

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 January 2003 (03.01.2003)

PCT

(10) International Publication Number  
**WO 03/000898 A1**

(51) International Patent Classification<sup>7</sup>: **C12N 15/29**,  
15/82, C12Q 1/68, A01H 5/00, G06F 17/00, C07K 14/415

(21) International Application Number: PCT/IB01/01105

(22) International Filing Date: 22 June 2001 (22.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **SYNGENTA PARTICIPATIONS AG** [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHANG, Hur-Song** [CN/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **CHEN, Wenqiong** [CN/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **COOPER, Bret** [US/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **GLAZEBROOK, Jane** [US/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **GOFF, Stephen, Arthur** [US/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **HOU, Yu-Ming** [CN/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **KATAGIRI, Fumiaki** [JP/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **QUAN, Sheng** [CN/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **TAO, Yi** [CN/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **WHITHAM, Steve**

[US/US]; 4025 Berkshire Avenue, Ames, IA 50010 (US). **XIE, Zhiyi** [CN/US]; Apartment 225, 8933 Lombard Place, San Diego, CA 92122 (US). **ZHU, Tong** [CN/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US). **ZOU, Guangzhou** [CN/US]; Torrey Mesa Research Institute, 3115 Merryfield Row, San Diego, CA 92121 (US).

(74) Agent: **BASTIAN, Werner**; c/o Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



**WO 03/000898 A1**

(54) Title: PLANT GENES INVOLVED IN DEFENSE AGAINST PATHOGENS

(57) Abstract: Methods to identify genes, the expression of which are altered in response to pathogen infection, are provided, as well as the genes identified thereby and their corresponding promoters.

## PLANT GENES INVOLVED IN DEFENSE AGAINST PATHOGENS

### Cross-Reference to Related Applications

This application claims the benefit of the filing date of U.S. application Serial No. 60/213,634, filed on June 23, 2000, U.S. application Serial No. 60/214,926, filed on June 23, 2000, U.S. application Serial No. 60/261,320, filed on January 12, 2001, U.S. application Serial No. 60/264,353, filed on January 26, 2001, and U.S. application Serial No. 60/273,879, filed on March 7, 2001 under 35 U.S.C. § 119(e).

### Field of the Invention

The present invention generally relates to the field of plant molecular biology, and more specifically to the regulation of gene expression in plants in response to pathogen exposure.

### Background of the Invention

Plants are capable of activating a large array of defense mechanisms in response to pathogen attack, some of which are preexisting and others are inducible. Pathogens must specialize to circumvent the defense mechanisms of the host, especially those biotrophic pathogens that derive their nutrition from an intimate association with living plant cells. If the pathogen can cause disease, the interaction is said to be compatible, but if the plant is resistant, the interaction is said to be incompatible. A crucial factor determining the success of these mechanisms is the speed of their activation. Consequently, there is considerable interest in understanding how plants recognize pathogen attack and control expression of defense mechanisms.

Some potential pathogens trigger a very rapid resistance response called gene-for-gene resistance. This occurs when the pathogen carries an avirulence (*avr*) gene that triggers specific recognition by a corresponding host resistance (*R*) gene. *R* gene specificity is generally quite narrow, in most cases only pathogens carrying a particular *avr* gene are recognized. Recognition is thought to be mediated by ligand-receptor binding. *R* genes have been studied extensively in recent years. For a review of *R* genes, see Ellis et al. (1998); Jones et al. (1997); and Ronald (1998).

One of the defense mechanisms triggered by gene-for-gene resistance is programmed cell death at the infection site. This is called the hypersensitive response, or HR. Pathogens that induce the HR, or cause cell death by other means, activate a systemic resistance response



called systemic acquired resistance (SAR). During SAR, levels of salicylic acid (SA) rise throughout the plant, defense genes such as pathogenesis related (PR) genes are expressed, and the plant becomes more resistant to pathogen attack. SA is a crucial component of this response. Plants that cannot accumulate SA due to the presence of a transgene that encodes an SA-degrading enzyme (*nahG*), develop a HR in response to challenge by avirulent pathogens, but do not exhibit systemic expression of defense genes and do not develop resistance to subsequent pathogen attack (Ryals et al., 1996). The nature of the systemic signal that triggers SAR is a subject of debate (Shulaev et al., 1995; Vernooji et al., 1994). SA clearly moves from the site of the HR to other parts of the plant, but if this is the signal, it must be effective at extremely low concentration (Willits et al., 1998).

SAR is quite similar to some reactions that occur locally in response to attack by virulent (those that cause disease) or avirulent (those that trigger gene-for-gene resistance) pathogens. In general, activation of defense gene expression occurs more slowly in response to virulent pathogens than in response to avirulent pathogens. Some pathogens trigger expression of defense genes through a different signaling pathway that requires components of the jasmonic acid (JA) and ethylene signaling pathways (Creelman et al., 1997).

One approach to understanding the signal transduction networks that control defense mechanisms is to use genetic methods to identify signaling components and determine their roles within the network. Considerable progress has been made using this approach in *Arabidopsis*-pathogen model systems.

#### **R gene signal transduction**

Genes such as *NDR1* and *EDS1*, as well as *DND1* and the lesion-mimic genes, likely act in signal transduction pathways downstream from *R-avr* recognition. *NDR1* and *EDS1* are required for gene-for-gene mediated resistance to avirulent strains of the bacterial pathogen *Pseudomonas syringae* and the oomycete pathogen *Peronospora parasitica*. Curiously, *ndr1* mutants are susceptible to one set of avirulent pathogens, whereas *eds1* mutants are susceptible to a non-overlapping set (Aarts et al., 1998). The five cloned *R* genes that require *EDS1* all belong to the subset of the nucleotide binding site-leucine rich repeat (NBS-LRR) class of *R* genes that contain sequences similar to the cytoplasmic domains of *Drosophila* Toll and mammalian interleukin 1 transmembrane receptors (TIR-NBS-LRR). The two genes that require *NDR1* belong to the leucine-zipper (LZ-NBS-LRR) subclass of NBS-LRR genes. There is another LZ-NBS-LRR gene, *RPP8*, that does not require *EDS1* or *NDR1*, so the correlation between *R* gene structure and requirement for *EDS1* or *NDR1* is not perfect.

Nevertheless, these results show that *R* genes differ in their requirements for downstream factors and that these differences are correlated with *R* gene structural type.

*NDR1* encodes a protein with two predicted transmembrane domains (Century et al. 1997). *RPM1*, which requires *NDR1* to mediate resistance, is membrane-associated, despite  
5 the fact that its primary sequence does not include any likely membrane-integral stretches (Boyes et al., 1998). It is possible that part of the function of *NDR1* is to hold *R* proteins close to the membrane. *EDS1* encodes a protein with blocks of homology to triacyl glycerol lipases (Falk et al., 1999). The significance of this homology is not known, but it is tempting to speculate that *EDS1* is involved in synthesis or degradation of a signal molecule. *EDS1*  
10 expression is inducible by SA and pathogen infection, suggesting that *EDS1* may be involved in signal amplification (Falk et al., 1999).

It has been extremely difficult to isolate mutations in genes other than the *R* genes that are required for gene-for-gene resistance. A selection procedure was devised (McNellis et al., 1998) on the basis of precisely controlled inducible expression of the *avr* gene *avrRpt2* in  
15 plants carrying the corresponding resistance gene *RPS2*. Expression of *avrRpt2* in this background is lethal, as it triggers a systemic HR. It is now possible to select for mutants with subtle defects in gene-for-gene signaling by requiring growth on a concentration of inducer slightly higher than the lethal dose.

Putative plant receptor proteins encoded by *RPP* genes (recognition of *P. parasitica*)  
20 mediate specific recognition of *Peronospora* isolates and trigger defense reactions. Recently, McDowell et al. (2000) reported that two members of this class, *RPP7* and *RPP8* (the latter of which encodes a LZ-NBS-LRR type R protein) were not significantly suppressed by mutations in either *EDS1* or *NDR1*, and that *RPP7* resistance was also not compromised by mutations in *EIN2*, *JAR1* or *COI1*, which affect ethylene or jasmonic acid signaling, or in *coil/npr1* or  
25 *coil/NahG* backgrounds. The authors suggested that *RPP7* initiates resistance through a novel signaling pathway that is independent of salicylic acid accumulation or jasmonic acid response components.

#### SA-dependent signaling

SA levels increase locally in response to pathogen attack, and systemically in response  
30 to the SAR-inducing signal. SA is necessary and sufficient for activation of PR gene expression and enhanced disease resistance. Physiological analyses and characterization of certain lesion-mimic mutants strongly suggest that there is a positive autoregulatory loop affecting SA concentrations (Shirasu et al., 1997; Hunt et al., 1997; Weymann et al., 1995).

Several mutants with defects in SA signaling have been characterized. These include *npr1*, in which expression of *PR* genes in response to SA is blocked; *cpr1*, *cpr5*, and *cpr6*, which constitutively express *PR* genes; the *npr1* suppressor *ssil*; *pad4*, which has a defect in SA accumulation; and *eds5*, which has a defect in *PR1* expression.

5           Expression of the defense genes *PR1*, *BG2*, and *PR5* in response to SA treatment requires a gene called *NPR1* or *NIM1*. Mutations in *npr1* abolish SAR, and cause enhanced susceptibility to infection by various pathogens (Cao et al., 1994; Delaney et al., 1995; Glazebrook et al., 1996; Shah et al., 1997). *NPR1* appears to be a positive regulator of *PR* gene expression that acts downstream from SA. *NPR1* encodes a novel protein that contains  
10   ankyrin repeats (which are often involved in protein-protein interactions (Cao et al., 1997; Ryals et al., 1997), and that is localized to the nucleus in the presence of SA (Dong et al., 1998). Consequently, it is unlikely that *NPR1* acts as a transcription factor to directly control *PR* gene expression, but its nuclear localization suggests that it may interact with such transcription factors.

15           *PAD4* appears to act upstream from SA. In *pad4* plants infected with a virulent *P. syringae* strain, SA levels, synthesis of the antimicrobial compound camalexin, and *PR1* expression are all reduced (Zhou et al., 1998). SA is necessary, but not sufficient, for activation of camalexin synthesis (Zhou et al., 1998; Zhao et al., 1996). The camalexin defect in *pad4* plants is reversible by exogenous SA (Zhou et al., 1998). Mutations in *pad4* do not  
20   affect SA levels, camalexin synthesis, or *PR1* when plants are infected with an avirulent *P. syringae* strain (Zhou et al., 1998). Taken together, these results suggest that *PAD4* is required for signal amplification to activate the SA pathway in response to pathogens that do not elicit a strong defense response (Zhou et al., 1998).

#### **JA-dependent signaling**

25           JA signaling affects diverse processes including fruit ripening, pollen development, root growth, and response to wounding (Creelman et al., 1997). The *jar1* and *coi1* mutants fail to respond to JA (Feys et al., 1994; Staswick et al., 1992). *COI1* has been cloned, and found to encode protein containing leucine-rich repeats and a degenerate F-box motif (Xie et al., 1998). These features are characteristic of proteins that function in complexes that ubiquitinate  
30   protein targeted for degradation.

In the past few years it has become apparent that JA plays an important role in regulation of pathogen defenses. For example, the induction of the defensin gene *PDF1.2* after

inoculation of *Arabidopsis* with the avirulent pathogen *Alternaria brassicicola* does not require SA or NPR1, but does require ethylene and JA signaling (Penninck et al., 1996).

SA signaling and JA signaling pathways are interconnected in complicated ways.

Studies in other systems have shown that SA signaling and JA signaling are mutually

- 5 inhibitory (Creelman et al., 1997; Harms et al., 1998). However, synthesis of camalexin in response to *P. syringae* infection is blocked in *nahG* (Zhou et al., 1998; Zhao et al., 1996) and *coi1* (Glazebrook, 1999) plants, strongly suggesting that camalexin synthesis requires both SA and JA signaling.

#### Induced systemic resistance (ISR)

- 10 Some rhizosphere-associated bacteria promote disease resistance (van Loon et al., 1998). This phenomenon, called ISR, has been studied using *Pseudomonas fluorescens* strain WCS417r to colonize *Arabidopsis* roots (Pieterse et al., 1996). Colonized plants are more resistant to infection by the fungal pathogen *Fusarium oxysporum* f sp *raphani* and *P. syringae* (Pieterse et al., 1996). ISR occurs in *nahG* plants, indicating that it is not a SA-dependent
- 15 phenomenon (Pieterse et al., 1996). Rather, ISR appears to be JA- and ethylene-dependent. The observation that ethylene can induce ISR in *jar1* mutants led to the hypothesis that ISR requires a JA signal followed by an ethylene signal (Pieterse et al., 1998). No changes in gene expression associated with ISR have been detected (Pieterse et al., 1998), suggesting that it is different from activation of *PDF1.2* expression by *A. brassicicola*.

- 20 Curiously, ISR requires *NPR1* (Pieterse et al., 1996). This was unexpected in light of the fact that NPR1 was previously known to be involved only in SA-dependent processes and ISR is SA-independent. If the SA-dependent signal is received, NPR1 mediates a resistance response characterized by *PR1* expression, whereas if the ISR signal is received, NPR1 mediates a different resistance response. It is difficult to imagine how this could occur, unless
- 25 NPR1 is interacting with different 'adapter' molecules to mediate the different signals. The ankyrin repeats found in NPR1 could function in protein-protein interactions between NPR1 and adapter proteins. Identification of proteins that interact with NPR1, and characterization of plants with loss-of-function mutations affecting those proteins, would be very helpful for understanding how NPR1 acts in each pathway. It would also be worthwhile to determine if
- 30 the *ssl1* or *cpr6* mutations suppress the ISR defect of *npr1* mutants.

#### Relevance to disease resistance

Characterization of the effects of various mutations on resistance to different pathogens has revealed that there is considerable variation in the extent to which pathogens are affected

by defense mechanisms. SAR is known to confer resistance to a wide array of pathogens, including bacteria, fungi, oomycetes, and viruses. JA signaling is important for limiting the growth of certain fungal pathogens. In *Arabidopsis*, the SA pathway mutants *npr1* and *pad4* show enhanced susceptibility to *P. syringae* and *P. parasitica* (Cao et al., 1994; Delaney et al., 5 1995; Shah et al., 1997; Zhou et al., 1998; Glazebrook et al., 1997).

Overexpression of rate-limiting defense response regulators may cause the signaling network to respond faster or more strongly to pathogen attack, thereby improving resistance. For example, overexpression of *NPR1* caused increased resistance to *P. syringae* and *P. parasitica* in a dosage dependent manner (Cao et al., 1998). Moreover, *NPR1*-overexpression 10 had no obvious deleterious effects on plant growth, in contrast to mutations that lead to constitutive overexpression of defense responses, which generally cause dwarfism.

#### **Promoters for gene expression of plant pathogen defense genes**

Promoters (and other regulatory components) from bacteria, viruses, fungi and plants have been used to control gene expression in plant cells. Numerous plant transformation 15 experiments using DNA constructs comprising various promoter sequences fused to various foreign genes (for example, bacterial marker genes) have led to the identification of useful promoter sequences. It has been demonstrated that sequences up to 500-1,000 bases in most instances are sufficient to allow for the regulated expression of foreign genes. However, it has also been shown that sequences much longer than 1 kb may have useful features which permit 20 high levels of gene expression in transgenic plants. The expression of genes encoding proteins that are useful for protecting plants from pathogen attack may have deleterious effects on plant growth if expressed constitutively. Consequently, it is desirable to have promoter sequences that control expression of these gene(s) in such a way that expression is absent or very low in the absence of pathogens, and high in the presence of pathogens.

25 Thus, what is needed is the identification of plant genes useful to confer resistance to a pathogen(s) and plant promoters, the expression of which is altered in response to pathogen attack.

#### **Summary of the Invention**

30 The invention generally provides an isolated nucleic acid molecule (polynucleotide) comprising a plant nucleotide sequence obtained or isolatable from a gene, the expression of which is altered, either increased or decreased, in response to pathogen infection. In one embodiment, the plant nucleotide sequence comprises an open reading frame, while in another

embodiment, the plant nucleotide sequence comprises a promoter. A promoter sequence of the invention directs transcription of a linked nucleic acid segment, e.g., a linked plant DNA comprising an open reading frame for a structural or regulatory gene, in a host cell, such as a plant cell, in response to pathogen infection of that cell. As used herein, a "pathogen" includes  
5 bacteria, fungi, oomycetes, viruses, nematodes and insects, e.g., aphids (see Hammond-Kosack and Jones, 1997). Moreover, the expression of a plant nucleotide sequence of the invention comprising a promoter may be altered in response to one or more species of bacteria, nematode, fungi, oomycete, virus, or insect. Likewise, the expression of a plant nucleotide sequence of the invention comprising an open reading frame may be useful to confer tolerance  
10 or resistance of a plant to one or more species of bacteria, nematode, fungi, oomycete, virus or insect.

The nucleotide sequence preferably is obtained or isolatable from plant DNA. In particular, the nucleotide sequence is obtained or isolatable from a gene encoding a polypeptide which is substantially similar, and preferably has at least 70%, e.g., 71%, 72%,  
15 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, and even 90% or more, e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, and 99%, amino acid sequence identity, to a polypeptide encoded by an *Arabidopsis* gene comprising any one of SEQ ID NOs: 1-953 and 2137-2661 or a fragment (portion) thereof which encodes a partial length polypeptide having substantially the same activity of the full-length  
20 polypeptide, a rice gene comprising one of SEQ ID NOs:2000-2129 or SEQ ID NOs:2662-6813, or a *Chenopodium* gene comprising one of SEQ ID NOs:1954-1966.

The present invention also provides an isolated nucleic acid molecule comprising a plant nucleotide sequence that directs transcription of a linked nucleic acid segment in a host cell, e.g., a plant cell. The nucleotide sequence preferably is obtained or isolatable from plant  
25 genomic DNA. In particular the plant DNA is obtained or isolatable from a gene encoding a polypeptide which is substantially similar, and preferably has at least 70%, e.g., 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, and even 90% or more, e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, and 99%, amino acid sequence identity, to a polypeptide encoded by an *Arabidopsis* gene comprising  
30 any one of SEQ ID NOs:1-953, a rice gene comprising one of SEQ ID NOs:2000-2129 or SEQ ID NOs:2662-4737, or a *Chenopodium* gene having any one of SEQ ID NOs:1954-1966, the expression of which is increased or decreased in response to pathogen infection. Preferred promoters comprise DNA obtained or isolatable from a gene encoding a polypeptide which is

substantially similar, and preferably has at least 70%, e.g., 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, and even 90% or more, e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, and 99%, amino acid sequence identity, to a polypeptide encoded by an *Arabidopsis* gene comprising a promoter according to  
5 SEQ ID NOs:2137-2661, a rice gene comprising a promoter according to SEQ ID NOs:4738-6813 or a fragment thereof (i.e., promoters isolatable from any one of SEQ ID NOs:2137-2661 or SEQ ID NOs:4738-6813) which increases or decreases transcription of a linked nucleic acid segment in response to pathogen infection.

The invention also provides uses for an isolated nucleic acid molecule, e.g., DNA or  
10 RNA, comprising a plant nucleotide sequence comprising an open reading frame encoding a polypeptide which is substantially similar, and preferably has at least 70%, e.g., 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, and even 90% or more, e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, and 99%, amino acid sequence identity, to a polypeptide encoded by an *Arabidopsis*, *Chenopodium* or  
15 rice gene comprising an open reading frame comprising any one of SEQ ID NOs:1-953, 1954-1966, 2000-2129, 2662-4737, or the complement thereof. For example, these open reading frames may be useful to prepare plants that over- or under-express the encoded product or to prepare knockout plants.

The promoters and open reading frames of the invention can be identified by any  
20 method. For example, they can be identified by employing an array of nucleic acid samples, e.g., each sample having a plurality of oligonucleotides, and each plurality corresponding to a different plant gene, on a solid substrate, e.g., a DNA chip, and probes corresponding to nucleic acid which is up- or down-regulated in response to pathogen infection in one or more ecotypes or species of plant relative to a control (e.g., a water control, nucleic acid from an  
25 uninfected plant or nucleic acid from a mutant plant). Thus, genes that are upregulated or downregulated in response to pathogen infection can be systematically identified.

As described herein, GeneChip® technology was utilized to discover a plurality of genes, the expression of which is altered after pathogen infection. The *Arabidopsis* oligonucleotide probe array consists of probes from about 8,100 unique *Arabidopsis* genes,  
30 which covers approximately one third of the genome. This genome array permits a broader, more complete and less biased analysis of gene expression. Using labeled cRNA probes, expression levels were determined by laser scanning and genes generally selected for expression levels that were > 2 fold over the control.

For example, using this approach, 953 genes were identified, the expression of which was altered after infection of wild-type *Arabidopsis* plants with a pathogen (SEQ ID NOs:1-953). In addition, 745 genes were identified, the expression which was increased after infection of wild-type *Arabidopsis* with *Pseudomonas syringae* (SEQ ID NOs: 2-6, 8-13, 16, 18, 22-23, 25, 28-29, 31-32, 35-37, 39-43, 45-47, 49-50, 52, 54-55, 57-58, 60-66, 70-72, 74, 76-77, 79, 81, 83, 85, 87-90, 92, 94, 97, 100-107, 111-115, 117-125, 127-135, 138-140, 142-153, 156-158, 160, 162-165, 168-170, 173-181, 183-184, 186-188, 190-198, 200-201, 203-211, 214-215, 218-224, 227-232, 234-249, 251-262, 264, 266-268, 270, 272-275, 277-281, 283, 286-294, 297-298, 302, 304-306, 308-326, 328-339, 341, 344-345, 347, 350-351, 353-358, 361-371, 373-377, 379-386, 388-390, 392, 394-400, 402-406, 408-410, 412-417, 419-427, 429-433, 435-443, 445-452, 454-457, 459-460, 462-464, 466-470, 473-475, 478-479, 481-482, 484-187, 489-494, 496-498, 500-501, 503-506, 508, 510, 512-515, 517-523, 526, 528-529, 531-538, 540, 544-548, 550-558, 560, 563-568, 570, 572-577, 579-580, 582-585, 588-594, 596, 598-600, 602-603, 605-606, 608-612, 614-617, 619-624, 626-630, 632-639, 642, 644, 646-651, 653-657, 659-665, 667-671, 673-678, 681-689, 691-693, 695-713, 715-717, 719, 721-727, 729-733, 736-738, 740, 742, 744, 746, 748-752, 755-756, 758-760, 762-769, 771, 774, 776-781, 783-788, 790-796, 798-799, 802, 804-808, 810-815, 817-831, 833-848, 850-855, 857-869, 871-880, 882-900, 903-907, 909, 911-915, 918-920, 922-925, 927, 929, 931-938, 940, 943-945, 947, and 950-953). Of the 745 genes, the expression of 530 of those genes was altered in at least one mutant *Arabidopsis* after infection with *Pseudomonas syringae* (SEQ ID NOs: 2, 4-6, 11-13, 18, 22-23, 28, 31, 36, 39-43, 45, 47, 49-50, 52, 54-55, 57-58, 60-61, 63-66, 71-72, 74, 77, 81, 83, 85, 87-89, 92, 97, 100-107, 111-112, 114-115, 117-120, 122, 125, 127-128, 134, 128-140, 143-144, 148-151, 153, 156-157, 160, 165, 168-170, 173-174, 176-180, 183, 187-188, 191, 193-194, 197-198, 200, 203-210, 214, 219-224, 227, 230-232, 235-237, 239-240, 243-246, 248-249, 251-254, 256-258, 261, 264, 266-268, 270, 272-275, 277-278, 280, 283, 286-287, 290-293, 297, 302, 305-306, 308-310, 312-316, 321-326, 328-331, 333, 336-339, 341, 345, 351, 353, 355-358, 361-363, 365-366, 368-371, 373, 375, 377, 379-381, 384-385, 388-390, 392, 394-400, 402-406, 410, 412, 415-416, 419-420, 422-425, 429-433, 435-439, 441-443, 445-452, 454, 459-460, 463, 466, 468-470, 473, 481-482, 485-486, 489, 491-494, 497-498, 500-501, 503, 505-506, 508, 510, 513-515, 517, 520-521, 523, 528-529, 531, 533-538, 540, 545-548, 550-551, 553-554, 556-558, 560, 566-567, 575, 580, 582-584, 588-593, 596, 598-600, 602-603, 605-606, 608-610, 612, 614, 616, 620-622, 627-629, 633-634, 636-639, 644, 646, 648-651, 654-657, 659, 661-663, 667, 669, 673-



674, 677, 682, 684-687, 689, 691-693, 697, 699, 701, 703-708, 713, 717, 719, 721-727, 730-733, 736, 740, 744, 746, 749-752, 755-756, 758-760, 762-764, 766-769, 774, 776-778, 780-781, 786, 788, 791-796, 799, 802, 804-808, 810-812, 815, 818-821, 823-825, 827-829, 831, 833-836, 838-843, 845, 847-848, 852-853, 855, 858, 860-869, 871-874, 876, 878-880, 884-887, 889, 892-894, 896-900, 904-907, 911-915, 918-920, 922-924, 931, 933, 938, 943-945, 947, and 950-952). Of the 530, 81 encode regulatory factors (SEQ ID NOs: 39, 52, 60, 63, 81, 83, 106, 107, 115, 117, 118, 168, 174, 176, 179, 204, 207, 208, 220, 221, 248, 258, 268, 275, 280, 309, 323, 326, 329, 351, 419, 422, 429, 430, 432, 459, 460, 468, 469, 473, 500, 505, 506, 508, 529, 531, 533, 535, 538, 545, 553, 602, 606, 608, 610, 614, 616, 634, 654, 655, 684, 686, 687, 691, 717, 751, 752, 766, 777, 815, 831, 834, 835, 839, 841, 847, 876, 884, 906, 920, and 924).

As also described herein, 333 genes were identified that are useful to confer improved resistance to plants to bacterial infection (SEQ ID NOs: 12-13, 18, 23, 36, 39-40, 43, 45, 50, 52, 57-58, 60-61, 64, 71-72, 81, 87-89, 97, 100, 102-105, 107, 111-112, 115, 119-120, 122, 125, 127-128, 140, 144, 148-150, 153, 165, 168-169, 176-177, 179, 183, 188, 191, 193-194, 197-198, 203-206, 208-209, 214, 219-222, 227, 230, 232, 237, 244-246, 248-249, 251-253, 258, 261, 264, 266, 268, 273-275, 283, 287, 290, 293, 297, 302, 305-306, 308, 312-315, 321-322, 324, 326, 330, 333, 338, 341, 345, 353, 356-358, 362-363, 366, 369, 371, 375, 377, 380, 384-385, 389, 392, 394-395, 398-399, 402-404, 406, 410, 415, 419, 422, 425, 429-430, 433, 435-439, 443, 445-452, 454, 463, 466, 468-470, 473, 486, 489, 491-492, 4894, 498, 500-501, 503, 508, 513-514, 517, 529, 533-538, 548, 550, 553-554, 4556-558, 566, 575, 580, 582-583, 590-591, 593, 600, 602, 609-610, 612, 614, 620-622, 627-629, 637-638, 644, 649, 654-657, 659, 663, 667, 669, 673-674, 677, 684-685, 689, 691-693, 699, 703-705, 708, 719, 721, 724-726, 730-732, 744, 746, 749-750, 752, 755-756, 758, 760, 762-764, 767, 769, 774, 780-781, 786, 788, 791-792, 794-796, 799, 804-808, 810-812, 815, 818-819, 823, 828-829, 833, 840841, 843, 847, 852-853, 858, 860, 862-865, 867-868, 872-874, 876, 885-887, 889, 892-894, 896-900, 904-905, 907, 911-914, 918-920, 922-924, 931, 933, 938, 947, 950, and 952).

Further, 296 genes were identified that are useful to confer improved resistance to plants to fungal infection (SEQ ID NOs: 2, 4, 6, 11-13, 18, 22-23, 31, 41-43, 49-50, 54, 57-58, 61, 64-66, 71-72, 74, 77, 85, 87, 89, 92, 97, 101, 103, 106-107, 112, 114, 117-119, 125, 128, 134, 138, 143, 149, 151, 156-157, 165, 169-170, 174, 176-180, 187-188, 191, 193, 206, 208, 219-220, 222, 224, 231, 236, 239, 243-245, 251-254, 256-257, 267, 272, 287, 290, 292, 297, 302, 312-313, 315-316, 321-322, 324-325, 328, 330, 345, 351, 353, 355-357, 362-363, 366,

368-371, 373, 375, 379, 381, 384, 388-390, 392, 395-400, 405, 410, 415-416, 419, 422, 424, 431-432, 435-436, 438-439, 447, 459-460, 470, 473, 481-482, 489, 491, 493-494, 500-501, 505-506, 513-514, 517, 520-521, 523, 528-529, 531, 535, 537-538, 540, 545-548, 551, 553-554, 557-558, 566, 575, 580, 582, 584, 589, 591, 593, 596, 598-599, 603, 605, 608-609, 612, 5 628, 633-634, 636-637, 639, 646, 648, 650-651, 656, 661, 663, 667, 674, 685-687, 689, 691, 693, 697, 699, 701, 705, 707, 713, 723-724, 726, 736, 740, 749, 751-752, 756, 758-759, 764, 766-768, 774, 776, 778, 780, 792-796, 799, 802, 806, 810-812, 818, 820-821, 825, 827-829, 833-836, 838-839, 841-843, 848, 855, 860-861, 866, 868-869, 871, 873-874, 876, 878-880, 889, 892, 898-900, 904-905, 907, 915, 918, 922, 924, 933, 943-945, 947, and 951).

10 In addition, 288 genes were identified that are useful to confer improved resistance to plants to infection with more than one pathogen, e.g., pathogens that include bacteria, oomycetes and viruses (SEQ ID NOs: 12-13, 18, 23, 36, 39-40, 43, 45, 50, 52, 57-58, 60-61, 64, 71-72, 81, 87-88, 100, 102-105, 107, 111-112, 115, 119-120, 122, 125, 127-128, 140, 148-150, 153, 168-169, 176-177, 188, 191, 193-194, 197-198, 203-206, 209, 219-222, 227, 232, 15 237, 244-246, 248-249, 251-253, 258, 261, 264, 266, 268, 273-275, 283, 287, 290, 293, 297, 302, 305-306, 308, 312-315, 324, 326, 330, 333, 341, 345, 353, 356, 358, 366, 371, 375, 377, 380, 385, 389, 392, 394, 398, 402-404, 406, 410, 415, 419, 425, 429-430, 433, 435-438, 443, 445-447, 449-452, 454, 463, 466, 468-470, 473, 486, 489, 492, 494, 498, 500-501, 503, 508, 513-514, 517, 533-538, 548, 550, 553-554, 57-558, 566, 575, 582-583, 590-591, 593, 600, 20 602, 609-610, 612, 620-622, 627-629, 637-638, 644, 649, 654-657, 659, 667, 669, 673, 677, 684, 689, 692-693, 703-705, 719, 721, 724-726, 730-732, 744, 746, 749-750, 752, 755-756, 760, 762-764, 767, 769, 774, 780-781, 786, 788, 791-792, 795-796, 805-808, 810-812, 815, 818-819, 823, 828, 833, 840-841, 843, 852-853, 858, 860, 862-865, 867-868, 872-874, 876, 887, 889, 893-894, 896-898, 900, 905, 907, 911-914, 918-920, 922-923, 931, 933, 938, 947, 25 950, and 952).

Using the same approach described above, 25 genes were identified (SEQ ID NOs: 1, 15, 19, 20, 24, 26, 27, 34, 38, 51, 56, 59, 67-69, 99, 116, 155, 159, 182, 212, 284, 372, 444, and 789), the expression of which was decreased at 6 hours in an *avr2* plant. Also identified were 33 genes (SEQ ID NOs: 17, 70, 76, 81, 84, 109, 123, 144, 160, 230, 265, 268, 269, 271, 30 323, 333, 385, 427, 428, 430, 457, 505, 569, 597, 602, 606, 616, 708, 730, 741, 812, 862, and 942), the expression of which was elevated in an incompatible or a compatible interaction in four *Arabidopsis* ecotypes infected with bacteria. Eight of the genes were upregulated by 3 hours in an incompatible interaction, 18 of the genes were upregulated by 6 hours, but not at 3

hours, in an incompatible interaction, and 6 of the genes were upregulated in a compatible interaction.

Further identified were 33 genes, the expression of which was induced early after infection (SEQ ID NOs:17, 21, 80, 81, 156, 174, 176, 221, 227, 296, 302, 303, 306, 333, 340, 360, 500, 505, 524, 575, 601, 602, 614, 628, 687, 733, 782, 811, 835, 862, 900, 905, and 912), 10 genes, the expression of which was decreased early after infection (SEQ ID NOs:30, 73, 282, 541, 640, 679, 761, 870, 917, and 930), and 135 genes, 107 of which were induced at 3 and/or 6 hours after infection, and 28 of which were decreased after infection (SEQ ID NOs:7, 21, 33, 44, 46, 60, 82, 86, 91, 93, 106, 110, 119, 122, 130, 131, 136, 141, 154, 161, 166-168, 171, 176, 185, 189, 199, 200, 202, 203, 213, 225, 227, 248, 261, 262, 266, 274, 285, 300, 301, 302, 320, 326, 341, 345, 348, 349, 360, 366, 378, 406, 409, 422, 425, 434, 441, 443, 446, 449, 454, 461, 471, 475, 476, 483, 485, 499, 500, 511, 512, 516, 527, 530, 533, 543, 545, 549, 550, 552, 567, 575, 578, 586, 590, 608, 611, 615, 618, 625, 631, 643, 656, 658, 659, 666, 668, 671, 680, 690, 694, 704, 706, 711, 714, 718, 721, 728, 734, 738, 757, 770, 772, 791, 807, 811, 813, 816, 827, 857, 864, 868, 875, 881, 893, 901, 905, 908, 912, 916, 939, 941, 951, and 952).

In a similar approach, 48 genes that were upregulated in response to infection, e.g., bacterial or fungal infection, as well as 46 of the corresponding promoter containing regions, were identified. Thirty-six of the genes were upregulated in response to bacterial, e.g., *Pseudomonas*, infection (the promoters for genes corresponding to SEQ ID NOs: 104-106, 119, 123, 129, 131, 151-152, 183, 191, 198, 200, 227, 249, 274, 302, 358, 415, 481, 547, 566, 582, 628, 633, 639, 656, 673, 793, 818, 827, 864, 874, 880, and 904-905), while 23 of the genes were upregulated in response to fungal, e.g., *Botrytis*, infection (SEQ ID NOs: 18, 71, 119, 123, 129, 151, 191, 244, 245, 302, 545, 547, 562, 566, 637, 653, 747, 756, 774, 793, 842, 864, and 905). Twenty-five of the genes were upregulated only in response to bacterial, e.g., *Pseudomonas*, infection (the promoters for genes corresponding to SEQ ID NOs: 104-106, 131, 152, 183, 198, 200, 227, 249, 274, 358, 415, 481, 582, 628, 633, 639, 656, 673, 818, 827, 874, 880, and 904 are provided in SEQ ID NOs:1001-1025), 10 of the genes were upregulated only in response to fungal, e.g., *Botrytis*, infection (the promoters for genes corresponding to SEQ ID NOs:18, 71, 244, 245, 545, 562, 637, 653, 747, 756, 774, and 842 are provided in SEQ ID NOs:1026-1035), and 11 genes were upregulated in response to both bacterial and fungal infection (the promoters for genes corresponding to SEQ ID NOs:119, 123, 129, 151, 191, 302, 547, 566, 793, 864, and 905 are provided in SEQ ID NOs:1036-1046).

As also described hereinbelow, 129 *Arabidopsis* genes (SEQ ID NOs: 3, 51, 54, 60, 61, 66, 75, 76, 78, 88, 95, 96, 101, 106, 108, 123, 126, 128, 129, 131, 137, 145-147, 150, 158, 169, 170, 172, 173, 197, 200, 216, 219, 224, 230, 233, 237, 249, 250, 263, 274, 275, 276, 299, 307, 323, 333, 342, 346, 359, 382, 383, 387, 391, 393, 401, 411, 415, 427, 442, 455, 459, 466, 5 477, 481, 485, 487, 502, 511, 515, 525, 534, 539, 542, 560, 571, 577, 579, 584, 587, 595, 600, 627, 638, 645, 654, 659, 668, 681, 688, 695, 696, 706, 708, 730, 742, 753, 775, 785, 786, 791, 797, 800, 801, 809, 817, 819, 820, 823, 827, 847, 856, 875, 885, 896, 902, 910, 921, 922, 923, 925, 926, 928, 946, and 952) were identified that were upregulated in response to viral infection, and 46 *Arabidopsis* genes were identified that were downregulated in response to viral infection (SEQ ID NOs: 14, 48, 53, 98, 217, 226, 295, 327, 343, 352, 369, 404, 407, 418, 10 453, 458, 465, 472, 480, 488, 495, 507, 509, 513, 514, 559, 561, 581, 604, 607, 613, 641, 652, 672, 720, 735, 739, 743, 745, 754, 773, 803, 832, 849, 948, and 949).

Also provided are nucleic acid molecules comprising a nucleotide sequence comprising an open reading frame expressed in response to pathogen infection comprising SEQ ID NOs: 209, 216, 262, 267, 317, 386, 425, 440 and 800. These sequences are useful to over- or 15 under-express the encoded product, or prepare knock-out plants which have an altered response to pathogen infection.

The invention therefore provides a method in which the open reading frame of a plant pathogen resistance gene, e.g., a gene that is associated with a response to pathogen infection, 20 which is altered in a plant in response to infection is identified and isolated. A transgene comprising the isolated open reading frame may be introduced to and expressed in a transgenic plant, e.g., prior to infection, e.g., constitutively, or early and/or rapidly after infection, or in regulatable (inducible) fashion, e.g., after exposure to a chemical or using a promoter that is upregulated after infection, so as to confer resistance to that transgenic plant to the pathogen 25 relative to a corresponding plant which does not have the transgene. The expression of the transgene is preferably at higher than normal levels, and under the regulation of a promoter that allows very fast and high induction in response to the presence of a pathogen or under cycling promoters (e.g., circadian clock regulated promoters), such that the encoded gene product(s) is maintained at sufficiently high levels to provide enhanced resistance or tolerance. 30 The invention further provides a method in which a gene in a plant which is downregulated in response to infection, is disrupted or the expression of that gene is further downregulated, e.g., using antisense expression, so as to result in a plant that has enhanced resistance to infection,

and which disruption or downregulation preferably has little or no detrimental effect(s) on the host plant.

As also described herein, it was found that the early incompatible response was similar to the late compatible response, suggesting that early expression of plant pathogen-resistance genes is important for resistance. Also, various plant strains were found to respond differently to the same pathogen, but there was also an identifiable global pattern of response. Thus, the comparison of the expression patterns in incompatible and compatible interactions in one or more ecotypes can lead to identifying subsets of key responding genes and clusters of genes that are key (early) responders. In addition, the observed global expression pattern indicated that the least resistant strain tested (Ws) had a low basal level of pathogen-upregulated genes and a high level of pathogen-downregulated genes compared to the most resistant strain (Ler). Thus, plant strains that are more resistant to pathogens have a gene expression phenotype in which genes that are upregulated in response to infection are already expressed at a higher than normal basal level, and those genes that are downregulated are expressed at a lower than normal basal level.

The genes and promoters described hereinabove can be used to identify orthologous genes and their promoters which are also likely useful to enhance resistance of plants to pathogens. Moreover, the orthologous promoters are useful to express linked open reading frames. In addition by aligning the promoters of these orthologs, novel *cis* elements can be identified that are useful to generate synthetic promoters.

Hence, the isolated nucleic acid molecules of the invention include the orthologs of the *Arabidopsis*, *Chenopodium* and rice sequences disclosed herein, i.e., the corresponding nucleic acid molecules in organisms other than *Arabidopsis*, *Chenopodium* and rice, including, but not limited to, plants other than *Arabidopsis*, *Chenopodium* and rice, preferably cereal plants, e.g., corn, wheat, rye, turfgrass, sorghum, millet, sugarcane, soybean, barley, alfalfa, sunflower, canola, soybean, cotton, peanut, tobacco, sugarbeet, or rice. An ortholog is a gene from a different species that encodes a product having the same function as the product encoded by a gene from a reference organism. Databases such GenBank or one found at <http://bioserver.myongji.ac.kr/rjce.html> (for rice) may be employed to identify sequences related to the *Arabidopsis* or *Chenopodium* sequences, e.g., orthologs in cereal crops such as rice. Alternatively, recombinant DNA techniques such as hybridization or PCR may be employed to identify sequences related to the *Arabidopsis* sequences. The encoded ortholog products likely have at least 70% sequence identity to each other. Hence, the invention

includes an isolated nucleic acid molecule comprising a nucleotide sequence encoding a polypeptide having at least 70% identity to a polypeptide encoded by one or more of the *Arabidopsis*, *Chenopodium* or rice sequences disclosed herein. For example, promoter sequences within the scope of the invention are those which direct expression of an open reading frame which encodes a polypeptide that is substantially similar to an *Arabidopsis* polypeptide encoded by a gene comprising SEQ ID NOs:1-953.

The genes and promoters described hereinabove can be used to identify orthologous genes and their promoters which are also likely expressed in a particular tissue and/or development manner. Moreover, the orthologous promoters are useful to express linked open reading frames. In addition, by aligning the promoters of these orthologs, novel *cis* elements can be identified that are useful to generate synthetic promoters. Hence, the isolated nucleic acid molecules of the invention include the orthologs of the *Arabidopsis* sequences disclosed herein, i.e., the corresponding nucleotide sequences in organisms other than *Arabidopsis*, including, but not limited to, plants other than *Arabidopsis*, preferably cereal plants, e.g., corn, wheat, rye, turfgrass, sorghum, millet, sugarcane, soybean, barley, alfalfa, sunflower, canola, soybean, cotton, peanut, tobacco, sugarbeet, or rice. An orthologous gene is a gene from a different species that encodes a product having the same or similar function, e.g., catalyzing the same reaction as a product encoded by a gene from a reference organism. Thus, an ortholog includes polypeptides having less than, e.g., 65% amino acid sequence identity, but which ortholog encodes a polypeptide having the same or similar function. Databases such as GenBank or one found at <http://bioserver.myongji.ac.kr/rjce.html> (for rice) may be employed to identify sequences related to the *Arabidopsis* sequences, e.g., orthologs in cereal crops such as rice, wheat, sunflower or alfalfa. SEQ ID NOs: 6286 and 4210, for example, are the rice promoter and open reading frame for rice peroxidase, the ortholog of the *Arabidopsis* gene comprising SEQ ID NO: 50. SEQ ID NOs: 3311, 5387, 3791 and 5867 are rice orthologs of the *Arabidopsis* gene comprising SEQ ID NO:609; SEQ ID NOs: 2699, 4775, 3463, 5539, 3584, 5660, 4451, 6527, 4595 and 6671 are rice orthologs of the *Arabidopsis* gene comprising SEQ ID NO: 139.

Preferably, the promoters of the invention include a consecutive stretch of about 25 to 2000, including 50 to 500 or 100 to 250, and up to 1000 or 1500, contiguous nucleotides, e.g., 40 to about 743, 60 to about 743, 125 to about 743, 250 to about 743, 400 to about 743, 600 to about 743, of any one of SEQ ID NOs:2137-2661, SEQ ID NOs:4738-6813 or the promoter orthologs thereof, which include the minimal promoter region. Preferably, the nucleotide

sequence that includes the promoter region includes at least one copy of a TATA box. Thus, the invention provides plant promoters, including orthologs of *Arabidopsis* promoters corresponding to genes comprising any one of SEQ ID NOs: 1-953. The present invention further provides a composition, an expression cassette or a recombinant vector containing the nucleic acid molecule of the invention, and host cells comprising the expression cassette or vector, e.g., comprising a plasmid. In particular, the present invention provides an expression cassette or a recombinant vector comprising a promoter of the invention linked to a nucleic acid segment which, when present in a plant, plant cell or plant tissue, results in transcription of the linked nucleic acid segment.

10 In its broadest sense, the term "substantially similar" when used herein with respect to a nucleotide sequence means that the nucleotide sequence is part of a gene which encodes a polypeptide having substantially the same structure and function as a polypeptide encoded by a gene for the reference nucleotide sequence, e.g., the nucleotide sequence comprises a promoter from a gene that is the ortholog of the gene corresponding to the reference nucleotide sequence, as well as promoter sequences that are structurally related the promoter sequences particularly exemplified herein, i.e., the substantially similar promoter sequences hybridize to the complement of the promoter sequences exemplified herein under high or very high stringency conditions. The term "substantially similar" thus includes nucleotide sequences wherein the sequence has been modified, for example, to optimize expression in particular cells, as well as nucleotide sequences encoding a variant polypeptide comprising one or more amino acid substitutions relative to the (unmodified) polypeptide encoded by the reference sequence, which substitution(s) does not alter the activity of the variant polypeptide relative to the unmodified polypeptide. In its broadest sense, the term "substantially similar" when used herein with respect to polypeptide means that the polypeptide has substantially the same structure and function as the reference polypeptide. The percentage of amino acid sequence identity between the substantially similar and the reference polypeptide is at least 65%, 66%, 67%, 68%, 69%, 70%, e.g., 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, and even 90% or more, e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, up to at least 99%, wherein the reference polypeptide is a polypeptide encoded by an *Arabidopsis* gene comprising any one of SEQ ID NOs:1-953, a *Chenopodium* gene comprising any one of SEQ ID NOs:1954-1966, or a rice gene comprising any one of SEQ ID NOs:2000-2129 or 2662-4737. One indication that two polypeptides are substantially similar to each other, besides having substantially the same function, is that an

agent, e.g., an antibody, which specifically binds to one of the polypeptides, specifically binds to the other.

Sequence comparisons may be carried out using a Smith-Waterman sequence alignment algorithm (see e.g., Waterman (1995) or <http://www.hto.usc.edu/software/seqaln/index.html>).

- 5 The localS program, version 1.16, is preferably used with following parameters: match: 1, mismatch penalty: 0.33, open-gap penalty: 2, extended-gap penalty: 2. Further, a nucleotide sequence that is "substantially similar" to a reference nucleotide sequence hybridizes to the reference nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 2X SSC, 0.1% SDS at 50°C, more desirably in 7% sodium
- 10 dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 1X SSC, 0.1% SDS at 50°C, more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.5X SSC, 0.1% SDS at 50°C, preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 50°C, more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM
- 15 EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 65°C.

- Hence, the present invention further provides an expression cassette or a vector containing the nucleic acid molecule comprising an open reading frame of the invention operably linked to a promoter, or comprising a promoter of the invention operably linked to an open reading frame or portion thereof, and the vector may be a plasmid. Such cassettes or
- 20 vectors, when present in a plant, plant cell or plant tissue result in transcription of the linked nucleic acid fragment in the plant. The expression cassettes or vectors of the invention may optionally include other regulatory sequences, e.g., transcription terminator sequences, operator, repressor binding site, transcription factor binding site, and/or an enhancer and may be contained in a host cell. The expression cassette or vector may augment the genome of a
  - 25 transformed plant or may be maintained extrachromosomally. The expression cassette or vector may further have a Ti plasmid and be contained in an *Agrobacterium tumefaciens* cell; it may be carried on a microparticle, wherein the microparticle is suitable for ballistic transformation of a plant cell; or it may be contained in a plant cell or protoplast. Further, the expression cassette can be contained in a transformed plant or cells thereof and the plant may
  - 30 be a dicot or a monocot. In particular, the plant may be a cereal plant.

The invention also provides sense and anti-sense nucleic acid molecules corresponding to the open reading frames identified herein as well as their orthologs. Also provided are expression cassettes, e.g., recombinant vectors, and host cells, comprising the nucleic acid



molecule of the invention, e.g., one which comprises a nucleotide sequence which encodes a polypeptide the expression of which is altered in response to pathogen infection.

The present invention further provides a method of augmenting a plant genome by contacting plant cells with a nucleic acid molecule of the invention, e.g., one isolatable or  
5 obtained from a plant gene encoding a polypeptide that is substantially similar to a polypeptide encoded by an *Arabidopsis*, *Chenopodium* or rice gene comprising a sequence comprising any one of SEQ ID NOs: 1-953, 1954-1966, 2000-2129 or 2662-4737 so as to yield transformed plant cells; and regenerating the transformed plant cells to provide a differentiated transformed plant, wherein the differentiated transformed plant expresses the nucleic acid molecule in the  
10 cells of the plant. The nucleic acid molecule may be present in the nucleus, chloroplast, mitochondria and/or plastid of the cells of the plant. The present invention also provides a transgenic plant prepared by this method, a seed from such a plant and progeny plants from such a plant including hybrids and inbreds. Preferred transgenic plants are transgenic maize, soybean, barley, alfalfa, sunflower, canola, soybean, cotton, peanut, sorghum, tobacco,  
15 sugarbeet, rice, wheat, rye, turfgrass, millet, sugarcane, tomato, or potato.

The invention also provides a method of plant breeding, e.g., to prepare a crossed fertile transgenic plant. The method comprises crossing a fertile transgenic plant comprising a particular nucleic acid molecule of the invention with itself or with a second plant, e.g., one lacking the particular nucleic acid molecule, to prepare the seed of a crossed fertile transgenic  
20 plant comprising the particular nucleic acid molecule. The seed is then planted to obtain a crossed fertile transgenic plant. The plant may be a monocot or a dicot. In a particular embodiment, the plant is a cereal plant.

The crossed fertile transgenic plant may have the particular nucleic acid molecule inherited through a female parent or through a male parent. The second plant may be an inbred  
25 plant. The crossed fertile transgenic may be a hybrid. Also included within the present invention are seeds of any of these crossed fertile transgenic plants.

The various breeding steps are characterized by well-defined human intervention such as selecting the lines to be crossed, directing pollination of the parental lines, or selecting appropriate progeny plants. Depending on the desired properties different breeding measures  
30 are taken. The relevant techniques are well known in the art and include but are not limited to hybridization, inbreeding, backcross breeding, multiline breeding, variety blend, interspecific hybridization, aneuploid techniques, etc. Hybridization techniques also include the sterilization of plants to yield male or female sterile plants by mechanical, chemical or biochemical means.

Cross pollination of a male sterile plant with pollen of a different line assures that the genome of the male sterile but female fertile plant will uniformly obtain properties of both parental lines. Thus, the transgenic plants according to the invention can be used for the breeding of improved plant lines that for example increase the effectiveness of conventional methods such as herbicide or pesticide treatment or allow to dispense with said methods due to their modified genetic properties. Alternatively new crops with improved stress tolerance can be obtained that, due to their optimized genetic "equipment", yield harvested product of better quality than products that were not able to tolerate comparable adverse developmental conditions.

The nucleic acid molecules of the invention, their encoded polypeptides and compositions thereof, are: for open reading frames, useful to provide resistance to pathogens to alter expression of a particular gene corresponding to the open reading frame by decreasing or eliminating expression of that plant gene or by overexpressing a particular gene product, and as a diagnostic for the presence or absence of the pathogen by correlating the expression level or pattern of expression of one or more of the nucleic acid molecules or polypeptides of the invention; and for promoters, useful to alter the expression of a linked open reading frame in response to pathogen infection. As one embodiment of the invention includes isolated nucleic acid molecules that have increased expression in response to pathogen infection, the invention further provides compositions and methods for enhancing resistance to pathogen infection.

The compositions of the invention include plant nucleic acid sequences and the amino acid sequences for the polypeptides or partial-length polypeptides encoded thereby which are described herein, or other plant nucleic acid sequences and the amino acid sequences for the polypeptides or partial-length polypeptides encoded thereby which are operably linked to a promoters are useful to provide tolerance or resistance to a plant to a pathogen, preferably by preventing or inhibiting pathogen infection. Methods of the invention involve stably transforming a plant with one or more of at least a portion of these nucleotide sequences which confer tolerance or resistance operably linked to a promoter capable of driving expression of that nucleotide sequence in a plant cell. By "portion" or "fragment", as it relates to a nucleic acid molecule, sequence or segment of the invention, when it is linked to other sequences for expression, is meant a sequence comprising at least 80 nucleotides, more preferably at least 150 nucleotides, and still more preferably at least 400 nucleotides. If not employed for expressing, a "portion" or "fragment" means at least 9, preferably 12, more preferably 15, even more preferably at least 20, consecutive nucleotides, e.g., probes and primers

(oligonucleotides), corresponding to the nucleotide sequence of the nucleic acid molecules of the invention. By "resistant" is meant a plant which exhibits substantially no phenotypic changes as a consequence of infection with the pathogen. By "tolerant" is meant a plant which, although it may exhibit some phenotypic changes as a consequence of infection, does not have a substantially decreased reproductive capacity or substantially altered metabolism.

A method of combating a pathogen in an agricultural crop is also provided. The method comprises introducing to a plant, plant cell, or plant tissue an expression cassette comprising a nucleic acid molecule of the invention comprising an open reading frame so as to yield a transformed differentiated plant, transformed cell or transformed tissue. Transformed cells or tissue can be regenerated to provide a transformed differentiated plant. The transformed differentiated plant preferably expresses the nucleic acid molecule in an amount that confers resistance to the transformed plant to pathogen infection relative to a corresponding nontransformed plant. The present invention also provides a transformed plant prepared by the method, progeny and seed thereof. Examples of plant viruses which may be combated by the present invention include single stranded RNA viruses (with and without envelope), double stranded RNA viruses, and single and double stranded DNA viruses such as (but not limited to) tobacco mosaic virus, cucumber mosaic virus, turnip mosaic virus, turnip vein clearing virus, oilseed rape mosaic virus, tobacco rattle virus, pea enation mosaic virus, barley stripe mosaic virus, potato viruses X and Y, carnation latent virus, beet yellows virus, maize chlorotic virus, tobacco necrosis virus, turnip yellow mosaic virus, tomato bushy stunt virus, southern bean mosaic virus, barley yellow dwarf virus, tomato spotted wilt virus, lettuce necrotic yellows virus, wound tumor virus, maize streak virus, and cauliflower mosaic virus. Other pathogens within the scope of the invention include, but are not limited to, fungi such as *Cochliobolus carbonum*, *Phytophthora infestans*, *Phytophthora sojae*, *Colletesichum*, *Melampsora lini*, *cladosporium fulvum*, *Heminthosporium maydia*, *Peronospora parasitica*, *Puccinia sorghi*, and *Puccinia polysora*; bacteria such as *Phynchosporium secalis*, *Pseudomonas glycinea*, *Xanthomonas oryzae* and, *Fusarium oxyaporium*; and nematodes such as *Globodera rostochiensis*.

For example, the invention provides a nucleic acid molecule comprising a plant nucleotide sequence comprising at least a portion of a key effector gene(s) responsible for host resistance to particular pathogens. To provide resistance or tolerance to a pathogen in a plant, this sequence may be overexpressed individually, in the sense or antisense orientation, or in combination with other sequences to confer improved disease resistance or tolerance to a plant

relative to a plant that does not comprise and/or express the sequence. The overexpression may be constitutive, or it may be preferable to express the effector gene(s) in a tissue-specific manner or from an inducible promoter including a promoter which is responsive to external stimuli, such as chemical application, or to pathogen infection, e.g., so as to avoid possible deleterious effects on plant growth if the effector gene(s) was constitutively expressed. In one embodiment of the invention, the promoter employed may be one that is rapidly and transiently and/or highly transcribed after pathogen infection.

A transformed (transgenic) plant of the invention includes plants, the genome of which is augmented by a nucleic acid molecule of the invention, or in which the corresponding gene has been disrupted, e.g., to result in a loss, a decrease or an alteration, in the function of the product encoded by the gene, which plant may also have increased yields, e.g., under conditions of pathogen infection, and/or produce a better-quality product than the corresponding wild-type plant. The nucleic acid molecules of the invention are thus useful for targeted gene disruption, as well as markers and probes.

For example, the invention includes a pathogen, e.g., virus, tolerant or resistant plant and seed thereof having stably integrated and expressed within its genome, a nucleic acid molecule of the invention. The normal fertile transformed (transgenic) plant may be selfed to yield a substantially homogenous line with respect to viral resistance or tolerance. Individuals of the line, or the progeny thereof, may be crossed with plants which optionally exhibit the trait. In a particular embodiment of the method, the selfing and selection steps are repeated at least five times in order to obtain the homogenous (isogenic) line. Thus, the invention also provides transgenic plants and the products of the transgenic plants.

The invention further includes a nucleotide sequence which is complementary to one (hereinafter "test" sequence) which hybridizes under low, moderate or stringent conditions with the nucleic acid molecules of the invention as well as RNA which is encoded by the nucleic acid molecule. When the hybridization is performed under stringent conditions, either the test or nucleic acid molecule of invention is preferably supported, e.g., on a membrane or DNA chip. Thus, either a denatured test or nucleic acid molecule of the invention is preferably first bound to a support and hybridization is effected for a specified period of time at a temperature of, e.g., between 55 and 70°C, in double strength citrate buffered saline (SC) containing 0.1% SDS followed by rinsing of the support at the same temperature but with a buffer having a reduced SC concentration. Depending upon the degree of stringency required

such reduced concentration buffers are typically single strength SC containing 0.1% SDS, half strength SC containing 0.1% SDS and one-tenth strength SC containing 0.1% SDS.

The invention further provides a method to identify an open reading frame in the genome of a plant cell, the expression of which is altered by pathogen infection of that cell.

- 5 The method comprises contacting a solid substrate comprising a plurality of samples comprising isolated plant nucleic acid of a probe comprising plant nucleic acid, e.g., cRNA, isolated from a pathogen infected plant so as to form a complex. Each individual sample comprises one or more nucleic acid sequences (e.g., oligonucleotides) corresponding to at least a portion of a plant gene. The method may be employed with nucleic acid samples and probes
- 10 from any organism, e.g., any prokaryotic or eukaryotic organism. Preferably, the nucleic acid sample and probes are from a plant, such as a dicot or monocot. More preferably the nucleic acid samples and probes are from a cereal plant. Even more preferably the nucleic acids and probes are from a crop plant. A second plurality of samples on a solid substrate, i.e., a DNA chip, each comprising a plurality of samples comprising isolated plant nucleic acid is contacted
- 15 with a probe comprising plant nucleic acid isolated from an uninfected or infected control (mutant) plant so as to form a complex. Then complex formation between the samples and probes comprising nucleic acid from infected or control cells compared. For example, potato virus X, tobacco mosaic virus, tobnavirus, cucumber mosaic virus and geminivirus are known to infect *Arabidopsis*. Thus, *Arabidopsis* genes, the expression of which is altered in response
- 20 to infection by any of these viruses, can be identified. Regions that are 5N to the start codon for the gene can then be identified and/or isolated.

- The invention further provides a method for identifying a plant cell infected with a pathogen. The method comprises contacting nucleic acid obtained from a plant cell suspected of being infected with a pathogen with oligonucleotides corresponding to a portion of a
- 25 plurality of sequences selected from SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737 under conditions effective to amplify those sequences. Then the presence of the amplified product is detected or determined. The presence of two or more amplified products, e.g., in an amount that is different than the amount of the corresponding amplified products from an uninfected plant, each corresponding to two or more SEQ ID NOs: 1-953, 1954-1966, 2000-
- 30 2129, or 2662-4737 is indicative of pathogen infection.

The invention further provides a method for identifying a plant cell infected with a pathogen. The method comprises contacting a protein sample obtained from a plant cell suspected of being infected with a pathogen with an agent that specifically binds a polypeptide

encoded by an open reading frame comprising SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737 so as to form a complex. Then the presence or amount of complex formation is detected or determined.

The invention provides an additional method for identifying a plant cell infected with a pathogen. The method comprises hybridizing a probe selected from SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737 to nucleic acid obtained from a plant cell suspected of being infected with a pathogen. The amount of the probe hybridized to nucleic acid obtained from a cell suspected of being infected with a virus is compared to hybridization of the probe to nucleic acid isolated from an uninfected cell. A change in the amount of at least two probes that hybridize to nucleic acid isolated from a cell suspected of being infected by a virus relative to hybridization of at least two probes to nucleic acid isolated from an uninfected cell is indicative of viral infection.

A method to shuffle the nucleic acids of the invention is provided. This method involves fragmentation of a nucleic acid corresponding to a nucleic acid sequence listed in SEQ ID NOs: 1-953, 1954-1966, 2000-2129 or 2662-4737, the orthologs thereof, and the corresponding genes, followed by religation. This method allows for the production of polypeptides having altered activity relative to the native form of the polypeptide. Accordingly, the invention provides cells and transgenic plants containing nucleic acid segments produced through shuffling that encode polypeptides having altered activity relative to the corresponding native polypeptide.

A computer readable medium containing the nucleic acid sequences of the invention as well as methods of use for the computer readable medium are provided. This medium allows a nucleic acid segment corresponding to a nucleic acid sequence listed in SEQ ID NOs:1-953, 2137-2661, 1954-1966, 2000-2129, 2662-4737 or 4738-6813 to be used as a reference sequence to search against databases. This medium also allows for computer-based manipulation of a nucleic acid sequence corresponding to a nucleic acid sequence listed in SEQ ID NOs :1-953, 2137-2661, 1954-1966, 2000-2129, 2662-4737 or 4738-6813, and the corresponding gene and polypeptide encoded by the nucleic acid sequence.

Therefore, another embodiment of the present invention provides a method of using known inducers or inhibitors of genes identified as being important in plant-pathogen interactions to induce genes that are important in resistance, or to inhibit genes that are downregulated in resistance.

Thus, some of the isolated nucleic acid molecules of the invention are useful in a method of combating a pathogen in an agricultural crop. The method comprises introducing to a plant an expression cassette comprising a nucleic acid molecule of the invention so as to yield a transformed differentiated plant. The transformed differentiated plant expresses the nucleic acid molecule in an amount that confers resistance to the transformed plant to infection relative to a corresponding nontransformed plant.

### **Detailed Description of the Invention**

#### **I. Definitions**

10

The term "gene" is used broadly to refer to any segment of nucleic acid associated with a biological function. Thus, genes include coding sequences and/or the regulatory sequences required for their expression. For example, gene refers to a nucleic acid fragment that expresses mRNA or functional RNA, or encodes a specific protein, and which includes regulatory sequences. Genes also include nonexpressed DNA segments that, for example, form recognition sequences for other proteins. Genes can be obtained from a variety of sources, including cloning from a source of interest or synthesizing from known or predicted sequence information, and may include sequences designed to have desired parameters.

The term "native" or "wild type" gene refers to a gene that is present in the genome of an untransformed cell, i.e., a cell not having a known mutation.

A "marker gene" encodes a selectable or screenable trait.

The term "chimeric gene" refers to any gene that contains 1) DNA sequences, including regulatory and coding sequences, that are not found together in nature, or 2) sequences encoding parts of proteins not naturally adjoined, or 3) parts of promoters that are not naturally adjoined. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or comprise regulatory sequences and coding sequences derived from the same source, but arranged in a manner different from that found in nature.

A "transgene" refers to a gene that has been introduced into the genome by transformation and is stably maintained. Transgenes may include, for example, genes that are either heterologous or homologous to the genes of a particular plant to be transformed. Additionally, transgenes may comprise native genes inserted into a non-native organism, or chimeric genes. The term "endogenous gene" refers to a native gene in its natural location in

the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism but that is introduced by gene transfer.

An "oligonucleotide" corresponding to a nucleotide sequence of the invention, e.g., for use in probing or amplification reactions, may be about 30 or fewer nucleotides in length (e.g., 9, 12, 15, 18, 20, 21 or 24, or any number between 9 and 30). Generally specific primers are upwards of 14 nucleotides in length. For optimum specificity and cost effectiveness, primers of 16 to 24 nucleotides in length may be preferred. Those skilled in the art are well versed in the design of primers for use processes such as PCR. If required, probing can be done with entire restriction fragments of the gene disclosed herein which may be 100's or even 1000's of nucleotides in length.

The terms "protein," "peptide" and "polypeptide" are used interchangeably herein.

The nucleotide sequences of the invention can be introduced into any plant. The genes to be introduced can be conveniently used in expression cassettes for introduction and expression in any plant of interest. Such expression cassettes will comprise the transcriptional initiation region of the invention linked to a nucleotide sequence of interest. Preferred promoters include constitutive, tissue-specific, developmental-specific, inducible and/or viral promoters. Such an expression cassette is provided with a plurality of restriction sites for insertion of the gene of interest to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes. The cassette will include in the 5'-3' direction of transcription, a transcriptional and translational initiation region, a DNA sequence of interest, and a transcriptional and translational termination region functional in plants. The termination region may be native with the transcriptional initiation region, may be native with the DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also, Guerineau et al., 1991; Proudfoot, 1991; Sanfacon et al., 1991; Mogen et al., 1990; Munroe et al., 1990; Ballas et al., 1989; Joshi et al., 1987.

"Coding sequence" refers to a DNA or RNA sequence that codes for a specific amino acid sequence and excludes the non-coding sequences. It may constitute an "uninterrupted coding sequence", i.e., lacking an intron, such as in a cDNA or it may include one or more introns bounded by appropriate splice junctions. An "intron" is a sequence of RNA which is contained in the primary transcript but which is removed through cleavage and re-ligation of the RNA within the cell to create the mature mRNA that can be translated into a protein.



The terms "open reading frame" and "ORF" refer to the amino acid sequence encoded between translation initiation and termination codons of a coding sequence. The terms "initiation codon" and "termination codon" refer to a unit of three adjacent nucleotides ('codon') in a coding sequence that specifies initiation and chain termination, respectively, of protein synthesis (mRNA translation).

A "functional RNA" refers to an antisense RNA, ribozyme, or other RNA that is not translated.

The term "RNA transcript" refers to the product resulting from RNA polymerase catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA" (mRNA) refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a single- or a double-stranded DNA that is complementary to and derived from mRNA.

"Regulatory sequences" and "suitable regulatory sequences" each refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences include enhancers, promoters, translation leader sequences, introns, and polyadenylation signal sequences. They include natural and synthetic sequences as well as sequences which may be a combination of synthetic and natural sequences. As is noted above, the term "suitable regulatory sequences" is not limited to promoters.

"5' non-coding sequence" refers to a nucleotide sequence located 5' (upstream) to the coding sequence. It is present in the fully processed mRNA upstream of the initiation codon and may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency (Turner et al., 1995).

"3' non-coding sequence" refers to nucleotide sequences located 3' (downstream) to a coding sequence and include polyadenylation signal sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al., 1989.

The term "translation leader sequence" refers to that DNA sequence portion of a gene

between the promoter and coding sequence that is transcribed into RNA and is present in the fully processed mRNA upstream (5') of the translation start codon. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency.

5           The term "mature" protein refers to a post-translationally processed polypeptide without its signal peptide. "Precursor" protein refers to the primary product of translation of an mRNA. "Signal peptide" refers to the amino terminal extension of a polypeptide, which is translated in conjunction with the polypeptide forming a precursor peptide and which is required for its entrance into the secretory pathway. The term "signal sequence" refers to a  
10       nucleotide sequence that encodes the signal peptide.

          The term "intracellular localization sequence" refers to a nucleotide sequence that encodes an intracellular targeting signal. An "intracellular targeting signal" is an amino acid sequence that is translated in conjunction with a protein and directs it to a particular sub-cellular compartment. "Endoplasmic reticulum (ER) stop transit signal" refers to a carboxy-  
15       terminal extension of a polypeptide, which is translated in conjunction with the polypeptide and causes a protein that enters the secretory pathway to be retained in the ER. "ER stop transit sequence" refers to a nucleotide sequence that encodes the ER targeting signal. Other intracellular targeting sequences encode targeting signals active in seeds and/or leaves and vacuolar targeting signals.

20           "Pathogen" as used herein includes but is not limited to bacteria, fungi, yeast, oomycetes and virus, e.g., American wheat striate mosaic virus mosaic (AWSMV), barley stripe mosaic virus (BSMV), barley yellow dwarf virus (BYDV), Brome mosaic virus (BMV), cereal chlorotic mottle virus (CCMV), corn chlorotic vein banding virus (CCVBV), maize chlorotic mottle virus (MCMV), maize dwarf mosaic virus (MDMV), A or B, wheat streak  
25       mosaic virus (WSMV), cucumber mosaic virus (CMV), cynodon chlorotic streak virus (CCSV), Johnsongrass mosaic virus (JGMV), maize bushy stunt or mycoplasma-like organism (NJILO), maize chlorotic dwarf virus (MCDV), maize chlorotic mottle virus (MCMV), maize dwarf mosaic virus (MDMV) strains A, D, E and F, maize leaf fleck virus (MLFV), maize line virus (NELV), maize mosaic virus (MMV), maize mottle and chlorotic stunt virus, maize  
30       pellucid ringspot virus (MPRV), maize raya gruesa virus (MRGV), maize rayado fino virus (MRFV), maize red leaf and red stripe virus (MRSV), maize ring mottle virus (MRMV), maize rio cuarto virus (MRCV), maize rough dwarf virus (MRDV), maize sterile stunt virus (strains of barley yellow striate virus), maize streak virus (MSV), maize chlorotic stripe, maize

hoja Maize stripe virus blanca, maize stunting virus, maize tassel abortion virus (MTAV), maize vein enation virus (MVEV), maize wallaby ear virus (MAVEV), maize white leaf virus, maize white line mosaic virus (NTVVLMV), millet red leaf virus (NMV), Northern cereal mosaic virus (NCMV), oat pseudorosette virus, oat sterile dwarf virus (OSDV), rice black-streaked dwarf virus (RBSDV), rice stripe virus (RSV), sorghum mosaic virus (SrMV),  
 5 formerly sugarcane mosaic virus (SCMV) strains H, I and M, sugarcane Fiji disease virus (FDV), sugarcane mosaic virus (SCMV) strains A, B, D, E, SC, BC, Sabi and NM vein enation virus, and wheat spot mosaic virus (WSMV).

Bacterial pathogens include but are not limited to *Pseudomonas avenae* subsp. *avenae*,  
 10 *Xanthomonas campestris* pv. *holcicola*, *Enterobacter dissolvens*, *Erwinia dissolvens*, *Erwinia carotovora* subsp. *carotovora*, *Erwinia chrysanthemi* pv. *zeae*, *Pseudomonas andropogonis*, *Pseudomonas syringae* pv. *coronafaciens*, *Clavibacter michiganensis* subsp., *Corynebacterium michiganense* pv. *nebraskense*, *Pseudomonas syringae* pv. *syringae*, *Herniparasitic bacteria* (see under fungi), *Bacillus subtilis*, *Erwinia stewartii*, and *Spiroplasma kunkelii*.

Fungal pathogens include but are not limited to *Collelotrichum graminicola*,  
*Glomerella graminicola* Politis, *Glomerella lucumanensis*, *Aspergillus flavus*, *Rhizoctonia solani* Kuhn, *Thanatephorus cucumeris*, *Acremonium strictum* W. Gams, *Cephalosporium acremonium* Auct. non Corda Black *Lasiodiplodia theobromae* = *BoIr odiplochia y theobromae*  
*Borde blanco Marasmiellus* sp., *Physoderma maydis*, *Cephalosporium Corticium sasakii*,  
 20 *Curvularia clavata*, *C. maculans*, *Cochhobolus eragrostidis*, *Curvularia inaequahs*, *C. intermedia* (teleomorph *Cochhobolus intermedius*), *Curvularia lunata* (teleomorph: *Cochliobolus lunatus*), *Curvularia pallescens* (teleomorph - *Cochliobolus pallescens*), *Curvularia senegalensis*, *C. luberculata* (teleomorph: *Cochliobolus tuberculatus*), *Didymella exitalis* *Diplodiaftumenti* (teleomorph - *Botryosphaeriafestucae*), *Diplodia maydis* =  
 25 *Stenocarpella maydis*, *Stenocarpella macrospora* = *Diplodia macrospora*, *Sclerophthora rayssiae* var. *zeae*, *Sclerophthora macrospora* = *Sclerospora macrospora*, *Sclerospora graminicola*, *Peronosclerospora maydis* = *Sclerospora maydis*, *Peronosclerospora philippinensis*, *Sclerospora philippinensis*, *Peronosclerospora sorghi* = *Sclerospora sorghi*, *Peronosclerospora spontanea* = *Sclerospora spontanea*, *Peronosclerospora sacchari* =  
 30 *Sclerospora sacchari*, *Nigrospora oryzae* (teleomorph: *Khuskia oryzae*) A. *Iternaria alternala* = *A. tenuis*, *Aspergillus glaucus*, *A. niger*, *Aspergillus spp.*, *Botrytis cinerea*, *Cunninghamella sp.*, *Curvulariapallescens*, *Doratomyces slemonitis* = *Cephalotrichum slemonitis*, *Fusarium culmorum*, *Gonatobotrys simplex*, *Pithomyces maydicus*, *Rhizopus microsporus* Tiegh., *R.*

- stolonifer* = *R. nigricans*, *Scopulariopsis brumptii*, *Claviceps gigantea* (anamorph: *Sphacelia* sp.) *Aureobasidium zeae* = *Kabatiella zeae*, *Fusarium subglutinans* = *F. moniliforme* var. *subglutinans*, *Fusarium moniliforme*, *Fusarium avenaceum* (teleomorph - *Gibberella avenacea*), *Botryosphaeria zeae* = *Physalospora zeae* (anamorph: *Allacrophoma zeae*),
- 5 *Cercospora sorghi* = *C. sorghi* var. *maydis*, *Helminthosporium pedicellatum* (teleomorph: *Selosphaeriapedicellata*), *Cladosporium cladosporioides* = *Hormodendrum cladosporioides*, *C. herbarum* (teleomorph - *Mycosphaerella tassiana*), *Cephalosporium maydis*, *A. Iternaria alternata*, *A. scochyta maydis*, *A. tritici*, *A. zeicola*, *Bipolaris victoriae*, *Helminthosporium victoriae* (teleomorph *Cochhoholus victoriae*), *C sativus* (anamorph: *Bipolaris sorokiniana* =
- 10 *H. sorokinianum* = *H. sativum*), *Epicoccum nigrum*, *Exserohilum prolatum* = *Drechslera prolata* (teleomorph: *Setosphaeriaprolata*), *Graphium penicillioides*, *Leptosphaeria maydis*, *Leptothyrium zeae*, *Ophiosphaerella herpotricha* (anamorph - *Scolecosporella* sp.), *Pataphaeosphaeria michotii*, *Phoma* sp., *Septoria zeae*, *S. zeicola*, *S. zeina* *Setosphaeria turcica*, *Exserohilum turcicum* = *Helminthosporium furcicum*, *Cochhoholus carbonum*,
- 15 *Bipolaris zeicola* = *Helminthosporium carbonum*, *Penicillium* spp., *P. chrysogenum*, *P. expansum*, *P. oxalicum*, *Phaeocytostroma ambiguum*, *Phaeocylosporella zeae*, *Phaeosphaeria maydis* = *Sphaerulina maydis*, *Botryosphaeriafestucae* = *Physalospora zeicola* (anamorph: *Diplodiafumensis*), *Herniparasitic bacteria and fungi* *Pyrenochaeta Phoma terrestris* = *Pyrenochaeta terrestris*, *Pythium* spp., *P. arrhenomanes*, *P. graminicola*, *Pythium*
- 20 *aphanidermatum* = *P. hutleri* L., *Rhizoctonia zeae* (teleomorph: *Waitea circinata*), *Rhizoctonia solani*, minor *A Iternaria alternata*, *Cercospora sorghi*, *Dictochaetafritilis*, *Fusarium acuminatum* (teleomorph *Gibberella acuminata*), *E. equiseti* (teleomorph: *G. intricans*), *E. oxysporum*, *E. pallidroseum*, *E. poae*, *E. roseum*, *G. cyanogena* (anamorph: *E. sulphureum*), *Microdochium holleyi*, *Mucor* sp., *Periconia circinata*, *Phytophthora cactorum*,
- 25 *P. drechsleri*, *P. nicotianae* var. *parasitica*, *Rhizopus arrhizus*, *Setosphaeria rostrata*, *Exserohilum rostratum* = *Helminthosporium rostratum*, *Puccinia sorghi*, *Physopella pallescens*, *P. zeae*, *Sclerotium rofsii* Sacc. (teleomorph- *Athelia rofsii*), *Bipolaris sorokiniana*, *B. zeicola* = *Helminthosporium carbonum*, *Diplodia maydis*, *Exserohilum pedicellatum*, *Exserohilum furcicum* = *Helminthosporium turcicum*, *Fusarium avenaceum*, *E.*
- 30 *culmorum*, *E. moniliforme*, *Gibberella zeae* (anamorph - *E. graminearum*), *Macrophominaphaseolina*, *Penicillium* spp., *Phomopsis* sp., *Pythium* spp., *Rhizoctonia solani*, *R. zeae*, *Sclerotium rolfsii*, *Spicaria* sp., *Selenophoma* sp., *Gaeumannomyces graminis*, *Myrothecium gramineum*, *Monascus purpureus*, *M. ruber* Smut, *Ustilago zeae* = *U. maydis*

*Smut, Ustilaginoidea virens Smut, Sphacelotheca reiliana = Sporisorium holci, Cochliobolus heterostrophus* (anamorph: *Bipolaris maydis = Helminthosporium maydis*), *Stenocarpella macrospora = Diplodia macrospora, Cercospora sorghi, Fusarium episphaeria, E. merismoides, F. oxysporum Schlechtend, E. poae, E. roseum, E. solani* (teleomorph: *Nectria haematococca*), *F. tricinturn, Mariannaea elegans, Mucor sp., Rhopoglyphus zeae, Spicaria sp., Aspergillus spp., Penicillium spp., Trichoderma viride = T lignorum teleomorph: Hypocrea sp., Stenocarpella maydis = Diplodia zeae, Ascochyta ischaemi, Phyllosticta maydis* (telomorph: *Mycosphaerella zeae-maydis*), and *Gloeocercospora sorghi*.

Parasitic nematodes include but are not limited to *Awl Dolichodorus spp., D. heterocephalus* Bulb and stem (Europe), *Ditylenchus dipsaci* Burrowing Radopholus similis Cyst Heterodera avenae, *H. zeae, Punctodera chaltoensis* Dagger *Xiphinema spp., X americanum, X mediterraneum* False root-knot *Nacobbus dorsalis* Lance, Columbia *Hoplolaimus columbus* Lance *Hoplolaimus spp., H. galeatus* Lesion *Pratylenchus spp., P. brachyurus, P. crenatus, P. hexincisus, P. neglectus, P. penetrans, P. scribneri, P. thornei, P. zeae* Needle Longidorus spp., *L. breviannulatus* Ring *Criconemella spp., C. ornata* Root-knot Meloidogyne spp., *M. chitwoodi, M. incognita, M. javanica* Spiral *Helicotylenchus spp., Belonolaimus spp., B. longicaudatus* Stubby-root *Paratrichodorus spp., P. christiei, P. minor, Quinisulcius aculus, and Trichodorus spp.*

"Promoter" refers to a nucleotide sequence, usually upstream (5') to its coding sequence, which controls the expression of the coding sequence by providing the recognition for RNA polymerase and other factors required for proper transcription. "Promoter" includes a minimal promoter that is a short DNA sequence comprised of a TATA box and other sequences that serve to specify the site of transcription initiation, to which regulatory elements are added for control of expression. "Promoter" also refers to a nucleotide sequence that includes a minimal promoter plus regulatory elements that is capable of controlling the expression of a coding sequence or functional RNA. This type of promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an "enhancer" is a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue specificity of a promoter. It is capable of operating in both orientations (normal or flipped), and is capable of functioning even when moved either upstream or downstream from the promoter. Both enhancers and other upstream promoter elements bind sequence-specific DNA-binding proteins that mediate their effects. Promoters

may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even be comprised of synthetic DNA segments. A promoter may also contain DNA sequences that are involved in the binding of protein factors which control the effectiveness of transcription initiation in response to  
5 physiological or developmental conditions.

The "initiation site" is the position surrounding the first nucleotide that is part of the transcribed sequence, which is also defined as position +1. With respect to this site all other sequences of the gene and its controlling regions are numbered. Downstream sequences (i.e., further protein encoding sequences in the 3' direction) are denominated positive, while  
10 upstream sequences (mostly of the controlling regions in the 5' direction) are denominated negative.

Promoter elements, particularly a TATA element, that are inactive or that have greatly reduced promoter activity in the absence of upstream activation are referred to as "minimal or core promoters." In the presence of a suitable transcription factor, the minimal promoter  
15 functions to permit transcription. A "minimal or core promoter" thus consists only of all basal elements needed for transcription initiation, e.g., a TATA box and/or an initiator.

"Constitutive expression" refers to expression using a constitutive or regulated promoter. "Conditional" and "regulated expression" refer to expression controlled by a regulated promoter.

20 "Constitutive promoter" refers to a promoter that is able to express the open reading frame (ORF) that it controls in all or nearly all of the plant tissues during all or nearly all developmental stages of the plant. Each of the transcription-activating elements do not exhibit an absolute tissue-specificity, but mediate transcriptional activation in most plant parts at a level of  $\geq 1\%$  of the level reached in the part of the plant in which transcription is most active.

25 "Regulated promoter" refers to promoters that direct gene expression not constitutively, but in a temporally- and/or spatially-regulated manner, and includes both tissue-specific and inducible promoters. It includes natural and synthetic sequences as well as sequences which may be a combination of synthetic and natural sequences. Different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or  
30 in response to different environmental conditions. New promoters of various types useful in plant cells are constantly being discovered, numerous examples may be found in the compilation by Okamuro et al. (1989). Typical regulated promoters useful in plants include but are not limited to safener-inducible promoters, promoters derived from the tetracycline-

inducible system, promoters derived from salicylate-inducible systems, promoters derived from alcohol-inducible systems, promoters derived from glucocorticoid-inducible system, promoters derived from pathogen-inducible systems, and promoters derived from ecdysome-inducible systems.

5 "Tissue-specific promoter" refers to regulated promoters that are not expressed in all plant cells but only in one or more cell types in specific organs (such as leaves or seeds), specific tissues (such as embryo or cotyledon), or specific cell types (such as leaf parenchyma or seed storage cells). These also include promoters that are temporally regulated, such as in early or late embryogenesis, during fruit ripening in developing seeds or fruit, in fully  
10 differentiated leaf, or at the onset of senescence.

"Inducible promoter" refers to those regulated promoters that can be turned on in one or more cell types by an external stimulus, such as a chemical, light, hormone, stress, or a pathogen.

"Operably-linked" refers to the association of nucleic acid sequences on single nucleic  
15 acid fragment so that the function of one is affected by the other. For example, a regulatory DNA sequence is said to be "operably linked to" or "associated with" a DNA sequence that codes for an RNA or a polypeptide if the two sequences are situated such that the regulatory DNA sequence affects expression of the coding DNA sequence (i.e., that the coding sequence or functional RNA is under the transcriptional control of the promoter). Coding sequences can  
20 be operably-linked to regulatory sequences in sense or antisense orientation.

"Expression" refers to the transcription and/or translation of an endogenous gene, ORF or portion thereof, or a transgene in plants. For example, in the case of antisense constructs, expression may refer to the transcription of the antisense DNA only. In addition, expression refers to the transcription and stable accumulation of sense (mRNA) or functional RNA.  
25 Expression may also refer to the production of protein.

"Specific expression" is the expression of gene products which is limited to one or a few plant tissues (spatial limitation) and/or to one or a few plant developmental stages (temporal limitation). It is acknowledged that hardly a true specificity exists: promoters seem to be preferably switch on in some tissues, while in other tissues there can be no or only little  
30 activity. This phenomenon is known as leaky expression. However, with specific expression in this invention is meant preferable expression in one or a few plant tissues.

The "expression pattern" of a promoter (with or without enhancer) is the pattern of expression levels which shows where in the plant and in what developmental stage

transcription is initiated by said promoter. Expression patterns of a set of promoters are said to be complementary when the expression pattern of one promoter shows little overlap with the expression pattern of the other promoter. The level of expression of a promoter can be determined by measuring the 'steady state' concentration of a standard transcribed reporter mRNA. This measurement is indirect since the concentration of the reporter mRNA is dependent not only on its synthesis rate, but also on the rate with which the mRNA is degraded. Therefore, the steady state level is the product of synthesis rates and degradation rates.

The rate of degradation can however be considered to proceed at a fixed rate when the transcribed sequences are identical, and thus this value can serve as a measure of synthesis rates. When promoters are compared in this way techniques available to those skilled in the art are hybridization S1-RNase analysis, northern blots and competitive RT-PCR. This list of techniques in no way represents all available techniques, but rather describes commonly used procedures used to analyze transcription activity and expression levels of mRNA.

The analysis of transcription start points in practically all promoters has revealed that there is usually no single base at which transcription starts, but rather a more or less clustered set of initiation sites, each of which accounts for some start points of the mRNA. Since this distribution varies from promoter to promoter the sequences of the reporter mRNA in each of the populations would differ from each other. Since each mRNA species is more or less prone to degradation, no single degradation rate can be expected for different reporter mRNAs. It has been shown for various eukaryotic promoter sequences that the sequence surrounding the initiation site ('initiator') plays an important role in determining the level of RNA expression directed by that specific promoter. This includes also part of the transcribed sequences. The direct fusion of promoter to reporter sequences would therefore lead to suboptimal levels of transcription.

A commonly used procedure to analyze expression patterns and levels is through determination of the 'steady state' level of protein accumulation in a cell. Commonly used candidates for the reporter gene, known to those skilled in the art are  $\beta$ -glucuronidase (GUS), chloramphenicol acetyl transferase (CAT) and proteins with fluorescent properties, such as green fluorescent protein (GFP) from *Aequora victoria*. In principle, however, many more proteins are suitable for this purpose, provided the protein does not interfere with essential plant functions. For quantification and determination of localization a number of tools are suited. Detection systems can readily be created or are available which are based on, e.g.,



immunochemical, enzymatic, fluorescent detection and quantification. Protein levels can be determined in plant tissue extracts or in intact tissue using *in situ* analysis of protein expression.

Generally, individual transformed lines with one chimeric promoter reporter construct will vary in their levels of expression of the reporter gene. Also frequently observed is the phenomenon that such transformants do not express any detectable product (RNA or protein). The variability in expression is commonly ascribed to 'position effects', although the molecular mechanisms underlying this inactivity are usually not clear.

The term "average expression" is used here as the average level of expression found in all lines that do express detectable amounts of reporter gene, so leaving out of the analysis plants that do not express any detectable reporter mRNA or protein.

"Root expression level" indicates the expression level found in protein extracts of complete plant roots. Likewise, leaf, and stem expression levels, are determined using whole extracts from leaves and stems. It is acknowledged however, that within each of the plant parts just described, cells with variable functions may exist, in which promoter activity may vary.

"Non-specific expression" refers to constitutive expression or low level, basal ('leaky') expression in undesired cells or tissues from a 'regulated promoter'.

"Altered levels" refers to the level of expression in transgenic organisms that differs from that of normal or untransformed organisms.

"Overexpression" refers to the level of expression in transgenic cells or organisms that exceeds levels of expression in normal or untransformed (nontransgenic) cells or organisms.

"Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of protein from an endogenous gene or a transgene.

"Co-suppression" and "transwitch" each refer to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar transgene or endogenous genes (U.S. Patent No. 5,231,020).

"Gene silencing" refers to homology-dependent suppression of viral genes, transgenes, or endogenous nuclear genes. Gene silencing may be transcriptional, when the suppression is due to decreased transcription of the affected genes, or post-transcriptional, when the suppression is due to increased turnover (degradation) of RNA species homologous to the affected genes (English et al., 1996). Gene silencing includes virus-induced gene silencing (Ruiz et al. 1998).

"Silencing suppressor" gene refers to a gene whose expression leads to counteracting gene silencing and enhanced expression of silenced genes. Silencing suppressor genes may be of plant, non-plant, or viral origin. Examples include, but are not limited to HC-Pro, P1-HC-Pro, and 2b proteins. Other examples include one or more genes in TGMV-B genome.

5       The terms "heterologous DNA sequence," "exogenous DNA segment" or "heterologous nucleic acid," as used herein, each refer to a sequence that originates from a source foreign to the particular host cell or, if from the same source, is modified from its original form. Thus, a heterologous gene in a host cell includes a gene that is endogenous to the particular host cell but has been modified through, for example, the use of DNA shuffling. The terms also include  
10 non-naturally occurring multiple copies of a naturally occurring DNA sequence. Thus, the terms refer to a DNA segment that is foreign or heterologous to the cell, or homologous to the cell but in a position within the host cell nucleic acid in which the element is not ordinarily found. Exogenous DNA segments are expressed to yield exogenous polypeptides. A "homologous" DNA sequence is a DNA sequence that is naturally associated with a host cell  
15 into which it is introduced.

"Homologous to" in the context of nucleotide sequence identity refers to the similarity between the nucleotide sequence of two nucleic acid molecules or between the amino acid sequences of two protein molecules. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood  
20 by those skilled in the art (as described in Haines and Higgins (eds.), *Nucleic Acid Hybridization*, IRL Press, Oxford, U.K.), or by the comparison of sequence similarity between two nucleic acids or proteins.

The term "substantially similar" refers to nucleotide and amino acid sequences that represent functional and/or structural equivalents of *Arabidopsis* sequences disclosed herein.  
25 For example, altered nucleotide sequences which simply reflect the degeneracy of the genetic code but nonetheless encode amino acid sequences that are identical to a particular amino acid sequence are substantially similar to the particular sequences. In addition, amino acid sequences that are substantially similar to a particular sequence are those wherein overall amino acid identity is at least 65% or greater to the instant sequences. Modifications that  
30 result in equivalent nucleotide or amino acid sequences are well within the routine skill in the art. Moreover, the skilled artisan recognizes that equivalent nucleotide sequences encompassed by this invention can also be defined by their ability to hybridize, under low, moderate and/or stringent conditions (e.g., 0.1X SSC, 0.1% SDS, 65°C), with the nucleotide

sequences that are within the literal scope of the instant claims.

"Target gene" refers to a gene on the replicon that expresses the desired target coding sequence, functional RNA, or protein. The target gene is not essential for replicon replication. Additionally, target genes may comprise native non-viral genes inserted into a non-native organism, or chimeric genes, and will be under the control of suitable regulatory sequences. Thus, the regulatory sequences in the target gene may come from any source, including the virus. Target genes may include coding sequences that are either heterologous or homologous to the genes of a particular plant to be transformed. However, target genes do not include native viral genes. Typical target genes include, but are not limited to genes encoding a structural protein, a seed storage protein, a protein that conveys herbicide resistance, and a protein that conveys insect resistance. Proteins encoded by target genes are known as "foreign proteins". The expression of a target gene in a plant will typically produce an altered plant trait.

The term "altered plant trait" means any phenotypic or genotypic change in a transgenic plant relative to the wild-type or non-transgenic plant host.

"Transcription Stop Fragment" refers to nucleotide sequences that contain one or more regulatory signals, such as polyadenylation signal sequences, capable of terminating transcription. Examples include the 3' non-regulatory regions of genes encoding nopaline synthase and the small subunit of ribulose biphosphate carboxylase.

"Replication gene" refers to a gene encoding a viral replication protein. In addition to the ORF of the replication protein, the replication gene may also contain other overlapping or non-overlapping ORF(s), as are found in viral sequences in nature. While not essential for replication, these additional ORFs may enhance replication and/or viral DNA accumulation. Examples of such additional ORFs are AC3 and AL3 in ACMV and TGMV geminiviruses, respectively.

"Chimeric *trans*-acting replication gene" refers either to a replication gene in which the coding sequence of a replication protein is under the control of a regulated plant promoter other than that in the native viral replication gene, or a modified native viral replication gene, for example, in which a site specific sequence(s) is inserted in the 5' transcribed but untranslated region. Such chimeric genes also include insertion of the known sites of

replication protein binding between the promoter and the transcription start site that attenuate transcription of viral replication protein gene.

"Chromosomally-integrated" refers to the integration of a foreign gene or DNA construct into the host DNA by covalent bonds. Where genes are not "chromosomally integrated" they may be "transiently expressed." Transient expression of a gene refers to the expression of a gene that is not integrated into the host chromosome but functions independently, either as part of an autonomously replicating plasmid or expression cassette, for example, or as part of another biological system such as a virus.

"Production tissue" refers to mature, harvestable tissue consisting of non-dividing, terminally-differentiated cells. It excludes young, growing tissue consisting of germline, meristematic, and not-fully-differentiated cells.

"Germline cells" refer to cells that are destined to be gametes and whose genetic material is heritable.

"*Trans*-activation" refers to switching on of gene expression or replicon replication by the expression of another (regulatory) gene in *trans*.

The term "transformation" refers to the transfer of a nucleic acid fragment into the genome of a host cell, resulting in genetically stable inheritance. Host cells containing the transformed nucleic acid fragments are referred to as "transgenic" cells, and organisms comprising transgenic cells are referred to as "transgenic organisms". Examples of methods of transformation of plants and plant cells include *Agrobacterium*-mediated transformation (De Blaere et al., 1987) and particle bombardment technology (Klein et al. 1987; U.S. Patent No. 4,945,050). Whole plants may be regenerated from transgenic cells by methods well known to the skilled artisan (see, for example, Fromm et al., 1990).

"Transformed," "transgenic," and "recombinant" refer to a host organism such as a bacterium or a plant into which a heterologous nucleic acid molecule has been introduced. The nucleic acid molecule can be stably integrated into the genome generally known in the art and are disclosed in Sambrook et al., 1989. See also Innis et al., 1995 and Gelfand, 1995; and Innis and Gelfand, 1999. Known methods of PCR include, but are not limited to, methods using paired primers, nested primers, single specific primers, degenerate primers, gene-specific primers, vector-specific primers, partially mismatched primers, and the like. For example, "transformed," "transformant," and "transgenic" plants or calli have been through the transformation process and contain a foreign gene integrated into their chromosome. The term "untransformed" refers to normal plants that have not been through the transformation process.

"Transiently transformed" refers to cells in which transgenes and foreign DNA have been introduced (for example, by such methods as *Agrobacterium*-mediated transformation or biolistic bombardment), but not selected for stable maintenance.

5 "Stably transformed" refers to cells that have been selected and regenerated on a selection media following transformation.

"Transient expression" refers to expression in cells in which a virus or a transgene is introduced by viral infection or by such methods as *Agrobacterium*-mediated transformation, electroporation, or biolistic bombardment, but not selected for its stable maintenance.

10 "Genetically stable" and "heritable" refer to chromosomally-integrated genetic elements that are stably maintained in the plant and stably inherited by progeny through successive generations.

"Primary transformant" and "T0 generation" refer to transgenic plants that are of the same genetic generation as the tissue which was initially transformed (i.e., not having gone through meiosis and fertilization since transformation).

15 "Secondary transformants" and the "T1, T2, T3, etc. generations" refer to transgenic plants derived from primary transformants through one or more meiotic and fertilization cycles. They may be derived by self-fertilization of primary or secondary transformants or crosses of primary or secondary transformants with other transformed or untransformed plants.

"Wild-type" refers to a virus or organism found in nature without any known mutation.

20 "Genome" refers to the complete genetic material of an organism.

The term "nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form, composed of monomers (nucleotides) containing a sugar, phosphate and a base which is either a purine or pyrimidine. Unless specifically limited, the term encompasses nucleic acids containing known analogs of natural  
25 nucleotides which have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions) and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may  
30 be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al., 1991; Ohtsuka et al., 1985; Rossolini et al. 1994). A "nucleic acid fragment" is a fraction of a given nucleic acid molecule. In higher plants, deoxyribonucleic acid (DNA) is the genetic material

while ribonucleic acid (RNA) is involved in the transfer of information contained within DNA into proteins. The term "nucleotide sequence" refers to a polymer of DNA or RNA which can be single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases capable of incorporation into DNA or RNA polymers. The terms "nucleic acid" or  
5 "nucleic acid sequence" may also be used interchangeably with gene, cDNA, DNA and RNA encoded by a gene.

The invention encompasses isolated or substantially purified nucleic acid or protein compositions. In the context of the present invention, an "isolated" or "purified" DNA molecule or an "isolated" or "purified" polypeptide is a DNA molecule or polypeptide that, by  
10 the hand of man, exists apart from its native environment and is therefore not a product of nature. An isolated DNA molecule or polypeptide may exist in a purified form or may exist in a non-native environment such as, for example, a transgenic host cell. For example, an "isolated" or "purified" nucleic acid molecule or protein, or biologically active portion thereof, is substantially free of other cellular material, or culture medium when produced by  
15 recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can  
20 contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. A protein that is substantially free of cellular material includes preparations of protein or polypeptide having less than about 30%, 20%, 10%, 5%, (by dry weight) of contaminating protein. When the protein of the invention, or biologically active  
25 portion thereof, is recombinantly produced, preferably culture medium represents less than about 30%, 20%, 10%, or 5% (by dry weight) of chemical precursors or non-protein of interest chemicals.

The nucleotide sequences of the invention include both the naturally occurring sequences as well as mutant (variant) forms. Such variants will continue to possess the desired  
30 activity, i.e., either promoter activity or the activity of the product encoded by the open reading frame of the non-variant nucleotide sequence.

Thus, by "variants" is intended substantially similar sequences. For nucleotide sequences comprising an open reading frame, variants include those sequences that, because of

the degeneracy of the genetic code, encode the identical amino acid sequence of the native protein. Naturally occurring allelic variants such as these can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridization techniques. Variant nucleotide sequences also include synthetically  
5 derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis and for open reading frames, encode the native protein, as well as those that encode a polypeptide having amino acid substitutions relative to the native protein. Generally, nucleotide sequence variants of the invention will have at least 40, 50, 60, to 70%, e.g., preferably 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, to 79%, generally at least 80%, e.g.,  
10 81%-84%, at least 85%, e.g., 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, to 98% and 99% nucleotide sequence identity to the native (wild type or endogenous) nucleotide sequence.

“Conservatively modified variations” of a particular nucleic acid sequence refers to those nucleic acid sequences that encode identical or essentially identical amino acid  
15 sequences, or where the nucleic acid sequence does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given polypeptide. For instance the codons CGT, CGC, CGA, CGG, AGA, and AGG all encode the amino acid arginine. Thus, at every position where an arginine is specified by a codon, the codon can be altered to any of the  
20 corresponding codons described without altering the encoded protein. Such nucleic acid variations are “silent variations” which are one species of “conservatively modified variations.” Every nucleic acid sequence described herein which encodes a polypeptide also describes every possible silent variation, except where otherwise noted. One of skill will recognize that each codon in a nucleic acid (except ATG, which is ordinarily the only codon  
25 for methionine) can be modified to yield a functionally identical molecule by standard techniques. Accordingly, each “silent variation” of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

The nucleic acid molecules of the invention can be “optimized” for enhanced expression in plants of interest. See, for example, EPA 035472; WO 91/16432; Perlak et al.,  
30 1991; and Murray et al., 1989. In this manner, the open reading frames in genes or gene fragments can be synthesized utilizing plant-preferred codons. See, for example, Campbell and Gowri, 1990 for a discussion of host-preferred codon usage. Thus, the nucleotide sequences can be optimized for expression in any plant. It is recognized that all or any part of

the gene sequence may be optimized or synthetic. That is, synthetic or partially optimized sequences may also be used. Variant nucleotide sequences and proteins also encompass sequences and protein derived from a mutagenic and recombinogenic procedure such as DNA shuffling. With such a procedure, one or more different coding sequences can be manipulated  
5 to create a new polypeptide possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined *in vitro* or *in vivo*. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer, 1994; Stemmer, 1994; Cramer et al., 1997; Moore et  
10 al., 1997; Zhang et al., 1997; Cramer et al., 1998; and U.S. Patent Nos. 5,605,793 and 5,837,458.

By "variant" polypeptide is intended a polypeptide derived from the native protein by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or more amino acids at one or  
15 more sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Such variants may result from, for example, genetic polymorphism or from human manipulation. Methods for such manipulations are generally known in the art.

Thus, the polypeptides may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are  
20 generally known in the art. For example, amino acid sequence variants of the polypeptides can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel, 1985; Kunkel et al., 1987; U. S. Patent No. 4,873,192; Walker and Gaastra, 1983 and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of  
25 interest may be found in the model of Dayhoff et al. (1978). Conservative substitutions, such as exchanging one amino acid with another having similar properties, are preferred.

Individual substitutions deletions or additions that alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 1%) in an encoded sequence are "conservatively modified variations," where the alterations result  
30 in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. The following five groups each contain amino acids that are conservative substitutions for one another: Aliphatic: Glycine (G), Alanine (A), Valine (V), Leucine (L), Isoleucine (I);



Aromatic: Phenylalanine (F), Tyrosine (Y), Tryptophan (W); Sulfur-containing: Methionine (M), Cysteine (C); Basic: Arginine (R), Lysine (K), Histidine (H); Acidic: Aspartic acid (D), Glutamic acid (E), Asparagine (N), Glutamine (Q). See also, Creighton, 1984. In addition, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids in an encoded sequence are also "conservatively modified variations."

"Expression cassette" as used herein means a DNA sequence capable of directing expression of a particular nucleotide sequence in an appropriate host cell, comprising a promoter operably linked to the nucleotide sequence of interest which is operably linked to termination signals. It also typically comprises sequences required for proper translation of the nucleotide sequence. The coding region usually codes for a protein of interest but may also code for a functional RNA of interest, for example antisense RNA or a nontranslated RNA, in the sense or antisense direction. The expression cassette comprising the nucleotide sequence of interest may be chimeric, meaning that at least one of its components is heterologous with respect to at least one of its other components. The expression cassette may also be one which is naturally occurring but has been obtained in a recombinant form useful for heterologous expression. The expression of the nucleotide sequence in the expression cassette may be under the control of a constitutive promoter or of an inducible promoter which initiates transcription only when the host cell is exposed to some particular external stimulus. In the case of a multicellular organism, the promoter can also be specific to a particular tissue or organ or stage of development.

"Vector" is defined to include, inter alia, any plasmid, cosmid, phage or *Agrobacterium* binary vector in double or single stranded linear or circular form which may or may not be self transmissible or mobilizable, and which can transform prokaryotic or eukaryotic host either by integration into the cellular genome or exist extrachromosomally (e.g. autonomous replicating plasmid with an origin of replication).

Specifically included are shuttle vectors by which is meant a DNA vehicle capable, naturally or by design, of replication in two different host organisms, which may be selected from actinomycetes and related species, bacteria and eukaryotic (e.g. higher plant, mammalian, yeast or fungal cells).

Preferably the nucleic acid in the vector is under the control of, and operably linked to, an appropriate promoter or other regulatory elements for transcription in a host cell such as a microbial, e.g. bacterial, or plant cell. The vector may be a bi-functional expression vector

which functions in multiple hosts. In the case of genomic DNA, this may contain its own promoter or other regulatory elements and in the case of cDNA this may be under the control of an appropriate promoter or other regulatory elements for expression in the host cell.

"Cloning vectors" typically contain one or a small number of restriction endonuclease recognition sites at which foreign DNA sequences can be inserted in a determinable fashion without loss of essential biological function of the vector, as well as a marker gene that is suitable for use in the identification and selection of cells transformed with the cloning vector. Marker genes typically include genes that provide tetracycline resistance, hygromycin resistance or ampicillin resistance.

10 A "transgenic plant" is a plant having one or more plant cells that contain an expression vector.

"Plant tissue" includes differentiated and undifferentiated tissues or plants, including but not limited to roots, stems, shoots, leaves, pollen, seeds, tumor tissue and various forms of cells and culture such as single cells, protoplast, embryos, and callus tissue. The plant tissue  
15 may be in plants or in organ, tissue or cell culture.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: (a) "reference sequence", (b) "comparison window", (c) "sequence identity", (d) "percentage of sequence identity", and (e) "substantial identity".

(a) As used herein, "reference sequence" is a defined sequence used as a basis for  
20 sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full length cDNA or gene sequence, or the complete cDNA or gene sequence.

(b) As used herein, "comparison window" makes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence in the  
25 comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally can be 30, 40, 50, 100, or longer. Those of skill in the art understand that to avoid a high similarity to a reference sequence due to inclusion of gaps in the polynucleotide  
30 sequence a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well known in the art. Thus, the determination of percent identity between any two sequences can be accomplished using a mathematical algorithm. Preferred, non-limiting examples of such mathematical algorithms

are the algorithm of Myers and Miller, 1988; the local homology algorithm of Smith et al. 1981; the homology alignment algorithm of Needleman and Wunsch 1970; the search-for-similarity-method of Pearson and Lipman 1988; the algorithm of Karlin and Altschul, 1990, modified as in Karlin and Altschul, 1993.

5 Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, California); the ALIGN program (Version 2.0) and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 8 (available from  
10 Genetics Computer Group (GCG), 575 Science Drive, Madison, Wisconsin, USA). Alignments using these programs can be performed using the default parameters. The CLUSTAL program is well described by Higgins et al. 1988; Higgins et al. 1989; Corpet et al. 1988; Huang et al. 1992; and Pearson et al. 1994. The ALIGN program is based on the algorithm of Myers and Miller, *supra*. The BLAST programs of Altschul et al., 1990, are  
15 based on the algorithm of Karlin and Altschul *supra*.

Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when  
20 aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., 1990). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the  
25 parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when the cumulative alignment score falls off by the quantity X from its maximum achieved value, the cumulative score goes to zero or below due to the accumulation of one or more  
30 negative-scoring residue alignments, or the end of either sequence is reached.

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul (1993)). One measure of similarity provided by the BLAST algorithm is the smallest

sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a test nucleic acid sequence is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid sequence to the reference nucleic acid sequence is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul et al. 1997. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al., *supra*. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g. BLASTN for nucleotide sequences, BLASTX for proteins) can be used. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, 1989). See <http://www.ncbi.nlm.nih.gov>. Alignment may also be performed manually by inspection.

For purposes of the present invention, comparison of nucleotide sequences for determination of percent sequence identity to the promoter sequences disclosed herein is preferably made using the BlastN program (version 1.4.7 or later) with its default parameters or any equivalent program. By "equivalent program" is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by the preferred program.

(c) As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the

substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity." Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California).

(d) As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

(e)(i) The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, or 79%, preferably at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, or 89%, more preferably at least 90%, 91%, 92%, 93%, or 94%, and most preferably at least 95%, 96%, 97%, 98%, or 99% sequence identity, compared to a reference sequence using one of the alignment programs described using standard parameters. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 70%, more preferably at least 80%, 90%, and most preferably at least 95%.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions (see below). Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. However, stringent conditions encompass temperatures in the range of about 1°C to about 20°C, depending upon the desired

degree of stringency as otherwise qualified herein. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides they encode are substantially identical. This may occur, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. One indication that two  
5 nucleic acid sequences are substantially identical is when the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid.

(e)(ii) The term "substantial identity" in the context of a peptide indicates that a peptide comprises a sequence with at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, or  
10 79%, preferably 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, or 89%, more preferably at least 90%, 91%, 92%, 93%, or 94%, or even more preferably, 95%, 96%, 97%, 98% or 99%, sequence identity to the reference sequence over a specified comparison window. Preferably, optimal alignment is conducted using the homology alignment algorithm of Needleman and Wunsch (1970). An indication that two peptide sequences are substantially  
15 identical is that one peptide is immunologically reactive with antibodies raised against the second peptide. Thus, a peptide is substantially identical to a second peptide, for example, where the two peptides differ only by a conservative substitution.

For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference  
20 sequences are input into a computer, subsequence coordinates are designated if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

As noted above, another indication that two nucleic acid sequences are substantially  
25 identical is that the two molecules hybridize to each other under stringent conditions. The phrase "hybridizing specifically to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular) DNA or RNA. "Bind(s) substantially" refers to complementary hybridization between a probe nucleic acid and a target  
30 nucleic acid and embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the desired detection of the target nucleic acid sequence.

“Stringent hybridization conditions” and “stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments such as Southern and Northern hybridization are sequence dependent, and are different under different environmental parameters. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the  $T_m$  can be approximated from the equation of Meinkoth and Wahl, 1984;  $T_m = 81.5^\circ\text{C} + 16.6 (\log M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$ ; where  $M$  is the molarity of monovalent cations,  $\%GC$  is the percentage of guanosine and cytosine nucleotides in the DNA,  $\% \text{ form}$  is the percentage of formamide in the hybridization solution, and  $L$  is the length of the hybrid in base pairs.  $T_m$  is reduced by about  $1^\circ\text{C}$  for each 1% of mismatching; thus,  $T_m$ , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with  $>90\%$  identity are sought, the  $T_m$  can be decreased  $10^\circ\text{C}$ . Generally, stringent conditions are selected to be about  $5^\circ\text{C}$  lower than the thermal melting point  $T_m$  for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or  $4^\circ\text{C}$  lower than the thermal melting point  $T_m$ ; moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or  $10^\circ\text{C}$  lower than the thermal melting point  $T_m$ ; low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or  $20^\circ\text{C}$  lower than the thermal melting point  $T_m$ . Using the equation, hybridization and wash compositions, and desired  $T$ , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a  $T$  of less than  $45^\circ\text{C}$  (aqueous solution) or  $32^\circ\text{C}$  (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen, 1993. Generally, highly stringent hybridization and wash conditions are selected to be about  $5^\circ\text{C}$  lower than the thermal melting point  $T_m$  for the specific sequence at a defined ionic strength and pH.

An example of highly stringent wash conditions is 0.15 M NaCl at  $72^\circ\text{C}$  for about 15 minutes. An example of stringent wash conditions is a 0.2X SSC wash at  $65^\circ\text{C}$  for 15 minutes (see, Sambrook, *infra*, for a description of SSC buffer). Often, a high stringency wash is preceded by a low stringency wash to remove background probe signal. An example medium

stringency wash for a duplex of, e.g., more than 100 nucleotides, is 1X SSC at 45°C for 15 minutes. An example low stringency wash for a duplex of, e.g., more than 100 nucleotides, is 4-6X SSC at 40°C for 15 minutes. For short probes (e.g., about 10 to 50 nucleotides), stringent conditions typically involve salt concentrations of less than about 1.5 M, more preferably about 0.01 to 1.0 M, Na ion concentration (or other salts) at pH 7.0 to 8.3, and the temperature is typically at least about 30°C and at least about 60°C for long robes (e.g., >50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. In general, a signal to noise ratio of 2X (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the proteins that they encode are substantially identical. This occurs, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

Very stringent conditions are selected to be equal to the  $T_m$  for a particular probe. An example of stringent conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or Northern blot is 50% formamide, e.g., hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37°C, and a wash in 0.5X to 1X SSC at 55 to 60°C.

The following are examples of sets of hybridization/wash conditions that may be used to clone orthologous nucleotide sequences that are substantially identical to reference nucleotide sequences of the present invention: a reference nucleotide sequence preferably hybridizes to the reference nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 2X SSC, 0.1% SDS at 50°C, more desirably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 1X SSC, 0.1% SDS at 50°C, more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.5X SSC, 0.1% SDS at 50°C, preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.1X



SSC, 0.1% SDS at 50°C, more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 65°C.

“DNA shuffling” is a method to introduce mutations or rearrangements, preferably randomly, in a DNA molecule or to generate exchanges of DNA sequences between two or more DNA molecules, preferably randomly. The DNA molecule resulting from DNA shuffling is a shuffled DNA molecule that is a non-naturally occurring DNA molecule derived from at least one template DNA molecule. The shuffled DNA preferably encodes a variant polypeptide modified with respect to the polypeptide encoded by the template DNA, and may have an altered biological activity with respect to the polypeptide encoded by the template DNA.

“Recombinant DNA molecule” is a combination of DNA sequences that are joined together using recombinant DNA technology and procedures used to join together DNA sequences as described, for example, in Sambrook et al., 1989.

The word “plant” refers to any plant, particularly to seed plant, and “plant cell” is a structural and physiological unit of the plant, which comprises a cell wall but may also refer to a protoplast. The plant cell may be in form of an isolated single cell or a cultured cell, or as a part of higher organized unit such as, for example, a plant tissue, or a plant organ.

“Significant increase” is an increase that is larger than the margin of error inherent in the measurement technique, preferably an increase by about 2-fold or greater.

“Significantly less” means that the decrease is larger than the margin of error inherent in the measurement technique, preferably a decrease by about 2-fold or greater.

## **II. DNA Sequences for Transformation**

Virtually any DNA composition may be used for delivery to recipient plant cells, e.g., monocotyledonous cells, to ultimately produce fertile transgenic plants in accordance with the present invention. For example, DNA segments in the form of vectors and plasmids, or linear DNA fragments, in some instances containing only the DNA element to be expressed in the plant, and the like, may be employed. The construction of vectors which may be employed in conjunction with the present invention will be known to those of skill of the art in light of the present disclosure (see, e.g., Sambrook et al., 1989; Gelvin et al., 1990).

Vectors, plasmids, cosmids, YACs (yeast artificial chromosomes), BACs (bacterial artificial chromosomes) and DNA segments for use in transforming such cells will, of course, generally comprise the cDNA, gene or genes which one desires to introduce into the cells. These DNA constructs can further include structures such as promoters, enhancers,

polylinkers, or even regulatory genes as desired. The DNA segment or gene chosen for cellular introduction will often encode a protein which will be expressed in the resultant recombinant cells, such as will result in a screenable or selectable trait and/or which will impart an improved phenotype to the regenerated plant. However, this may not always be the case, and the present invention also encompasses transgenic plants incorporating non-expressed transgenes.

In certain embodiments, it is contemplated that one may wish to employ replication-competent viral vectors in monocot transformation. Such vectors include, for example, wheat dwarf virus (WDV) "shuttle" vectors, such as pW1-11 and PW1-GUS (Ugaki et al., 1991). These vectors are capable of autonomous replication in maize cells as well as *E. coli*, and as such may provide increased sensitivity for detecting DNA delivered to transgenic cells. A replicating vector may also be useful for delivery of genes flanked by DNA sequences from transposable elements such as Ac, Ds, or Mu. It has been proposed (Laufs et al., 1990) that transposition of these elements within the maize genome requires DNA replication. It is also contemplated that transposable elements would be useful for introducing DNA fragments lacking elements necessary for selection and maintenance of the plasmid vector in bacteria, e.g., antibiotic resistance genes and origins of DNA replication. It is also proposed that use of a transposable element such as Ac, Ds, or Mu would actively promote integration of the desired DNA and hence increase the frequency of stably transformed cells. The use of a transposable element such as Ac, Ds, or Mu may actively promote integration of the DNA of interest and hence increase the frequency of stably transformed cells. Transposable elements may be useful to allow separation of genes of interest from elements necessary for selection and maintenance of a plasmid vector in bacteria or selection of a transformant. By use of a transposable element, desirable and undesirable DNA sequences may be transposed apart from each other in the genome, such that through genetic segregation in progeny, one may identify plants with either the desirable undesirable DNA sequences.

DNA useful for introduction into plant cells includes that which has been derived or isolated from any source, that may be subsequently characterized as to structure, size and/or function, chemically altered, and later introduced into plants. An example of DNA "derived" from a source, would be a DNA sequence that is identified as a useful fragment within a given organism, and which is then chemically synthesized in essentially pure form. An example of such DNA "isolated" from a source would be a useful DNA sequence that is excised or removed from said source by chemical means, e.g., by the use of restriction endonucleases, so

that it can be further manipulated, e.g., amplified, for use in the invention, by the methodology of genetic engineering. Such DNA is commonly referred to as "recombinant DNA."

Therefore useful DNA includes completely synthetic DNA, semi-synthetic DNA, DNA isolated from biological sources, and DNA derived from introduced RNA. Generally, the introduced DNA is not originally resident in the plant genotype which is the recipient of the DNA, but it is within the scope of the invention to isolate a gene from a given plant genotype, and to subsequently introduce multiple copies of the gene into the same genotype, e.g., to enhance production of a given gene product such as a storage protein or a protein that confers tolerance or resistance to water deficit.

The introduced DNA includes but is not limited to, DNA from plant genes, and non-plant genes such as those from bacteria, yeasts, animals or viruses. The introduced DNA can include modified genes, portions of genes, or chimeric genes, including genes from the same or different maize genotype. The term "chimeric gene" or "chimeric DNA" is defined as a gene or DNA sequence or segment comprising at least two DNA sequences or segments from species which do not combine DNA under natural conditions, or which DNA sequences or segments are positioned or linked in a manner which does not normally occur in the native genome of untransformed plant.

The introduced DNA used for transformation herein may be circular or linear, double-stranded or single-stranded. Generally, the DNA is in the form of chimeric DNA, such as plasmid DNA, that can also contain coding regions flanked by regulatory sequences which promote the expression of the recombinant DNA present in the resultant plant. For example, the DNA may itself comprise or consist of a promoter that is active in a plant which is derived from a source other than that plant, or may utilize a promoter already present in a plant genotype that is the transformation target.

Generally, the introduced DNA will be relatively small, i.e., less than about 30 kb to minimize any susceptibility to physical, chemical, or enzymatic degradation which is known to increase as the size of the DNA increases. As noted above, the number of proteins, RNA transcripts or mixtures thereof which is introduced into the plant genome is preferably preselected and defined, e.g., from one to about 5-10 such products of the introduced DNA may be formed.

Two principal methods for the control of expression are known, viz.: overexpression and underexpression. Overexpression can be achieved by insertion of one or more than one extra copy of the selected gene. It is, however, not unknown for plants or their progeny,

originally transformed with one or more than one extra copy of a nucleotide sequence, to exhibit the effects of underexpression as well as overexpression. For underexpression there are two principle methods which are commonly referred to in the art as "antisense downregulation" and "sense downregulation" (sense downregulation is also referred to as "cosuppression"). Generically these processes are referred to as "gene silencing". Both of these methods lead to an inhibition of expression of the target gene.

Obtaining sufficient levels of transgene expression in the appropriate plant tissues is an important aspect in the production of genetically engineered crops. Expression of heterologous DNA sequences in a plant host is dependent upon the presence of an operably linked promoter that is functional within the plant host. Choice of the promoter sequence will determine when and where within the organism the heterologous DNA sequence is expressed.

Furthermore, it is contemplated that promoters combining elements from more than one promoter may be useful. For example, U.S. Patent No. 5,491,288 discloses combining a Cauliflower Mosaic Virus promoter with a histone promoter. Thus, the elements from the promoters disclosed herein may be combined with elements from other promoters.

Promoters which are useful for plant transgene expression include those that are inducible, viral, synthetic, constitutive (Odell et al., 1985), temporally regulated, spatially regulated, tissue-specific, and spatio-temporally regulated.

Where expression in specific tissues or organs is desired, tissue-specific promoters may be used. In contrast, where gene expression in response to a stimulus is desired, inducible promoters are the regulatory elements of choice. Where continuous expression is desired throughout the cells of a plant, constitutive promoters are utilized. Additional regulatory sequences upstream and/or downstream from the core promoter sequence may be included in expression constructs of transformation vectors to bring about varying levels of expression of heterologous nucleotide sequences in a transgenic plant.

#### **A. Transcription Regulatory Sequences**

##### **1. Promoters**

The choice of promoter will vary depending on the temporal and spatial requirements for expression, and also depending on the target species. In some cases, expression in multiple tissues is desirable. While in others, tissue-specific, e.g., leaf-specific, seed-specific, petal-specific, anther-specific, or pith-specific, expression is desirable. Although many promoters from dicotyledons have been shown to be operational in monocotyledons and *vice versa*, ideally dicotyledonous promoters are selected for expression

in dicotyledons, and monocotyledonous promoters for expression in monocotyledons.

However, there is no restriction to the provenance of selected promoters; it is sufficient that they are operational in driving the expression of the nucleotide sequences in the desired cell.

These promoters include, but are not limited to, constitutive, inducible, temporally  
5 regulated, developmentally regulated, spatially-regulated, chemically regulated, stress-  
responsive, tissue-specific, viral and synthetic promoters. Promoter sequences are known to be  
strong or weak. A strong promoter provides for a high level of gene expression, whereas a  
weak promoter provides for a very low level of gene expression. An inducible promoter is a  
promoter that provides for the turning on and off of gene expression in response to an  
10 exogenously added agent, or to an environmental or developmental stimulus. A bacterial  
promoter such as the  $P_{lac}$  promoter can be induced to varying levels of gene expression  
depending on the level of isothiopyrogalactoside added to the transformed bacterial cells.  
An isolated promoter sequence that is a strong promoter for heterologous nucleic acid is  
advantageous because it provides for a sufficient level of gene expression to allow for easy  
15 detection and selection of transformed cells and provides for a high level of gene expression  
when desired.

Within a plant promoter region there are several domains that are necessary for full  
function of the promoter. The first of these domains lies immediately upstream of the  
structural gene and forms the "core promoter region" containing consensus sequences,  
20 normally 70 base pairs immediately upstream of the gene. The core promoter region contains  
the characteristic CAAT and TATA boxes plus surrounding sequences, and represents a  
transcription initiation sequence that defines the transcription start point for the structural gene.

The presence of the core promoter region defines a sequence as being a promoter: if the  
region is absent, the promoter is non-functional. Furthermore, the core promoter region is  
25 insufficient to provide full promoter activity. A series of regulatory sequences upstream of the  
core constitute the remainder of the promoter. The regulatory sequences determine expression  
level, the spatial and temporal pattern of expression and, for an important subset of promoters,  
expression under inductive conditions (regulation by external factors such as light,  
temperature, chemicals, hormones).

30 A range of naturally-occurring promoters are known to be operative in plants and have  
been used to drive the expression of heterologous (both foreign and endogenous) genes in  
plants: for example, the constitutive 35S cauliflower mosaic virus (CaMV) promoter, the  
ripening-enhanced tomato polygalacturonase promoter (Bird et al., 1988), the E8 promoter

(Diekman & Fischer, 1988) and the fruit specific 2A1 promoter (Pear et al., 1989) and many others, e.g., U2 and U5 snRNA promoters from maize, the promoter from alcohol dehydrogenase, the Z4 promoter from a gene encoding the Z4 22 kD zein protein, the Z10 promoter from a gene encoding a 10 kD zein protein, a Z27 promoter from a gene encoding a 27 kD zein protein, the A20 promoter from the gene encoding a 19 kD -zein protein, inducible promoters, such as the light inducible promoter derived from the pea *rbcS* gene and the actin promoter from rice, e.g., the actin 2 promoter (WO 00/70067); seed specific promoters, such as the phaseolin promoter from beans, may also be used. The nucleotide sequences of this invention can also be expressed under the regulation of promoters that are chemically regulated. This enables the nucleic acid sequence or encoded polypeptide to be synthesized only when the crop plants are treated with the inducing chemicals. Chemical induction of gene expression is detailed in EP 0 332 104 (to Ciba-Geigy) and U.S. Patent 5,614,395. A preferred promoter for chemical induction is the tobacco PR-1a promoter.

Examples of some constitutive promoters which have been described include the rice actin 1 (Wang et al., 1992; U.S. Patent No. 5,641,876), CaMV 35S (Odell et al., 1985), CaMV 19S (Lawton et al., 1987), *nos*, *Adh*, sucrose synthase; and the ubiquitin promoters.

Examples of tissue specific promoters which have been described include the lectin (Vodkin, 1983; Lindstrom et al., 1990) corn alcohol dehydrogenase 1 (Vogel et al., 1989; Dennis et al., 1984), corn light harvesting complex (Simpson, 1986; Bansal et al., 1992), corn heat shock protein (Odell et al., 1985), pea small subunit RuBP carboxylase (Poulsen et al., 1986), Ti plasmid mannopine synthase (Langridge et al., 1989), Ti plasmid nopaline synthase (Langridge et al., 1989), petunia chalcone isomerase (vanTunen et al., 1988), bean glycine rich protein 1 (Keller et al., 1989), truncated CaMV 35s (Odell et al., 1985), potato patatin (Wenzler et al., 1989), root cell (Yamamoto et al., 1990), maize zein (Reina et al., 1990; Kriz et al., 1987; Wandelt et al., 1989; Langridge et al., 1983; Reina et al., 1990), globulin-1 (Belanger et al., 1991),  $\alpha$ -tubulin, *cab* (Sullivan et al., 1989), PEPCase (Hudspeth & Grula, 1989), R gene complex-associated promoters (Chandler et al., 1989), histone, and chalcone synthase promoters (Franken et al., 1991). Tissue specific enhancers are described in Fromm et al. (1989).

Inducible promoters that have been described include the ABA- and turgor-inducible promoters, the promoter of the auxin-binding protein gene (Schwob et al., 1993), the UDP glucose flavonoid glycosyl-transferase gene promoter (Ralston et al., 1988), the MPI

proteinase inhibitor promoter (Cordero et al., 1994), and the glyceraldehyde-3-phosphate dehydrogenase gene promoter (Kohler et al., 1995; Quigley et al., 1989; Martinez et al., 1989).

Several other tissue-specific regulated genes and/or promoters have been reported in plants. These include genes encoding the seed storage proteins (such as napin, cruciferin, beta-conglycinin, and phaseolin) zein or oil body proteins (such as oleosin), or genes involved in fatty acid biosynthesis (including acyl carrier protein, stearoyl-ACP desaturase. And fatty acid desaturases (fad 2-1)), and other genes expressed during embryo development (such as Bce4, see, for example, EP 255378 and Kridl et al., 1991). Particularly useful for seed-specific expression is the pea vicilin promoter (Czako et al., 1992). (See also U.S. Pat. No. 5,625,136, herein incorporated by reference.) Other useful promoters for expression in mature leaves are those that are switched on at the onset of senescence, such as the SAG promoter from *Arabidopsis* (Gan et al., 1995).

A class of fruit-specific promoters expressed at or during antithesis through fruit development, at least until the beginning of ripening, is discussed in U.S. 4,943,674. cDNA clones that are preferentially expressed in cotton fiber have been isolated (John et al., 1992). cDNA clones from tomato displaying differential expression during fruit development have been isolated and characterized (Mansson et al., 1985, Slater et al., 1985). The promoter for polygalacturonase gene is active in fruit ripening. The polygalacturonase gene is described in U.S. Patent No. 4,535,060, U.S. Patent No. 4,769,061, U.S. Patent No. 4,801,590, and U.S. Patent No. 5,107,065, which disclosures are incorporated herein by reference.

Other examples of tissue-specific promoters include those that direct expression in leaf cells following damage to the leaf (for example, from chewing insects), in tubers (for example, patatin gene promoter), and in fiber cells (an example of a developmentally-regulated fiber cell protein is E6 (John et al., 1992). The E6 gene is most active in fiber, although low levels of transcripts are found in leaf, ovule and flower.

The tissue-specificity of some "tissue-specific" promoters may not be absolute and may be tested by one skilled in the art using the diphtheria toxin sequence. One can also achieve tissue-specific expression with "leaky" expression by a combination of different tissue-specific promoters (Beals et al., 1997). Other tissue-specific promoters can be isolated by one skilled in the art (see U.S. 5,589,379). Several inducible promoters ("gene switches") have been reported. Many are described in the review by Gatz (1996) and Gatz (1997). These include tetracycline repressor system, *Lac* repressor system, copper-inducible systems, salicylate-inducible systems (such as the PR1a system), glucocorticoid- (Aoyama et al., 1997) and

ecdysome-inducible systems. Also included are the benzene sulphonamide- (U.S. Patent No. 5,364,780) and alcohol-(WO 97/06269 and WO 97/06268) inducible systems and glutathione S-transferase promoters. Other studies have focused on genes inducibly regulated in response to environmental stress or stimuli such as increased salinity. Drought, pathogen and wounding.  
5 (Graham et al., 1985; Graham et al., 1985, Smith et al., 1986). Accumulation of metallocarboxypeptidase-inhibitor protein has been reported in leaves of wounded potato plants (Graham et al., 1981). Other plant genes have been reported to be induced methyl jasmonate, elicitors, heat-shock, anaerobic stress, or herbicide safeners.

Regulated expression of the chimeric transacting viral replication protein can be further  
10 regulated by other genetic strategies. For example, *Cre*-mediated gene activation as described by Odell et al. 1990. Thus, a DNA fragment containing 3' regulatory sequence bound by lox sites between the promoter and the replication protein coding sequence that blocks the expression of a chimeric replication gene from the promoter can be removed by *Cre*-mediated excision and result in the expression of the *trans*-acting replication gene. In this case, the  
15 chimeric *Cre* gene, the chimeric *trans*-acting replication gene, or both can be under the control of tissue- and developmental- specific or inducible promoters. An alternate genetic strategy is the use of tRNA suppressor gene. For example, the regulated expression of a tRNA suppressor gene can conditionally control expression of a *trans*-acting replication protein coding sequence containing an appropriate termination codon as described by Ulmasov et al.  
20 1997. Again, either the chimeric tRNA suppressor gene, the chimeric transacting replication gene, or both can be under the control of tissue- and developmental-specific or inducible promoters.

Frequently it is desirable to have continuous or inducible expression of a DNA sequence throughout the cells of an organism in a tissue-independent manner. For example,  
25 increased resistance of a plant to infection by soil- and airborne-pathogens might be accomplished by genetic manipulation of the plant's genome to comprise a continuous promoter operably linked to a heterologous pathogen-resistance gene such that pathogen-resistance proteins are continuously expressed throughout the plant's tissues.

Alternatively, it might be desirable to inhibit expression of a native DNA sequence  
30 within a plant's tissues to achieve a desired phenotype. In this case, such inhibition might be accomplished with transformation of the plant to comprise a constitutive, tissue-independent promoter operably linked to an antisense nucleotide sequence, such that constitutive



expression of the antisense sequence produces an RNA transcript that interferes with translation of the mRNA of the native DNA sequence.

To define a minimal promoter region, a DNA segment representing the promoter region is removed from the 5' region of the gene of interest and operably linked to the coding sequence of a marker (reporter) gene by recombinant DNA techniques well known to the art. The reporter gene is operably linked downstream of the promoter, so that transcripts initiating at the promoter proceed through the reporter gene. Reporter genes generally encode proteins which are easily measured, including, but not limited to, chloramphenicol acetyl transferase (CAT), beta-glucuronidase (GUS), green fluorescent protein (GFP), beta-galactosidase (beta-GAL), and luciferase.

The construct containing the reporter gene under the control of the promoter is then introduced into an appropriate cell type by transfection techniques well known to the art. To assay for the reporter protein, cell lysates are prepared and appropriate assays, which are well known in the art, for the reporter protein are performed. For example, if CAT were the reporter gene of choice, the lysates from cells transfected with constructs containing CAT under the control of a promoter under study are mixed with isotopically labeled chloramphenicol and acetyl-coenzyme A (acetyl-CoA). The CAT enzyme transfers the acetyl group from acetyl-CoA to the 2- or 3-position of chloramphenicol. The reaction is monitored by thin-layer chromatography, which separates acetylated chloramphenicol from unreacted material. The reaction products are then visualized by autoradiography.

The level of enzyme activity corresponds to the amount of enzyme that was made, which in turn reveals the level of expression from the promoter of interest. This level of expression can be compared to other promoters to determine the relative strength of the promoter under study. In order to be sure that the level of expression is determined by the promoter, rather than by the stability of the mRNA, the level of the reporter mRNA can be measured directly, such as by Northern blot analysis.

Once activity is detected, mutational and/or deletional analyses may be employed to determine the minimal region and/or sequences required to initiate transcription. Thus, sequences can be deleted at the 5' end of the promoter region and/or at the 3' end of the promoter region, and nucleotide substitutions introduced. These constructs are then introduced to cells and their activity determined.

In one embodiment, the promoter may be a gamma zein promoter, an oleosin ole16 promoter, a globulinI promoter, an actin I promoter, an actin cl promoter, a sucrose synthetase

promoter, an INOPS promoter, an EXM5 promoter, a globulin2 promoter, a b-32, ADPG-pyrophosphorylase promoter, an LtpI promoter, an Ltp2 promoter, an oleosin ole17 promoter, an oleosin ole18 promoter, an actin 2 promoter, a pollen-specific protein promoter, a pollen-specific pectate lyase promoter, an anther-specific protein promoter (Huffman), an anther-specific gene RTS2 promoter, a pollen-specific gene promoter, a tapetum-specific gene promoter, tapetum-specific gene RAB24 promoter, a anthranilate synthase alpha subunit promoter, an alpha zein promoter, an anthranilate synthase beta subunit promoter, a dihydrodipicolinate synthase promoter, a Thil promoter, an alcohol dehydrogenase promoter, a cab binding protein promoter, an H3C4 promoter, a RUBISCO SS starch branching enzyme promoter, an ACCase promoter, an actin3 promoter, an actin7 promoter, a regulatory protein GF14-12 promoter, a ribosomal protein L9 promoter, a cellulose biosynthetic enzyme promoter, an S-adenosyl-L-homocysteine hydrolase promoter, a superoxide dismutase promoter, a C-kinase receptor promoter, a phosphoglycerate mutase promoter, a root-specific RCc3 mRNA promoter, a glucose-6 phosphate isomerase promoter, a pyrophosphate-fructose 6-phosphatetphosphotransferase promoter, an ubiquitin promoter, a beta-ketoacyl-ACP synthase promoter, a 33 kDa photosystem 11 promoter, an oxygen evolving protein promoter, a 69 kDa vacuolar ATPase subunit promoter, a metallothionein-like protein promoter, a glyceraldehyde-3-phosphate dehydrogenase promoter, an ABA- and ripening-inducible-like protein promoter, a phenylalanine ammonia lyase promoter, an adenosine triphosphatase S-adenosyl-L-homocysteine hydrolase promoter, an a-tubulin promoter, a cab promoter, a PEPCase promoter, an R gene promoter, a lectin promoter, a light harvesting complex promoter, a heat shock protein promoter, a chalcone synthase promoter, a zein promoter, a globulin-1 promoter, an ABA promoter, an auxin-binding protein promoter, a UDP glucose flavonoid glycosyl-transferase gene promoter, an NTI promoter, an actin promoter, an opaque 2 promoter, a b70 promoter, an oleosin promoter, a CaMV 35S promoter, a CaMV 19S promoter, a histone promoter, a turgor-inducible promoter, a pea small subunit RuBP carboxylase promoter, a Ti plasmid mannopine synthase promoter, Ti plasmid nopaline synthase promoter, a petunia chalcone isomerase promoter, a bean glycine rich protein I promoter, a CaMV 35S transcript promoter, a potato patatin promoter, or a S-E9 small subunit RuBP carboxylase promoter.

## 2. Other Regulatory Elements

In addition to promoters, a variety of 5N and 3N transcriptional regulatory sequences are also available for use in the present invention. Transcriptional terminators are responsible

for the termination of transcription and correct mRNA polyadenylation. The 3N nontranslated regulatory DNA sequence preferably includes from about 50 to about 1,000, more preferably about 100 to about 1,000, nucleotide base pairs and contains plant transcriptional and translational termination sequences. Appropriate transcriptional terminators and those which are known to function in plants include the CaMV 35S terminator, the *tml* terminator, the nopaline synthase terminator, the pea *rbcS* E9 terminator, the terminator for the T7 transcript from the octopine synthase gene of *Agrobacterium tumefaciens*, and the 3N end of the protease inhibitor I or II genes from potato or tomato, although other 3N elements known to those of skill in the art can also be employed. Alternatively, one also could use a gamma coixin, oleosin 3 or other terminator from the genus *Coix*.

Preferred 3' elements include those from the nopaline synthase gene of *Agrobacterium tumefaciens* (Bevan et al., 1983), the terminator for the T7 transcript from the octopine synthase gene of *Agrobacterium tumefaciens*, and the 3' end of the protease inhibitor I or II genes from potato or tomato.

As the DNA sequence between the transcription initiation site and the start of the coding sequence, i.e., the untranslated leader sequence, can influence gene expression, one may also wish to employ a particular leader sequence. Preferred leader sequences are contemplated to include those which include sequences predicted to direct optimum expression of the attached gene, i.e., to include a preferred consensus leader sequence which may increase or maintain mRNA stability and prevent inappropriate initiation of translation. The choice of such sequences will be known to those of skill in the art in light of the present disclosure. Sequences that are derived from genes that are highly expressed in plants will be most preferred.

Other sequences that have been found to enhance gene expression in transgenic plants include intron sequences (e.g., from *Adh1*, *bronzel*, *actin1*, *actin 2* (WO 00/760067), or the sucrose synthase intron) and viral leader sequences (e.g., from TMV, MCMV and AMV). For example, a number of non-translated leader sequences derived from viruses are known to enhance expression. Specifically, leader sequences from Tobacco Mosaic Virus (TMV), Maize Chlorotic Mottle Virus (MCMV), and Alfalfa Mosaic Virus (AMV) have been shown to be effective in enhancing expression (e.g., Gallie et al., 1987; Skuzeski et al., 1990). Other leaders known in the art include but are not limited to: Picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5 noncoding region) (Elroy-Stein et al., 1989); Potyvirus leaders, for example, TEV leader (Tobacco Etch Virus); MDMV leader (Maize Dwarf Mosaic

Virus); Human immunoglobulin heavy-chain binding protein (BiP) leader, (Macejak et al., 1991); Untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4), (Jobling et al., 1987; Tobacco mosaic virus leader (TMV), (Gallie et al., 1989; and Maize Chlorotic Mottle Virus leader (MCMV) (Lommel et al., 1991. See also, Della-Cioppa et al., 5 1987.

Regulatory elements such as Adh intron 1 (Callis et al., 1987), sucrose synthase intron (Vasil et al., 1989) or TMV omega element (Gallie, et al., 1989), may further be included where desired.

Examples of enhancers include elements from the CaMV 35S promoter, octopine 10 synthase genes (Ellis et al., 1987), the rice actin I gene, the maize alcohol dehydrogenase gene (Callis et al., 1987), the maize shrunken I gene (Vasil et al., 1989), TMV Omega element (Gallie et al., 1989) and promoters from non-plant eukaryotes (e.g. yeast; Ma et al., 1988).

Vectors for use in accordance with the present invention may be constructed to include the ocs enhancer element. This element was first identified as a 16 bp palindromic enhancer 15 from the octopine synthase (ocs) gene of *utilane* (Ellis et al., 1987), and is present in at least 10 other promoters (Bouchez et al., 1989). The use of an enhancer element, such as the ocs element and particularly multiple copies of the element, will act to increase the level of transcription from adjacent promoters when applied in the context of monocot transformation.

Ultimately, the most desirable DNA segments for introduction into for example a 20 monocot genome may be homologous genes or gene families which encode a desired trait (e.g., increased yield per acre) and which are introduced under the control of novel promoters or enhancers, etc., or perhaps even homologous or tissue specific (e.g., root-, collar/sheath-, whorl-, stalk-, earshank-, kernel- or leaf-specific) promoters or control elements. Indeed, it is envisioned that a particular use of the present invention will be the targeting of a gene in a 25 constitutive manner or a root-specific manner. For example, insect resistant genes may be expressed specifically in the whorl and collar/sheath tissues which are targets for the first and second broods, respectively, of ECB. Likewise, genes encoding proteins with particular activity against rootworm may be targeted directly to root tissues.

Vectors for use in tissue-specific targeting of genes in transgenic plants will typically 30 include tissue-specific promoters and may also include other tissue-specific control elements such as enhancer sequences. Promoters which direct specific or enhanced expression in certain plant tissues will be known to those of skill in the art in light of the present disclosure. These include, for example, the *rbcS* promoter, specific for green tissue; the ocs, nos and mas

promoters which have higher activity in roots or wounded leaf tissue; a truncated (-90 to +8) 35S promoter which directs enhanced expression in roots, an alpha-tubulin gene that directs expression in roots and promoters derived from zein storage protein genes which direct expression in endosperm. It is particularly contemplated that one may advantageously use the  
5 16 bp ocs enhancer element from the octopine synthase (ocs) gene (Ellis et al., 1987; Bouchez et al., 1989), especially when present in multiple copies, to achieve enhanced expression in roots.

Tissue specific expression may be functionally accomplished by introducing a constitutively expressed gene (all tissues) in combination with an antisense gene that is  
10 expressed only in those tissues where the gene product is not desired. For example, a gene coding for the crystal toxin protein from *B. thuringiensis* (Bt) may be introduced such that it is expressed in all tissues using the 35S promoter from Cauliflower Mosaic Virus. Expression of an antisense transcript of the Bt gene in a maize kernel, using for example a zein promoter, would prevent accumulation of the Bt protein in seed. Hence the protein encoded by the  
15 introduced gene would be present in all tissues except the kernel.

Expression of some genes in transgenic plants will be desired only under specified conditions. For example, it is proposed that expression of certain genes that confer resistance to environmental stress factors such as drought will be desired only under actual stress conditions. It is contemplated that expression of such genes throughout a plants development  
20 may have detrimental effects. It is known that a large number of genes exist that respond to the environment. For example, expression of some genes such as *rbcS*, encoding the small subunit of ribulose biphosphate carboxylase, is regulated by light as mediated through phytochrome. Other genes are induced by secondary stimuli. For example, synthesis of abscisic acid (ABA) is induced by certain environmental factors, including but not limited to  
25 water stress. A number of genes have been shown to be induced by ABA (Skriver and Mundy, 1990). It is also anticipated that expression of genes conferring resistance to insect predation would be desired only under conditions of actual insect infestation. Therefore, for some desired traits inducible expression of genes in transgenic plants will be desired.

Expression of a gene in a transgenic plant will be desired only in a certain time period  
30 during the development of the plant. Developmental timing is frequently correlated with tissue specific gene expression. For example, expression of zein storage proteins is initiated in the endosperm about 15 days after pollination.

Additionally, vectors may be constructed and employed in the intracellular targeting of a specific gene product within the cells of a transgenic plant or in directing a protein to the extracellular environment. This will generally be achieved by joining a DNA sequence encoding a transit or signal peptide sequence to the coding sequence of a particular gene. The resultant transit, or signal, peptide will transport the protein to a particular intracellular, or extracellular destination, respectively, and will then be post-translationally removed. Transit or signal peptides act by facilitating the transport of proteins through intracellular membranes, e.g., vacuole, vesicle, plastid and mitochondrial membranes, whereas signal peptides direct proteins through the extracellular membrane.

10 A particular example of such a use concerns the direction of a herbicide resistance gene, such as the EPSPS gene, to a particular organelle such as the chloroplast rather than to the cytoplasm. This is exemplified by the use of the rbcS transit peptide which confers plastid-specific targeting of proteins. In addition, it is proposed that it may be desirable to target certain genes responsible for male sterility to the mitochondria, or to target certain genes for resistance to phytopathogenic organisms to the extracellular spaces, or to target proteins to the vacuole.

By facilitating the transport of the protein into compartments inside and outside the cell, these sequences may increase the accumulation of gene product protecting them from proteolytic degradation. These sequences also allow for additional mRNA sequences from highly expressed genes to be attached to the coding sequence of the genes. Since mRNA being translated by ribosomes is more stable than naked mRNA, the presence of translatable mRNA in front of the gene may increase the overall stability of the mRNA transcript from the gene and thereby increase synthesis of the gene product. Since transit and signal sequences are usually post- translationally removed from the initial translation product, the use of these sequences allows for the addition of extra translated sequences that may not appear on the final polypeptide. Targeting of certain proteins may be desirable in order to enhance the stability of the protein (U.S. Patent No. 5,545,818).

It may be useful to target DNA itself within a cell. For example, it may be useful to target introduced DNA to the nucleus as this may increase the frequency of transformation. Within the nucleus itself it would be useful to target a gene in order to achieve site specific integration. For example, it would be useful to have an gene introduced through transformation replace an existing gene in the cell.

Other elements include those that can be regulated by endogenous or exogenous agents, e.g., by zinc finger proteins, including naturally occurring zinc finger proteins or chimeric zinc finger proteins (see, e.g., U.S. Patent No. 5,789,538, WO 99/48909; WO 99/45132; WO 98/53060; WO 98/53057; WO 98/53058; WO 00/23464; WO 95/19431; and WO 98/54311) or myb-like transcription factors. For example, a chimeric zinc finger protein may include amino acid sequences which bind to a specific DNA sequence (the zinc finger) and amino acid sequences that activate (e.g., GAL 4 sequences) or repress the transcription of the sequences linked to the specific DNA sequence.

### 3. Preferred Nucleic Acid Molecules of the Invention

The invention relates to an isolated plant, e.g., *Arabidopsis*, *Chenopodium* and rice, nucleic acid molecule comprising a gene having an open reading frame, the expression of which is altered in response to pathogen infection, as well as the endogenous plant promoters for those genes. However, the expression of these genes may also be altered in response to non-pathogens, e.g., in response to environmental stimuli. The nucleic acid molecules can be used in pathogen control strategies, e.g., by overexpressing nucleic acid molecules which can confer tolerance to a cell, or by altering the expression of host genes which are required for pathogen infection, e.g., by "knocking out" the expression of at least one genomic copy of the gene. Plants having genetic disruptions in host genes may be less susceptible to infection, e.g., due to a decrease or absence of a host protein needed for infection, or, alternatively, hypersusceptible to infection. Plants that are hypersusceptible to infection may be useful to prepare transgenic plants as the expression of the gene(s) which was disrupted may be related to gene silencing.

Preferred sources from which the nucleic acid molecules of the invention can be obtained or isolated include, but are not limited to, corn (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*)), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Cofea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*),

tea (*Camellia sinensis*), banana (*Musa* spp.), avocado (*Persea utilis*), fig (*Ficus casica*),  
 guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica*  
*papaya*), cashew (*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond  
 (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, duckweed  
 5 (*Lemna*), barley, vegetables, ornamentals, and conifers.

Duckweed (*Lemna*, see WO 00/07210) includes members of the family *Lemnaceae*.  
 There are known four genera and 34 species of duckweed as follows: genus *Lemna* (*L.*  
*aequinoctialis*, *L. disperma*, *L. ecuadoriensis*, *L. gibba*, *L. japonica*, *L. minor*, *L. miniscula*, *L.*  
*obscura*, *L. perpusilla*, *L. tenera*, *L. trisulca*, *L. turionifera*, *L. valdiviana*); genus *Spirodela* (*S.*  
 10 *intermedia*, *S. polyrrhiza*, *S. punctata*); genus *Woffia* (*Wa. Angusta*, *Wa. Arrhiza*, *Wa.*  
*Australina*, *Wa. Borealis*, *Wa. Brasiliensis*, *Wa. Columbiana*, *Wa. Elongata*, *Wa. Globosa*, *Wa.*  
*Microscopica*, *Wa. Neglecta*) and genus *Wofsiella* (*Wl. utila*, *Wl. utilis*, *Wl. gladiata*, *Wl.*  
*utila*, *Wl. lingulata*, *Wl. repunda*, *Wl. rotunda*, and *Wl. neotropica*). Any other genera or  
 species of *Lemnaceae*, if they exist, are also aspects of the present invention. *Lemna gibba*,  
 15 *Lemna minor*, and *Lemna miniscula* are preferred, with *Lemna minor* and *Lemna miniscula*  
 being most preferred. *Lemna* species can be classified using the taxonomic scheme described  
 by Landolt, Biosystematic Investigation on the Family of Duckweeds: The family of  
*Lemnaceae* – A Monograph Study. Geobotanischen Institut ETH, Stiftung Rubel, Zurich  
 (1986)).

20 Vegetables from which to obtain or isolate the nucleic acid molecules of the invention  
 include, but are not limited to, tomatoes (*Lycopersicon esculentum*), lettuce (e.g., *Lactuca*  
*sativa*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus*  
 spp.), and members of the genus *Cucumis* such as cucumber (*C. sativus*), cantaloupe (*C.*  
*cantalupensis*), and musk melon (*C. melo*). Ornamentals from which to obtain or isolate the  
 25 nucleic acid molecules of the invention include, but are not limited to, azalea (*Rhododendron*  
 spp.), hydrangea (*Macrophylla hydrangea*), hibiscus (*Hibiscus rosasanensis*), roses (*Rosa*  
 spp.), tulips (*Tulipa* spp.), daffodils (*Narcissus* spp.), petunias (*Petunia hybrida*), carnation  
 (*Dianthus caryophyllus*), poinsettia (*Euphorbia pulcherrima*), and chrysanthemum. Conifers  
 that may be employed in practicing the present invention include, for example, pines such as  
 30 loblolly pine (*Pinus taeda*), slash pine (*Pinus elliotii*), ponderosa pine (*Pinus ponderosa*),  
 lodgepole pine (*Pinus contorta*), and Monterey pine (*Pinus radiata*), Douglas-fir (*Pseudotsuga*  
*menziesii*); Western hemlock (*Tsuga utilis*); Sitka spruce (*Picea glauca*); redwood (*Sequoia*  
*sempervirens*); true firs such as silver fir (*Abies amabilis*) and balsam fir (*Abies balsamea*);



and cedars such as Western red cedar (*Thuja plicata*) and Alaska yellow-cedar (*Chamaecyparis nootkatensis*). Leguminous plants from which the nucleic acid molecules of the invention can be isolated or obtained include, but are not limited to, beans and peas. Beans include guar, locust bean, fenugreek, soybean, garden beans, cowpea, mungbean, lima bean, fava bean, lentils, chickpea, and the like. Legumes include, but are not limited to, *Arachis*, e.g., peanuts, *Vicia*, e.g., crown vetch, hairy vetch, adzuki bean, mung bean, and chickpea, *Lupinus*, e.g., lupine, trifolium, *Phaseolus*, e.g., common bean and lima bean, *Pisum*, e.g., field bean, *Melilotus*, e.g., clover, *Medicago*, e.g., alfalfa, *Lotus*, e.g., trefoil, lens, e.g., lentil, and false indigo.

10 Papaya, garlic, pea, peach, pepper, petunia, sorghum, sweet potato, turnip, safflower, corn, pea, endive, gourd, grape, snap bean, chicory, cotton, tobacco, aubergine, beet, buckwheat, broad bean, nectarine, avocado, mango, banana, groundnut, potato, peanut, lettuce, pineapple, spinach, squash, sugarbeet, sugarcane, sweet corn, chrysanthemum.

Other preferred sources of the nucleic acid molecules of the invention include *Acacia*,  
 15 aneth, artichoke, arugula, blackberry, canola, cilantro, clementines, escarole, eucalyptus, fennel, grapefruit, honey dew, jicama, kiwifruit, lemon, lime, mushroom, nut, okra, orange, parsley, persimmon, plantain, pomegranate, poplar, radiata pine, radicchio, Southern pine, sweetgum, tangerine, triticale, vine, yams, apple, pear, quince, cherry, apricot, melon, hemp, buckwheat, grape, raspberry, chenopodium, blueberry, nectarine, peach, plum, strawberry,  
 20 watermelon, eggplant, pepper, cauliflower, Brassica, e.g., broccoli, cabbage, utilian sprouts, onion, carrot, leek, beet, broad bean, celery, radish, pumpkin, endive, gourd, garlic, snapbean, spinach, squash, turnip, utilane, and zucchini.

Yet other sources of nucleic acid molecules are ornamental plants including, but not limited to, impatiens, Begonia, Pelargonium, Viola, Cyclamen, Verbena, Vinca, Tagetes,  
 25 Primula, Saint Paulia, Agertum, Amaranthus, Antihirrhinum, Aquilegia, Cineraria, Clover, Cosmo, Cowpea, Dahlia, Datura, Delphinium, Gerbera, Gladiolus, Gloxinia, Hippeastrum, Mesembryanthemum, Salpiglossos, and Zinnia, and plants such as those shown in Table 1.

Table 1

| FAMILY | LATIN NAME | COMMON NAME | MAP REFERENCES<br>RESOURCES | LINKS |
|--------|------------|-------------|-----------------------------|-------|
|        |            |             |                             |       |

| FAMILY        | LATIN NAME                | COMMON NAME        | MAP REFERENCES<br>RESOURCES | LINKS   |
|---------------|---------------------------|--------------------|-----------------------------|---|
| Cucurbitaceae | <i>Cucumis sativus</i>    | Cucumber           |                             | <a href="http://www.cucurbit.org/">http://www.cucurbit.org/</a>                               |
|               | <i>Cucumis melo</i>       | Melon              |                             | <a href="http://genome.cornell.edu/cgi/c/">http://genome.cornell.edu/cgi/c/</a>               |
|               | <i>Citrullus lanatus</i>  | Watermelon         |                             |   |
|               | <i>Cucurbita pepo</i>     | Squash – summer    |                             |   |
|               | <i>Cucurbita maxima</i>   | Squash – winter    |                             |   |
|               | <i>Cucurbita moschata</i> | Pumpkin /butternut |                             |   |
| Total         |                           |                    |                             | <a href="http://www.nal.usda.gov/pgdic/Map_proj/">http://www.nal.usda.gov/pgdic/Map_proj/</a> |
|               |                           |                    |                             |   |

| FAMILY     | LATIN NAME                         | COMMON NAME  | MAP REFERENCES<br>RESOURCES  | LINKS   |
|------------|------------------------------------|--------------|--|---|
| Solanaceae | <i>Lycopersicon<br/>esculentum</i> | Tomato       | <ul style="list-style-type: none"> <li>• 15x BAC on variety Heinz 1706 order from Clemson Genome center (www.genome.clemson.edu)</li> <li>• 11.6x BAC of L. cheesmanii (originates from J. Giovannoni) available from Clemson genome center (www.genome.clemson.edu)</li> <li>• EST collection from TIGR (www.tigr.org/tdb/lgi/index.html)</li> <li>• EST collection from Clemson Genome Center (www.genome.clemson.edu)</li> <li>• TAG 99:254-271, 1999 (esculentum x pennelli)</li> <li>• TAG 89:1007-1013, 1994 (peruvianum)</li> <li>• Plant Cell Reports 12:293-297, 1993 (RAPDs)</li> <li>• Genetics 132:1141-1160, 1992 (potato x tomato)</li> <li>• Genetics 120:1095-1105, 1988 (RFLP potato and tomato)</li> <li>• Genetics 115:387-393, 1986 (esculentum x pennelli isozyme and cDNAs)</li> </ul> | <a href="http://genome.cornell.edu/solgenes">genome.cornell.edu/solgenes</a><br><a href="http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=solgenes">http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=solgenes</a><br><a href="http://genome.cornell.edu/tgc/">http://genome.cornell.edu/tgc/</a><br><a href="http://tgrc.ucdavis.edu/">http://tgrc.ucdavis.edu/</a> |
|            | <i>Capsicum<br/>annuum</i>         | Pepper       |  | <a href="http://neptune.netimages.com/~chile/science.html">http://neptune.netimages.com/~chile/science.html</a>   |
|            | <i>Capsicum<br/>frutescens</i>     | Chile pepper |  |   |
|            | <i>Solanum<br/>melongena</i>       | Eggplant     |  |   |

| FAMILY       | LATIN NAME                                   | COMMON NAME     | MAP REFERENCES<br>RESOURCES  | LINKS  |
|--------------|--|-----------------|--|--|
|              | <i>(Nicotiana tabacum)</i>                   | (Tobacco)       |  |  |
|              | <i>(Solanum tuberosum)</i>                   | (Potato)        |  |  |
|              | <i>(Petunia x hybrida hort. Ex E. Vilm.)</i> | (Petunia)       | 4x BAC of Petunia hybrida 7984 available from Clemson genome center (www.genome.clemson.edu)     |  |
| Total        |  |                 |  | <a href="http://www.nal.usda.gov/pgdic/Map_proj/">http://www.nal.usda.gov/pgdic/Map_proj/</a>  |
| Brassicaceae | <i>Brassica oleracea L. var. italica</i>     | Broccoli        |  | <a href="http://res.agr.cornell.edu/ecorc/cwmt/crucifer/traits/index.htm">http://res.agr.cornell.edu/ecorc/cwmt/crucifer/traits/index.htm</a><br><a href="http://geneous.cit.cornell.edu/cabbage/aboutcab.html">http://geneous.cit.cornell.edu/cabbage/aboutcab.html</a> |
|              | <i>Brassica oleracea L. var. capitata</i>    | Cabbage         |  |  |
|              | <i>Brassica rapa</i>                         | Chinese Cabbage |  |  |
|              | <i>Brassica oleracea L. var. botrytis</i>    | Cauliflower     |  |  |
|              | <i>Raphanus sativus var. niger</i>           | Daikon          |  |  |
|              | <i>(Brassica napus)</i>                      | (Oilseed rape)  |  | <a href="http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=brassicadb">http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=brassicadb</a>  |
|              |  | Arabidopsis     | 12x and 6x BACs on Columbia strain available from Clemson genome center (www.genome.clemson.edu) | <a href="http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=agr">http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=agr</a>  |

| FAMILY         | LATIN NAME                | COMMON NAME  | MAP REFERENCES<br>RESOURCES  | LINKS   |
|----------------|---------------------------|--------------|--|---|
| Total          |                           |              |  | <a href="http://www.nal.usda.gov/pgdic/Map_proj/">http://www.nal.usda.gov/pgdic/Map_proj/</a>   |
| Umbelliferae   | <i>Daucus carota</i>      | Carrot       |  |   |
| Compositae     | <i>Lactuca sativa</i>     | Lettuce      |  |   |
|                | <i>Helianthus annuus</i>  | (Sunflower)  |  |   |
| Total          |                           |              |  |   |
| Chenopodiaceae | <i>Spinacia oleracea</i>  | Spinach      |  |   |
|                | ( <i>Beta vulgaris</i> )  | (Sugar Beet) |  |   |
| Total          |                           |              |  |   |
| Leguminosae    | <i>Phaseolus vulgaris</i> | Bean         | 4.3x BAC available from Clemson genome center ( <a href="http://www.genome.clemson.edu">www.genome.clemson.edu</a> )                             | <a href="http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=beanengenes">http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=beanengenes</a> |
|                | <i>Pisum sativum</i>      | Pea          |  |   |
|                | ( <i>Glycine max</i> )    | (Soybean)    | 7.5x and 7.9x BACs available from Clemson genome center ( <a href="http://www.genome.clemson.edu">www.genome.clemson.edu</a> )                   | <a href="http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=soybase">http://ars-genome.cornell.edu/cgi-bin/WebAce/webace?db=soybase</a>         |
| Total          |                           |              | <a href="http://www.nal.usda.gov/pgdic/Map_proj/">http://www.nal.usda.gov/pgdic/Map_proj/</a>  |   |
| Gramineae      | <i>Zea mays</i>           | Sweet Corn   | Novartis BACs for Mo17 and B73 have been donated to Clemson Genome Center ( <a href="http://www.genome.clemson.edu">www.genome.clemson.edu</a> ) |   |
|                | ( <i>Zea mays</i> )       | (Field Corn) |  | <a href="http://www.agron.missouri.edu/mnl/">http://www.agron.missouri.edu/mnl/</a>   |

| FAMILY    | LATIN NAME         | COMMON NAME | MAP REFERENCES<br>RESOURCES   | LINKS |
|-----------|--------------------|-------------|---|-------|
| Total     |                    |             | <a href="http://www.nal.usda.gov/pgdic/Map_proj/">http://www.nal.usda.gov/pgdic/Map_proj/</a> |       |
| Liliaceae | <i>Allium cepa</i> | Onion       |   |       |
|           |                    | Leek        |   |       |
|           |                    | (Garlic)    |   |       |
|           |                    | (Asparagus) |   |       |
| Total     |                    |             | <a href="http://www.nal.usda.gov/pgdic/Map_proj/">http://www.nal.usda.gov/pgdic/Map_proj/</a> |       |

Preferred forage and turf grass nucleic acid sources for the nucleic acid molecules of the invention include, but are not limited to, alfalfa, orchard grass, tall fescue, perennial ryegrass, creeping bent grass, and redtop. Yet other preferred sources include, but are not limited to, crop plants and in particular cereals (for example, corn, alfalfa, sunflower, rice, *Brassica*, canola, soybean, barley, soybean, sugarbeet, cotton, safflower, peanut, sorghum, oat, rye, rape, wheat, millet, tobacco, and the like), and even more preferably corn, rice and soybean.

According to one embodiment, the present invention is directed to a nucleic acid molecule comprising a nucleotide sequence isolated or obtained from any plant which encodes a polypeptide having at least 70% amino acid sequence identity to a polypeptide encoded by a gene comprising any one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737, or a gene comprising SEQ ID NOs:2137-2661 or 4738-6813. Based on the *Arabidopsis*, *Chenopodium* and rice nucleic acid sequences of the present invention, orthologs may be identified or isolated from the genome of any desired organism, preferably from another plant, according to well known techniques based on their sequence similarity to the *Arabidopsis*, *Chenopodium* and rice nucleic acid sequences, e.g., hybridization, PCR or computer generated sequence comparisons. For example, all or a portion of a particular *Arabidopsis*, *Chenopodium* and rice nucleic acid sequence is used as a probe that selectively hybridizes to other gene sequences present in a population of cloned genomic DNA fragments or cDNA fragments (i.e., genomic or cDNA libraries) from a chosen source organism. Further, suitable genomic and cDNA libraries may be prepared from any cell or tissue of an organism. Such techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, e.g., Sambrook et al., 1989) and amplification by PCR using oligonucleotide primers

preferably corresponding to sequence domains conserved among related polypeptide or subsequences of the nucleotide sequences provided herein (see, e.g., Innis et al., 1990). These methods are particularly well suited to the isolation of gene sequences from organisms closely related to the organism from which the probe sequence is derived. The application of these methods using the *Arabidopsis* sequences as probes is well suited for the isolation of gene sequences from any source organism, preferably other plant species. In a PCR approach, oligonucleotide primers can be designed for use in PCR reactions to amplify corresponding DNA sequences from cDNA or genomic DNA extracted from any plant of interest. Methods for designing PCR primers and PCR cloning are generally known in the art.

In hybridization techniques, all or part of a known nucleotide sequence is used as a probe that selectively hybridizes to other corresponding nucleotide sequences present in a population of cloned genomic DNA fragments or cDNA fragments (i.e., genomic or cDNA libraries) from a chosen organism. The hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments, or other oligonucleotides, and may be labeled with a detectable group such as  $^{32}\text{P}$ , or any other detectable marker. Thus, for example, probes for hybridization can be made by labeling synthetic oligonucleotides based on the sequence of the invention. Methods for preparation of probes for hybridization and for construction of cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook et al. (1989). In general, sequences that hybridize to the sequences disclosed herein will have at least 40% to 50%, about 60% to 70% and even about 80% 85%, 90%, 95% to 98% or more identity with the disclosed sequences. That is, the sequence similarity of sequences may range, sharing at least about 40% to 50%, about 60% to 70%, and even about 80%, 85%, 90%, 95% to 98% sequence similarity.

The nucleic acid molecules of the invention can also be identified by, for example, a search of known databases for genes encoding polypeptides having a specified amino acid sequence identity or DNA having a specified nucleotide sequence identity. Methods of alignment of sequences for comparison are well known in the art and are described hereinabove.

#### 4. Methods for Mutagenizing DNA

It is specifically contemplated by the inventors that one could mutagenize DNA having a promoter or open reading frame to, for example, potentially improve the utility of the DNA for expression of transgenes in plants. The mutagenesis can be carried out at random and the mutagenized sequences screened for activity in a trial-by-error procedure. Alternatively,

particular sequences which provide the promoter with desirable expression characteristics, or a promoter with expression enhancement activity, could be identified and these or similar sequences introduced into the sequences via mutation. It is further contemplated that one could mutagenize these sequences in order to enhance their expression of transgenes in a particular species.

The means for mutagenizing a DNA segment of the current invention are well-known to those of skill in the art. As indicated, modifications may be made by random or site-specific mutagenesis procedures. The DNA may be modified by altering its structure through the addition or deletion of one or more nucleotides from the sequence which encodes the corresponding un-modified sequences.

Mutagenesis may be performed in accordance with any of the techniques known in the art, such as, and not limited to, synthesizing an oligonucleotide having one or more mutations within the sequence of a particular regulatory region. In particular, site-specific mutagenesis is a technique useful in the preparation of promoter mutants, through specific mutagenesis of the underlying DNA. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to about 75 nucleotides or more in length is preferred, with about 10 to about 25 or more residues on both sides of the junction of the sequence being altered.

In general, the technique of site-specific mutagenesis is well known in the art, as exemplified by various publications. As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially available and their use is generally well known to those skilled in the art.

Double stranded plasmids also are routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double stranded vector which includes within its sequence a DNA sequence which encodes the promoter. An



oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation.

This heteroduplex vector is then used to transform or transfect appropriate cells, such as *E. coli* cells, and cells are selected which include recombinant vectors bearing the mutated sequence arrangement. Vector DNA can then be isolated from these cells and used for plant transformation. A genetic selection scheme was devised by Kunkel et al. (1987) to enrich for clones incorporating mutagenic oligonucleotides. Alternatively, the use of PCR with commercially available thermostable enzymes such as Taq polymerase may be used to incorporate a mutagenic oligonucleotide primer into an amplified DNA fragment that can then be cloned into an appropriate cloning or expression vector. The PCR-mediated mutagenesis procedures of Tomic et al. (1990) and Upender et al. (1995) provide two examples of such protocols. A PCR employing a thermostable ligase in addition to a thermostable polymerase also may be used to incorporate a phosphorylated mutagenic oligonucleotide into an amplified DNA fragment that may then be cloned into an appropriate cloning or expression vector. The mutagenesis procedure described by Michael (1994) provides an example of one such protocol.

The preparation of sequence variants of DNA segments using site-directed mutagenesis is provided as a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of DNA sequences may be obtained. For example, recombinant vectors encoding the desired promoter sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.

In addition, an unmodified or modified nucleotide sequence of the present invention can be varied by shuffling the sequence of the invention. To test for a function of variant DNA sequences according to the invention, the sequence of interest is operably linked to a selectable or screenable marker gene and expression of the marker gene is tested in transient expression assays with protoplasts or in stably transformed plants. It is known to the skilled artisan that DNA sequences capable of driving expression of an associated nucleotide sequence are build in a modular way. Accordingly, expression levels from shorter DNA fragments may be different than the one from the longest fragment and may be different from each other. For

example, deletion of a down-regulating upstream element will lead to an increase in the expression levels of the associated nucleotide sequence while deletion of an up-regulating element will decrease the expression levels of the associated nucleotide sequence. It is also known to the skilled artisan that deletion of development-specific or a tissue-specific element will lead to a temporally or spatially altered expression profile of the associated nucleotide sequence.

As used herein, the term "oligonucleotide directed mutagenesis procedure" refers to template-dependent processes and vector-mediated propagation which result in an increase in the concentration of a specific nucleic acid molecule relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed mutagenesis procedure" also is intended to refer to a process that involves the template-dependent extension of a primer molecule. The term template-dependent process refers to nucleic acid synthesis of an RNA or a DNA molecule wherein the sequence of the newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson and Rarnstad, 1987). Typically, vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of the amplified nucleic acid fragment. Examples of such methodologies are provided by U.S. Patent No. 4,237,224. A number of template dependent processes are available to amplify the target sequences of interest present in a sample, such methods being well known in the art and specifically disclosed herein below.

Where a clone comprising a promoter has been isolated in accordance with the instant invention, one may wish to delimit the essential promoter regions within the clone. One efficient, targeted means for preparing mutagenizing promoters relies upon the identification of putative regulatory elements within the promoter sequence. This can be initiated by comparison with promoter sequences known to be expressed in similar tissue-specific or developmentally unique manner. Sequences which are shared among promoters with similar expression patterns are likely candidates for the binding of transcription factors and are thus likely elements which confer expression patterns. Confirmation of these putative regulatory elements can be achieved by deletion analysis of each putative regulatory region followed by functional analysis of each deletion construct by assay of a reporter gene which is functionally

attached to each construct. As such, once a starting promoter sequence is provided, any of a number of different deletion mutants of the starting promoter could be readily prepared.

As indicated above, deletion mutants, deletion mutants of the promoter of the invention also could be randomly prepared and then assayed. With this strategy, a series of constructs are prepared, each containing a different portion of the clone (a subclone), and these constructs are then screened for activity. A suitable means for screening for activity is to attach a deleted promoter or intron construct which contains a deleted segment to a selectable or screenable marker, and to isolate only those cells expressing the marker gene. In this way, a number of different, deleted promoter constructs are identified which still retain the desired, or even enhanced, activity. The smallest segment which is required for activity is thereby identified through comparison of the selected constructs. This segment may then be used for the construction of vectors for the expression of exogenous genes.

### **B. Marker Genes**

In order to improve the ability to identify transformants, one may desire to employ a selectable or screenable marker gene as, or in addition to, the expressible gene of interest. "Marker genes" are genes that impart a distinct phenotype to cells expressing the marker gene and thus allow such transformed cells to be distinguished from cells that do not have the marker. Such genes may encode either a selectable or screenable marker, depending on whether the marker confers a trait which one can 'select' for by chemical means, i.e., through the use of a selective agent (e.g., a herbicide, antibiotic, or the like), or whether it is simply a trait that one can identify through observation or testing, i.e., by 'screening' (e.g., the R-locus trait, the green fluorescent protein (GFP)). Of course, many examples of suitable marker genes are known to the art and can be employed in the practice of the invention.

Included within the terms selectable or screenable marker genes are also genes which encode a "secretable marker" whose secretion can be detected as a means of identifying or selecting for transformed cells. Examples include markers which encode a secretable antigen that can be identified by antibody interaction, or even secretable enzymes which can be detected by their catalytic activity. Secretable proteins fall into a number of classes, including small, diffusible proteins detectable, e.g., by ELISA; small active enzymes detectable in extracellular solution (e.g., alpha-amylase, beta-lactamase, phosphinothricin acetyltransferase); and proteins that are inserted or trapped in the cell wall (e.g., proteins that include a leader sequence such as that found in the expression unit of extensin or tobacco PR-S).

With regard to selectable secretable markers, the use of a gene that encodes a protein that becomes sequestered in the cell wall, and which protein includes a unique epitope is considered to be particularly advantageous. Such a secreted antigen marker would ideally employ an epitope sequence that would provide low background in plant tissue, a promoter-  
5 leader sequence that would impart efficient expression and targeting across the plasma membrane, and would produce protein that is bound in the cell wall and yet accessible to antibodies. A normally secreted wall protein modified to include a unique epitope would satisfy all such requirements.

One example of a protein suitable for modification in this manner is extensin, or  
10 hydroxyproline rich glycoprotein (HPRG). For example, the maize HPRG (Steifel et al., 1990) molecule is well characterized in terms of molecular biology, expression and protein structure. However, any one of a variety of utilane and/or glycine-rich wall proteins (Keller et al., 1989) could be modified by the addition of an antigenic site to create a screenable marker.

One exemplary embodiment of a secretable screenable marker concerns the use of a  
15 maize sequence encoding the wall protein HPRG, modified to include a 15 residue epitope from the pro-region of murine interleukin, however, virtually any detectable epitope may be employed in such embodiments, as selected from the extremely wide variety of antigen-antibody combinations known to those of skill in the art. The unique extracellular epitope can then be straightforwardly detected using antibody labeling in conjunction with chromogenic or  
20 fluorescent adjuncts.

Elements of the present disclosure may be exemplified in detail through the use of the bar and/or GUS genes, and also through the use of various other markers. Of course, in light of this disclosure, numerous other possible selectable and/or screenable marker genes will be apparent to those of skill in the art in addition to the one set forth hereinbelow. Therefore, it  
25 will be understood that the following discussion is exemplary rather than exhaustive. In light of the techniques disclosed herein and the general recombinant techniques which are known in the art, the present invention renders possible the introduction of any gene, including marker genes, into a recipient cell to generate a transformed plant.

#### 1. Selectable Markers

30 Possible selectable markers for use in connection with the present invention include, but are not limited to, a *neo* gene which codes for kanamycin resistance and can be selected for using kanamycin, G418, paromomycin, and the like; a *bar* gene which codes for bialaphos or phosphinothricin resistance; a gene which encodes an altered EPSP synthase protein (Hinchee

et al., 1988) thus conferring glyphosate resistance; a nitrilase gene such as *bxn* from *Klebsiella*  
*ozaenae* which confers resistance to bromoxynil (Stalker et al., 1988); a mutant acetolactate  
synthase gene (ALS) which confers resistance to imidazolinone, sulfonyleurea or other ALS-  
inhibiting chemicals (European Patent Application 154,204, 1985); a methotrexate-resistant  
5 DHFR gene (Thillet et al., 1988); a dalapon dehalogenase gene that confers resistance to the  
herbicide dalapon; a mutated anthranilate synthase gene that confers resistance to 5-methyl  
tryptophan. Preferred selectable marker genes encode phosphinothricin acetyltransferase;  
glyphosate resistant EPSPS, aminoglycoside phosphotransferase; hygromycin  
phosphotransferase, or neomycin phosphotransferase. Where a mutant EPSP synthase gene is  
10 employed, additional benefit may be realized through the incorporation of a suitable  
chloroplast transit peptide, CTP (European Patent Application 0,218,571, 1987).

An illustrative embodiment of a selectable marker gene capable of being used in  
systems to select transformants is the genes that encode the enzyme phosphinothricin  
acetyltransferase, such as the *bar* gene from *Streptomyces hygroscopicus* or the *pat* gene from  
15 *Streptomyces viridochromogenes*. The enzyme phosphinothricin acetyl transferase (PAT)  
inactivates the active ingredient in the herbicide bialaphos, phosphinothricin (PPT). PPT  
inhibits glutamine synthetase, (Murakami et al., 1986; Twell et al., 1989) causing rapid  
accumulation of ammonia and cell death. The success in using this selective system in  
conjunction with monocots was particularly surprising because of the major difficulties which  
20 have been reported in transformation of cereals.

Where one desires to employ a bialaphos resistance gene in the practice of the  
invention, a particularly useful gene for this purpose is the *bar* or *pat* genes obtainable from  
species of *Streptomyces* (e.g., ATCC No. 21,705). The cloning of the *bar* gene has been  
described (Murakami et al., 1986; Thompson et al., 1987) as has the use of the *bar* gene in the  
25 context of plants other than monocots (De Block et al., 1987; De Block et al., 1989).

Selection markers resulting in positive selection, such as a phosphomannose  
isomerase gene, as described in patent application WO 93/05163, may also be used.  
Alternative genes to be used for positive selection are described in WO 94/20627 and encode  
xyloisomerases and phosphomanno-isomerases such as mannose-6-phosphate isomerase and  
30 mannose-1-phosphate isomerase; phosphomanno mutase; mannose epimerases such as those  
which convert carbohydrates to mannose or mannose to carbohydrates such as glucose or  
galactose; phosphatases such as mannose or xylose phosphatase, mannose-6-phosphatase and  
mannose-1-phosphatase, and permeases which are involved in the transport of mannose, or a

derivative, or a precursor thereof into the cell. Transformed cells are identified without damaging or killing the non-transformed cells in the population and without co-introduction of antibiotic or herbicide resistance genes. As described in WO 93/05163, in addition to the fact that the need for antibiotic or herbicide resistance genes is eliminated, it has been shown that the positive selection method is often far more efficient than traditional negative selection.

## 2. Screenable Markers

Screenable markers that may be employed include, but are not limited to, a beta-glucuronidase (GUS) or *uidA* gene which encodes an enzyme for which various chromogenic substrates are known; an R-locus gene, which encodes a product that regulates the production of anthocyanin pigments (red color) in plant tissues (Dellaporta et al., 1988); a beta-lactamase gene (Sutcliffe, 1978), which encodes an enzyme for which various chromogenic substrates are known (e.g., PADAC, a chromogenic cephalosporin); a *xylE* gene (Zukowsky et al., 1983) which encodes a catechol dioxygenase that can convert chromogenic catechols; an  $\alpha$ -amylase gene (Ikuta et al., 1990); a tyrosinase gene (Katz et al., 1983) which encodes an enzyme capable of oxidizing tyrosine to DOPA and dopaquinone which in turn condenses to form the easily detectable compound melanin; a  $\beta$ -galactosidase gene, which encodes an enzyme for which there are chromogenic substrates; a luciferase (*lux*) gene (Ow et al., 1986), which allows for bioluminescence detection; or even an aequorin gene (Prasher et al., 1985), which may be employed in calcium-sensitive bioluminescence detection, or a green fluorescent protein gene (Niedz et al., 1995).

Genes from the maize R gene complex are contemplated to be particularly useful as screenable markers. The R gene complex in maize encodes a protein that acts to regulate the production of anthocyanin pigments in most seed and plant tissue. A gene from the R gene complex was applied to maize transformation, because the expression of this gene in transformed cells does not harm the cells. Thus, an R gene introduced into such cells will cause the expression of a red pigment and, if stably incorporated, can be visually scored as a red sector. If a maize line is carries dominant *utila* for genes encoding the enzymatic intermediates in the anthocyanin biosynthetic pathway (C2, A1, A2, Bz1 and Bz2) (Roth et al., 1990), but carries a recessive allele at the R locus, transformation of any cell from that line with R will result in red pigment formation. Exemplary lines include Wisconsin 22 which contains the *rg*-Stadler allele and TR112, a K55 derivative which is *r-g*, *b*, *P1*. Alternatively any genotype of maize can be utilized if the C1 and R alleles are introduced together.

It is further proposed that R gene regulatory regions may be employed in chimeric constructs in order to provide mechanisms for controlling the expression of chimeric genes. More diversity of phenotypic expression is known at the R locus than at any other locus (Coe et al., 1988). It is contemplated that regulatory regions obtained from regions 5' to the structural R gene would be valuable in directing the expression of genes, e.g., insect resistance, drought resistance, herbicide tolerance or other protein coding regions. For the purposes of the present invention, it is believed that any of the various R gene family members may be successfully employed (e.g., P, S, Lc, etc.). However, the most preferred will generally be Sn (particularly Sn:bol3). Sn is a dominant member of the R gene complex and is functionally similar to the R and B loci in that Sn controls the tissue specific deposition of anthocyanin pigments in certain seedling and plant cells, therefore, its phenotype is similar to R.

A further screenable marker contemplated for use in the present invention is firefly luciferase, encoded by the *lux* gene. The presence of the *lux* gene in transformed cells may be detected using, for example, X-ray film, scintillation counting, fluorescent spectrophotometry, low-light video cameras, photon counting cameras or multiwell luminometry. It is also envisioned that this system may be developed for populational screening for bioluminescence, such as on tissue culture plates, or even for whole plant screening. Where use of a screenable marker gene such as *lux* or GFP is desired, benefit may be realized by creating a gene fusion between the screenable marker gene and a selectable marker gene, for example, a GFP-NPTII gene fusion. This could allow, for example, selection of transformed cells followed by screening of transgenic plants or seeds.

### C. Exogenous Genes for Modification of Plant Phenotypes

Genes of interest are reflective of the commercial markets and interests of those involved in the development of the crop. Crops and markets of interest changes, and as developing nations open up world markets, new crops and technologies will also emerge. In addition, as the understanding of agronomic traits and characteristics such as yield and heterosis increase, the choice of genes for transformation will change accordingly. General categories of genes of interest include, for example, those genes involved in information, such as zinc fingers, those involved in communication, such as kinases, and those involved in housekeeping, such as heat shock proteins. More specific categories of transgenes, for example, include genes encoding important traits for agronomics, insect resistance, disease resistance, herbicide resistance, sterility, grain characteristics, and commercial products. Genes of interest include, generally, those involved in starch, oil, carbohydrate, or nutrient

metabolism, as well as those affecting kernel size, sucrose loading, zinc finger proteins, see, e.g., U.S. Patent No. 5,789,538, WO 99/48909; WO 99/45132; WO 98/53060; WO 98/53057; WO 98/53058; WO 00/23464; WO 95/19431; and WO 98/54311, and the like.

One skilled in the art recognizes that the expression level and regulation of a transgene  
5 in a plant can vary significantly from line to line. Thus, one has to test several lines to find one with the desired expression level and regulation. Once a line is identified with the desired regulation specificity of a chimeric Cre transgene, it can be crossed with lines carrying different inactive replicons or inactive transgene for activation.

Other sequences which may be linked to the gene of interest which encodes a  
10 polypeptide are those which can target to a specific organelle, e.g., to the mitochondria, nucleus, or plastid, within the plant cell. Targeting can be achieved by providing the polypeptide with an appropriate targeting peptide sequence, such as a secretory signal peptide (for secretion or cell wall or membrane targeting, a plastid transit peptide, a chloroplast transit peptide, e.g., the chlorophyll a/b binding protein, a mitochondrial target peptide, a vacuole  
15 targeting peptide, or a nuclear targeting peptide, and the like. For example, the small subunit of ribulose biphosphate carboxylase transit peptide, the EPSPS transit peptide or the dihydrodipicolinic acid synthase transit peptide may be used. For examples of plastid organelle targeting sequences (see WO 00/12732). Plastids are a class of plant organelles derived from proplastids and include chloroplasts, leucoplasts, aravloplast, and chromoplasts.  
20 The plastids are major sites of biosynthesis in plants. In addition to photosynthesis in the chloroplast, plastids are also sites of lipid biosynthesis, nitrate reduction to ammonium, and starch storage. And while plastids contain their own circular genome, most of the proteins localized to the plastids are encoded by the nuclear genome and are imported into the organelle from the cytoplasm.

25 Transgenes used with the present invention will often be genes that direct the expression of a particular protein or polypeptide product, but they may also be non-expressible DNA segments, e.g., transposons such as Ds that do not direct their own transposition. As used herein, an "expressible gene" is any gene that is capable of being transcribed into RNA (e.g., mRNA, antisense RNA, etc.) or translated into a protein, expressed as a trait of interest, or the  
30 like, etc., and is not limited to selectable, screenable or non-selectable marker genes. The invention also contemplates that, where both an expressible gene that is not necessarily a marker gene is employed in combination with a marker gene, one may employ the separate



genes on either the same or different DNA segments for transformation. In the latter case, the different vectors are delivered concurrently to recipient cells to maximize cotransformation.

The choice of the particular DNA segments to be delivered to the recipient cells will often depend on the purpose of the transformation. One of the major purposes of transformation of crop plants is to add some commercially desirable, agronomically important traits to the plant. Such traits include, but are not limited to, herbicide resistance or tolerance; insect resistance or tolerance; disease resistance or tolerance (viral, bacterial, fungal, nematode); stress tolerance and/or resistance, as exemplified by resistance or tolerance to drought, heat, chilling, freezing, excessive moisture, salt stress; oxidative stress; increased yields; food content and makeup; physical appearance; male sterility; drydown; standability; prolificacy; starch properties; oil quantity and quality; and the like. One may desire to incorporate one or more genes conferring any such desirable trait or traits, such as, for example, a gene or genes encoding pathogen resistance.

In certain embodiments, the present invention contemplates the transformation of a recipient cell with more than one advantageous transgene. Two or more transgenes can be supplied in a single transformation event using either distinct transgene-encoding vectors, or using a single vector incorporating two or more gene coding sequences. For example, plasmids bearing the *bar* and *aroA* expression units in either convergent, divergent, or colinear orientation, are considered to be particularly useful. Further preferred combinations are those of an insect resistance gene, such as a *Bt* gene, along with a protease inhibitor gene such as *pinII*, or the use of *bar* in combination with either of the above genes. Of course, any two or more transgenes of any description, such as those conferring herbicide, insect, disease (viral, bacterial, fungal, nematode) or drought resistance, male sterility, drydown, standability, prolificacy, starch properties, oil quantity and quality, or those increasing yield or nutritional quality may be employed as desired.

#### 1. Herbicide Resistance

The genes encoding phosphinothricin acetyltransferase (*bar* and *pat*), glyphosate tolerant EPSP synthase genes, the glyphosate degradative enzyme gene *gox* encoding glyphosate oxidoreductase, *deh* (encoding a dehalogenase enzyme that inactivates dalapon), herbicide resistant (e.g., sulfonyleurea and imidazolinone) acetolactate synthase, and *bxn* genes (encoding a nitrilase enzyme that degrades bromoxynil) are good examples of herbicide resistant genes for use in transformation. The *bar* and *pat* genes code for an enzyme, phosphinothricin acetyltransferase (PAT), which inactivates the herbicide phosphinothricin

and prevents this compound from inhibiting glutamine synthetase enzymes. The enzyme 5-enolpyruvylshikimate 3-phosphate synthase (EPSP Synthase), is normally inhibited by the herbicide N-(phosphonomethyl)glycine (glyphosate). However, genes are known that encode glyphosate-resistant EPSP Synthase enzymes.

- 5        These genes are particularly contemplated for use in monocot transformation. The *deh* gene encodes the enzyme dalapon dehalogenase and confers resistance to the herbicide dalapon. The *bxn* gene codes for a specific nitrilase enzyme that converts bromoxynil to a non-herbicidal degradation product.

## 2. Insect Resistance

- 10        An important aspect of the present invention concerns the introduction of insect resistance-conferring genes into plants. Potential insect resistance genes which can be introduced include *Bacillus thuringiensis* crystal toxin genes or *Bt* genes (Watrud et al., 1985). *Bt* genes may provide resistance to lepidopteran or coleopteran pests such as European Corn Borer (ECB) and corn rootworm (CRW). Preferred *Bt* toxin genes for use in such
- 15        embodiments include the CryIA(b) and CryIA(c) genes. Endotoxin genes from other species of *B. thuringiensis* which affect insect growth or development may also be employed in this regard.

- The poor expression of *Bt* toxin genes in plants is a well-documented phenomenon, and the use of different promoters, fusion proteins, and leader sequences has not led to significant
- 20        increases in *Bt* protein expression (Vaeck et al., 1989; Barton et al., 1987). It is therefore contemplated that the most advantageous *Bt* genes for use in the transformation protocols disclosed herein will be those in which the coding sequence has been modified to effect increased expression in plants, and more particularly, those in which maize preferred codons have been used. Examples of such modified *Bt* toxin genes include the variant *Bt* CryIA(b)
- 25        gene termed Iab6 (Perlak et al., 1991) and the synthetic CryIA(c) genes termed 1800a and 1800b.

- Protease inhibitors may also provide insect resistance (Johnson et al., 1989), and will thus have utility in plant transformation. The use of a protease inhibitor II gene, *pinII*, from tomato or potato is envisioned to be particularly useful. Even more advantageous is the use of
- 30        a *pinII* gene in combination with a *Bt* toxin gene, the combined effect of which has been discovered by the present inventors to produce synergistic insecticidal activity. Other genes which encode inhibitors of the insects' digestive system, or those that encode enzymes or co-

factors that facilitate the production of inhibitors, may also be useful. This group may be exemplified by oryzacystatin and amylase inhibitors, such as those from wheat and barley.

Also, genes encoding lectins may confer additional or alternative insecticide properties. Lectins (originally termed phytohemagglutinins) are multivalent carbohydrate-binding proteins which have the ability to agglutinate red blood cells from a range of species. Lectins have been identified recently as insecticidal agents with activity against weevils, ECB and rootworm (Murdock et al., 1990; Czapla and Lang, 1990). Lectin genes contemplated to be useful include, for example, barley and wheat germ agglutinin (WGA) and rice lectins (Gatehouse et al., 1984), with WGA being preferred.

Genes controlling the production of large or small polypeptides active against insects when introduced into the insect pests, such as, e.g., lytic peptides, peptide hormones and toxins and venoms, form another aspect of the invention. For example, it is contemplated that the expression of juvenile hormone esterase, directed towards specific insect pests, may also result in insecticidal activity, or perhaps cause cessation of metamorphosis (Hammock et al., 1990).

Transgenic plants expressing genes which encode enzymes that affect the integrity of the insect cuticle form yet another aspect of the invention. Such genes include those encoding, e.g., chitinase, proteases, lipases and also genes for the production of nikkomycin, a compound that inhibits chitin synthesis, the introduction of any of which is contemplated to produce insect resistant maize plants. Genes that code for activities that affect insect molting, such those affecting the production of ecdysteroid UDP-glucosyl transferase, also fall within the scope of the useful transgenes of the present invention.

Genes that code for enzymes that facilitate the production of compounds that reduce the nutritional quality of the host plant to insect pests are also encompassed by the present invention. It may be possible, for instance, to confer insecticidal activity on a plant by altering its sterol composition. Sterols are obtained by insects from their diet and are used for hormone synthesis and membrane stability. Therefore alterations in plant sterol composition by expression of novel genes, e.g., those that directly promote the production of undesirable sterols or those that convert desirable sterols into undesirable forms, could have a negative effect on insect growth and/or development and hence endow the plant with insecticidal activity. Lipxygenases are naturally occurring plant enzymes that have been shown to exhibit anti-nutritional effects on insects and to reduce the nutritional quality of their diet. Therefore, further embodiments of the invention concern transgenic plants with enhanced lipxygenase activity which may be resistant to insect feeding.

The present invention also provides methods and compositions by which to achieve qualitative or quantitative changes in plant secondary metabolites. One example concerns transforming plants to produce DIMBOA which, it is contemplated, will confer resistance to European corn borer, rootworm and several other maize insect pests. Candidate genes that are particularly considered for use in this regard include those genes at the *bx* locus known to be involved in the synthetic DIMBOA pathway (Dunn et al., 1981). The introduction of genes that can regulate the production of maysin, and genes involved in the production of dhurrin in sorghum, is also contemplated to be of use in facilitating resistance to earworm and rootworm, respectively.

*Tripsacum dactyloides* is a species of grass that is resistant to certain insects, including corn root worm. It is anticipated that genes encoding proteins that are toxic to insects or are involved in the biosynthesis of compounds toxic to insects will be isolated from *Tripsacum* and that these novel genes will be useful in conferring resistance to insects. It is known that the basis of insect resistance in *Tripsacum* is genetic, because said resistance has been transferred to *Zea mays* via sexual crosses (Branson and Guss, 1972).

Further genes encoding proteins characterized as having potential insecticidal activity may also be used as transgenes in accordance herewith. Such genes include, for example, the cowpea trypsin inhibitor (CpTI; Hilder et al., 1987) which may be used as a rootworm deterrent; genes encoding avermectin (Campbell, 1989; Ikeda et al., 1987) which may prove particularly useful as a corn rootworm deterrent; ribosome inactivating protein genes; and even genes that regulate plant structures. Transgenic maize including anti-insect antibody genes and genes that code for enzymes that can covert a non-toxic insecticide (pro-insecticide) applied to the outside of the plant into an insecticide inside the plant are also contemplated.

### 3. Environment or Stress Resistance

Improvement of a plant's ability to tolerate various environmental stresses such as, but not limited to, drought, excess moisture, chilling, freezing, high temperature, salt, and oxidative stress, can also be effected through expression of heterologous, or overexpression of homologous genes. Benefits may be realized in terms of increased resistance to freezing temperatures through the introduction of an "antifreeze" protein such as that of the Winter Flounder (Cutler et al., 1989) or synthetic gene derivatives thereof. Improved chilling tolerance may also be conferred through increased expression of glycerol-3-phosphate acetyltransferase in chloroplasts (Murata et al., 1992; Wolter et al., 1992). Resistance to oxidative stress (often exacerbated by conditions such as chilling temperatures in combination

with high light intensities) can be conferred by expression of superoxide dismutase (Gupta et al., 1993), and may be improved by glutathione reductase (Bowler et al., 1992). Such strategies may allow for tolerance to freezing in newly emerged fields as well as extending later maturity higher yielding varieties to earlier relative maturity zones.

5           Expression of novel genes that favorably effect plant water content, total water potential, osmotic potential, and turgor can enhance the ability of the plant to tolerate drought. As used herein, the terms "drought resistance" and "drought tolerance" are used to refer to a plants increased resistance or tolerance to stress induced by a reduction in water availability, as compared to normal circumstances, and the ability of the plant to function and survive in  
10 lower-water environments, and perform in a relatively superior manner. In this aspect of the invention it is proposed, for example, that the expression of a gene encoding the biosynthesis of osmotically-active solutes can impart protection against drought. Within this class of genes are DNAs encoding mannitol dehydrogenase (Lee and Saier, 1982) and trehalose-6-phosphate synthase (Kaasen et al., 1992). Through the subsequent action of native phosphatases in the  
15 cell or by the introduction and coexpression of a specific phosphatase, these introduced genes will result in the accumulation of either mannitol or trehalose, respectively, both of which have been well documented as protective compounds able to mitigate the effects of stress. Mannitol accumulation in transgenic tobacco has been verified and preliminary results indicate that plants expressing high levels of this metabolite are able to tolerate an applied osmotic stress  
20 (Tarczynski et al., cited supra (1992), 1993).

          Similarly, the efficacy of other metabolites in protecting either enzyme function (e.g. alanopine or propionic acid) or membrane integrity (e.g., alanopine) has been documented (Loomis et al., 1989), and therefore expression of gene encoding the biosynthesis of these compounds can confer drought resistance in a manner similar to or complimentary to mannitol.  
25 Other examples of naturally occurring metabolites that are osmotically active and/or provide some direct protective effect during drought and/or desiccation include sugars and sugar derivatives such as fructose, erythritol (Coxson et al., 1992), sorbitol, dulcitol (Karsten et al., 1992), glucosylglycerol (Reed et al., 1984; Erdmann et al., 1992), sucrose, stachyose (Koster and Leopold, 1988; Blackman et al., 1992), ononitol and pinitol (Vernon and Bohnert, 1992),  
30 and raffinose (Bernal-Lugo and Leopold, 1992). Other osmotically active solutes which are not sugars include, but are not limited to, proline and glycine-betaine (Wyn-Jones and Storey, 1981). Continued canopy growth and increased reproductive fitness during times of stress can be augmented by introduction and expression of genes such as those controlling the

osmotically active compounds discussed above and other such compounds, as represented in one exemplary embodiment by the enzyme myoinositol 0-methyltransferase.

It is contemplated that the expression of specific proteins may also increase drought tolerance. Three classes of Late Embryogenic Proteins have been assigned based on structural similarities (see Dure et al., 1989). All three classes of these proteins have been demonstrated in maturing (i.e., desiccating) seeds. Within these 3 types of proteins, the Type-II (dehydrin-type) have generally been implicated in drought and/or desiccation tolerance in vegetative plant parts (i.e. Mundy and Chua, 1988; Piatkowski et al., 1990; Yamaguchi-Shinozaki et al., 1992). Recently, expression of a Type-III LEA (HVA-1) in tobacco was found to influence plant height, maturity and drought tolerance (Fitzpatrick, 1993). Expression of structural genes from all three groups may therefore confer drought tolerance. Other types of proteins induced during water stress include thiol proteases, aldolases and transmembrane transporters (Guerrero et al., 1990), which may confer various protective and/or repair-type functions during drought stress. The expression of a gene that effects lipid biosynthesis and hence membrane composition can also be useful in conferring drought resistance on the plant.

Many genes that improve drought resistance have complementary modes of action. Thus, combinations of these genes might have additive and/or synergistic effects in improving drought resistance in maize. Many of these genes also improve freezing tolerance (or resistance); the physical stresses incurred during freezing and drought are similar in nature and may be mitigated in similar fashion. Benefit may be conferred via constitutive expression of these genes, but the preferred means of expressing these novel genes may be through the use of a turgor-induced promoter (such as the promoters for the turgor-induced genes described in Guerrero et al. 1990 and Shagan et al., 1993). Spatial and temporal expression patterns of these genes may enable maize to better withstand stress.

Expression of genes that are involved with specific morphological traits that allow for increased water extractions from drying soil would be of benefit. For example, introduction and expression of genes that alter root characteristics may enhance water uptake. Expression of genes that enhance reproductive fitness during times of stress would be of significant value. For example, expression of DNAs that improve the synchrony of pollen shed and receptiveness of the female flower parts, i.e., silks, would be of benefit. In addition, expression of genes that minimize kernel abortion during times of stress would increase the amount of grain to be harvested and hence be of value. Regulation of cytokinin levels in monocots, such as maize, by introduction and expression of an isopentenyl transferase gene with appropriate regulatory

sequences can improve monocot stress resistance and yield (Gan et al., Science, 270:1986 (1995)).

Given the overall role of water in determining yield, it is contemplated that enabling plants to utilize water more efficiently, through the introduction and expression of novel genes, will improve overall performance even when soil water availability is not limiting. By introducing genes that improve the ability of plants to maximize water usage across a full range of stresses relating to water availability, yield stability or consistency of yield performance may be realized.

#### 4. Disease Resistance

It is proposed that increased resistance to diseases may be realized through introduction of genes into plants period. It is possible to produce resistance to diseases caused by viruses, bacteria, fungi, root pathogens, insects and nematodes. It is also contemplated that control of mycotoxin producing organisms may be realized through expression of introduced genes.

Resistance to viruses may be produced through expression of novel genes. For example, it has been demonstrated that expression of a viral coat protein in a transgenic plant can impart resistance to infection of the plant by that virus and perhaps other closely related viruses (Cuozzo et al., 1988, Hemenway et al., 1988, Abel et al., 1986). It is contemplated that expression of antisense genes targeted at essential viral functions may impart resistance to said virus. For example, an antisense gene targeted at the gene responsible for replication of viral nucleic acid may inhibit said replication and lead to resistance to the virus. It is believed that interference with other viral functions through the use of antisense genes may also increase resistance to viruses. Further it is proposed that it may be possible to achieve resistance to viruses through other approaches, including, but not limited to the use of satellite viruses.

It is proposed that increased resistance to diseases caused by bacteria and fungi may be realized through introduction of novel genes. It is contemplated that genes encoding so-called "peptide antibiotics," pathogenesis related (PR) proteins, toxin resistance, and proteins affecting host-pathogen interactions such as morphological characteristics will be useful. Peptide antibiotics are polypeptide sequences which are inhibitory to growth of bacteria and other microorganisms. For example, the classes of peptides referred to as cecropins and magainins inhibit growth of many species of bacteria and fungi. It is proposed that expression of PR proteins in plants may be useful in conferring resistance to bacterial disease. These genes are induced following pathogen attack on a host plant and have been divided into at least five classes of proteins (Bol et al., 1990). Included amongst the PR proteins are beta-1,3-

glucanases, chitinases, and osmotin and other proteins that are believed to function in plant resistance to disease organisms. Other genes have been identified that have antifungal properties, e.g., UDA (stinging nettle lectin) and hevein (Broakgert et al., 1989; Barkai-Golan et al., 1978). It is known that certain plant diseases are caused by the production of  
5 phytotoxins. Resistance to these diseases could be achieved through expression of a novel gene that encodes an enzyme capable of degrading or otherwise inactivating the phytotoxin. Expression novel genes that alter the interactions between the host plant and pathogen may be useful in reducing the ability the disease organism to invade the tissues of the host plant, e.g., an increase in the waxiness of the leaf cuticle or other morphological characteristics.

10 Plant parasitic nematodes are a cause of disease in many plants. It is proposed that it would be possible to make the plant resistant to these organisms through the expression of novel genes. It is anticipated that control of nematode infestations would be accomplished by altering the ability of the nematode to recognize or attach to a host plant and/or enabling the plant to produce nematocidal compounds, including but not limited to proteins.

#### 15 5. Mycotoxin Reduction/Elimination

Production of mycotoxins, including aflatoxin and fumonisin, by fungi associated with plants is a significant factor in rendering the grain not useful. These fungal organisms do not cause disease symptoms and/or interfere with the growth of the plant, but they produce chemicals (mycotoxins) that are toxic to animals. Inhibition of the growth of these fungi  
20 would reduce the synthesis of these toxic substances and, therefore, reduce grain losses due to mycotoxin contamination. Novel genes may be introduced into plants that would inhibit synthesis of the mycotoxin without interfering with fungal growth. Expression of a novel gene which encodes an enzyme capable of rendering the mycotoxin nontoxic would be useful in order to achieve reduced mycotoxin contamination of grain. The result of any of the above  
25 mechanisms would be a reduced presence of mycotoxins on grain.

#### 6. Grain Composition or Quality

Genes may be introduced into plants, particularly commercially important cereals such as maize, wheat or rice, to improve the grain for which the cereal is primarily grown. A wide range of novel transgenic plants produced in this manner may be envisioned depending on the  
30 particular end use of the grain.

For example, the largest use of maize grain is for feed or food. Introduction of genes that alter the composition of the grain may greatly enhance the feed or food value. The primary components of maize grain are starch, protein, and oil. Each of these primary



components of maize grain may be improved by altering its level or composition. Several examples may be mentioned for illustrative purposes but in no way provide an exhaustive list of possibilities.

The protein of many cereal grains is suboptimal for feed and food purposes especially when fed to pigs, poultry, and humans. The protein is deficient in several amino acids that are essential in the diet of these species, requiring the addition of supplements to the grain. Limiting essential amino acids may include lysine, methionine, tryptophan, threonine, valine, arginine, and histidine. Some amino acids become limiting only after the grain is supplemented with other inputs for feed formulations. For example, when the grain is supplemented with soybean meal to meet lysine requirements, methionine becomes limiting. The levels of these essential amino acids in seeds and grain may be elevated by mechanisms which include, but are not limited to, the introduction of genes to increase the biosynthesis of the amino acids, decrease the degradation of the amino acids, increase the storage of the amino acids in proteins, or increase transport of the amino acids to the seeds or grain.

One mechanism for increasing the biosynthesis of the amino acids is to introduce genes that deregulate the amino acid biosynthetic pathways such that the plant can no longer adequately control the levels that are produced. This may be done by deregulating or bypassing steps in the amino acid biosynthetic pathway which are normally regulated by levels of the amino acid end product of the pathway. Examples include the introduction of genes that encode deregulated versions of the enzymes aspartokinase or dihydrodipicolinic acid (DHDP)-synthase for increasing lysine and threonine production, and anthranilate synthase for increasing tryptophan production. Reduction of the catabolism of the amino acids may be accomplished by introduction of DNA sequences that reduce or eliminate the expression of genes encoding enzymes that catalyse steps in the catabolic pathways such as the enzyme lysine-ketoglutarate reductase.

The protein composition of the grain may be altered to improve the balance of amino acids in a variety of ways including elevating expression of native proteins, decreasing expression of those with poor composition, changing the composition of native proteins, or introducing genes encoding entirely new proteins possessing superior composition. DNA may be introduced that decreases the expression of members of the zein family of storage proteins. This DNA may encode ribozymes or antisense sequences directed to impairing expression of zein proteins or expression of regulators of zein expression such as the opaque-2 gene product. The protein composition of the grain may be modified through the phenomenon of

cosuppression, i.e., inhibition of expression of an endogenous gene through the expression of an identical structural gene or gene fragment introduced through transformation (Goring et al., 1991). Additionally, the introduced DNA may encode enzymes which degrade seines. The decreases in zein expression that are achieved may be accompanied by increases in proteins  
5 with more desirable amino acid composition or increases in other major seed constituents such as starch. Alternatively, a chimeric gene may be introduced that comprises a coding sequence for a native protein of adequate amino acid composition such as for one of the globulin proteins or 10 kD zein of maize and a promoter or other regulatory sequence designed to elevate expression of said protein. The coding sequence of said gene may include additional or  
10 replacement codons for essential amino acids. Further, a coding sequence obtained from another species, or, a partially or completely synthetic sequence encoding a completely unique peptide sequence designed to enhance the amino acid composition of the seed may be employed.

The introduction of genes that alter the oil content of the grain may be of value.  
15 Increases in oil content may result in increases in metabolizable energy content and density of the seeds for uses in feed and food. The introduced genes may encode enzymes that remove or reduce rate-limitations or regulated steps in fatty acid or lipid biosynthesis. Such genes may include, but are not limited to, those that encode acetyl-CoA carboxylase, ACP-acyltransferase, beta-ketoacyl-ACP synthase, plus other well known fatty acid biosynthetic activities. Other  
20 possibilities are genes that encode proteins that do not possess enzymatic activity such as acyl carrier protein. Additional examples include 2-acetyltransferase, oleosin pyruvate dehydrogenase complex, acetyl CoA synthetase, ATP citrate lyase, ADP-glucose pyrophosphorylase and genes of the carnitine-CoA- acetyl-CoA shuttles. It is anticipated that expression of genes related to oil biosynthesis will be targeted to the plastid, using a plastid  
25 transit peptide sequence and preferably expressed in the seed embryo. Genes may be introduced that alter the balance of fatty acids present in the oil providing a more healthful or nutritive feedstuff. The introduced DNA may also encode sequences that block expression of enzymes involved in fatty acid biosynthesis, altering the proportions of fatty acids present in the grain such as described below.

30 Genes may be introduced that enhance the nutritive value of the starch component of the grain, for example by increasing the degree of branching, resulting in improved utilization of the starch in cows by delaying its metabolism.

Besides affecting the major constituents of the grain, genes may be introduced that affect a variety of other nutritive, processing, or other quality aspects of the grain as used for feed or food. For example, pigmentation of the grain may be increased or decreased. Enhancement and stability of yellow pigmentation is desirable in some animal feeds and may be achieved by introduction of genes that result in enhanced production of xanthophylls and carotenes by eliminating rate-limiting steps in their production. Such genes may encode altered forms of the enzymes phytoene synthase, phytoene desaturase, or lycopene synthase. Alternatively, unpigmented white corn is desirable for production of many food products and may be produced by the introduction of DNA which blocks or eliminates steps in pigment production pathways.

Feed or food comprising some cereal grains possesses insufficient quantities of vitamins and must be supplemented to provide adequate nutritive value. Introduction of genes that enhance vitamin biosynthesis in seeds may be envisioned including, for example, vitamins A, E, B<sub>12</sub>, choline, and the like. For example, maize grain also does not possess sufficient mineral content for optimal nutritive value. Genes that affect the accumulation or availability of compounds containing phosphorus, sulfur, calcium, manganese, zinc, and iron among others would be valuable. An example may be the introduction of a gene that reduced phytic acid production or encoded the enzyme phytase which enhances phytic acid breakdown. These genes would increase levels of available phosphate in the diet, reducing the need for supplementation with mineral phosphate.

Numerous other examples of improvement of cereals for feed and food purposes might be described. The improvements may not even necessarily involve the grain, but may, for example, improve the value of the grain for silage. Introduction of DNA to accomplish this might include sequences that alter lignin production such as those that result in the "brown midrib" phenotype associated with superior feed value for cattle.

In addition to direct improvements in feed or food value, genes may also be introduced which improve the processing of grain and improve the value of the products resulting from the processing. The primary method of processing certain grains such as maize is via wetmilling. Maize may be improved through the expression of novel genes that increase the efficiency and reduce the cost of processing such as by decreasing steeping time.

Improving the value of wetmilling products may include altering the quantity or quality of starch, oil, corn gluten meal, or the components of corn gluten feed. Elevation of starch may be achieved through the identification and elimination of rate limiting steps in starch

biosynthesis or by decreasing levels of the other components of the grain resulting in proportional increases in starch. An example of the former may be the introduction of genes encoding ADP-glucose pyrophosphorylase enzymes with altered regulatory activity or which are expressed at higher level. Examples of the latter may include selective inhibitors of, for  
5 example, protein or oil biosynthesis expressed during later stages of kernel development.

The properties of starch may be beneficially altered by changing the ratio of amylose to amylopectin, the size of the starch molecules, or their branching pattern. Through these changes a broad range of properties may be modified which include, but are not limited to, changes in gelatinization temperature, heat of gelatinization, clarity of films and pastes,  
10 Theological properties, and the like. To accomplish these changes in properties, genes that encode granule-bound or soluble starch synthase activity or branching enzyme activity may be introduced alone or combination. DNA such as antisense constructs may also be used to decrease levels of endogenous activity of these enzymes. The introduced genes or constructs may possess regulatory sequences that time their expression to specific intervals in starch  
15 biosynthesis and starch granule development. Furthermore, it may be advisable to introduce and express genes that result in the *in vivo* derivatization, or other modification, of the glucose moieties of the starch molecule. The covalent attachment of any molecule may be envisioned, limited only by the existence of enzymes that catalyze the derivatizations and the accessibility of appropriate substrates in the starch granule. Examples of important derivations may include  
20 the addition of functional groups such as amines, carboxyls, or phosphate groups which provide sites for subsequent *in vitro* derivatizations or affect starch properties through the introduction of ionic charges. Examples of other modifications may include direct changes of the glucose units such as loss of hydroxyl groups or their oxidation to aldehyde or carboxyl groups.

25 Oil is another product of wetmilling of corn and other grains, the value of which may be improved by introduction and expression of genes. The quantity of oil that can be extracted by wetmilling may be elevated by approaches as described for feed and food above. Oil properties may also be altered to improve its performance in the production and use of cooking oil, shortenings, lubricants or other oil-derived products or improvement of its health attributes  
30 when used in the food-related applications. Novel fatty acids may also be synthesized which upon extraction can serve as starting materials for chemical syntheses. The changes in oil properties may be achieved by altering the type, level, or lipid arrangement of the fatty acids present in the oil. This in turn may be accomplished by the addition of genes that encode

enzymes that catalyze the synthesis of novel fatty acids and the lipids possessing them or by increasing levels of native fatty acids while possibly reducing levels of precursors.

Alternatively DNA sequences may be introduced which slow or block steps in fatty acid biosynthesis resulting in the increase in precursor fatty acid intermediates. Genes that might be added include desaturases, epoxidases, hydratases, dehydratases, and other enzymes that catalyze reactions involving fatty acid intermediates. Representative examples of catalytic steps that might be blocked include the desaturations from stearic to oleic acid and oleic to linolenic acid resulting in the respective accumulations of stearic and oleic acids.

Improvements in the other major cereal wetmilling products, gluten meal and gluten feed, may also be achieved by the introduction of genes to obtain novel plants. Representative possibilities include but are not limited to those described above for improvement of food and feed value.

In addition it may further be considered that the plant be used for the production or manufacturing of useful biological compounds that were either not produced at all, or not produced at the same level, in the plant previously. The novel plants producing these compounds are made possible by the introduction and expression of genes by transformation methods. The possibilities include, but are not limited to, any biological compound which is presently produced by any organism such as proteins, nucleic acids, primary and intermediary metabolites, carbohydrate polymers, etc. The compounds may be produced by the plant, extracted upon harvest and/or processing, and used for any presently recognized useful purpose such as pharmaceuticals, fragrances, industrial enzymes to name a few.

Further possibilities to exemplify the range of grain traits or properties potentially encoded by introduced genes in transgenic plants include grain with less breakage susceptibility for export purposes or larger grit size when processed by dry milling through introduction of genes that enhance gamma-zein synthesis, popcorn with improved popping quality and expansion volume through genes that increase pericarp thickness, corn with whiter grain for food uses through introduction of genes that effectively block expression of enzymes involved in pigment production pathways, and improved quality of alcoholic beverages or sweet corn through introduction of genes which affect flavor such as the shrunken gene (encoding sucrose synthase) for sweet corn.

#### 7. Plant Agronomic Characteristics

Two of the factors determining where plants can be grown are the average daily temperature during the growing season and the length of time between frosts. Within the areas

where it is possible to grow a particular plant, there are varying limitations on the maximal time it is allowed to grow to maturity and be harvested. The plant to be grown in a particular area is selected for its ability to mature and dry down to harvestable moisture content within the required period of time with maximum possible yield. Therefore, plant of varying

5 maturities are developed for different growing locations. Apart from the need to dry down sufficiently to permit harvest is the desirability of having maximal drying take place in the field to minimize the amount of energy required for additional drying post-harvest. Also the more readily the grain can dry down, the more time there is available for growth and kernel fill. Genes that influence maturity and/or dry down can be identified and introduced into plant

10 lines using transformation techniques to create new varieties adapted to different growing locations or the same growing location but having improved yield to moisture ratio at harvest. Expression of genes that are involved in regulation of plant development may be especially useful, e.g., the liguleless and rough sheath genes that have been identified in plants.

Genes may be introduced into plants that would improve standability and other plant

15 growth characteristics. For example, expression of novel genes which confer stronger stalks, improved root systems, or prevent or reduce ear droppage would be of great value to the corn farmer. Introduction and expression of genes that increase the total amount of photoassimilate available by, for example, increasing light distribution and/or interception would be advantageous. In addition the expression of genes that increase the efficiency of

20 photosynthesis and/or the leaf canopy would further increase gains in productivity. Such approaches would allow for increased plant populations in the field.

Delay of late season vegetative senescence would increase the flow of assimilate into the grain and thus increase yield. Overexpression of genes within plants that are associated with "stay green" or the expression of any gene that delays senescence would achieve be

25 advantageous. For example, a non-yellowing mutant has been identified in *Festuca pratensis* (Davies et al., 1990). Expression of this gene as well as others may prevent premature breakdown of chlorophyll and thus maintain canopy function.

#### 8. Nutrient Utilization

The ability to utilize available nutrients and minerals may be a limiting factor in growth

30 of many plants. It is proposed that it would be possible to alter nutrient uptake, tolerate pH extremes, mobilization through the plant, storage pools, and availability for metabolic activities by the introduction of novel genes. These modifications would allow a plant to more efficiently utilize available nutrients. It is contemplated that an increase in the activity of, for

example, an enzyme that is normally present in the plant and involved in nutrient utilization would increase the availability of a nutrient. An example of such an enzyme would be phytase. It is also contemplated that expression of a novel gene may make a nutrient source available that was previously not accessible, e.g., an enzyme that releases a component of nutrient value from a more complex molecule, perhaps a macromolecule.

#### 9. Male Sterility

Male sterility is useful in the production of hybrid seed. It is proposed that male sterility may be produced through expression of novel genes. For example, it has been shown that expression of genes that encode proteins that interfere with development of the male inflorescence and/or gametophyte result in male sterility. Chimeric ribonuclease genes that express in the anthers of transgenic tobacco and oilseed rape have been demonstrated to lead to male sterility (Mariani et al, 1990).

For example, a number of mutations were discovered in maize that confer cytoplasmic male sterility. One mutation in particular, referred to as T cytoplasm, also correlates with sensitivity to Southern corn leaf blight. A DNA sequence, designated TURF-13 (Levings, 1990), was identified that correlates with T cytoplasm. It would be possible through the introduction of TURF-13 via transformation to separate male sterility from disease sensitivity. As it is necessary to be able to restore male fertility for breeding purposes and for grain production, it is proposed that genes encoding restoration of male fertility may also be introduced.

#### 10. Negative Selectable Markers

Introduction of genes encoding traits that can be selected against may be useful for eliminating undesirable linked genes. When two or more genes are introduced together by cotransformation, the genes will be linked together on the host chromosome. For example, a gene encoding a *Bt* gene that confers insect resistance on the plant may be introduced into a plant together with a *bar* gene that is useful as a selectable marker and confers resistance to the herbicide Ignite® on the plant. However, it may not be desirable to have an insect resistant plant that is also resistant to the herbicide Ignite®. It is proposed that one could also introduce an antisense *bar* gene that is expressed in those tissues where one does not want expression of the *bar* gene, e.g., in whole plant parts. Hence, although the *bar* gene is expressed and is useful as a selectable marker, it is not useful to confer herbicide resistance on the whole plant. The *bar* antisense gene is a negative selectable marker.

Negative selection is necessary in order to screen a population of transformants for rare homologous recombinants generated through gene targeting. For example, a homologous recombinant may be identified through the inactivation of a gene that was previously expressed in that cell. The antisense gene to neomycin phosphotransferase II (*nptII*) has been investigated as a negative selectable marker in tobacco (*Nicotiana tabacum*) and *Arabidopsis thaliana* (Xiang and Guerra, 1993). In this example both sense and antisense *nptII* genes are introduced into a plant through transformation and the resultant plants are sensitive to the antibiotic kanamycin. An introduced gene that integrates into the host cell chromosome at the site of the antisense *nptII* gene, and inactivates the antisense gene, will make the plant resistant to kanamycin and other aminoglycoside antibiotics. Therefore, rare site specific recombinants may be identified by screening for antibiotic resistance. Similarly, any gene, native to the plant or introduced through transformation, that when inactivated confers resistance to a compound, may be useful as a negative selectable marker.

It is contemplated that negative selectable markers may also be useful in other ways. One application is to construct transgenic lines in which one could select for transposition to unlinked sites. In the process of tagging it is most common for the transposable element to move to a genetically linked site on the same chromosome. A selectable marker for recovery of rare plants in which transposition has occurred to an unlinked locus would be useful. For example, the enzyme cytosine deaminase may be useful for this purpose (Stouggard, 1993). In the presence of this enzyme the compound 5-fluorocytosine is converted to 5-fluoruracil which is toxic to plant and animal cells. If a transposable element is linked to the gene for the enzyme cytosine deaminase, one may select for transposition to unlinked sites by selecting for transposition events in which the resultant plant is now resistant to 5-fluorocytosine. The parental plants and plants containing transpositions to linked sites will remain sensitive to 5-fluorocytosine. Resistance to 5-fluorocytosine is due to loss of the cytosine deaminase gene through genetic segregation of the transposable element and the cytosine deaminase gene. Other genes that encode proteins that render the plant sensitive to a certain compound will also be useful in this context. For example, T-DNA gene 2 from *Agrobacterium tumefaciens* encodes a protein that catalyzes the conversion of alpha-naphthalene acetamide (NAM) to alpha-naphthalene acetic acid (NAA) renders plant cells sensitive to high concentrations of NAM (Depicker et al., 1988).

It is also contemplated that negative selectable markers may be useful in the construction of transposon tagging lines. For example, by marking an autonomous



transposable element such as Ac, Master Mu, or En/Spn with a negative selectable marker, one could select for transformants in which the autonomous element is not stably integrated into the genome. This would be desirable, for example, when transient expression of the autonomous element is desired to activate in *trans* the transposition of a defective transposable element, such as Ds, but stable integration of the autonomous element is not desired. The presence of the autonomous element may not be desired in order to stabilize the defective element, i.e., prevent it from further transposing. However, it is proposed that if stable integration of an autonomous transposable element is desired in a plant the presence of a negative selectable marker may make it possible to eliminate the autonomous element during the breeding process.

#### 11. Non-Protein-Expressing Sequences

##### a. RNA-Expressing

DNA may be introduced into plants for the purpose of expressing RNA transcripts that function to affect plant phenotype yet are not translated into protein. Two examples are antisense RNA and RNA with ribozyme activity. Both may serve possible functions in reducing or eliminating expression of native or introduced plant genes.

Genes may be constructed or isolated, which when transcribed, produce antisense RNA that is complementary to all or part(s) of a targeted messenger RNA(s). The antisense RNA reduces production of the polypeptide product of the messenger RNA. The polypeptide product may be any protein encoded by the plant genome. The aforementioned genes will be referred to as antisense genes. An antisense gene may thus be introduced into a plant by transformation methods to produce a novel transgenic plant with reduced expression of a selected protein of interest. For example, the protein may be an enzyme that catalyzes a reaction in the plant. Reduction of the enzyme activity may reduce or eliminate products of the reaction which include any enzymatically synthesized compound in the plant such as fatty acids, amino acids, carbohydrates, nucleic acids and the like. Alternatively, the protein may be a storage protein, such as a zein, or a structural protein, the decreased expression of which may lead to changes in seed amino acid composition or plant morphological changes respectively. The possibilities cited above are provided only by way of example and do not represent the full range of applications.

Genes may also be constructed or isolated, which when transcribed produce RNA enzymes, or ribozymes, which can act as endoribonucleases and catalyze the cleavage of RNA molecules with selected sequences. The cleavage of selected messenger RNA's can result in

the reduced production of their encoded polypeptide products. These genes may be used to prepare novel transgenic plants which possess them. The transgenic plants may possess reduced levels of polypeptides including but not limited to the polypeptides cited above that may be affected by antisense RNA.

- 5 It is also possible that genes may be introduced to produce novel transgenic plants which have reduced expression of a native gene product by a mechanism of cosuppression. It has been demonstrated in tobacco, tomato, and petunia (Goring et al, 1991; Smith et al., 1990; Napoli et al., 1990; van der Krol et al., 1990) that expression of the sense transcript of a native gene will reduce or eliminate expression of the native gene in a manner similar to that  
10 observed for antisense genes. The introduced gene may encode all or part of the targeted native protein but its translation may not be required for reduction of levels of that native protein.

#### b. Non-RNA-Expressing

- For example, DNA elements including those of transposable elements such as Ds, Ac,  
15 or Mu, may be inserted into a gene and cause mutations. These DNA elements may be inserted in order to inactivate (or activate) a gene and thereby "tag" a particular trait. In this instance the transposable element does not cause instability of the tagged mutation, because the utility of the element does not depend on its ability to move in the genome. Once a desired trait is tagged, the introduced DNA sequence may be used to clone the corresponding gene,  
20 e.g., using the introduced DNA sequence as a PCR primer together with PCR gene cloning techniques (Shapiro, 1983; Dellaporta et al., 1988). Once identified, the entire gene(s) for the particular trait, including control or regulatory regions where desired may be isolated, cloned and manipulated as desired. The utility of DNA elements introduced into an organism for purposed of gene tagging is independent of the DNA sequence and does not depend on any  
25 biological activity of the DNA sequence, i.e., transcription into RNA or translation into protein. The sole function of the DNA element is to disrupt the DNA sequence of a gene.

- It is contemplated that unexpressed DNA sequences, including novel synthetic sequences could be introduced into cells as proprietary "labels" of those cells and plants and seeds thereof. It would not be necessary for a label DNA element to disrupt the function of a  
30 gene endogenous to the host organism, as the sole function of this DNA would be to identify the origin of the organism. For example, one could introduce a unique DNA sequence into a plant and this DNA element would identify all cells, plants, and progeny of these cells as having arisen from that labeled source. It is proposed that inclusion of label DNAs would

enable one to distinguish proprietary germplasm or germplasm derived from such, from unlabelled germplasm.

Another possible element which may be introduced is a matrix attachment region element (MAR), such as the chicken lysozyme A element (Stief et al., 1989), which can be  
5 positioned around an expressible gene of interest to effect an increase in overall expression of the gene and diminish position dependant effects upon incorporation into the plant genome (Stief et al., 1989; Phi-Van et al., 1990).

### **III. Transformed (Transgenic) Plants of the Invention and Methods of Preparation**

Plant species may be transformed with the DNA construct of the present invention by  
10 the DNA-mediated transformation of plant cell protoplasts and subsequent regeneration of the plant from the transformed protoplasts in accordance with procedures well known in the art.

Any plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a vector of the present invention. The term "organogenesis," as used herein, means a process by which shoots and roots are developed  
15 sequentially from meristematic centers; the term "embryogenesis," as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or gametes. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons,  
20 hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g., apical meristems, axillary buds, and root meristems), and induced meristem tissue (e.g., cotyledon meristem and utilane meristem).

Plants of the present invention may take a variety of forms. The plants may be chimeras of transformed cells and non-transformed cells; the plants may be clonal  
25 transformants (e.g., all cells transformed to contain the expression cassette); the plants may comprise grafts of transformed and untransformed tissues (e.g., a transformed root stock grafted to an untransformed scion in citrus species). The transformed plants may be propagated by a variety of means, such as by clonal propagation or classical breeding techniques. For example, first generation (or T1) transformed plants may be selfed to give  
30 homozygous second generation (or T2) transformed plants, and the T2 plants further propagated through classical breeding techniques. A dominant selectable marker (such as npt II) can be associated with the expression cassette to assist in breeding.

Thus, the present invention provides a transformed (transgenic) plant cell, *in planta* or *ex planta*, including a transformed plastid or other organelle, e.g., nucleus, mitochondria or chloroplast. The present invention may be used for transformation of any plant species, including, but not limited to, cells from corn (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*)), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Cofea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa* spp.), avocado (*Persea utilane*), fig (*Ficus casica*), guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica papaya*), cashew (*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, duckweed (*Lemna*), barley, vegetables, ornamentals, and conifers.

Duckweed (*Lemna*, see WO 00/07210) includes members of the family *Lemnaceae*.

There are known four genera and 34 species of duckweed as follows: genus *Lemna* (*L. aequinoctialis*, *L. disperma*, *L. ecuadoriensis*, *L. gibba*, *L. japonica*, *L. minor*, *L. miniscula*, *L. obscura*, *L. perpusilla*, *L. tenera*, *L. trisulca*, *L. turionifera*, *L. valdiviana*); genus *Spirodela* (*S. intermedia*, *S. polyrrhiza*, *S. punctata*); genus *Woffia* (*Wa. Angusta*, *Wa. Arrhiza*, *Wa. Australina*, *Wa. Borealis*, *Wa. Brasiliensis*, *Wa. Columbiana*, *Wa. Elongata*, *Wa. Globosa*, *Wa. Microscopica*, *Wa. Neglecta*) and genus *Wofieldia* (*W1. ultila*, *W1. utilanen*, *W1. gladiata*, *W1. ultila*, *W1. lingulata*, *W1. repunda*, *W1. rotunda*, and *W1. neotropica*). Any other genera or species of *Lemnaceae*, if they exist, are also aspects of the present invention. *Lemna gibba*, *Lemna minor*, and *Lemna miniscula* are preferred, with *Lemna minor* and *Lemna miniscula* being most preferred. *Lemna* species can be classified using the taxonomic scheme described by Landolt, Biosystematic Investigation on the Family of Duckweeds: The family of Lemnaceae – A Monograph Study. Geobotanischen Institut ETH, Stiftung Rubel, Zurich (1986)).

Vegetables within the scope of the invention include tomatoes (*Lycopersicon esculentum*), lettuce (e.g., *Lactuca sativa*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus* spp.), and members of the genus *Cucumis* such as cucumber (*C. sativus*), cantaloupe (*C. cantalupensis*), and musk melon (*C. melo*).

- 5 Ornamentals include azalea (*Rhododendron* spp.), hydrangea (*Macrophylla hydrangea*), hibiscus (*Hibiscus rosasanensis*), roses (*Rosa* spp.), tulips (*Tulipa* spp.), daffodils (*Narcissus* spp.), petunias (*Petunia hybrida*), carnation (*Dianthus caryophyllus*), poinsettia (*Euphorbia pulcherrima*), and chrysanthemum. Conifers that may be employed in practicing the present invention include, for example, pines such as loblolly pine (*Pinus taeda*), slash pine (*Pinus*
- 10 *elliotii*), ponderosa pine (*Pinus ponderosa*), lodgepole pine (*Pinus contorta*), and Monterey pine (*Pinus radiata*), Douglas-fir (*Pseudotsuga menziesii*); Western hemlock (*Tsuga ultilane*); Sitka spruce (*Picea glauca*); redwood (*Sequoia sempervirens*); true firs such as silver fir (*Abies amabilis*) and balsam fir (*Abies balsamea*); and cedars such as Western red cedar (*Thuja plicata*) and Alaska yellow-cedar (*Chamaecyparis nootkatensis*). Leguminous plants
- 15 include beans and peas. Beans include guar, locust bean, fenugreek, soybean, garden beans, cowpea, mungbean, lima bean, fava bean, lentils, chickpea, etc. Legumes include, but are not limited to, *Arachis*, e.g., peanuts, *Vicia*, e.g., crown vetch, hairy vetch, adzuki bean, mung bean, and chickpea, *Lupinus*, e.g., lupine, trifolium, *Phaseolus*, e.g., common bean and lima bean, *Pisum*, e.g., field bean, *Melilotus*, e.g., clover, *Medicago*, e.g., alfalfa, Lotus, e.g., trefoil,
- 20 lens, e.g., lentil, and false indigo. Preferred forage and turf grass for use in the methods of the invention include alfalfa, orchard grass, tall fescue, perennial ryegrass, creeping bent grass, and redtop.

- Papaya, garlic, pea, peach, pepper, petunia, strawberry, sorghum, sweet potato, turnip, safflower, corn, pea, endive, gourd, grape, snap bean, chicory, cotton, tobacco, aubergine, beet,
- 25 buckwheat, broad bean, nectarine, avocado, mango, banana, groundnut, potato, peanut, lettuce, pineapple, spinach, squash, sugarbeet, sugarcane, sweet corn, chrysanthemum.

- Other plants within the scope of the invention include *Acacia*, aneth, artichoke, arugula, blackberry, canola, cilantro, clementines, escarole, eucalyptus, fennel, grapefruit, honey dew, jicama, kiwifruit, lemon, lime, mushroom, nut, okra, orange, parsley, persimmon,
- 30 plantain, pomegranate, poplar, radiata pine, radicchio, Southern pine, sweetgum, tangerine, triticale, vine, yams, apple, pear, quince, cherry, apricot, melon, hemp, buckwheat, grape, raspberry, chenopodium, blueberry, nectarine, peach, plum, strawberry, watermelon, eggplant, pepper, cauliflower, Brassica, e.g., broccoli, cabbage, utilan sprouts, onion, carrot, leek, beet,

broad bean, celery, radish, pumpkin, endive, gourd, garlic, snapbean, spinach, squash, turnip, utilane, and zucchini.

Ornamental plants within the scope of the invention include impatiens, Begonia, Pelargonium, Viola, Cyclamen, Verbena, Vinca, Tagetes, Primula, Saint Paulia, Agertum, 5 Amaranthus, Antihirrhinum, Aquilegia, Cineraria, Clover, Cosmo, Cowpea, Dahlia, Datura, Delphinium, Gerbera, Gladiolus, Gloxinia, Hippeastrum, Mesembryanthemum, Salpiglossos, and Zinnia. Other plants within the scope of the invention are shown in Table 1 (above).

Preferably, transgenic plants of the present invention are crop plants and in particular cereals (for example, corn, alfalfa, sunflower, rice, *Brassica*, canola, soybean, barley, soybean, 10 sugarbeet, cotton, safflower, peanut, sorghum, wheat, millet, tobacco, etc.), and even more preferably corn, rice and soybean.

Transformation of plants can be undertaken with a single DNA molecule or multiple DNA molecules (i.e., co-transformation), and both these techniques are suitable for use with the expression cassettes of the present invention. Numerous transformation vectors are 15 available for plant transformation, and the expression cassettes of this invention can be used in conjunction with any such vectors. The selection of vector will depend upon the preferred transformation technique and the target species for transformation.

A variety of techniques are available and known to those skilled in the art for introduction of constructs into a plant cell host. These techniques generally include 20 transformation with DNA employing *A. tumefaciens* or *A. rhizogenes* as the transforming agent, liposomes, PEG precipitation, electroporation, DNA injection, direct DNA uptake, microprojectile bombardment, particle acceleration, and the like (See, for example, EP 295959 and EP 138341) (see below). However, cells other than plant cells may be transformed with the expression cassettes of the invention. The general descriptions of plant expression vectors 25 and reporter genes, and *Agrobacterium* and *Agrobacterium*-mediated gene transfer, can be found in Gruber et al. (1993).

Expression vectors containing genomic or synthetic fragments can be introduced into protoplasts or into intact tissues or isolated cells. Preferably expression vectors are introduced into intact tissue. General methods of culturing plant tissues are provided for example by 30 Maki et al., (1993); and by Phillips et al. (1988). Preferably, expression vectors are introduced into maize or other plant tissues using a direct gene transfer method such as microprojectile-mediated delivery, DNA injection, electroporation and the like. More preferably expression vectors are introduced into plant tissues using the microprojectile media delivery with the

biolistic device. See, for example, Tomes et al. (1995). The vectors of the invention can not only be used for expression of structural genes but may also be used in exon-trap cloning, or promoter trap procedures to detect differential gene expression in varieties of tissues, (Lindsey et al., 1993; Auch & Reth et al.).

5 It is particularly preferred to use the binary type vectors of Ti and Ri plasmids of *Agrobacterium spp.* Ti-derived vectors transform a wide variety of higher plants, including monocotyledonous and dicotyledonous plants, such as soybean, cotton, rape, tobacco, and rice (Pacciotti et al., 1985; Byrne et al., 1987; Sukhapinda et al., 1987; Park et al., 1985; Hiei et al., 1994). The use of T-DNA to transform plant cells has received extensive study and is amply  
10 described (EP 120516; Hoekema, 1985; Knauf, et al., 1983; and An et al., 1985). For introduction into plants, the chimeric genes of the invention can be inserted into binary vectors as described in the examples.

Other transformation methods are available to those skilled in the art, such as direct uptake of foreign DNA constructs (see EP 295959), techniques of electroporation (Fromm et al., 1986) or high velocity ballistic bombardment with metal particles coated with the nucleic  
15 acid constructs (Kline et al., 1987, and U.S. Patent No. 4,945,050). Once transformed, the cells can be regenerated by those skilled in the art. Of particular relevance are the recently described methods to transform foreign genes into commercially important crops, such as rapeseed (De Block et al., 1989), sunflower (Everett et al., 1987), soybean (McCabe et al.,  
20 1988; Hinchee et al., 1988; Chee et al., 1989; Christou et al., 1989; EP 301749), rice (Hiei et al., 1994), and corn (Gordon Kamm et al., 1990; Fromm et al., 1990).

Those skilled in the art will appreciate that the choice of method might depend on the type of plant, i.e., monocotyledonous or dicotyledonous, targeted for transformation. Suitable methods of transforming plant cells include, but are not limited to, microinjection (Crossway  
25 et al., 1986), electroporation (Riggs et al., 1986), *Agrobacterium*-mediated transformation (Hinchee et al., 1988), direct gene transfer (Paszkowski et al., 1984), and ballistic particle acceleration using devices available from Agracetus, Inc., Madison, Wis. And BioRad, Hercules, Calif. (see, for example, Sanford et al., U.S. Pat. No. 4,945,050; and McCabe et al.,  
1988). Also see, Weissinger et al., 1988; Sanford et al., 1987 (onion); Christou et al., 1988  
30 (soybean); McCabe et al., 1988 (soybean); Datta et al., 1990 (rice); Klein et al., 1988 (maize); Klein et al., 1988 (maize); Klein et al., 1988 (maize); Fromm et al., 1990 (maize); and Gordon-Kamm et al., 1990 (maize); Svab et al., 1990 (tobacco chloroplast); Koziel et al., 1993 (maize); Shimamoto et al., 1989 (rice); Christou et al., 1991 (rice); European Patent

Application EP 0 332 581 (orchardgrass and other Pooideae); Vasil et al., 1993 (wheat); Weeks et al., 1993 (wheat). In one embodiment, the protoplast transformation method for maize is employed (European Patent Application EP 0 292 435, U. S. Pat. No. 5,350,689).

In another embodiment, a nucleotide sequence of the present invention is directly  
5 transformed into the plastid genome. Plastid transformation technology is extensively described in U.S. Patent Nos. 5,451,513, 5,545,817, and 5,545,818, in PCT application no. WO 95/16783, and in McBride et al., 1994. The basic technique for chloroplast transformation involves introducing regions of cloned plastid DNA flanking a selectable marker together with  
10 the gene of interest into a suitable target tissue, e.g., using biolistics or protoplast transformation (e.g., calcium chloride or PEG mediated transformation). The 1 to 1.5 kb flanking regions, termed targeting sequences, facilitate orthologous recombination with the plastid genome and thus allow the replacement or modification of specific regions of the plastome. Initially, point mutations in the chloroplast 16S rRNA and *rps12* genes conferring resistance to spectinomycin and/or streptomycin are utilized as selectable markers for  
15 transformation (Svab et al., 1990; Staub et al., 1992). This resulted in stable homoplasmic transformants at a frequency of approximately one per 100 bombardments of target leaves. The presence of cloning sites between these markers allowed creation of a plastid targeting vector for introduction of foreign genes (Staub et al., 1993). Substantial increases in transformation frequency are obtained by replacement of the recessive rRNA or r-protein  
20 antibiotic resistance genes with a dominant selectable marker, the bacterial *aadA* gene encoding the spectinomycin-detoxifying enzyme aminoglycoside-3N-adenyltransferase (Svab et al., 1993). Other selectable markers useful for plastid transformation are known in the art and encompassed within the scope of the invention. Typically, approximately 15-20 cell division cycles following transformation are required to reach a homoplastidic state. Plastid  
25 expression, in which genes are inserted by orthologous recombination into all of the several thousand copies of the circular plastid genome present in each plant cell, takes advantage of the enormous copy number advantage over nuclear-expressed genes to permit expression levels that can readily exceed 10% of the total soluble plant protein. In a preferred  
embodiment, a nucleotide sequence of the present invention is inserted into a plastid targeting  
30 vector and transformed into the plastid genome of a desired plant host. Plants homoplasmic for plastid genomes containing a nucleotide sequence of the present invention are obtained, and are preferentially capable of high expression of the nucleotide sequence.



*Agrobacterium tumefaciens* cells containing a vector comprising an expression cassette of the present invention, wherein the vector comprises a Ti plasmid, are useful in methods of making transformed plants. Plant cells are infected with an *Agrobacterium tumefaciens* as described above to produce a transformed plant cell, and then a plant is regenerated from the transformed plant cell. Numerous *Agrobacterium* vector systems useful in carrying out the present invention are known.

For example, vectors are available for transformation using *Agrobacterium tumefaciens*. These typically carry at least one T-DNA border sequence and include vectors such as pBIN19 (Bevan, 1984). In one preferred embodiment, the expression cassettes of the present invention may be inserted into either of the binary vectors pCIB200 and pCIB2001 for use with *Agrobacterium*. These vector cassettes for *Agrobacterium*-mediated transformation wear constructed in the following manner. PTJS75kan was created by NarI digestion of pTJS75 (Schmidhauser & Helinski, 1985) allowing excision of the tetracycline-resistance gene, followed by insertion of an AccI fragment from pUC4K carrying an NPTII (Messing & Vierra, 1982; Bevan et al., 1983; McBride et al., 1990). XhoI linkers were ligated to the EcoRV fragment of pCIB7 which contains the left and right T-DNA borders, a plant selectable nos/nptII chimeric gene and the pUC polylinker (Rothstein et al., 1987), and the XhoI-digested fragment was cloned into SalI-digested pTJS75kan to create pCIB200 (see also EP 0 332 104, example 19). PCIB200 contains the following unique polylinker restriction sites: EcoRI, SstI, KpnI, BglII, XbaI, and SalI. The plasmid pCIB2001 is a derivative of pCIB200 which was created by the insertion into the polylinker of additional restriction sites. Unique restriction sites in the polylinker of pCIB2001 are EcoRI, SstI, KpnI, BglII, XbaI, SalI, MluI, BclI, AvrII, ApaI, HpaI, and StuI. PCIB2001, in addition to containing these unique restriction sites also has plant and bacterial kanamycin selection, left and right T-DNA borders for *Agrobacterium*-mediated transformation, the RK2-derived trfA function for mobilization between *E. coli* and other hosts, and the OriT and OriV functions also from RK2. The pCIB2001 polylinker is suitable for the cloning of plant expression cassettes containing their own regulatory signals.

An additional vector useful for *Agrobacterium*-mediated transformation is the binary vector pCIB 10, which contains a gene encoding kanamycin resistance for selection in plants, T-DNA right and left border sequences and incorporates sequences from the wide host-range plasmid pRK252 allowing it to replicate in both *E. coli* and *Agrobacterium*. Its construction is described by Rothstein et al., 1987. Various derivatives of pCIB10 have been constructed

which incorporate the gene for hygromycin B phosphotransferase described by Gritz et al., 1983. These derivatives enable selection of transgenic plant cells on hygromycin only (pCIB743), or hygromycin and kanamycin (pCIB715, pCIB717).

5 Methods using either a form of direct gene transfer or *Agrobacterium*-mediated transfer usually, but not necessarily, are undertaken with a selectable marker which may provide resistance to an antibiotic (e.g., kanamycin, hygromycin or methotrexate) or a herbicide (e.g., phosphinothricin). The choice of selectable marker for plant transformation is not, however, critical to the invention.

10 For certain plant species, different antibiotic or herbicide selection markers may be preferred. Selection markers used routinely in transformation include the nptII gene which confers resistance to kanamycin and related antibiotics (Messing & Vierra, 1982; Bevan et al., 1983), the bar gene which confers resistance to the herbicide phosphinothricin (White et al., 1990, Spencer et al., 1990), the hph gene which confers resistance to the antibiotic hygromycin (Blochinger & Diggelmann), and the dhfr gene, which confers resistance to methotrexate  
15 (Bourouis et al., 1983).

One such vector useful for direct gene transfer techniques in combination with selection by the herbicide Basta (or phosphinothricin) is pCIB3064. This vector is based on the plasmid pCIB246, which comprises the CaMV 35S promoter in operational fusion to the *E. coli* GUS gene and the CaMV 35S transcriptional terminator and is described in the PCT  
20 published application WO 93/07278, herein incorporated by reference. One gene useful for conferring resistance to phosphinothricin is the bar gene from *Streptomyces viridochromogenes* (Thompson et al., 1987). This vector is suitable for the cloning of plant expression cassettes containing their own regulatory signals.

An additional transformation vector is pSOG35 which utilizes the *E. coli* gene  
25 dihydrofolate reductase (DHFR) as a selectable marker conferring resistance to methotrexate. PCR was used to amplify the 35S promoter (about 800 bp), intron 6 from the maize Adh1 gene (about 550 bp) and 18 bp of the GUS untranslated leader sequence from pSOG10. A 250 bp fragment encoding the *E. coli* dihydrofolate reductase type II gene was also amplified by PCR and these two PCR fragments were assembled with a SacI-PstI fragment from pBI221  
30 (Clontech) which comprised the pUC19 vector backbone and the nopaline synthase terminator. Assembly of these fragments generated pSOG19 which contains the 35S promoter in fusion with the intron 6 sequence, the GUS leader, the DHFR gene and the nopaline synthase terminator. Replacement of the GUS leader in pSOG19 with the leader sequence from Maize

Chlorotic Mottle Virus check (MCMV) generated the vector pSOG35. pSOG19 and pSOG35 carry the pUC-derived gene for ampicillin resistance and have HindIII, SphI, PstI and EcoRI sites available for the cloning of foreign sequences.

#### **IV. Production and Characterization of Stably Transformed Plants**

5 Transgenic plant cells are then placed in an appropriate selective medium for selection of transgenic cells which are then grown to callus. Shoots are grown from callus and plantlets generated from the shoot by growing in rooting medium. The various constructs normally will be joined to a marker for selection in plant cells. Conveniently, the marker may be resistance to a biocide (particularly an antibiotic, such as kanamycin, G418, bleomycin, hygromycin,  
10 chloramphenicol, herbicide, or the like). The particular marker used will allow for selection of transformed cells as compared to cells lacking the DNA which has been introduced. Components of DNA constructs including transcription cassettes of this invention may be prepared from sequences which are native (endogenous) or foreign (exogenous) to the host. By "foreign" it is meant that the sequence is not found in the wild-type host into which the  
15 construct is introduced. Heterologous constructs will contain at least one region which is not native to the gene from which the transcription-initiation-region is derived.

To confirm the presence of the transgenes in transgenic cells and plants, a variety of assays may be performed. Such assays include, for example, "molecular biological" assays well known to those of skill in the art, such as Southern and Northern blotting, *in situ*  
20 hybridization and nucleic acid-based amplification methods such as PCR or RT-PCR; "biochemical" assays, such as detecting the presence of a protein product, e.g., by immunological means (ELISAs and Western blots) or by enzymatic function; plant part assays, such as leaf or root assays; and also, by analyzing the phenotype of the whole regenerated plant, e.g., for disease or pest resistance.

25 DNA may be isolated from cell lines or any plant parts to determine the presence of the preselected nucleic acid segment through the use of techniques well known to those skilled in the art. Note that intact sequences will not always be present, presumably due to rearrangement or deletion of sequences in the cell.

The presence of nucleic acid elements introduced through the methods of this invention  
30 may be determined by polymerase chain reaction (PCR). Using this technique discreet fragments of nucleic acid are amplified and detected by gel electrophoresis. This type of analysis permits one to determine whether a preselected nucleic acid segment is present in a stable transformant, but does not prove integration of the introduced preselected nucleic acid

segment into the host cell genome. In addition, it is not possible using PCR techniques to determine whether transformants have exogenous genes introduced into different sites in the genome, i.e., whether transformants are of independent origin. It is contemplated that using PCR techniques it would be possible to clone fragments of the host genomic DNA adjacent to an introduced preselected DNA segment.

Positive proof of DNA integration into the host genome and the independent identities of transformants may be determined using the technique of Southern hybridization. Using this technique specific DNA sequences that were introduced into the host genome and flanking host DNA sequences can be identified. Hence the Southern hybridization pattern of a given transformant serves as an identifying characteristic of that transformant. In addition it is possible through Southern hybridization to demonstrate the presence of introduced preselected DNA segments in high molecular weight DNA, i.e., confirm that the introduced preselected DNA segment has been integrated into the host cell genome. The technique of Southern hybridization provides information that is obtained using PCR, e.g., the presence of a preselected DNA segment, but also demonstrates integration into the genome and characterizes each individual transformant.

It is contemplated that using the techniques of dot or slot blot hybridization which are modifications of Southern hybridization techniques one could obtain the same information that is derived from PCR, e.g., the presence of a preselected DNA segment.

Both PCR and Southern hybridization techniques can be used to demonstrate transmission of a preselected DNA segment to progeny. In most instances the characteristic Southern hybridization pattern for a given transformant will segregate in progeny as one or more Mendelian genes (Spencer et al., 1992); Laursen et al., 1994) indicating stable inheritance of the gene. The nonchimeric nature of the callus and the parental transformants ( $R_0$ ) was suggested by germline transmission and the identical Southern blot hybridization patterns and intensities of the transforming DNA in callus,  $R_0$  plants and  $R_1$  progeny that segregated for the transformed gene.

Whereas DNA analysis techniques may be conducted using DNA isolated from any part of a plant, RNA may only be expressed in particular cells or tissue types and hence it will be necessary to prepare RNA for analysis from these tissues. PCR techniques may also be used for detection and quantitation of RNA produced from introduced preselected DNA segments. In this application of PCR it is first necessary to reverse transcribe RNA into DNA, using enzymes such as reverse transcriptase, and then through the use of conventional PCR

techniques amplify the DNA. In most instances PCR techniques, while useful, will not demonstrate integrity of the RNA product. Further information about the nature of the RNA product may be obtained by Northern blotting. This technique will demonstrate the presence of an RNA species and give information about the integrity of that RNA. The presence or  
5 absence of an RNA species can also be determined using dot or slot blot Northern hybridizations. These techniques are modifications of Northern blotting and will only demonstrate the presence or absence of an RNA species.

While Southern blotting and PCR may be used to detect the preselected DNA segment in question, they do not provide information as to whether the preselected DNA segment is  
10 being expressed. Expression may be evaluated by specifically identifying the protein products of the introduced preselected DNA segments or evaluating the phenotypic changes brought about by their expression.

Assays for the production and identification of specific proteins may make use of physical-chemical, structural, functional, or other properties of the proteins. Unique physical-  
15 chemical or structural properties allow the proteins to be separated and identified by electrophoretic procedures, such as native or denaturing gel electrophoresis or isoelectric focusing, or by chromatographic techniques such as ion exchange or gel exclusion chromatography. The unique structures of individual proteins offer opportunities for use of specific antibodies to detect their presence in formats such as an ELISA assay. Combinations  
20 of approaches may be employed with even greater specificity such as Western blotting in which antibodies are used to locate individual gene products that have been separated by electrophoretic techniques. Additional techniques may be employed to absolutely confirm the identity of the product of interest such as evaluation by amino acid sequencing following purification. Although these are among the most commonly employed, other procedures may  
25 be additionally used.

Assay procedures may also be used to identify the expression of proteins by their functionality, especially the ability of enzymes to catalyze specific chemical reactions involving specific substrates and products. These reactions may be followed by providing and quantifying the loss of substrates or the generation of products of the reactions by physical or  
30 chemical procedures. Examples are as varied as the enzyme to be analyzed.

Very frequently the expression of a gene product is determined by evaluating the phenotypic results of its expression. These assays also may take many forms including but not limited to analyzing changes in the chemical composition, morphology, or physiological

properties of the plant. Morphological changes may include greater stature or thicker stalks. Most often changes in response of plants or plant parts to imposed treatments are evaluated under carefully controlled conditions termed bioassays.

#### **V. Uses of Transgenic Plants**

5           Once an expression cassette of the invention has been transformed into a particular plant species, it may be propagated in that species or moved into other varieties of the same species, particularly including commercial varieties, using traditional breeding techniques. Particularly preferred plants of the invention include the agronomically important crops listed above. The genetic properties engineered into the transgenic seeds and plants described above  
10   are passed on by sexual reproduction and can thus be maintained and propagated in progeny plants. The present invention also relates to a transgenic plant cell, tissue, organ, seed or plant part obtained from the transgenic plant. Also included within the invention are transgenic descendants of the plant as well as transgenic plant cells, tissues, organs, seeds and plant parts obtained from the descendants.

15           Preferably, the expression cassette in the transgenic plant is sexually transmitted. In one preferred embodiment, the coding sequence is sexually transmitted through a complete normal sexual cycle of the R0 plant to the R1 generation. Additionally preferred, the expression cassette is expressed in the cells, tissues, seeds or plant of a transgenic plant in an amount that is different than the amount in the cells, tissues, seeds or plant of a plant which  
20   only differs in that the expression cassette is absent.

          The transgenic plants produced herein are thus expected to be useful for a variety of commercial and research purposes. Transgenic plants can be created for use in traditional agriculture to possess traits beneficial to the grower (e.g., agronomic traits such as resistance to water deficit, pest resistance, herbicide resistance or increased yield), beneficial to the  
25   consumer of the grain harvested from the plant (e.g., improved nutritive content in human food or animal feed; increased vitamin, amino acid, and antioxidant content; the production of antibodies (passive immunization) and nutraceuticals), or beneficial to the food processor (e.g., improved processing traits). In such uses, the plants are generally grown for the use of their grain in human or animal foods. Additionally, the use of root-specific promoters in transgenic  
30   plants can provide beneficial traits that are localized in the consumable (by animals and humans) roots of plants such as carrots, parsnips, and beets. However, other parts of the plants, including stalks, husks, vegetative parts, and the like, may also have utility, including use as part of animal silage or for ornamental purposes. Often, chemical constituents (e.g., oils

or starches) of maize and other crops are extracted for foods or industrial use and transgenic plants may be created which have enhanced or modified levels of such components.

Transgenic plants may also find use in the commercial manufacture of proteins or other molecules, where the molecule of interest is extracted or purified from plant parts, seeds, and the like. Cells or tissue from the plants may also be cultured, grown *in vitro*, or fermented to manufacture such molecules.

The transgenic plants may also be used in commercial breeding programs, or may be crossed or bred to plants of related crop species. Improvements encoded by the expression cassette may be transferred, e.g., from maize cells to cells of other species, e.g., by protoplast fusion.

The transgenic plants may have many uses in research or breeding, including creation of new mutant plants through insertional mutagenesis, in order to identify beneficial mutants that might later be created by traditional mutation and selection. An example would be the introduction of a recombinant DNA sequence encoding a transposable element that may be used for generating genetic variation. The methods of the invention may also be used to create plants having unique "signature sequences" or other marker sequences which can be used to identify proprietary lines or varieties.

Thus, the transgenic plants and seeds according to the invention can be used in plant breeding which aims at the development of plants with improved properties conferred by the expression cassette, such as tolerance of drought, disease, or other stresses. The various breeding steps are characterized by well-defined human intervention such as selecting the lines to be crossed, directing pollination of the parental lines, or selecting appropriate descendant plants. Depending on the desired properties different breeding measures are taken. The relevant techniques are well known in the art and include but are not limited to hybridization, inbreeding, backcross breeding, utilane breeding, variety blend, interspecific hybridization, aneuploid techniques, etc. Hybridization techniques also include the sterilization of plants to yield male or female sterile plants by mechanical, chemical or biochemical means. Cross pollination of a male sterile plant with pollen of a different line assures that the genome of the male sterile but female fertile plant will uniformly obtain properties of both parental lines. Thus, the transgenic seeds and plants according to the invention can be used for the breeding of improved plant lines which for example increase the effectiveness of conventional methods such as herbicide or pesticide treatment or allow to dispense with said methods due to their modified genetic properties. Alternatively new crops with improved stress tolerance can be

obtained which, due to their optimized genetic "equipment", yield harvested product of better quality than products which were not able to tolerate comparable adverse developmental conditions.

#### **VI. A Computer Readable Medium**

- 5           The invention also provides a computer readable medium having stored thereon a data structure containing nucleic acid sequences having at least 70% sequence identity to a nucleic acid sequence selected from those listed in SEQ ID Nos: 1-953, 2137-2661, 1954-1966, 2000-2129, 2662-4737 and 4738-6813, as well as complementary, ortholog, and variant sequences thereof. Storage and use of nucleic acid sequences on a computer readable medium is well
- 10   known in the art. (See for example U.S. Patent Nos. 6,023,659; 5,867,402; 5,795,716) Examples of such medium include, but are not limited to, magnetic tape, optical disk, CD-ROM, random access memory, volatile memory, non-volatile memory and bubble memory. Accordingly, the nucleic acid sequences contained on the computer readable medium may be compared through use of a module that receives the sequence information and compares it to
- 15   other sequence information. Examples of other sequences to which the nucleic acid sequences of the invention may be compared include those maintained by the National Center for Biotechnology Information (NCBI)(<http://www.ncbi.nlm.nih.gov/>) and the Swiss Protein Data Bank. A computer is an example of such a module that can read and compare nucleic acid sequence information. Accordingly, the invention also provides the method of comparing a
- 20   nucleic acid sequence of the invention to another sequence. For example, a sequence of the invention may be submitted to the NCBI for a Blast search as described herein where the sequence is compared to sequence information contained within the NCBI database and a comparison is returned. The invention also provides nucleic acid sequence information in a computer readable medium that allows the encoded polypeptide to be optimized for a desired
- 25   property. Examples of such properties include, but are not limited to, increased or decreased: thermal stability, chemical stability, hydrophylicity, hydrophobicity, and the like. Methods for the use of computers to model polypeptides and polynucleotides having altered activities are well known in the art and have been reviewed. (Lesyng et al., 1993; Surles et al., 1994; Koehl et al., 1996; Rossi et al., 2001).
- 30           The invention will be further described by the following non-limiting examples.



### Example 1

#### GeneChip Standard Protocol

##### Quantitation of total RNA

Total RNA from plant tissue is extracted and quantified.

- 5        1. Quantify total RNA using GeneQuant
- 1OD<sub>260</sub>=40 mg RNA/ml; A260/A280=1.9 to about 2.1
2. Run gel to check the integrity and purity of the extracted RNA

##### Synthesis of double-stranded cDNA

- 10        Gibco/BRL SuperScript Choice System for cDNA Synthesis (Cat#1B090-019) was employed to prepare cDNAs. T7-(dT)<sub>24</sub> oligonucleotides were prepared and purified by HPLC. (5'-GGCCAGTGAATTGTAATACGACTCACTATAGGGAGGCGG-(dT)<sub>24</sub>-3' SEQ ID NO:2136).

##### Step 1. Primer hybridization:

- 15        Incubate at 70°C for 10 minutes
- Quick spin and put on ice briefly

##### Step 2. Temperature adjustment:

          Incubate at 42°C for 2 minutes

##### Step 3. First strand synthesis:

- 20        DEPC-water- 1 µl
- RNA (10 µg final)-10 µl
- T7=(dT)<sub>24</sub> Primer (100 pmol final)-1 µl pmol
- 5X 1st strand cDNA buffer-4 µl
- 0.1M DTT (10 mM final)- 2 µl
- 25        10 mM dNTP mix (500 µM final)-1 µl
- Superscript II RT 200 U/µl- 1 µl
- Total of 20 µl
- Mix well
- Incubate at 42°C for 1 hour

##### Step 4. Second strand synthesis:

30        Place reactions on ice, quick spin

          DEPC-water- 91 µl

          5X 2nd strand cDNA buffer- 30 µl

- mM dNTP mix (250 mM final) - 3  $\mu$ l  
*E. coli* DNA ligase (10 U/ $\mu$ l)-1  $\mu$ l  
*E. coli* DNA polymerase 1-10 U/ $\mu$ l- 4  $\mu$ l  
 RnaseH 2U/ $\mu$ l -1  $\mu$ l  
 5 T4 DNA polymerase 5 U/ $\mu$ l-2  $\mu$ l  
 0.5 M EDTA (0.5 M final)--10  $\mu$ l  
 Total 162  $\mu$ l  
 Mix/spin down/incubate 16°C for 2 hours

Step 5. Completing the reaction:

- 10 Incubate at 16°C for 5 minutes

Purification of double stranded cDNA

1. Centrifuge PLG (Phase Lock Gel, Eppendorf 5 Prime, Inc., PI-188233) at 14,000X, transfer 162  $\mu$ l of cDNA to PLG
- 15 2. Add 162  $\mu$ l of Phenol:Chloroform:Isoamyl alcohol (pH 8.0), centrifuge 2 minutes
3. Transfer the supernatant to a fresh 1.5 ml tube, add
 

|                                      |     |
|--------------------------------------|-----|
| Glycogen (5 mg/ml)                   | 2   |
| 0.5 M NH <sub>4</sub> OAC (0.75xVol) | 120 |
| ETOH (2.5xVol, -20 C)                | 400 |
- 20 4. Mix well and centrifuge at 14,000X for 20 minutes
5. Remove supernatant, add 0.5 ml 80% EtOH (-20°C)
6. Centrifuge for 5 minutes, air dry or by speed vac for 5-10 minutes
7. Add 44  $\mu$ l DEPC H<sub>2</sub>O

Analyze of quantity and size distribution of cDNA

- 25 Run a gel using 1  $\mu$ l of the double-stranded synthesis product

Synthesis of biotinylated cRNA

(use Enzo BioArray High Yield RNA Transcript Labeling Kit Cat#900182)

- |    |                            |            |
|----|----------------------------|------------|
|    | Purified cDNA              | 22 $\mu$ l |
| 30 | 10X Hy buffer              | 4 $\mu$ l  |
|    | 10X biotin ribonucleotides | 4 $\mu$ l  |
|    | 10X DTT                    | 4 $\mu$ l  |
|    | 10X Rnase inhibitor mix    | 4 $\mu$ l  |

|                       |           |
|-----------------------|-----------|
| 20X T7 RNA polymerase | 2 $\mu$ l |
|-----------------------|-----------|

|       |            |
|-------|------------|
| Total | 40 $\mu$ l |
|-------|------------|

Centrifuge 5 seconds, and incubate for 4 hours at 37EC

Gently mix every 30-45 minutes

5

#### Purification and quantification of cRNA

(use Qiagen Rneasy Mini kit Cat# 74103)

Determine concentration and dilute to 1  $\mu$ g/ $\mu$ l concentration

#### 10 Fragmentation of cRNA

|                           |            |
|---------------------------|------------|
| cRNA (1 $\mu$ g/ $\mu$ l) | 15 $\mu$ l |
|---------------------------|------------|

|                          |           |
|--------------------------|-----------|
| 5X Fragmentation Buffer* | 6 $\mu$ l |
|--------------------------|-----------|

|                       |                            |
|-----------------------|----------------------------|
| DEPC H <sub>2</sub> O | <u>9 <math>\mu</math>l</u> |
|-----------------------|----------------------------|

|            |
|------------|
| 30 $\mu$ l |
|------------|

15

#### \*5x Fragmentation Buffer

|                 |        |
|-----------------|--------|
| 1M Tris (pH8.1) | 4.0 ml |
|-----------------|--------|

|       |        |
|-------|--------|
| MgOAc | 0.64 g |
|-------|--------|

|      |        |
|------|--------|
| KOAC | 0.98 g |
|------|--------|

20

|                       |  |
|-----------------------|--|
| DEPC H <sub>2</sub> O |  |
|-----------------------|--|

|       |       |
|-------|-------|
| Total | 20 ml |
|-------|-------|

Filter Sterilize

#### Array wash and staining

25 Stringent Wash Buffer\*\*

Non-Stringent Wash Buffer\*\*\*

SAPE Stain\*\*\*\*

Antibody Stain\*\*\*\*\*

30 Wash on fluidics station using the appropriate antibody amplification protocol

\*\*Stringent Buffer: 12X MES 83.3 ml, 5 M NaCl 5.2 ml, 10% Tween 1.0 ml, H<sub>2</sub>O 910 ml;

Filter Sterilize

\*\*\*Non-Stringent Buffer: 20X SSPE 300 ml, 10% Tween 1.0 ml, H<sub>2</sub>O 698 ml, Filter Sterilize, Antifoam 1.0.

\*\*\*\*SAPE stain: 2X Stain Buffer 600 µl, BSA 48 µl, SAPE 12µl, H<sub>2</sub>O 540 µl.

\*\*\*\*\*Antibody Stain: 2X Stain Buffer 300 µl, H<sub>2</sub>O 266.4 µl, BSA 24 ul, Goat IgG 6 µl,

5 Biotinylated Ab 3.6 µl

#### Image analysis and data mining

1. Two text files are included in the analysis:

- a. One with Absolute analysis: giving the status of each gene, either absent or present in the samples
- 10 b. The other with Comparison analysis: comparing gene expression levels between two samples

### Example 2

#### Analysis of the RPS2 Mediated Interaction in *Arabidopsis*

15 The identification and cloning of resistance genes is extremely important for the treatment of crops. For example, bacterial blight disease caused by *Xanthomonas spp.* infects virtually all crop plants and leads to extensive crop losses worldwide. Therefore, it is of interest to identify diverse and abundant plant resistance genes for use as future crop treatments for pathogen resistance, e.g., to identify particular pathogen resistance (*R*) genes in  
20 a plant.

Differential gene expression analysis was used to identify pathogen resistance (*R*) genes in a plant. This method takes advantage of the HR-associated disease resistance. One model plant-pathogen interaction is that of *Arabidopsis thaliana* and *Pseudomonas syringae* pv *tomato*. There are four possible genetic interactions of a *P. syringae* infection of  
25 *Arabidopsis* when analyzing HR-associated disease resistance (Table 2). However, there are only two possible outcomes: a compatible outcome occurs when there is disease, and an incompatible outcome occurs when there is no disease. An incompatible outcome, or disease resistance, occurs only when the plant possesses the resistance gene, e.g., *RPS2*, and the pathogen possesses the corresponding *avr* gene, e.g., *avrRpt2*. *RPS2* belongs to the NBS-LRR  
30 class of *R* genes, which can confer resistance to a wide variety of phytopathogens. It has been suggested that AvrRpt2 is delivered to the plant via the bacteria's type III secretion system and recognized by a surveillance system involving *RPS2* inside the plant cell. The plant response during an incompatible interaction includes a change in ion flux across the plasma membrane,

generation of reactive oxygen species, induction of defense genes, induction of HR, fortification of the cell wall, accumulation of salicylic acid, and anti-microbial compounds.

Table 2

| Number | Plant       | Pathogen       | Outcome    |              |
|--------|-------------|----------------|------------|--------------|
| 1      | <i>RPS2</i> | no <i>avr</i>  | Disease    | Compatible   |
| 2      | <i>RPS2</i> | <i>avrRpt2</i> | No disease | Incompatible |
| 3      | <i>rps2</i> | no <i>avr</i>  | Disease    | Compatible   |
| 4      | <i>rps2</i> | <i>avrRpt2</i> | Disease    | Compatible   |

5

### Methods

#### Differential Expression

Analysis of differential gene expression is a classic and very powerful tool in experimental biology not only to study large trends in gene regulation but also small differences among similar responses. Historically, methods for analysis only allowed the comparison of a very few genes in each experiment. However, with new methods to identify and quantitate differential mRNA profiles, such as long distance differential display PCR, cDNA microarrays, and gene chips, one can much more quickly and comprehensively identify and analyze differentially expressed genes.

By analyzing and comparing the expression profile of genes in the above 4-way matrix, a number of types of genes can be identified that are involved in the resistance pathway. Resistance genes would be highly expressed or strongly downregulated in outcome number 2 in the four way matrix and less oppositely expressed in outcome numbers 1, 3, and 4. Genes that are highly expressed or strongly downregulated in outcome numbers 1 and 2 and oppositely expressed or not expressed above baseline in outcome numbers 3 and 4 are of interest as being associated with the reaction of a plant having resistance genes to a bacterial infection, regardless of the *avr* genotype of the bacterium. Such a comparison is very useful in identifying strong candidates for different roles in plant/pathogen interactions, as are numerous other kinds of outcomes in the four-way plant/pathogen interaction analysis of gene expression. Such genes include those involved in recognition of pathogen (unrelated to virulence status); genes involved in recognition of pathogen having a virulence or avirulence gene (regardless of the status of the corresponding plant); genes related to the status of the

plant, regardless of the status of the pathogen; and genes that do not change expression during plant-pathogen interaction.

#### Use of a Gene Chip to Study Gene Regulation in *Arabidopsis* in Response to Exposure to Pathogen

5 Initially isogenic strains of *Arabidopsis thaliana* ecotype Col-0 were used, one having the wild type *RPS2* gene that confers resistance, and one having the *rps2* mutant that confers susceptibility to attack by *Pseudomonas syringae* pathovar *tomato* (Pst). Subsequently, comparisons between ecotypes, mutant *Arabidopsis*, and infection with different pathogens were made. After infection, the RNA was isolated and a probe produced using the Affymetrix  
10 GeneChip™ protocol. A gene array representing approximately 8,100 *Arabidopsis thaliana* genes was used to carry out global gene expression profiling in response to exposure to a particular pathogen.

Initially, the analysis involved comparing all four of the interactions to a water control (plants "infected" with water). In the initial analysis, the mRNA levels of approximately 1,600  
15 genes were significantly affected (> 2.5-fold change in expression) by exposure to the bacterial pathogen. This suggested a dramatic change in the molecular biology of the cell and a more detailed analysis was performed.

#### Results

##### A. Comparison Of Compatible To Incompatible Infections

20 Two different types of interactions between *Arabidopsis* and *Pseudomonas syringae* were analyzed. In one type of experiment, a gene for gene interaction conditioned by the plant resistance (*R*) gene *RPS2* and the bacterial avirulence gene *avrRpt2* at a relatively early stage was analyzed. When the pathogen has an *avr* gene and the plant has the corresponding *R* gene, the plant is resistant to the pathogen and the interaction is called incompatible. When the  
25 plant-pathogen system lacks either or both genes, the plant is susceptible to the pathogen and the interaction is called compatible. A hypersensitive response (HR, localized rapid cell death of the plant) is one aspect of resistance.

Isogenic strains of *Arabidopsis thaliana* ecotype Col-0 were used, one having the wild type *RPS2* gene that confers resistance, and one having the mutant *rps2* mutant that confers  
30 susceptibility to attack by *Pseudomonas syringae* pathovar *tomato* (Pst) carrying *avrRpt2*. Two strains of *Pseudomonas syringae* were used, one having the *avr* gene *avrRpt2* and the other having no *avr*. The *avr* gene is carried on a plasmid.

A gene array having 8,700 probe sets representing approximately 8,100 *Arabidopsis thaliana* genes was used to carry out global gene expression profiling of each of the infection outcomes. The pairings were as follows:

1. RPS2 WT plant; *P. syringae* (no *avr*)
- 5      2. RPS2 WT plant; *P. syringae/avrRpt2*
3. *rps2-101C* mutant plant; *P. syringae* (no *avr*)
4. *rps2-101C* mutant plant; *P. syringae/avrRpt2*

Additionally, two controls were used:

5. RPS2 WT plant; water control
- 10      6. *Rps2-101C* mutant plant; water control

Data were processed such that genes having a difference in mRNA levels that was greater than 2.5-fold increased or reduced, compared with controls were selected. The fold change for each gene was log-scaled and normalized.

1. Data analysis: identification of expression clusters

15      Data analysis was carried out by comparing expression of each gene in interactions 1-4 (Table 2), plotting that expression level, and identifying the genes of interest, i.e., those that show more than a 2.5X change in expression (about 1,600 genes). Classification of patterns, or expression clusters were as follows:

- a) Genes strongly induced (> 2.5X change in expression level) only in the resistant  
20 (incompatible) response;
- b) Genes responding weakly only in the resistance response, but strongly induced in the compatible response;
- c) Genes that show a high level of expression in all outcomes;
- d) Genes that show a high level of repression in all outcomes;
- 25      e) Genes that show a very high level of repression only when the bacterial *avr* is expressed; and
- f) Genes that show a very different level of expression in the presence of the plant resistance compared to the level in the absence of the plant resistance (the mutant *rps2*).

30      Genes that fall within groups 1a and 1b, i.e., those that are differentially expressed only when an incompatible interaction occurs, include genes directly involved in resistance to pathogens. These genes show a peak (either up or down) only during plant-pathogen interaction 2. The differential expression can be of two types: upregulated (increased expression of this gene is potentially important in the incompatible interaction) or

downregulated (decreased expression of this gene is potentially important in the incompatible interaction).

## 2. Heat shock proteins and transcription factors

All major heat shock proteins (HSPs) were identified to be upregulated only during the incompatible interaction. Heat shock factors (HSFs) are transcription factors which control the transcription of the HSP genes. Eight HSF genes are known in *Arabidopsis*. HSF4 and HSF21 were identified as being upregulated when the plant was infected with *P. syringae*. HSF4 showed strong induction that was restricted to resistance, and HSF4 was the only HSF specifically upregulated during the incompatible interaction. The data suggests that the upregulation of HSPs is downstream of upregulation of HSF4.

To analyze whether the response was a more general one, or specific to a given ecotype, expression of HSF4 was analysed in two different *Arabidopsis* ecotypes, *A. thaliana*, ecotypes Col-0 and Ws. HSF4 was also upregulated in the response of Ws ecotype to infection and, specifically, was upregulated during an incompatible response. HSF21 is thus a preferred protein for resistance applications, and HSF4, a protein which is expressed in all plants, is especially preferred for engineering resistance.

A transgene containing the ACT2 promoter and the HSF4 open reading frame was introduced to *Arabidopsis* and transgenic HSF4 *Arabidopsis* lines generated to overexpress and underexpress HSF4. The expression of HSF4 during pathogen infection may cause lower general resistance to *P. syringae*.

Conditional overexpression lines were also generated using the estradiol-inducible promoter system. Infiltration of 20  $\mu$ M estradiol into the intercellular space of the leaves of transgenic plants induced expression of HSF4 mRNA for a short time (down by 4 hours). Addition of 20  $\mu$ M estradiol to the hydroponic medium yielded sustained HSF4 mRNA accumulation.

## B. Genes Involved in *Arabidopsis* Responses to Pathogens

A number of mutations in *Arabidopsis thaliana* that disrupt expression of pathogen-induced genes and cause enhanced disease susceptibility have been identified. Pathogen-induced genes whose expression is altered in these enhanced disease susceptibility mutants are likely to play important roles in conferring disease resistance.

To identify such genes, wild type and various mutant plants were infected with strain *Psm* ES4326 at a dose of 10,000 colony forming units per square centimeter of leaf tissue. Control plants were mock-infected. After thirty hours, tissue samples were collected and used



to prepare RNA. Three sets of experiments were carried out. Each set of experiments included three independent replicate experiments. RNA from replicate experiments was pooled to reduce errors arising from the effects of variations in environmental conditions. Each RNA sample was used to prepare a fluorescently-labelled probe which was applied to an Affymetrix GeneChip™, allowing the expression level of each gene represented on the GeneChip™ to be determined for each sample. The plant genotypes included in each experiment were as follows:

#### Experiment #1

- 10 Wild-type (ecotype Columbia)  
     *nahG*  
     *pad4-1*  
     *eds5-1*  
     *eds4*  
     *pad2-1*  
     *npr1-1*  
     *npr1-3*

#### Experiment #2

- 20 Wild-type (ecotype Columbia)  
     *coil*  
     *ein2*  
     *pad1*  
     *FN1-3*  
     *eds3*  
     *eds8*

#### Experiment #3

- 30 Wild-type (ecotype Columbia)  
     *pad4-1*  
     *nahG*  
     *sid2*  
     *eds5-3*  
     *FN1-9*  
     *FN3-2*

#### 1. Data analysis

- Expression values that were less than 5 were set to five. This ensures that no values are 0 or negative. Such values interfere with subsequent analysis steps. To obtain a list of pathogen-induced genes, the ratios of infected wild-type to mock infected wild type were calculated for each experiment. Then genes were selected in which expression levels were infected wild-type/mock wild-type > 2.5, and infected wild-type > 50 for at least 2 of 3

experiments. The ratio of 2.5 was chosen because the false positive rate for the GeneChip™ is essentially 0 at this level of stringency, and the absolute value of 50 was chosen to eliminate expression values below the detection limit of the GeneChip™. The result of this analysis was a list of 745 probe sets representing genes that are induced by infection in wild-type plants

5 (note that some genes are represented by more than one probe set, so the number of different genes is somewhat fewer) (see Table 3 below). Hence, the expression of genes comprising SEQ ID NOs:2-6, 8-13, 16, 18, 22-23, 25, 28-29, 31-32, 35-37, 39-43, 45-47, 49-50, 52, 54-55, 57-58, 60-66, 70-72, 74, 76-77, 79, 81, 83, 85, 87-90, 92, 94, 97, 100-107, 111-115, 117-125, 127-135, 138-140, 142-153, 156-158, 160, 162-165, 168-170, 173-181, 183-184, 186-188, 190-198, 200-201, 203-211, 214-215, 218-224, 227-232, 234-249, 251-262, 264, 266-268, 270, 272-275, 277-281, 283, 286-294, 297-298, 302, 304-306, 308-326, 328-339, 341, 344-345, 347, 350-351, 353-358, 361-371, 373-377, 379-386, 388-390, 392, 394-400, 402-406, 408-410, 412-417, 419-427, 429-433, 435-443, 445-452, 454-457, 459-460, 462-464, 466-470, 473-475, 478-479, 481-482, 484-187, 489-494, 496-498, 500-501, 503-506, 508,

15 510, 512-515, 517-523, 526, 528-529, 531-538, 540, 544-548, 550-558, 560, 563-568, 570, 572-577, 579-580, 582-585, 588-594, 596, 598-600, 602-603, 605-606, 608-612, 614-617, 619-624, 626-630, 632-639, 642, 644, 646-651, 653-657, 659-665, 667-671, 673-678, 681-689, 691-693, 695-713, 715-717, 719, 721-727, 729-733, 736-738, 740, 742, 744, 746, 748-752, 755-756, 758-760, 762-769, 771, 774, 776-781, 783-788, 790-796, 798-799, 802, 804-

20 808, 810-815, 817-831, 833-848, 850-855, 857-869, 871-880, 882-900, 903-907, 909, 911-915, 918-920, 922-925, 927, 929, 931-938, 940, 943-945, 947, and 950-953 is increased after infection of wild-type *Arabidopsis* with *Pseudomonas syringae*.

To identify pathogen-induced genes whose expression is affected by the mutations, genes for which the ratio of infected mutant/infected wild-type was  $< 0.5$  or  $> 2$  for at least one

25 mutant were selected from the list of 745 pathogen-inducible probe sets. The limits of 5 and 2 were chosen because changes of at least 2-fold are likely to be significant for impact on disease resistance, and because the false positive rate for the GeneChip™ at 2-fold is 0.2%. This selection yielded a list of 530 probe sets corresponding to genes, the expression of which is induced by *Pseudomonas* infection in wild-type plants and perturbed in at least one mutant

30 plant (see Tables 4a and 4b below). Thus, the expression of genes comprising SEQ ID NOs:2, 4-6, 11-13, 18, 22-23, 28, 31, 36, 39-43, 45, 47, 49-50, 52, 54-55, 57-58, 60-61, 63-66, 71-72, 74, 77, 81, 83, 85, 87-89, 92, 97, 100-107, 111-112, 114-115, 117-120, 122, 125, 127-128, 134, 138-140, 143-144, 148-151, 153, 156-157, 160, 165, 168-170, 173-174, 176-180, 183,

187-188, 191, 193-194, 197-198, 200, 203-210, 214, 219-224, 227, 230-232, 235-237, 239-240, 243-246, 248-249, 251-254, 256-258, 261, 264, 266-268, 270, 272-275, 277-278, 280, 283, 286-287, 290-293, 297, 302, 305-306, 308-310, 312-316, 321-326, 328-331, 333, 336-339, 341, 345, 351, 353, 355-358, 361-363, 365-366, 368-371, 373, 375, 377, 379-381, 384-385, 388-390, 392, 394-400, 402-406, 410, 412, 415-416, 419-420, 422-425, 429-433, 435-439, 441-443, 445-452, 454, 459-460, 463, 466, 468-470, 473, 481-482, 485-486, 489, 491-494, 497-498, 500-501, 503, 505-506, 508, 510, 513-515, 517, 520-521, 523, 528-529, 531, 533-538, 540, 545-548, 550-551, 553-554, 556-558, 560, 566-567, 575, 580, 582-584, 588-593, 596, 598-600, 602-603, 605-606, 608-610, 612, 614, 616, 620-622, 627-629, 633-634, 636-639, 644, 646, 648-651, 654-657, 659, 661-663, 667, 669, 673-674, 677, 682, 684-687, 689, 691-693, 697, 699, 701, 703-708, 713, 717, 719, 721-727, 730-733, 736, 740, 744, 746, 749-752, 755-756, 758-760, 762-764, 766-769, 774, 776-778, 780-781, 786, 788, 791-796, 799, 802, 804-808, 810-812, 815, 818-821, 823-825, 827-829, 831, 833-836, 838-843, 845, 847-848, 852-853, 855, 858, 860-869, 871-874, 876, 878-880, 884-887, 889, 892-894, 896-900, 904-907, 911-915, 918-920, 922-924, 931, 933, 938, 943-945, 947, and 950-952 is increased after infection of wild-type *Arabidopsis*, and altered after infection of at least one mutant *Arabidopsis*, with *Pseudomonas syringae*.

## 2. Data interpretation

Genes that encode regulatory proteins such as transcription factors, protein kinases, calcium binding proteins and the like, are likely to play important roles in disease resistance, as they are likely to affect the expression of multiple defense effector genes. The list of 530 probe sets include 81 that correspond to genes encoding regulatory factors. These are likely to be useful for engineering plants to respond more quickly to pathogen attack by activating expression of defense responses (see Table 5 below). Thus, the expression of genes comprising SEQ ID NOs:39, 52, 60, 63, 81, 83, 106, 107, 115, 117, 118, 168, 174, 176, 179, 204, 207, 208, 220, 221, 248, 258, 268, 275, 280, 309, 323, 326, 329, 351, 419, 422, 429, 430, 432, 459, 460, 468, 469, 473, 500, 505, 506, 508, 529, 531, 533, 535, 538, 545, 553, 602, 606, 608, 610, 614, 616, 634, 654, 655, 684, 686, 687, 691, 717, 751, 752, 766, 777, 815, 831, 834, 835, 839, 841, 847, 876, 884, 906, 920, and 924 is increased after infection of wild-type *Arabidopsis*, and altered after infection of at least one mutant *Arabidopsis*, with *Pseudomonas syringae*.

The mutations *nahG*, *pad4-1*, *eds5-1*, *eds4*, *pad2-1*, *npr1-1*, *npr1-3*, *pad1*, *FN1-3*, *eds3*, *eds8*, *sid2*, *eds5-3*, *FN1-3* and *FN3-2* cause enhanced susceptibility to *Pseudomonas syringae*. Consequently, pathogen-inducible genes whose expression is reduced by one of these

mutations are likely to be important for resistance to *Pseudomonas syringae* and possibly other bacterial pathogens. These 333 probe sets are shown in Table 6 (below). Therefore, the expression of genes comprising SEQ ID NOs:12-13, 18, 23, 36, 39-40, 43, 45, 50, 52, 57-58, 60-61, 64, 71-72, 81, 87-89, 97, 100, 102-105, 107, 111-112, 115, 119-120, 122, 125, 127-128, 140, 144, 148-150, 153, 165, 168-169, 176-177, 179, 183, 188, 191, 193-194, 197-198, 203-206, 208-209, 214, 219-222, 227, 230, 232, 237, 244-246, 248-249, 251-253, 258, 261, 264, 266, 268, 273-275, 283, 287, 290, 293, 297, 302, 305-306, 308, 312-315, 321-322, 324, 326, 330, 333, 338, 341, 345, 353, 356-358, 362-363, 366, 369, 371, 375, 377, 380, 384-385, 389, 392, 394-395, 398-399, 402-404, 406, 410, 415, 419, 422, 425, 429-430, 433, 435-439, 443, 445-452, 454, 463, 466, 468-470, 473, 486, 489, 491-492, 494, 498, 500-501, 503, 508, 513-514, 517, 529, 533-538, 548, 550, 553-554, 4556-558, 566, 575, 580, 582-583, 590-591, 593, 600, 602, 609-610, 612, 614, 620-622, 627-629, 637-638, 644, 649, 654-657, 659, 663, 667, 669, 673-674, 677, 684-685, 689, 691-693, 699, 703-705, 708, 719, 721, 724-726, 730-732, 744, 746, 749-750, 752, 755-756, 758, 760, 762-764, 767, 769, 774, 780-781, 786, 788, 791-792, 794-796, 799, 804-808, 810-812, 815, 818-819, 823, 828-829, 833, 840-841, 843, 847, 852-853, 858, 860, 862-865, 867-868, 872-874, 876, 885-887, 889, 892-894, 896-900, 904-905, 907, 911-914, 918-920, 922-924, 931, 933, 938, 947, 950, and 952 is increased after infection of wild-type *Arabidopsis*, and altered after infection of at least one mutant *Arabidopsis* having a mutation that results in enhanced susceptibility to *Pseudomonas* (*nahG*, *pad 4-1*, *eds 5-1*, *eds4*, *pad2-1*, *np4 1-1*, *npr 1-3*, *pad1*, *FN1-3*, *eds3*, *eds8*, *sid2*, *eds5-3*, *NF1-3* and *FN3-2*).

The mutations *coil* and *ein2* block jasmonate and ethylene signaling, respectively. Jasmonate and ethylene-dependent disease resistance responses are known to be important for resistance to the fungal pathogens *Alternaria brassicicola* and *Botrytis cinerea*, and may also be important for resistance to other necrotrophic fungal pathogens. *Alternaria* and *Botrytis* are distantly related, yet plant resistance to these fungi is controlled similarly, suggesting that jasmonate- and ethylene-dependent responses function to limit growth of a wide range of fungal pathogens. Consequently, pathogen-induced genes whose expression is reduced in *coil* and *ein2* mutants are likely to be important for resistance to these necrotrophic fungal pathogens. These 296 probe sets are shown in Table 7 (see below). Hence, the expression of genes comprising SEQ ID NOs:2, 4, 6, 11-13, 18, 22-23, 31, 41-43, 49-50, 54, 57-58, 61, 64-66, 71-72, 74, 77, 85, 87, 89, 92, 97, 101, 103, 106-107, 112, 114, 117-119, 125, 128, 134, 138, 143, 149, 151, 156-157, 165, 169-170, 174, 176-180, 187-188, 191, 193, 206, 208, 219-

220, 222, 224, 231, 236, 239, 243-245, 251-254, 256-257, 267, 272, 287, 290, 292, 297, 302, 312-313, 315-316, 321-322, 324-325, 328, 330, 345, 351, 353, 355-357, 362-363, 366, 368-371, 373, 375, 379, 381, 384, 388-390, 392, 395-400, 405, 410, 415-416, 419, 422, 424, 431-432, 435-436, 438-439, 447, 459-460, 470, 473, 481-482, 489, 491, 493-494, 500-501, 505-506, 513-514, 517, 520-521, 523, 528-529, 531, 535, 537-538, 540, 545-548, 551, 553-554, 557-558, 566, 575, 580, 582, 584, 589, 591, 593, 596, 598-599, 603, 605, 608-609, 612, 628, 633-634, 636-637, 639, 646, 648, 650-651, 656, 661, 663, 667, 674, 685-687, 689, 691, 693, 697, 699, 701, 705, 707, 713, 723-724, 726, 736, 740, 749, 751-752, 756, 758-759, 764, 766-768, 774, 776, 778, 780, 792-796, 799, 802, 806, 810-812, 818, 820-821, 825, 827-829, 833-836, 838-839, 841-843, 848, 855, 860-861, 866, 868-869, 871, 873-874, 876, 878-880, 889, 892, 898-900, 904-905, 907, 915, 918, 922, 924, 933, 943-945, 947, and 951 is increased after infection of wild-type *Arabidopsis*, and altered after infection of at least one mutant *Arabidopsis* having a mutation in a gene whose expression is important for resistance to necrotrophic fungi (a mutation that blocks or interferes with jasmonate and ethylene signaling such as *col1* and *ein2*). Accordingly, these genes are useful to improve the resistance of plants to fungal infection.

The mutations *nahG*, *pad4-1*, *sid2*, *eds5-1*, *eds5-3*, and *eds4* are known to interfere with salicylic acid dependent signaling. Such signaling is known to be important for resistance to the bacterial pathogen *Pseudomonas syringae*, the oomycete pathogen *Peronospora parasitica*, the viral pathogen tobacco mosaic virus, as well as various other plant pathogens. Consequently, pathogen-induced genes whose expression is reduced by one of the mutations that block salicylate signaling are likely to be important for disease resistance, and useful for engineering improved disease resistance. These 288 probesets are shown in Table 8 (see below). Therefore, the expression of genes comprising SEQ ID NOs: 12-13, 18, 23, 36, 39-40, 43, 45, 50, 52, 57-58, 60-61, 64, 71-72, 81, 87-88, 100, 102-105, 107, 111-112, 115, 119-120, 122, 125, 127-128, 140, 148-150, 153, 168-169, 176-177, 188, 191, 193-194, 197-198, 203-206, 209, 219-222, 227, 232, 237, 244-246, 248-249, 251-253, 258, 261, 264, 266, 268, 273-275, 283, 287, 290, 293, 297, 302, 305-306, 308, 312-315, 324, 326, 330, 333, 341, 345, 353, 356, 358, 366, 371, 375, 377, 380, 385, 389, 392, 394, 398, 402-404, 406, 410, 415, 419, 425, 429-430, 433, 435-438, 443, 445-447, 449-452, 454, 463, 466, 468-470, 473, 486, 489, 492, 494, 498, 500-501, 503, 508, 513-514, 517, 533-538, 548, 550, 553-554, 57-558, 566, 575, 582-583, 590-591, 593, 600, 602, 609-610, 612, 620-622, 627-629, 637-638, 644, 649, 654-657, 659, 667, 669, 673, 677, 684, 689, 692-693, 703-705, 719, 721, 724-726, 730-732, 744,

746, 749-750, 752, 755-756, 760, 762-764, 767, 769, 774, 780-781, 786, 788, 791-792, 795-796, 805-808, 810-812, 815, 818-819, 823, 828, 833, 840-841, 843, 852-853, 858, 860, 862-865, 867-868, 872-874, 876, 887, 889, 893-894, 896-898, 900, 905, 907, 911-914, 918-920, 922-923, 931, 933, 938, 947, 950, and 952 which is increased after infection of wild-type *Arabidopsis* and altered after infection of at least one mutant *Arabidopsis* having a mutation in a gene that interferes with salicylic acid dependent signaling (*nahG*, *pad4-1*, *sid2*, *eds5-1*, *eds5-3* and *eds4*). Thus, these genes are particularly useful to improve the resistance of plants to infection by more than one pathogen including bacteria, oomycetes and viruses, such as TMV.

10

### Example 3

#### Further Analysis of the Pathogen Response and Comparison of the Response in Different Ecotypes

##### Materials and Methods

15 *Arabidopsis* ecotypes (or accessions) (the wild-types of all the *Arabidopsis* ecotypes used here have wild-type alleles of *RPS2* and *RPM1*).

Col, Columbia-0

Ler, Landsberg *erecta*

Ws, Wassilewskija

20 *Arabidopsis* mutants and transgenics

Col rps2-101C, a loss-of-function mutant of the resistance gene *RPS2* in Col background.

NahG, transgene for salicylic acid hydroxylase (inactivating salicylic acid). Col background.

25 ndr1-1, null mutant allele of *NDRI* (non-race specific disease resistance).

The mutation strongly affects *RPS2*-mediated resistance and partially affects *RPM1*-mediated resistance. Col background.

##### Bacterial strains

30 Pst, *Pseudomonas syringae* pv. *tomato* DC3000 (virulent strain of *Arabidopsis*)

Psm, *P. syringae* pv. *maculicola* ES4326 (another virulent strain of *Arabidopsis*)

Psp, *P. syringae* pv. *phaseolicola* NPS3121 (very weak pathogen of

*Arabidopsis*)Avirulence (*avr*) genes of *P. syringae*

*avrRpt2*: corresponding to the *Arabidopsis* resistance (*R*) gene *RPS2*

*avrB*: corresponding to the *Arabidopsis* resistance (*R*) gene *RPM1*

5 Experimental ProtocolsA. Gene for gene resistance (6 hours after treatment)

|    | <u>plant</u>          | <u>treatment</u>    |
|----|-----------------------|---------------------|
|    | Col WT                | H <sub>2</sub> O    |
|    | Col WT                | Pst                 |
| 10 | Col WT                | Pst/ <i>avrRpt2</i> |
|    | Col <i>rps2</i> -101C | H <sub>2</sub> O    |
|    | Col <i>rps2</i> -101C | Pst                 |
|    | Col <i>rps2</i> -101C | Pst/ <i>avrRpt2</i> |
|    | Ws WT                 | H <sub>2</sub> O    |
| 15 | Ws WT                 | Pst                 |
|    | Ws WT                 | Pst/ <i>avrRpt2</i> |

B. Differences in the response to bacterial pathogens among ecotypes (3, 6, and 9 hours after treatment)

|    | <u>Plant</u> | <u>treatment</u>    |
|----|--------------|---------------------|
| 20 | Col          | H <sub>2</sub> O    |
|    | Col          | Pst                 |
|    | Col          | Pst/ <i>avrRpt2</i> |
|    | Ler          | H <sub>2</sub> O    |
|    | Ler          | Pst                 |
| 25 | Ler          | Pst/ <i>avrRpt2</i> |
|    | Ws           | H <sub>2</sub> O    |
|    | Ws           | Pst                 |
|    | Ws           | Pst/ <i>avrRpt2</i> |

Note that overall results for Cvi were very similar to Ler.

30 C. Genetic factors that affect the plant response to incompatible interactions (3, 6, and 9 hours after treatment)

| <u>plant</u> | <u>treatment</u> |
|--------------|------------------|
| Col          | H <sub>2</sub> O |

|    |            |                           |
|----|------------|---------------------------|
|    | Col        | Pst                       |
|    | Col        | Pst/avrRpt2               |
|    | Col        | Pst/avrB                  |
|    | Col        | Psm                       |
| 5  | Col        | Psm/avrRpt2               |
|    | Col        | Psp (not 9 hours)         |
|    | Col        | Psp/avrRpt2 (not 9 hours) |
|    | Col NahG   | Pst                       |
|    | Col NahG   | Pst/avrRpt2               |
| 10 | Col NahG   | Pst/avrB                  |
|    | Col ndr1-1 | Pst                       |
|    | Col ndr1-1 | Pst/avrRpt2               |
|    | Col ndr1-1 | Pst/avrB                  |

### Results

Four hundred sixty-five genes were specifically/preferentially induced in the incompatible interaction (WT and Pst/avrRpt2), and 616 genes were specifically/preferentially repressed in the incompatible interaction. Examples of these genes are provided in Tables 10 and 13. Gene expression patterns in the incompatible interaction in Col and Ws were significantly different, indicating that the genetic diversity among ecotypes can affect gene regulation during the incompatible interaction significantly. In comparison, a relatively small number of genes (314 genes for induction, 167 genes for repression) were affected at this time point during the compatible interactions (but not preferential to the incompatible interactions). A comparison of the results in three genetically different compatible interactions (WT and Pst, rps2 and Pst, rps2 and avrRpt2) revealed that 25 genes were repressed in an avrRpt2-dependent manner (see Table 9). Thus, the expression of genes comprising SEQ ID NOs:1, 15, 19, 20, 24, 26, 27, 34, 38, 51, 56, 59, 67-69, 99, 116, 155, 159, 182, 212, 284, 372, 444, and 789 is downregulated (repressed) in an avrRpt2-dependent manner in *Arabidopsis*. These genes are good candidates to be involved in avrRpt2 virulence functions (in rps2 plants).

Genes that were induced in rps2 plants after infection irrespective of avrRpt2 indicate a function of RPS2 other than an interaction with avrRpt2. Thus, global gene expression profiling can identify large and minor trends in gene regulation and is useful in gene discovery.

One general phenomenon when plants are resistant to a pathogen is the early response of pathogen-responsive (induced or repressed) genes compared to plants that are susceptible to



infection. This has been proposed based on observing expression of a very limited number of genes, but it has not been proven as a global trend. To examine the results from early incompatible interactions and late compatible interactions, 4 week old Col-0 plants with well expanded leaves were infected with a high dose ( $OD_{600} = 0.02$ ) or low dose ( $OD_{600} = 0.002$ ) of *P. syringae* and samples collected at 6 or 30 hours, respectively. The two expression patterns were similar. The correlation values between the late compatible and incompatible interaction at either 6 hours, 9 hours or the average of 3-9 hour time points was 0.71, 0.72 and 0.75, respectively.

The majority of genes that did not respond within 9 hours after infection of a virulent strain but that responded in 30 hours (Pst or Psm, for *Pseudomonas syringae* pv. tomato DC3000 and *Pseudomonas syringae* pv. *maculicola* ES4326, respectively; the plant is susceptible to these strains) responded within 6 hours after infection of an avirulent strain (Pst/avrRpt2; Pst carrying the avirulence gene *avrRpt2*; the plant is resistant to this strain). This strongly suggests that early response of the pathogen-responsive genes is crucial for the plant to be resistant.

A comparison of the differences in the expression patterns of the 2 primary ecotypes of *Arabidopsis* response to infection provides a further way to identify which genes have a more universal role (unchanged expression pattern) and which may be very specific to a particular plant ecotype involved in a very specific gene-for-gene interaction. For example, responses that are common between two ecotypes may be important for resistance. Genes that show the same pattern in both ecotypes may be part of more universal, or commonly-used, mechanisms involved in plant-pathogen interactions. Responses that are different may indicate that the two ecotypes use different combinations of responses to achieve resistance. This implies that a variety of genes can participate in plant-pathogen interactions. Nevertheless, ecotype-specific responses are expected to have counterparts in other plant species.

The differences in resistance response between ecotypes can be used for improving resistance in plants. In responses that are different between ecotypes, using the methods and compounds of the invention, such a response can be added to (induced or repressed) the response seen in the ecotype which does not normally use that response. This will likely give the plant a more robust or a wider range of resistance.

Table shows a comparison of gene expression in 4 ecotypes, i.e., Col-0, Ws-2, Cvi and Ler in response to infection. Table 10A shows the expression data for 9 probe sets corresponding to genes that are specifically induced at 3 hours after incompatible infection of

four different ecotypes of *Arabidopsis* with *P. syringae* pv. tomato DC3000. Table 10B shows expression data for 18 probe sets corresponding to genes that are induced by 6 hours but not at 3 hours after incompatible infection of four different ecotypes of *Arabidopsis* with three different bacterial strains, i.e., *P. syringae* pv. tomato DC3000. Table 10C illustrates the expression data for 6 probe sets corresponding to genes that are activated by *P. syringae* at 6 hours post-infection. Most of the genes are compatible interaction-specific or -preferential.

Four week old plants with fully expanded leaves were infected and samples collected at 3 or 6 hours post-infection ( $OD_{600} = 0.02$ ). Some common patterns were observed. At 3 hours after infection of an avirulent strain, Pst/avrRpt2, the overall qualitative gene expression patterns were very similar for all the ecotypes tested. Common responses to Pst/avrRpt2 could be important for gene-for-gene resistance and so may be useful to identify targets for reverse genetics. Quantitative and qualitative differences in the response were noted, indicating that there are qualitative and/or quantitative differences in the signal transduction mechanisms that regulate the response among the ecotypes. Such signal transduction mechanism differences are attributed to genetic differences among the ecotypes.

In particular, early inducible genes (3 hours) in the incompatible interaction were identified (70 genes are common in all the ecotypes, and 360 genes if selected for induced in at least one ecotype). One group of the early genes (38 genes in Col) were repressed to the control level by 6 hours. These genes did not respond in the compatible interaction at 3 hours and were repressed below the control level in the compatible interaction by 6 hours. This suggests that shutting down these genes in the incompatible interaction by 6 hours could be caused by defense response inhibiting factor(s) delivered by bacteria. Another group of the early genes were expressed even higher at 6 hours in the incompatible interaction. One hundred eighty-eight genes showed significant induction or repression at 3 hours in the compatible interaction in at least one of the ecotypes. Of these, 3 induced genes and 3 repressed genes were induced or repressed in all three ecotypes.

At 3 hours in the incompatible interaction, a major difference among the ecotypes was quantitative; overall expression patterns were very similar, but overall fold change amplitudes were clearly in the order of  $Ws > Col > Ler$ . Thus, in this type of analysis it is not appropriate to analyze datasets by comparing the genes from different datasets that are selected by a certain cut-off value (e.g., 2.5-fold difference). This fold change difference was mainly caused by differences in the basal expression of these genes. In fact, a strong negative correlation in each gene was found between the relative basal expression level in *Ws* (relative to the other

ecotypes; Pearson correlation  $-0.78$ ) and response in the incompatible interaction (especially at 3 hours) and a moderate positive correlation between the relative basal expression level in Ler and response in the incompatible interaction (Person correlation  $0.38$ ) (almost no correlation for the relative basal expression level in Col; Person correlation  $0.10$ ). These observations  
5 indicate that Ws has the tightest regulation of these incompatible interaction-responsive genes, and Ler has the loosest. Another interesting observation is that the relative susceptibility to a virulent strain (Pst) was in the order of Ws>Col>Ler. Although it is unknown whether these two phenomena are controlled by same gene(s), it is conceivable that leaky expression of early response genes (in Ler) confers relative resistance to a virulent strain. At 6 hours in the  
10 incompatible interaction, the gene expression pattern for Col was significantly different from the other ecotypes.

Moreover, different ecotypes may use a different but overlapping set of responses to achieve resistance against the same pathogen. Gene expression profiling can thus reveal ecotype differences. Therefore, it is possible to isolate the genes responsible for these  
15 differences in regulatory mechanisms using ecotype differences in gene expression as a phenotype, by a map-based cloning approach.

For example, a majority of the incompatible response-inducible genes have lower basal levels in ecotype Ws and higher basal levels in ecotype Ler. Among the numerous genes, a few genes that display large differences in the basal level in two ecotypes are chosen. The  
20 large differences in expression level constitute easy-to-score phenotypic markers. Ws and Ler are crossed to obtain F2 populations. The larger the F2 population is, the better resolution in the map position can be obtained. For each of the F2 plants, expression levels of the chosen phenotypic marker genes are measured and physical markers that distinguish these ecotype genomes are scored. The map position of the responsible gene is determined by analyzing the  
25 linkage between the phenotype and the physical markers. If more than a single gene is responsible for the ecotype difference and each of the genes has a quantitative effect on the phenotype, quantitative trait locus (QTL) analysis can be used for mapping. Instead of using F2 populations, the use of recombinant inbred lines (RILs) between the ecotypes of interest may facilitate the analysis, especially using RILs that are already mapped for recombination  
30 points. Once the gene(s) responsible for the phenotype is mapped, a combination of increasing the map resolution, sequencing the chromosomal region identified by mapping in both ecotypes, and gene transfer from one ecotype to the other leads to isolation of the gene.

If the phenotype of interest in gene expression depends on bacterial infection, such as expression of ecotype Col-specific inducible genes at 6 hours after infection of Pst/avrRpt2, expression of the corresponding phenotypic marker genes (e.g., genes that show good difference in induction between Col and Ler) can be measured at an appropriate time after  
 5 bacterial infection.

Differences in gene expression patterns between two virulent strain backgrounds (Pst and Psm) are relatively small. Gene expression patterns for Pst/avrRpt2 and Pst/avrB were quite similar at 3 hours, but the difference increased at 6 hours. Psp (no avr) shows similar expression pattern to incompatible bacteria although the amplitude of fold difference was  
 10 smaller in general. This suggests that Psp, which does not induce the HR in the plant, is still recognized by the plant and induce major part of the defense response seen during the incompatible interaction. It also suggests that plants monitor the effect of the defense response and that if it seems effective (bacteria do not grow like Psp), the plant does not go for a full-blown defense response.

#### 15 Preferred Genes

Preferred early inducible genes were selected as induced > 2.5 fold (except for 2 fold for Psp at 6 hours) in all of the following datasets: Pst/avrRpt2 at 3 hours in Col, Ws, and Ler; Pst/avrRpt2 at 3 hours, Psm/avrRpt2 at 3 hours, Psp at 6 hours, and Pst/avrB at 3 hours, relative to the water control, as well as estradiol-inducible (avrRpm1 at 0, 45, and 120 minutes  
 20 and avrRpt2 at 0, 45, and 120 minutes, where the fold change was relative to the appropriate resistance gene mutant carrying the same transgenes. Among these genes, the genes were ranked according to genes that are not induced by SA or BTH and not induced in late time points with Psm. Regulatory genes were given higher rankings (see Table 11). Hence, the expression of genes comprising SEQ ID NOs:17, 70, 76, 81, 84, 109, 123, 144, 160, 230, 265,  
 25 268, 269, 271, 323, 333, 385, 427, 428, 430, 457, 505, 569, 597, 602, 606, 616, 708, 730, 741, 812, 862, and 942 is induced early after infection of different *Arabidopsis* ecotypes with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121 (at 3 or 6 hours) or is estradiol inducible (at 45 or 120 minutes).

Preferred early repressible genes were selected as repressed > 2.5 fold (except for > 2 fold for Psp at 6 hours) in all of the following datasets: Pst/avrRpt2 at 3 hours, Psm/avrRpt2 at 3 hours, Psp at 6 hours, and Pst/avrB at 3 hours) and Pst/avrRpt2 at 3 hours in Col (the fold change was relative to the appropriate water controls). Among them, the genes were ranked in order of expression (highest to lower levels of expression) (see Table 12). Thus, the  
 30

expression for genes comprising SEQ ID NOs:30, 73, 282, 541, 640, 679, 761, 870, 917, and 930 is repressed early after infection of *Arabidopsis* with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121.

Other genes are induced/repressed during incompatible interactions at 3 and/or 6 hours after inoculation of bacteria. Preferred genes in this group were selected as induced/repressed > 2.5 fold in the incompatible interaction compared to water inoculated control and 2 > fold compared to the corresponding compatible interaction at 3 and/or 6 hours after inoculation with Pst/avrRpt2 and Pst/avrB, and Psm/avrRpt2 and Pst/avrRpt2, in all four ecotypes (see Tables 13a and 13b). Hence, the expression of genes comprising SEQ ID NOs:21, 44, 46, 60, 86, 91, 93, 106, 110, 119, 122, 130, 131, 161, 166, 167, 168, 171, 176, 200, 203, 213, 225, 227, 248, 261, 262, 266, 274, 285, 300, 301, 302, 320, 326, 341, 345, 348, 349, 360, 366, 378, 615, 618, 406, 409, 422, 425, 441, 443, 446, 449, 454, 461, 475, 476, 485, 500, 511, 512, 527, 533, 543, 545, 549, 550, 552, 567, 575, 590, 608, 611, 625, 643, 656, 659, 666, 668, 671, 680, 690, 704, 706, 711, 721, 728, 738, 757, 791, 807, 811, 813, 827, 857, 864, 868, 875, 881, 893, 901, 905, 908, 912, 939, 941, 951, and 952 is induced in an incompatible interaction at 3 and/or 6 hours after infection of four *Arabidopsis* ecotypes with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121, while the expression of genes comprising SEQ ID NOs:7, 33, 82, 136, 141, 154, 185, 189, 199, 202, 434, 471, 483, 499, 516, 530, 578, 586, 631, 658, 694, 714, 718, 734, 770, 772, 816, and 916 is decreased in an incompatible interaction at 3 and/or 6 hours after infection of four *Arabidopsis* ecotypes with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121.

Garlic T-DNA insertion lines corresponding to these genes are searched by BLAST. Global expression profiling after infection with one of two different pathogens (*P. syringae* and *Alternaria brassicicola*) may be employed as a phenotyping method. Transgenic plants for overexpression, underexpression, and conditional overexpression of selected genes are also prepared.

#### Example 4

##### Promoters of Genes Responsive to Pathogen Infection

In many cases the major outcomes of plant-pathogen interactions are largely determined by how plants react in an early stage. Therefore, it is useful to isolate promoters

that rapidly react to pathogen attack for use in expressing proteins that provide tolerance or resistance to pathogen attack.

Genes were selected according to the conditions described below based on the results of a GeneChip™ analysis. These genes were particularly selected for a high level of induction in the *avrRpt2-RPS2* interaction and for a very low mRNA level in the absence of pathogen attack among four *Arabidopsis* ecotypes tested (Col, Ws, Ler, and Cvi). The genes were also analyzed to determine if their expression was similar in other combinations of incompatible interactions (three different bacterial strain backgrounds: *P. syringae* pv. tomato DC3000, *P. syringae* pv. *maculicola* ES4326, and *P. syringae* pv. *phaseolicola* NP3121; three different avirulence genes: *avrRpt2*, *avrB*, and *avrRpm1*; and direct expression of avirulence genes in plants using an estradiol-inducible system). For each gene, the 1.2-kb sequence upstream of the initiation codon is provided in SEQ ID NOs: 1047-1095.

#### Preferred Highly Inducible Promoters

Promoters were selected that had low basal expression level (i.e., uninduced level) in all the ecotypes (Col, Ler, Ws, and Cvi) and high inducibility in Col. Five such promoters of genes represented by the probe sets in Table 14 were identified: the promoters of germin precursor-like oxalate oxidase gene, extra-large G protein gene, PR-1, EREBP5 gene, and a C2H2-type zinc finger protein gene were chosen. The promoters for the germin-precursor like oxalate oxidase gene and PR-1 gene are relatively slow response promoters (no induction 3 hours after infection), but have high induction by 6 hours. The extralarge G protein gene is an intermediate in terms of response time, but maintains high expression over time. The other two are useful as early transient response promoters (good induction by 3 hours, but shut down by 6 hours) in the incompatible interaction (wild type plant infected with Pst/*avrRpt2*).

Promoter sequences comprising SEQ ID NOs:1046-1095 and 1047-1055 correspond to genes comprising one of SEQ ID Nos: 17, 21, 80, 81, 109, 156, 174, 176, 221, 227, 296, 302, 303, 306, 333, 340, 360, 500, 505, 524, 575, 601, 602, 614, 628, 687, 733, 782, 811, 835, 862, 900, 905, 912, and 109, 306, 524, 600, 875, 912, 913, 941 and 942, respectively. Promoter-LUC reporter fusions are prepared and tested in a transient expression system using biolistic co-bombardment of *avrRpt2* gene.

#### Promoters Responsive to Particular Pathogens

Proteins that are useful for protecting plants from pathogen attack may have deleterious effects on plant growth if expressed constitutively. Consequently, it is desirable to have

promoter sequences that control gene expression in such a way that expression is absent or very low in the absence of pathogens, and high in the presence of pathogens.

Wild-type *Arabidopsis* plants (ecotype Columbia) were either mock-infected or infected with the bacterial pathogen *Pseudomonas syringae* pv. *maculiola* strain ES4326 (2 x 10<sup>4</sup> cfu per square centimeter of leaf). After 30 hours, samples were collected, and RNA was purified. This procedure was repeated three times independently, and the RNAs from corresponding samples were pooled, in order to reduce the impact of variation due to uncontrolled variables. The two pools of RNA representing mock-infected and infected plants were then used for gene expression profiling using an *Arabidopsis* GeneChip®. This entire procedure was repeated three times, yielding three sets of GeneChip® data representing a total of nine independent experiments.

To identify promoter sequences that are likely to be useful for driving expression of transgenes in plants in response to pathogen attack, genes were selected whose expression level was less than 40 in all of the mock-infected samples and whose expression level was greater than 400 in all of the infected samples. The value of 40 was chosen arbitrarily as a low expression level and the value of 400 was chosen arbitrarily as a reasonably high expression level. Thirty-seven genes met these criteria and promoter sequences could be identified for 36 of them. Table 15 indicates the identifying probe set number for these 36 genes, the corresponding *Arabidopsis* gene, the mean expression level of each gene in mock-infected plants, the mean expression level of each gene in infected plants, and the fold induction in expression of each gene after infection. For 11 genes, expression in mock-infected plants was undetectable, so it was not possible to calculate fold induction. Therefore, the expression of genes comprising SEQ ID NOs:104-106, 119, 123, 129, 131, 151-152, 183, 191, 198, 200, 227, 249, 274, 302, 358, 415, 481, 547, 566, 582, 628, 633, 639, 656, 673, 793, 818, 827, 864, 874, 880, and 904-905 is induced in *Pseudomonas syringae* pv. *maculiola*-infected *Arabidopsis*.

It is possible that promoters that strongly activate gene expression in response to infection by a bacterial pathogen might be different from promoters that strongly activate gene expression in response to infection by a fungal pathogen. To test this possibility, a second GeneChip® experiment was conducted, in which wild-type *Arabidopsis* plants (ecotype Columbia) were mock-infected or infected with the fungus *Botrytis cinerea*. Samples were collected at 0, 12, 36, 60, and 84 hours after infection, RNA was purified and used for expression profiling using an *Arabidopsis* GeneChip®. To identify useful promoters, genes

were selected whose expression level was less than 40 in mock-infected samples from all time points and whose expression level was greater than 400 in infected plants at 84 hours after infection. Twenty-three genes met these criteria, and promoter sequences could be identified for 21 of them. These genes are described in Table 16, with their identifying probe set  
 5 number, the corresponding *Arabidopsis* gene, the mean expression level of each gene in mock-infected plants, and the expression level of each gene in infected plants at various times after infection. Among these 23 genes, 11 genes were previously identified in the search for genes whose expression was strongly induced by *Pseudomonas syringae* infection. These 11 genes correspond to identifying codes 12989, 13015, 13100, 13215, 13565, 14609, 16649, 16914,  
 10 19284, 19991, and 20356. Hence, the expression of genes comprising SEQ ID NOs:18, 71, 119, 123, 129, 151, 191, 244, 245, 302, 545, 547, 562, 566, 637, 653, 747, 756, 774, 793, 842, 864, and 905 is induced in *Botrytis cinerea*-infected *Arabidopsis*.

The promoter sequences for the 25 genes that were only identified in the *P. syringae* data set are shown in SEQ ID NOs:1001-1025. The promoter sequences for the 10 genes that  
 15 were only identified in the *B. cinerea* data set are listed in SEQ ID NOs:1026-1035) The promoter sequences of the 11 genes that were identified in both data sets are listed in SEQ ID NOs:1036-1046. The 11 promoter sequences that were identified in both data sets are most likely to be useful for driving expression of transgenes in response to attacks by various pathogens, as these promoters are activated in response to attack by either *Pseudomonas*  
 20 *syringae* or *Botrytis cinerea*, two very different pathogens. The other promoters may also be useful for driving expression of transgenes that are efficiently expressed in response to infection by certain types of pathogens.

Further, orthologs of the *Arabidopsis* promoters are also useful to drive expression of transgenes. To identify the orthologous promoter, a BLAST search for orthologous genes was  
 25 conducted. To identify the ortholog, the alignments from the BLAST search are used to determine the range of nucleotides showing homology to the *Arabidopsis* gene. The coding sequences shown at the beginning of each search result that contain regions corresponding to the nucleotides showing homology are likely orthologous genes. Orthologous promoter sequences may be isolated by any method known to the art, e.g., cloning of genomic DNA 5'  
 30 to the ATG in orthologous genes identified in a computer assisted database search or hybridization of a probe comprising any one of SEQ ID NOs:1001-1046 to genomic plant DNA.



### Example 6

#### Genes the Expression of Which Are Altered by Viral Infection

To identify host genes that are commonly up or down regulated during local RNA or DNA virus infection, gene expression profiling was employed. The host genes may include host factors that are induced by viral infection, e.g., activated host defense genes, suppressed by viral infection, e.g., suppressed host defense genes, genes involved in symptom development, as well as genes regulated by virus inducible promoters. Once the genes are identified, the function of each is then determined. Reverse genetics is then employed to examine the effect of mutations on these genes during virus infection.

#### 10 Experimental Procedure

*Arabidopsis thaliana* (Columbia-0 (Col-0)) were grown in a Conviron growth chamber to 4 weeks of age. The growth conditions were 22°C, 12 hour day length and 75% relative humidity. At least four rosette leaves of twenty plants were inoculated with one of five viruses or a mock control (120 plants total). The viruses were turnip vein clearing virus (TVCV), a tobamovirus, an oil seed rape mosaic virus (ORMV), a tobamovirus, tobacco rattle tobravirus (TRV), a tobamovirus, cucumber mosaic virus strain Y (CMV-Y), a cucumovirus, and turnip mosaic virus (TuMV), a potyvirus. Each virus was diluted to approximately 0.5 to 1.0 µg/ml in 10 mM potassium phosphate buffer pH 7.2 (or 20 mM Tris-HCl pH 8.0 for the TuMV). The phosphate buffer was used as the mock infection control for the experiments. Inoculated Col-0 leaves were first dusted with carborundum then 10 µl of virus solution or phosphate buffer were pipetted onto the leaf surface. The virus solution or phosphate buffer alone were then rubbed into the leaf surface using a gloved finger, and the leaf surfaces were washed with distilled water at about 10 minutes post inoculation.

Inoculated leaf tissue was removed from each plant at 1, 2, 4 and 5 days post inoculation (dpi), weighed, snap frozen in liquid nitrogen and stored at -80°C. Total RNA was extracted from leaf tissue by the *RNAwiz* method (Ambion, Inc.) and further purified using the *RNeasy* method (Qiagen, Inc.). RNA was diluted to 1 µg/ml and labeled as a probe for Affymetrix GeneChip hybridization according to Affymetrix protocol for synthesizing labeled copy RNA (cRNA) (see Example 1). Labeled cRNA for each virus or mock treatment was hybridized to an Affymetrix GeneChip containing sequences corresponding to 8775 *Arabidopsis* genes. The hybridization data was then analyzed using Affymetrix GeneChip software.

*Arabidopsis* genes that were induced by at least 2-fold in all virus treatments were identified by importing the data into Microsoft Excel and then subjecting the data to selection criteria. Within each time point, the expression level of a gene exceeded 25 and the fold change was greater than 2 by comparison with the mock-infected treatment. Thus, for genes  
 5 that were induced by all five viruses, the expression level exceeded 25 and the fold change was greater than 2 for all five viruses. For genes that were repressed by at least 2-fold, the expression level of the gene must exceed 25 in the mock-infected treatment and the fold change must be less than 2 in all of the five virus treatments.

### Results

10 A gene chip from Affymetrix having oligonucleotides corresponding to approximately 8,100 *Arabidopsis* genes was used with labeled cRNA obtained from plant cells infected with a selected viruses at different days post-infection (dpi). For example, for *Arabidopsis*, the RNA may be obtained from *Arabidopsis* infected with potyvirus, tobamovirus, tobnavirus, cucumovirus or geminivirus. After hybridization, laser scanning is employed to detect  
 15 expression levels and the data obtained is then analyzed. For genes that are induced in response to viral infection, genes that are expressed at levels greater than, for example, 2 fold over control, are selected. Alternatively, for genes that are suppressed in response to viral infection, genes that are expressed at levels lower than control are selected. The advantages of a gene chip in such an analysis include a global gene expression analysis, quantitative results, a  
 20 highly reproducible system, and a higher sensitivity than Northern blot analyses. Moreover, a gene chip with *Arabidopsis* DNA has a further advantage in that the *Arabidopsis* genome is well characterized.

Data obtained from probe sets which correspond to genes upregulated or downregulated in response to infection by all 5 viruses revealed forty-six genes that were  
 25 downregulated and 126 that were upregulated in response to viral infection (Tables 17 and 18). Once the induced and/or suppressed genes are identified, the functions of the genes are then characterized by standard methodology.

Therefore, the expression of genes comprising SEQ ID NOs:14, 48, 53, 98, 217, 226, 295, 327, 343, 352, 369, 404, 407, 418, 453, 458, 465, 472, 480, 488, 495, 507, 509, 513, 514,  
 30 559, 561, 581, 604, 607, 613, 641, 652, 672, 720, 735, 739, 743, 745, 754, 773, 803, 832, 849, 948, and 949 is downregulated after viral infection, and the expression of genes comprising SEQ ID NOs:3, 51, 54, 60, 61, 66, 75, 76, 78, 88, 95, 96, 101, 106, 108, 123, 126, 128, 129, 131, 137, 145-147, 150, 158, 169, 170, 172, 173, 197, 200, 216, 219, 224, 230, 233, 237, 249,

250, 263, 274, 275, 276, 299, 307, 323, 333, 342, 346, 359, 382, 383, 387, 391, 393, 401, 411, 415, 427, 442, 455, 459, 466, 477, 481, 485, 487, 502, 511, 515, 525, 534, 539, 542, 560, 571, 577, 579, 584, 587, 595, 600, 627, 638, 645, 654, 659, 668, 681, 688, 695, 696, 706, 708, 730, 742, 753, 775, 785, 786, 792, 797, 800, 801, 809, 817, 819, 820, 823, 827, 847, 856, 875, 885, 896, 902, 910, 921, 922, 923, 925, 926, 928, 946, and 952 is upregulated after viral infection.

Rice and other plant orthologs of these *Arabidopsis* sequences were identified as described in Example 14 below. The results are summarized in Tables 22 and 23.

A correlation of the SEQ ID NOs:1-953 and the probe set designations corresponding to genes, the expression of which is altered after infection of *Arabidopsis* with a pathogen is shown in Table 19.

### Example 7

#### Identification of Gene Products that are Modulated upon Infection

##### of a *Chenopodium* Cell with a Virus

Of the many disease resistance mechanisms that can be studied, the HR (hypersensitive resistance) system of *Chenopodium* spp. is attractive because of the broad-spectrum virus resistance it confers. This is shown by the ability of members of the bromo-, como-, cucumo-, ilar-, alfamo-, nepo-, sobemo-, tombus-, tymo-, carla-, clostero-, hordei-, potex-, poty-, tobra- and tobamovirus groups to elicit local lesion HR on *Chenopodium* spp. (CMI/AAB Description of Plant Viruses, 1984; Cooper et al., (1995)). In many instances, the HR completely blocks viral spread. However, certain viruses can break through the hypersensitive response and move from one species of *Chenopodium* to another. The ability of some viruses to infect more than one species of *Chenopodium* provides an opportunity to isolate genes that provide a cell with resistance to viral infection.

The genetic mechanisms of *Chenopodium* spp. HR involve a number of factors. These factors can be studied to further understand the hypersensitive response and the mechanism through which the response acts. There are some similarities between the products of *Chenopodium* spp. genes and gene products involved in common defense signaling pathways in other plants. These similarities allow comparisons to be made between *Chenopodium* and these other plants. One example includes genes that are induced upon viral infection during HR in *C. foetidum* (Visedo et al., (1990)).

Additionally, some circumstantial experimental evidence suggests that *Chenopodium* HR may be somewhat similar to tobacco *N* gene HR (Whitham et al., 1994). Movement defective tobacco mosaic tobamovirus (TMV) replicates within an inoculated cell of a tobacco plant with an *N* gene, but fails to move from cell to cell (Cooper et al., 1996). Hypersensitivity is not induced, thus replication alone is not sufficient to induce HR despite the *N* gene elicitor being mapped to the replicase gene of TMV (Padgett and Beachy, 1993). Therefore, the process of virus movement may trigger hypersensitivity, which implicates intercellular signaling in this type of HR. Support for this position comes from experiments in which cell-to-cell contacts were disrupted in *N* gene tobacco which resulted in the prevention of necrotic lesion formation in infected leaves (Gulyas and Farkas, 1978). Likewise, TMV will not induce HR cell death in NN tobacco protoplasts where plasmodesmata are not intact (Otsuki et al., 1972), although HR does occur in callus cultures where plasmodesmata are intact (Beachy and Murakishi, 1971). By comparison in *C. quinoa*, movement defective brome mosaic bromovirus (BMV) replicates but fails to move from cell to cell. Initial infection is not sufficient to induce HR since local lesions do not form (Schmitz and Rao, 1996). Similarly, in *C. amaranticolor*, cucumber mosaic cucumovirus (CMV) lacking a movement protein replicates within inoculated cells, fails to move and does not elicit cell death (Canto and Palukaitis, 1999). Therefore, like TMV on *N* gene tobacco, the process of viral spread of BMV and CMV in *C. quinoa* and *C. amaranticolor* may induce HR.

## 20 Methods and Materials

### Inoculation of Plants

Leaves of 10-week old *C. amaranticolor* or *C. quinoa* were inoculated with *in vitro* transcripts of TMV-MGFus (Heinlein et al., 1995), TMV virions, tobacco rattle tobravirus (TRV), or they were mock-inoculated. TMV-MGFus encodes GFP (green fluorescent protein) fused to the viral movement protein. Infectious spread can be monitored through the detection of GFP. Using an Olympus stereomicroscope fitted with a U-ULH Olympus lamp, infected *C. amaranticolor* tissue accumulating GFP was excised at 4, 7 and 11 days after inoculation (dai). Leaves inoculated with TRV or TMV were collected at 4 dai, at which point local lesions were forming. Mock-inoculated tissue was collected at the same time. Tissue was frozen in liquid nitrogen and total RNA was purified from it. Three separate sets of plants were inoculated with TMV-MGFus and yielded three independent preparations of RNA.

cDNA-AFLP (complementary DNA-amplified fragment length polymorphism)

Poly-A+ RNA was isolated from TMV-MGfus infected *C. amaranticolor* using Qiagen's Oligotex mRNA purification system (Qiagen, Valencia, CA) and cDNA was generated using cDNA synthesis reagents from Life Technologies (Rockville, MD). cDNA was used to generate AFLP fragments with the AFLP reagents from Life Technologies and reactions were performed according to the manufacturer's instructions. cDNA made from one microgram of poly-A+ RNA was digested with EcoRI and MseI and the supplied compatible linkers were ligated to the ends of the digested molecules. A few modifications were introduced. EcoRI-NN primers (GACTGCGTACCAATTCNN; SEQ ID NO:2134), rather than EcoRI-NNN, were used with the 5' fluorescent label NED (Applied Biosystems, Foster City, CA) and MseI-N and MseI-NN [GATGAGTCCTGAGTAAN(N); SEQ ID NO:2135], rather than MseI-NNN, primers were used (Genosys, The Woodlands, TX), to reduce the complexity of the primer sets evaluated. All possible primer combinations (256 + 64) were used for PCR amplification and products were separated on polyacrylamide gels and visualized using a Genomix SC fluorescent scanner (Beckman Coulter, Fullerton, CA). Gene fragments that appeared to be upregulated in infected tissues compared to mock-inoculated tissues were tested to see if they were also upregulated by the same primers from a second preparation of cDNA from RNA from a second set of infected plants. Gene fragments that were upregulated in both RNA preparations were excised from the gel, eluted from the gel in water and reamplified by PCR using the appropriate MseI and EcoRI primers and sequenced with 377 ABI sequencers (Applied Biosystems) using dideoxysequencing methods.

#### Quantitative RT-PCR

DNase treated total RNA (2 ng per reaction) from the third independent preparation of TMV-MGfus infected *C. amaranticolor*, the first preparation of TRV infected *C. amaranticolor*, or the first preparation of TMV *C. quinoa*, was used with TaqMan One-Step RT-PCR reagents for quantitative analysis in an ABI 7700 (Applied Biosystems). Reactions were performed according to the manufacturer's instructions. Primers and 6-FAM 5' end-labeled probes (6-carboxyfluorescein, Applied Biosystems or Genosys) were designed from the sequences from the *C. amaranticolor* upregulated gene fragments using Primer Express software (Applied Biosystems) and are listed in SEQ ID Nos:954-1000 and 2130-2135. Expression levels were interpolated from standard curves with a correlation coefficient of 0.99 or greater and the quantities were normalized to the expression level of actin in each sample.

## Results

The interaction of the elicitor and the R gene product establishes a cascade of reactions and signaling events that is then manifested in a phenotypic HR. In essence, HR is the end result of disease activated signaling events. In order to detect the early expression of genes induced by viral infection, it was necessary to isolate infected tissue before the onset of local lesion formation. Therefore, *C. amaranticolor* was infected with RNA transcripts of TMV-MGfus that express GFP (green fluorescent protein) in infected cells. This allowed the spread of viral infection to be monitored over time. Infection foci comprising over 100 cells could be detected at 4 dai and foci of more than 500 cells could be detected at 7 dai. There was no visible appearances of cell death or chlorotic local lesion formation at the infection foci at 4 and 7 dai. By 11 dai, the infection foci were associated with chlorotic local lesions. Virus infected tissue was excised from leaves at each time point and RNA was purified from the tissue and used for cDNA-AFLP as previously described.

cDNA-AFLP fragments were separated on polyacrylamide sequencing gels and imaged with a fluorescent scanner. Samples derived from mock-inoculated tissue at 7 dai were run next to samples derived from TMV-MGfus infected tissue at 7 dai for comparison. Ninety-eight bands having intensity in the TMV-MGfus lanes that was greater than that of analogous bands in the mock lanes were easily detected. Thirty out of the 98 bands were also upregulated in an independent set of experiments designed to reduce biological variation between experiments. These bands were excised from the gel, reamplified, and sequenced.

The hypothetical protein sequences derived from the reamplified fragments (SEQ ID NOs:1954-1966) translated from all six reading frames were compared to sequences in the GenBank protein sequence database. The results of the BLASTX search (Altschul et al., 1997) are summarized in Table 20a. The results of a BLAST search similar to the one described for Arabidopsis ORFs in Example 6 above is summarized in Table 24. To confirm that the expression levels of DESCA genes were upregulated in infected tissue compared to mock inoculated tissue, the relative amount of DESCA and actin transcript in a third independent set of samples at 4 dai, 7dai, and 11 dai was quantitatively measured (Table 20b).

The expression level of DESCA1 increased the most in the TMV-MGfus infected plants. The expression level of DESCA1 increased 200 times by 4 dai but tapered off drastically by 11 dai. DESCA1 is unrelated to any protein known at this time.

Two sequences, DESCA4 and DESCA10, are both related to pumps found in *Arabidopsis* and yeast (Sanchez-Fernandez et al., 1998; Smart and Fleming, 1996). DESCA4

is expressed highly at 4 dai but the expression drops off over time whereas DESCA10 is only moderately induced and its expression returns to normal by the time of the visible appearance of local lesions in *C. amaranticolor*.

DESCA7 is similar to a salicylate-induced glucosyltransferase gene in tobacco  
5 (Horvath and Chua, 1996). DESCA9 is similar to cytochrome P450-like proteins which can produce cytotoxic compounds including phytoalexins that are deployed by a plant to defend against invading microbes. DESCA12 is related to a proanthranilate benzoyltransferase from carnation that plays a direct role in the phytoalexin biosynthesis in carnation (Yang et al., 1998). DESCA11 is similar to the tryptophan biosynthetic enzyme phosphoribosylanthranilate  
10 transferase whose gene expression is induced in the presence of ozone in *Arabidopsis* (Conklin and Last, 1995).

DESCA3 is similar to endo-1,4-beta-glucanases that have a role in fruit ripening, abscission, and cell elongation (Lashbrook et al., 1994). DESCA3 is highly expressed in the infected *C. amaranticolor* and remains highly expressed during the appearance of local lesions  
15 and necrosis.

Many disease responses are mediated by positive regulators such as transcription factors or kinases that initiate signaling cascades for the activation of defense responses. One gene, DESCA5, is loosely similar to a yeast potential transcriptional regulator. DESCA5 expression is twice as high at the early stages of infection compared to the late stages of  
20 infection illustrating an important role played by gene regulation at the early stages of infection. DESCA6 is related to kinases of *Arabidopsis*. Kinases have essential roles in programmed cell death during viral infection (Dunigan and Madlener, 1995). DESCA2 is the most highly expressed of the group suggesting that it is an important regulator at the onset of infection. It is similar to a receptor-like protein kinase in bean that responds to *Fusarium*  
25 *solani* attack (Lange et al., 1999).

Some R genes have kinase-like regions that may function in initiating a signal cascade during the onset of HR (Song et al.; 1995, Zhou et al., 1997). Global amino acid sequence alignment (Henikoff and Henikoff, 1992) of DESCA2 with Pto or Xa21, R genes with ser/thr kinase domains, reveals a 37% similarity. DESCA8 has a nucleotide binding site and a  
30 leucine-rich repeat that is common for many R genes. (Meyers et al., 1999; Leister et al., 1998).

To link DESCA genes to a multivirus resistance pathway, *C. amaranticolor* was inoculated with TRV (tobacco rattle virus), a virus that is taxonomically distinct from TMV.

Local lesions appeared by 4 dai and RNA was purified from the infected leaves. DESCA gene expression levels in infected tissue were compared to mock inoculated tissue by quantitative RT-PCR and revealed that the same DESCA genes upregulated during a TMV infection are also upregulated during a TRV infection (Table 20b).

5       The gene expression levels in TMV infected *C. quinoa* were measured using the same *C. amaranticolor*-derived primers in quantitative PCR to determine if DESCA genes were up-regulated during HR in another *Chenopodium* species. Most of the DESCA genes were upregulated in *C. quinoa* and were expressed at levels many times higher than in *C. amaranticolor* (Table 3). This may be a result of the infection of *C. quinoa* with the  
10       aggressive wild-type virus rather than slower moving TMV-MGfus.

      The experimental procedure presented here can detect any similar gene involved in the aforementioned signaling pathways such as SA signaling. Except for DESCA1, whose expression is increased the most at 200+ fold, many of the fragments have homology to other genes that have been placed in disease resistance pathways in other plants. DESCA12 and  
15       DESCA9 are respectively similar to hypersensitivity related gene 201, possibly a proanthranilate benzoyltransferase, and p450 monooxygenases, both which are expressed during the hypersensitive response in tobacco upon infection with *Pseudomonas solanacearum* but are not regulated by SA (Czemiec et al., 1996). DESCA7 is similar to a salicylate-induced glucosyltransferase gene in tobacco (Horvath and Chua, 1996). Thus, the disease resistance  
20       response in *C. amaranticolor* involves pathways both dependent and independent of SA signaling.

      The surprising discovery of DESCA4, DESCA7, DESCA9, DESCA10, and DESCA12, reveal the underpinnings of an endogenous detoxification system. Briefly, the activation phase involves cytochrome P450 monooxygenases introducing functional groups  
25       (e.g. aromatic rings) to potential toxins. The conjugation phase in plants involves the linking of glutathione or glucose to the toxin at which point the conjugated molecule can be recognized by an ATP-binding cassette transporter and pumped into the vacuole, or possibly the neighboring cells, during the elimination phase. The final phase includes either storage or breakdown of such molecules. DESCA9, similar to cytochrome P450, and DESCA12, similar  
30       to a gene associated with the production of phytoalexin, may produce potential toxins. In fact, *C. amaranticolor* produces many such compounds that are antiviral to TMV. DESCA7, similar to a glucosyltransferase, may conjugate such toxins to be transported by the ABC-transporters encoded by DESCA4 or DESCA10. In this particular case, the transported



compound could then be deployed by the infected plant cell as an antiviral agent or cytotoxic compound, stored by noninfected cells in anticipation of infection, or eliminated by noninfected cells neighboring infected cells. Since all of these genes are induced by TMV and TRV in *C. amaranticolor*, their induced expressions are a result of a specific or general  
5 multiviral or disease resistance pathway.

Possessing the R genes that allow *C. amaranticolor* to initially recognize multiple viruses provides an opportunity to use these genes, and the regulatory elements associated with these genes, to transfer viral resistance to other plants. In addition, possession of genes that produce and transport antiviral and cytotoxic products allows for the transfer of viral resistance  
10 through a mechanism involving induced cell death upon viral infection.

Two genes that may be used for early recognition of viral infection are DESCA8 and DESCA2, as these genes may act as signaling components to initiate the resistance cascade. DESCA8 has a nucleotide binding site and a leucine-rich repeat that is common for many R genes and that can be found in other plants (Meyers et al., 1999; Leister et al., 1998).  
15 DESCA2 is induced in both *Chenopodium* species and is similar to other R genes, Xa21 and Pto, which have similar ser/thr kinase domains.

Resistance to viral spread may be transferred between *Chenopodium* spp. For example, BMV (brome mosaic virus) induces local lesions in the green variety of *C. hybridum*, however lesion formation does not limit the systemic spread of the virus (Verduin, 1978). The  
20 systematic spread of the BMV virus may be restricted in the green variety of *C. hybridum* by transformation with a gene from the purple variety that does limit spread (Komari, 1990). Thus, genes that confer viral resistance may be used for complementation, reverse genetics, overexpression, and gene silencing. Furthermore, as indicated by the functionality of the R genes N and Pto after being transferred into heterologous species, (Whitham et al., 1996;  
25 Rommens et al., 1995), the *Chenopodium* genes may function to initiate hypersensitivity in crops, *Arabidopsis* or other useful plants.

### Example 8

#### Other Plant-Pathogen Interactions

The methods set out hereinabove can be used for any type of comparable resistance interaction. For example any of the following plant/pathogen interactions will be produced as compatible and incompatible interactions. The RNA from such an interaction is isolated and subject to a protocol such as one outlined in Example 1, e.g., using a Genechip with a specific plant's genes or microarray, differential display PCR or cDNA-ALFP (Example 7). A four-way analysis is performed and genes which are expressed differently are identified. The plant/pathogen interactions in Table 21 are well known in the art. However, any type of plant/pathogen interaction that involves this type of resistance can be used.

Genes that are upregulated and cause resistance in a wide variety of plants are particularly useful in methods which upregulate or overexpress the gene. One method is to add an exogenous copy, thus providing more of the gene product or allowing for a different induction from that used by the plant. Alternatively, the endogenous gene can be upregulated using a known inducer or using artificial methods such as using an artificial induction signal in the endogenous promoter. Examples of the two methods are provided in Examples 9 and 10.

Accordingly, embodiments of the invention provide the sequences disclosed herein, which sequences can be used in genetic engineering of crops, as probes and markers to study the dynamics of plant/pathogen interactions, and as markers in marker-assisted breeding protocols to identify plants carrying particularly useful combinations of genes associated with pathogen resistance, as well as in plant defense.

### Example 9

#### Transformation of Resistance Genes into Plants

To produce resistant plants, resistance genes such as those identified herein can be introduced into plant cells to generate transgenic plants having enhanced resistance. While HSF4 is any preferred gene for this embodiment of the invention, the invention can be employed with other genes, alone or in combination, whose regulation is strongly responsive to plant/pathogen interactions, such as the genes identified herein. Since some genes are strongly induced and others are strongly repressed in plant/pathogen interactions, and since some genes that are strongly induced in one ecotype can be strongly repressed in another, the invention contemplates use of any of the genes and sequences, or fragments thereof, disclosed

herein, in a construct adapted to cause overexpression, repression, or knock out, of the genes in a transgenic plant.

Transgenic downregulation of genes associated with pathogen resistance can have several useful applications. In one embodiment, transgenic downregulation of genes that are strongly repressed in resistance interactions can enhance resistance. Such transgenic downregulation can employ the genes disclosed herein, or fragments thereof, in an antisense orientation to interfere with accumulation of the products of those genes. Likewise, any other methodology capable of lowering expression of such genes is also included within these embodiments of the invention.

Plant transformation can be carried out by conventional means, and can include *Agrobacterium*-mediated transformation, electroporation, particle acceleration, abrasion, and any other useful means leading to expression of a transgene in a plant of interest. Transformed plant cell are then used to regenerate one or more plants in tissue culture. Subsequent generations of transgenic plants can be used directly or bred with other lines to generate plants having enhanced pathogen resistance.

#### Example 10

##### Upregulation of Resistance Genes in Crops

Because many or most *Arabidopsis* genes have orthologs in other plants, the genes and sequences disclosed herein are generally useful in constructs to be up-regulated and cause resistance in a wide variety of plants. As examples, the heat shock proteins, and particularly HSF4, are found throughout the plant kingdom. For many such regulatory and responsive genes it is well known that there exist substances that can induce expression. Chemicals such as dexamethasone have been found to induce mammalian HSF proteins. Likewise, a chemical induction of key plant defense genes can be chemically induced. High throughput screening for chemical inducers of the plant HSF4 or other resistance gene is performed. Potentially useful substances are then tested on crop plants and eventually used as a soil additive or sprayed onto plants when needed to induce resistance. Accordingly, embodiments of the invention usefully employ the genes disclosed herein, or fragments thereof, for screening to identify useful chemical inducers and/or repressors of gene responsive to pathogenic infections.

### Example 11

#### Identification of Inducers and Repressors of Resistance Genes

The yeast two-hybrid method and many methods which use its basic idea, provide a technique to identify proteins which interact with a protein of interest. The method relies on the fact that a protein contains domains which can be separated. Thus the protein of interest is fused to the GAL4 DNA binding region of a known protein. The GAL4 (or another) activation signal is fused in a library to produce a library of fused proteins. If one of the proteins from the library interacts with the protein of interest the protein binds and a signal protein is produced, such as luciferase. There are a number of such systems presently, some of which can be used in mammalian cells, allowing for correct processing and folding of certain proteins and others which allow the interaction to occur in the cytoplasm allowing for the identification of other types of proteins.

cDNA from HSF4 and any other protein of interest is cloned in fusion to the yeast GAL4 DNA binding domain on a vector. A library containing cDNA from Arabidopsis is fused to the GAL4 or an activation domain of choice. Expression of luciferase correlates with identification of an interacting protein. This protein is then analyzed as to its action as an inducer or repressor.

### Example 12

#### Determination of the Minimal Promoter Fragment

A promoter sequence as given in SEQ ID Nos: 2137-2661 and 4738-6813, preferably a promoter sequence of one of the genes defined by SEQ ID NOs 536-579 such as SEQ ID Nos: 536; 537; 539-542; 548; 550-553; 555-558; 560; 565-568; 571-576, 578 and 579, or a promoter ortholog thereof is fused to the  $\beta$ -glucuronidase (GUS) gene at the native ATG to obtain a chimeric gene cloned into plasmid DNA. The plasmid DNA is then digested with restriction enzymes to release a fragment comprising the full-length promoter sequence and the GUS gene, which is then used to construct the binary vector. This binary vector is transformed into *Agrobacterium tumefaciens*, which is in turn used to transform *Arabidopsis* plants.

The above plasmid can also be used to form a series of 5' end deletion mutants having increasingly shorter promoter fragments fused to the GUS gene at the native ATG. Various restriction enzymes are used to digest the plasmid DNA to obtain the binary vectors with different lengths of promoter fragments. In particular, a binary vector 1 is

constructed with a 1,900-bp long promoter fragment; a binary vector 2 is constructed with a 1,300-bp long promoter fragment; a binary vector 3 is constructed with a 1000-bp long promoter fragment; a binary vector 4 is constructed with a 800-bp long promoter fragment; a binary vector 5 is constructed with a 700-bp long promoter fragment; a binary vector 6 is constructed with a 600-bp long promoter fragment; and a binary vector 7 is constructed with a 100-bp long promoter fragment. Like the binary vector comprising the full-length promoter fragment, these 5' end deletion mutants are also transformed into *Agrobacterium tumefaciens* and, in turn, *Arabidopsis* plants (for further details of *Arabidopsis* transformation and promoter assay procedures see example 5 above).

The presence of the correct hybrid construct in the transgenic lines is confirmed by PCR amplification.

By using the above protocol it can be determined, which portion of the promoter or the promoter orthologs thereof is required for gene expression.

Minimal promoter fragments having lengths substantially less than the full-length promoter can therefore be operatively linked to coding sequences to form smaller constructs than can be formed using the full-length promoter. As noted earlier, shorter DNA fragments are often more amenable to manipulation than longer fragments. The chimeric gene constructs thus formed can then be transformed into hosts such as crop plants to enable at-will regulation of coding sequences in the hosts.

### Example 13

#### Determination of Promoter Motifs

While a deletion analysis characterizes regions in a promoter that are required overall for its regulation, linker-scanning mutagenesis allows for the identification of short defined motifs whose mutation alters the promoter activity. Accordingly, a set of linker-scanning mutant promoters fused to the coding sequence of the GUS reporter gene are constructed. Each of them contains a 8-10-bp mutation located between defined positions and included in a promoter fragment as given in SEQ ID Nos: 2137-2661 and 4738-6813, preferably a promoter sequence of one of the genes defined by SEQ ID NOs 536-579 such as SEQ ID Nos: 536; 537; 539-542; 548; 550-553; 555-558; 560; 565-568; 571-576, 578 and 579 or the promoter orthologs thereof.

Each construct is transformed into *Arabidopsis* and GUS activity is assayed for 19 to 30 independent transgenic lines. The presence of the correct hybrid construct in transgenic lines is confirmed by PCR amplification of all lines containing the mutant constructs and by random sampling of lines containing the other constructs. Amplified fragments are digested with restriction enzyme (e.g. XbaI) and separated on high resolution agarose gels to distinguish between the different mutant constructs. The effect of each mutation on promoter activity is compared to an equivalent number of transgenic lines containing the unmutated construct. Two repetitions resulting from independent plating of seeds are carried out in every case.

The sequences mutated in the linker-scanning constructs, in particular those that showed marked differences from the control construct, are then examined more closely.

#### Example 14

##### Identifying Orthologs

Orthologs were identified through use of BLAST and SCAN software with some additional filters. For the *Arabidopsis* search, a BLAST database was created that was a subset of GenBank ver 123.0 (released April 15, 2001) that contained all of the plant translated regions excluding *Arabidopsis thaliana* sequences. The subset was created with PERL script. A BLAST search with all of the peptide sequences was performed against the GenBank subset. Each query was executed using the "blastall" command with the parameters "-p blastp", "-v 50", "-b 50", "-F F". The BLAST search results were then processed with SCAN (Sequence Comparison Analysis program, version 1.0k, Los Alamos National Laboratories) using default settings and the orthologs were identified following implementation of an E-value cutoff of  $\leq 1e-4$ . The candidate orthologs were further filtered by comparing words in the description to the text of the annotation fields: product, function and note. The sequence was considered to have the same or similar function if any of the words matched. Words excluded from the filter included: the, like, protein, related, unknown, subunit, hypothetical, and, putative, precursor, clone, homolog, small, beta, class, dna, rna, alpha, gamma, has, not, been, from, to, by, long, type and induced.

For the rice search, amino acid sequences were used that resulted from FGENESH (version 1.C) gene prediction results. The peptide sequences were obtained from gene predictions and formatted into a BLAST database. A BLASTP comparison was then performed against the *Arabidopsis* sequences. The BLASTP results were then filtered through

use of SCAN with the following parameters: "-a 60 60" with an E-value cutoff of  $1e-4$ . This produced orthologs having 60 or more identities and where 60% of the alignments were made up of identities.

- 5 The following pages compile Tables 3 to 24 referred to in the examples above.

**Table 3** Probe Sets corresponding to genes, the expression of which is increased after infection of wild-type *Arabidopsis* with *Pseudomonas syringae*

| ProbeSet                      | Description  | Blast Score | EC # | Family |
|-------------------------------|--|-------------|------|--------|
| 11997_at<br>(AC005967.4_AT)   | gb AAD03372.1  (AC005967)<br>unknown protein [Arabidopsis<br>thaliana]   | 0           |      |        |
| 12002_at<br>(AF069442.47_AT)  | gb AAC79112.1  (AF069442)<br>hypothetical protein<br>[Arabidopsis thaliana]  | 0           |      |        |
| 12004_at<br>(AL022023.132_AT) | emb CAA17771.1 <br>(AL022023) putative protein<br>[Arabidopsis thaliana]   | 8E-86       |      |        |
| 12007_at<br>(Z99708.249_AT)   | emb CAB16829.1  (Z99708)<br>putative protein [Arabidopsis<br>thaliana]   | 1E-168      |      |        |
| 12037_at<br>(AC004005.174_AT) | gb AAC23417.1  (AC004005)<br>unknown protein [Arabidopsis<br>thaliana]   | 0           |      |        |
| 12051_at<br>(AL021889.94_AT)  | emb CAA17133.1 <br>(AL021889) putative protein<br>[Arabidopsis thaliana]   | 1E-143      |      |        |
| 12062_at<br>(AC006069.147_AT) | gb AAD12706.1  (AC006069)<br>unknown protein [Arabidopsis<br>thaliana]   | 0           |      |        |
| 12068_at<br>(AF118223.24_AT)  | gb AAD03449.1  (AF118223)<br>contains similarity to<br>Methanobacterium<br>thermoautotrophicum<br>transcriptional regulator<br>(GB:AE000850) [Arabidopsis<br>thaliana] | 1E-162      |      |        |
| 12072_at<br>(AL035396.4_AT)   | emb CAA23058.1 <br>(AL035396) putative protein<br>[Arabidopsis thaliana]   | 1E-158      |      |        |
| 12079_s_at<br>(A71597.1_S_AT) | emb CAB42594.1  (A71597)<br>unnamed protein product<br>[Arabidopsis thaliana]  | 5E-64       |      |        |
| 12081_at<br>(AC001645.140_AT) | gb AAB63644.1  (AC001645)<br>unknown protein [Arabidopsis<br>thaliana]   | 1E-117      |      |        |
| 12092_at<br>(AC004793.13_AT)  | gb AAD21694.1  (AC004793)<br>ESTs gb T20423,<br>gb AA712864, gb H76323 and<br>gb Z25560 come from this<br>gene. [Arabidopsis thaliana]                                 | 1E-150      |      |        |



| ProbeSet                      | Description   | Blast Score | EC #       | Family     |
|-------------------------------|---|-------------|------------|------------|
| 12115_at<br>(AL033545.26_AT)  | emb CAA22152.1 <br>(AL033545) extensin-like<br>protein [Arabidopsis thaliana]   | 1E-87       |            |            |
| 12136_at<br>(AC007591.60_AT)  | gb AAD39663.1 AC007591_2<br>8 (AC007591) ESTs<br>gb R65145, gb N96612 and<br>gb R90096 come from this<br>gene. [Arabidopsis thaliana] | 1E-60       |            |            |
| 12150_at<br>(AC004005.151_AT) | gb AAC23415.1  (AC004005)<br>unknown protein [Arabidopsis<br>thaliana]  | 5E-32       |            |            |
| 12187_at<br>(AC005489.31_AT)  | gb AAD32893.1 AC005489_3<br>1 (AC005489) F14N23.31<br>[Arabidopsis thaliana]  | 0           |            |            |
| 12198_at<br>(AC006954.90_AT)  | gb AAD23890.1 AC006954_1<br>1 (AC006954) unknown<br>protein [Arabidopsis thaliana]  | 1E-70       |            |            |
| 12203_at<br>(AL021710.268_AT) | emb CAA16738.1 <br>(AL021710) hypothetical<br>protein [Arabidopsis thaliana]  | 7E-55       |            |            |
| 12216_at<br>(AC007119.56_AT)  | gb AAD23641.1 AC007119_7<br>(AC007119) unknown protein<br>[Arabidopsis thaliana]  | 8E-55       |            |            |
| 12217_at<br>(AJ223804.1_AT)   | gb AAF34796.1 AF228640_1<br>(AF228640) lipoamide<br>dehydrogenase precursor<br>[Arabidopsis thaliana]                                 | 0           |            |            |
| 12227_at<br>(AC007576.18_AT)  | gb AAD39285.1 AC007576_8<br>(AC007576) Unknown protein<br>[Arabidopsis thaliana]  | 1E-102      |            |            |
| 12233_at<br>(AJ001807.1_AT)   | emb CAA05023.1 <br>(AJ001807) succinyl-CoA-<br>ligase alpha subunit<br>[Arabidopsis thaliana]   | 0           | EC_6.2.1.5 | synthetase |
| 12278_at<br>(AJ011674.2_AT)   | emb CAA09731.1 <br>(AJ011674) receptor-like<br>protein kinase, RLK3<br>[Arabidopsis thaliana]   | 0           |            | kinase     |
| 12307_at<br>(AC002392.162_AT) | gb AAD12037.1  (AC002392)<br>putative receptor-like protein<br>kinase [Arabidopsis thaliana]  | 0           |            | kinase     |
| 12314_at<br>(AC001229.28_AT)  | gb AAB60922.1  (AC001229)<br>F5I14.14 [Arabidopsis<br>thaliana]   | 0           |            |            |

| ProbeSet                              | Description  | Blast Score | EC #       | Family                |
|---------------------------------------|--|-------------|------------|-----------------------|
| 12317_at<br>(AC004138.27_AT)          | gb AAC32907.1  (AC004138)<br>putative sucrose-proton<br>symporter [Arabidopsis<br>thaliana]                  | 0           |            |                       |
| 12323_at<br>(AC002333.18_AT)          | gb AAB64019.1  (AC002333)<br>putative polygalacturonase<br>[Arabidopsis thaliana]                            | 0           |            | Polygalactur<br>onase |
| 12332_s_at<br>(AB023448.2_S_AT)       | dbj BAA82810.1  (AB023448)<br>basic endochitinase<br>[Arabidopsis thaliana]                                  | 0           |            |                       |
| 12341_s_at<br>(AL021637.176_S_A<br>T) | emb CAA16619.1 <br>(AL021637) vacuolar sorting<br>receptor-like protein<br>[Arabidopsis thaliana]            | 0           |            |                       |
| 12347_at<br>(AC007258.28_AT)          | gb AAD39325.1  (AC007258_1<br>4 (AC007258) Putative<br>ATPase [Arabidopsis thaliana]                         | 0           | EC_3.6.1.- | ATPase                |
| 12349_s_at<br>(X84728.6_S_AT)         | gb AAA17993.1  (M91192)<br>phenylalanine ammonia-lyase<br>[Trifolium subterraneum]                           | 0           |            |                       |
| 12369_at<br>(AC002535.59_AT)          | gb AAC62871.1  (AC002535)<br>putative Na <sup>+</sup> /Ca <sup>2+</sup> antiporter<br>[Arabidopsis thaliana] | 0           |            |                       |
| 12400_at<br>(X98453.1_AT)             | emb CAA67092.1  (X98453)<br>peroxidase [Arabidopsis<br>thaliana]   | 0           |            | peroxidase            |
| 12438_at<br>(AL021710.83_AT)          | emb CAA16723.1 <br>(AL021710) membrane-bound<br>small GTP-binding - like<br>protein [Arabidopsis thaliana]   | 1E-122      |            |                       |
| 12449_s_at<br>(AC002343.179_S_A<br>T) | gb AAB63623.1  (AC002343)<br>cellulose synthase isolog<br>[Arabidopsis thaliana]                             | 0           |            |                       |
| 12454_at<br>(AC006232.164_AT)         | gb AAD15602.1  (AC006232)<br>putative ferredoxin<br>[Arabidopsis thaliana]                                   | 5E-85       |            |                       |
| 12475_at<br>(Y11794.1_AT)             | emb CAA72490.1  (Y11794)<br>peroxidase ATP29a<br>[Arabidopsis thaliana]                                      | 3E-67       |            | peroxidase            |
| 12487_at<br>(AC004411.126_AT)         | gb AAC34225.1  (AC004411)<br>putative ABC transporter<br>[Arabidopsis thaliana]                              | 0           |            |                       |

| ProbeSet                             | Description   | Blast Score | EC # | Family              |
|--------------------------------------|---|-------------|------|---------------------|
| 12497_at<br>(AC006533.51_AT)         | gb AAD32284.1 AC006533_8<br>(AC006533) putative<br>receptor-like protein kinase<br>[Arabidopsis thaliana] | 0           |      |                     |
| 12500_s_at<br>(AF081067.3_S_AT)      | gb AAC32192.1  (AF081067)<br>IAA-Ala hydrolase; IAA-<br>amino acid hydrolase<br>[Arabidopsis thaliana]    | 0           |      |                     |
| 12521_at<br>(AF049236.28_AT)         | gb AAC14413.1  (AF049236)<br>unknown [Arabidopsis<br>thaliana]  | 0           |      |                     |
| 12525_at<br>(AC006587.85_AT)         | gb AAD21486.1  (AC006587)<br>putative DOF zinc finger<br>protein [Arabidopsis thaliana]                   | 1E-132      |      |                     |
| 12530_at<br>(Z99707.184_AT)          | emb CAB16760.1  (Z99707)<br>hydroxynitrile lyase like<br>protein [Arabidopsis thaliana]                   | 1E-150      |      | LYASE               |
| 12535_at<br>(AL035538.156_AT)        | emb CAB37540.1 <br>(AL035538) putative protein<br>[Arabidopsis thaliana]                                  | 1E-132      |      |                     |
| 12538_at<br>(AF033205.2_AT)          | gb AAC02973.1  (AF033205)<br>putative pectin methylesterase<br>[Arabidopsis thaliana]                     | 6E-38       |      | methyl-<br>esterase |
| 12571_s_at<br>(AF149413.18_S_AT<br>) | gb AAD40138.1 AF149413_1<br>9 (AF149413) Arabidopsis<br>thaliana ferrochelatase-I<br>(SW:P42043); Pfam    | 0           |      |                     |
| 12574_at<br>(X82624.2_AT)            | emb CAA57944.1  (X82624)<br>SRG2At [Arabidopsis<br>thaliana]  | 1E-25       |      |                     |
| 12584_at<br>(AC004521.233_AT)        | gb AAC16096.1  (AC004521)<br>similar to axi 1 protein from<br>Nicotiana tabacum<br>[Arabidopsis thaliana] | 0           |      |                     |
| 12626_at<br>(AC006234.95_AT)         | gb AAD20931.1  (AC006234)<br>putative diacylglycerol kinase<br>[Arabidopsis thaliana]                     | 0           |      | kinase              |
| 12642_at<br>(AC006920.138_AT)        | gb AAD22285.1 AC006920_9<br>(AC006920) unknown protein<br>[Arabidopsis thaliana]                          | 0           |      |                     |
| 12645_at<br>(AL021712.56_AT)         | emb CAA16774.1 <br>(AL021712) fibrillin<br>precursor-like protein<br>[Arabidopsis thaliana]               | 1E-150      |      |                     |

| ProbeSet                             | Description  | Blast Score | EC # | Family |
|--------------------------------------|--|-------------|------|--------|
| 12698_at<br>(AC000106.42_AT)         | gb AAB70413.1  (AC000106)<br>Similar to Beta integral<br>membrane protein<br>(gb U43629). EST gb W43122<br>comes from this gene.<br>[Arabidopsis thaliana]   | 0           |      |        |
| 12712_f_at<br>(Z95774_F_AT)          | emb CAB09206.1  (Z95774)<br>R2R3-MYB transcription<br>factor [Arabidopsis thaliana]  | 6E-21       |      |        |
| 12736_f_at<br>(Z97048_F_AT)          | emb CAA90748.1  (Z50869)<br>MYB-related protein<br>[Arabidopsis thaliana]  | 4E-21       |      |        |
| 12744_at<br>(AC001645.15_AT)         | gb AAB63630.1  (AC001645)<br>jasmonate inducible protein<br>isolog [Arabidopsis thaliana]  | 0           |      |        |
| 12760_g_at<br>(AC005278.32_G_A<br>T) | gb AAC72120.1  (AC005278)<br>Strong similarity to<br>gb D14550 extracellular<br>dermal glycoprotein (EDGP)<br>precursor from Daucus carota.<br>ESTs gb 84105 and<br>gb AI100071 come from this<br>gene. [Arabidopsis thaliana]   | 0           |      |        |
| 12764_f_at<br>(AC004138.69_F_A<br>T) | gb AAC32912.1  (AC004138)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]  | 1E-111      |      |        |
| 12772_at<br>(AC005278.34_AT)         | gb AAC72119.1  (AC005278)<br>Strong similarity to<br>gb D14550 extracellular<br>dermal glycoprotein (EDGP)<br>precursor from Daucus carota.<br>ESTs gb H37281, gb T44167,<br>gb T21813, gb N38437,<br>gb Z26470, gb R65072,<br>gb N76373, gb F15470,<br>gb Z35182, gb H76373,<br>gb Z34678 a | 0           |      |        |
| 12776_at<br>(AL021811.156_AT)        | emb CAA16969.1 <br>(AL021811) putative protein<br>[Arabidopsis thaliana]   | 0           |      |        |
| 12797_s_at<br>(AC007138.25_S_A<br>T) | gb AAD22647.1 AC007138_1<br>1 (AC007138) S-<br>adenosylmethionine synthase<br>2 [Arabidopsis thaliana]   | 0           |      |        |

| ProbeSet                            | Description   | Blast Score | EC # | Family   |
|-------------------------------------|---|-------------|------|----------|
| 12802_at<br>(AL022373.153_AT)       | emb CAA18498.1 <br>(AL022373) DnaJ-like protein<br>[Arabidopsis thaliana]                               | 2E-74       |      |          |
| 12851_s_at<br>(ACCSYN1_S_AT)        | emb CAA78260.1  (Z12614)<br>1-aminocyclopropane 1-<br>carboxylate synthase<br>[Arabidopsis thaliana]    | 3E-29       |      |          |
| 12879_s_at<br>(AIG1_S_AT)           | gb AAC49282.1  (U40856)<br>AIG1 [Arabidopsis thaliana]  | 1E-150      |      |          |
| 12880_s_at<br>(AIG2_S_AT)           | gb AAC49283.1  (U40857)<br>AIG2 [Arabidopsis thaliana]  | 3E-97       |      |          |
| 12883_s_at<br>(APX_S_AT)            | emb CAA67425.1  (X98925)<br>stromal ascorbate peroxidase<br>[Arabidopsis thaliana]                      | 1E-161      |      |          |
| 12889_s_at<br>(ASA1_S_AT)           | gb AAA32738.1  (M92353)<br>anthranilate synthase alpha<br>subunit [Arabidopsis thaliana]                | 0           |      |          |
| 12891_at<br>(ATACS6_AT)             | gb AAC63850.1  (U73786)<br>ACC synthase [Arabidopsis<br>thaliana]                                       | 0           |      | synthase |
| 12892_g_at<br>(ATACS6_G_AT)         | gb AAC63850.1  (U73786)<br>ACC synthase [Arabidopsis<br>thaliana]                                       | 0           |      |          |
| 12904_s_at<br>(ATERF1_S_AT)         | dbj BAA32418.1  (AB008103)<br>ethylene responsive element<br>binding factor 1 [Arabidopsis<br>thaliana] | 1E-126      |      |          |
| 12905_s_at<br>(ATERF2_S_AT)         | dbj BAA32419.1  (AB008104)<br>ethylene responsive element<br>binding factor 2 [Arabidopsis<br>thaliana] | 1E-112      |      |          |
| 12911_s_at<br>(ATG6PDHE5_S_A<br>T)  | emb CAA59011.1  (X84229)<br>glucose-6-phosphate 1-<br>dehydrogenase [Arabidopsis<br>thaliana]           | 0           |      |          |
| 12921_s_at<br>(ATHHMGCOAR_S<br>_AT) | emb CAA33139.1  (X15032)<br>hydroxy methylglutaryl CoA<br>reductase (AA 1-592)                          | 0           |      |          |
| 12930_s_at<br>(ATLLS1_S_AT)         | gb AAC49679.1  (U77347)<br>lethal leaf-spot 1 homolog<br>[Arabidopsis thaliana]                         | 0           |      |          |
| 12951_at<br>(AC005489.5_AT)         | gb AAD32867.1 AC005489_5<br>(AC005489) F14N23.5<br>[Arabidopsis thaliana]                               | 0           |      |          |

| ProbeSet                              | Description  | Blast Score | EC # | Family             |
|---------------------------------------|--|-------------|------|--------------------|
| 12958_at<br>(AC002332.249_AT)         | gb AAB80675.1  (AC002332)<br>putative protein kinase<br>[Arabidopsis thaliana]                 | 0           |      | kinase             |
| 12965_at<br>(AL021711.118_AT)         | emb CAA16752.1 <br>(AL021711) protein kinase-<br>like protein [Arabidopsis<br>thaliana]        | 0           |      | kinase             |
| 12966_s_at<br>(AL023094.197_S_A<br>T) | emb CAA18838.1 <br>(AL023094) bZIP<br>transcription factor ATB2<br>[Arabidopsis thaliana]      | 2E-67       |      |                    |
| 12989_s_at<br>(AC004077.149_S_A<br>T) | gb AAC26690.1  (AC004077)<br>putative cytochrome P450<br>[Arabidopsis thaliana]                | 0           |      |                    |
| 13003_s_at<br>(AB021936.1_S_AT)       | dbj BAA74591.1  (AB021936)<br>nicotianamine synthase<br>[Arabidopsis thaliana]                 | 0           |      |                    |
| 13005_at<br>(AC004683.61_AT)          | gb AAC28763.1  (AC004683)<br>unknown protein [Arabidopsis<br>thaliana]                         | 0           |      |                    |
| 13014_at<br>(U93215.87_AT)            | gb AAB63082.1  (U93215)<br>putative lipase [Arabidopsis<br>thaliana]                           | 0           |      | lipase             |
| 13015_s_at<br>(X98673.2_S_AT)         | emb CAA67232.1  (X98674)<br>zinc finger protein<br>[Arabidopsis thaliana]                      | 7E-90       |      |                    |
| 13025_at<br>(AL050400.20_AT)          | emb CAB43695.1 <br>(AL050400) putative protein<br>[Arabidopsis thaliana]                       | 0           |      |                    |
| 13040_at<br>(AC002392.134_AT)         | gb AAD12039.1  (AC002392)<br>unknown protein [Arabidopsis<br>thaliana]                         | 0           |      |                    |
| 13070_at<br>(AC006919.171_AT)         | gb AAD24640.1 AC006919_2<br>0 (AC006919) putative<br>pyruvate kinase [Arabidopsis<br>thaliana] | 0           |      | kinase             |
| 13094_at<br>(AL035523.163_AT)         | emb CAB36745.1 <br>(AL035523) putative protein<br>[Arabidopsis thaliana]                       | 1E-172      |      |                    |
| 13100_at<br>(AC003680.50_AT)          | gb AAC06158.1  (AC003680)<br>putative cytochrome P450<br>[Arabidopsis thaliana]                | 0           |      | mono-<br>oxygenase |

| ProbeSet                              | Description   | Blast Score | EC #        | Family       |
|---------------------------------------|---|-------------|-------------|--------------|
| 13110_at<br>(AF074021.34_AT)          | gb AAD29776.1 AF074021_8<br>(AF074021) putative<br>symbiosis-related protein<br>[Arabidopsis thaliana]  | 1E-65       |             |              |
| 13115_at<br>(AC000375.44_AT)          | gb AAB60774.1  (AC000375)<br>ESTs<br>gb U75592,gb T13956,gb T43<br>869 come from from this<br>gene. [Arabidopsis thaliana]  | 0           |             |              |
| 13119_at<br>(AC007260.23_AT)          | gb AAD30579.1 AC007260_1<br>0 (AC007260) Similar to<br>dTDP-D-glucose 4,6-<br>dehydratase [Arabidopsis<br>thaliana]   | 0           | EC_4.2.1.46 | dehydratase  |
| 13128_at<br>(AL049607.47_AT)          | emb CAB40756.1 <br>(AL049607) protein<br>phosphatase 2C-like protein<br>[Arabidopsis thaliana]  | 0           |             |              |
| 13134_s_at<br>(AC002337.9_S_AT)       | gb AAB63818.1  (AC002337)<br>putative galactinol synthase<br>[Arabidopsis thaliana]   | 0           |             |              |
| 13137_at<br>(AC007169.86_AT)          | gb AAD26480.1 AC007169_1<br>2 (AC007169) putative<br>fructokinase [Arabidopsis<br>thaliana]   | 0           |             | fructokinase |
| 13152_s_at<br>(AC005322.24_S_A<br>T)  | gb AAC97998.1  (AC005322)<br>Identical to 1-<br>aminocyclopropane-1-<br>carboxylate oxidase (ACC<br>oxidase) gb X66719 (EAT1).<br>ESTs gb T43073, gb T5714,<br>gb R90435, gb R44023,<br>gb AA597926, gb AI099676,<br>gb AA650810 and gb 29725<br>come from this gene.<br>[Arabidopsis thaliana] | 1E-177      |             |              |
| 13154_s_at<br>(AC002333.210_S_A<br>T) | gb AAB64047.1  (AC002333)<br>putative endochitinase<br>[Arabidopsis thaliana]   | 1E-148      |             |              |
| 13157_at<br>(AC002409.35_AT)          | gb AAB86449.1  (AC002409)<br>putative cytochrome P450<br>[Arabidopsis thaliana]   | 0           |             |              |
| 13163_s_at<br>(AC005560.223_S_A<br>T) | gb AAD12692.2  (AC006069)<br>unknown protein [Arabidopsis<br>thaliana]  | 0           |             |              |

| ProbeSet   | Description   | Blast Score | EC # | Family |
|--|---|-------------|------|--------|
| 13176_at<br>(AL031394.56_AT)                         | emb CAA20567.1 <br>(AL031394) putative protein<br>[Arabidopsis thaliana]  | 0           |      |        |
| 13177_at<br>(AL049640.42_AT)                         | emb CAB40989.1 <br>(AL049640) growth factor<br>like protein [Arabidopsis<br>thaliana]   | 1E-164      |      |        |
| 13187_i_at<br>(ATTHIREDA4_I_AT)                      | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]  | 2E-63       |      |        |
| 13188_r_at<br>(ATTHIREDA4_R_AT<br>)                  | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]  | 2E-63       |      |        |
| 13189_s_at<br>(ATTHIREDA4_S_AT<br>)                  | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]  | 2E-63       |      |        |
| 13190_s_at<br>(ATTHIREDA4_S_A<br>T)                  | emb CAA80655.1  (Z23108)<br>NADPH thioredoxin reductase<br>[Arabidopsis thaliana]   | 1E-174      |      |        |
| 13211_s_at<br>(BCH1_S_AT)                            | dbj BAA82816.1  (AB023454)<br>basic endochitinase<br>[Arabidopsis thaliana]   | 2E-72       |      |        |
| 13212_s_at<br>(BGL2_S_AT)                            | emb CAB68132.1 <br>(AL137080) beta-1, 3-<br>glucanase 2 (BG2)<br>[Arabidopsis thaliana]   | 0           |      |        |
| 13215_s_at<br>(CAFFEROYLCOA<br>METHYLTRANS_S<br>_AT) | gb AAF16576.1 AC012563_2<br>9 (AC012563) putative S-<br>adenosyl-L-methionine:trans-<br>caffeoyl-Coenzyme A 3-O-<br>methyltransferase<br>[Arabidopsis thaliana] | 1E-121      |      |        |
| 13217_s_at<br>(CALMODULINLIK<br>E_S_AT)              | emb CAB42906.1 <br>(AL049862) calmodulin-like<br>protein [Arabidopsis thaliana]   | 6E-99       |      |        |
| 13219_s_at<br>(CHI4_S_AT)                            | emb CAA74930.1  (Y14590)<br>class IV chitinase<br>[Arabidopsis thaliana]  | 5E-78       |      |        |
| 13243_r_at<br>(ELI32_R_AT)                           | emb CAB37539.1 <br>(AL035538) cinnamyl-alcohol<br>dehydrogenase ELI3-2<br>[Arabidopsis  | 0           |      |        |



| ProbeSet  | Description   | Blast Score | EC #       | Family                           |
|---|---|-------------|------------|----------------------------------|
| 13244_s_at<br>(ELI32_S_AT)                          | emb CAB37539.1 <br>(AL035538) cinnamyl-alcohol<br>dehydrogenase ELI3-2<br>[Arabidopsis]             | 0           |            |                                  |
| 13246_at<br>(ERECTAL_AT)                            | gb AAF26067.1 AC012562_2<br>(AC012562) putative<br>protein kinase [Arabidopsis<br>thaliana]         | 1E-40       |            | kinase                           |
| 13255_i_at<br>(GAMMAGLUTAM<br>YLTRANSPEPTI_I<br>AT) | emb CAA89206.1  (Z49240)<br>gamma-glutamyl<br>transpeptidase [Arabidopsis<br>thaliana]              | 0           | EC_2.3.2.2 | glutamyl-<br>trans-<br>peptidase |
| 13259_s_at<br>(GLUTATHIONEPE<br>ROXIDASE1_S_AT)     | gb AAD24836.1 AC007071_8<br>(AC007071) putative<br>glutathione peroxidase<br>[Arabidopsis thaliana] | 4E-95       |            |                                  |
| 13261_s_at<br>(GLUTATHIONERE<br>DUCTASE1_S_AT)      | gb AAB67841.1  (U37697)<br>glutathione reductase<br>[Arabidopsis thaliana]                          | 0           |            |                                  |
| 13263_s_at<br>(GST1_RC_S_AT)                        | emb CAA10060.1 <br>(AJ012571) glutathione<br>transferase [Arabidopsis<br>thaliana]                  | 3E-49       |            |                                  |
| 13266_s_at<br>(GST4_S_AT)                           | emb CAB51026.1 <br>(AJ243812) glutathione<br>synthetase [Arabidopsis<br>thaliana]                   | 0           |            |                                  |
| 13273_s_at<br>(HSF4_S_AT)                           | gb AAC31756.1  (U68017)<br>heat shock transcription factor<br>4 [Arabidopsis thaliana]              | 1E-113      |            |                                  |
| 13275_f_at<br>(HSP174_F_AT)                         | emb CAA35182.1  (X17293)<br>heat shock protein (AA 1 -<br>156) [Arabidopsis thaliana]               | 8E-80       |            |                                  |
| 13277_i_at<br>(HSP176A_I_AT)                        | emb CAA74399.1  (Y14070)<br>Heat Shock Protein 17.6A<br>[Arabidopsis thaliana]                      | 2E-75       |            |                                  |
| 13285_s_at<br>(HSP83_S_AT)                          | gb AAA32822.1  (M62984)<br>heat shock protein 83<br>[Arabidopsis thaliana]                          | 0           |            |                                  |
| 13312_at<br>(AC006223.75_AT)                        | gb AAD15391.1  (AC006223)<br>putative disease resistance<br>protein [Arabidopsis thaliana]          | 1E-126      |            | disease                          |

| ProbeSet                      | Description  | Blast Score | EC #       | Family              |
|-------------------------------|--|-------------|------------|---------------------|
| 13367_at<br>(AC004680.97_AT)  | gb AAC31853.1  (AC004680)<br>putative NADH<br>dehydrogenase (ubiquinone<br>oxidoreductase) [Arabidopsis<br>thaliana]   | 0           | EC_1.6.5.3 | oxido-<br>reductase |
| 13370_at<br>(AC005322.4_AT)   | gb AAC97990.1  (AC005322)<br>Strong similarity to Dsor1<br>protein kinase gb D13782<br>from Drosophila<br>melanogaster. [Arabidopsis<br>thaliana]                            | 0           |            | kinase              |
| 13381_at<br>(AC006580.8_AT)   | gb AAD22369.1 AC006580_1<br>(AC006580) NAM (no apical<br>meristem)-like protein<br>[Arabidopsis thaliana]  | 1E-169      |            |                     |
| 13395_at<br>(AL035528.202_AT) | emb CAB36843.1 <br>(AL035528) SAUR-AC-like<br>protein (small auxin up RNA)<br>[Arabidopsis thaliana]   | 1E-48       |            |                     |
| 13435_at<br>(AF003102.3_AT)   | gb AAC49775.1  (AF003102)<br>AP2 domain containing<br>protein RAP2.9 [Arabidopsis<br>thaliana]   | 1E-33       |            |                     |
| 13437_at<br>(AF096371.8_AT)   | gb AAC62791.1  (AF096371)<br>contains similarity to D-<br>isomer specific 2-hydroxyacid<br>dehydrogenases (Pfam: 2-<br>Hacid_DH.hmm, score: 19.11)<br>[Arabidopsis thaliana] | 0           |            |                     |
| 13450_at<br>(AL049657.33_AT)  | emb CAB41122.1 <br>(AL049657) putative<br>proteasome regulatory subunit<br>[Arabidopsis thaliana]  | 0           |            |                     |
| 13467_at<br>(AL096860.198_AT) | emb CAB51214.1 <br>(AL096860) putative protein<br>[Arabidopsis thaliana]   | 0           |            | permease            |
| 13480_at<br>(AC005223.15_AT)  | gb AAD10644.1  (AC005223)<br>40409 [Arabidopsis thaliana]  | 1E-169      |            |                     |
| 13534_at<br>(AF149413.36_AT)  | gb AAD40124.1 AF149413_5<br>(AF149413) contains<br>similarity to soybean early<br>nodulin 93 (N-93)<br>(SW:Q02921) [Arabidopsis<br>thaliana]                                 | 2E-38       |            |                     |

| ProbeSet                      | Description  | Blast Score | EC #       | Family             |
|-------------------------------|--|-------------|------------|--------------------|
| 13536_at<br>(AL021636.47_AT)  | emb CAA16575.1 <br>(AL021636) putative protein<br>[Arabidopsis thaliana]   | 0           |            |                    |
| 13538_at<br>(AL080254.75_AT)  | emb CAB45844.1 <br>(AL080254) calcium-binding<br>protein-like [Arabidopsis<br>thaliana]  | 1E-105      |            |                    |
| 13564_at<br>(AC005312.113_AT) | gb AAC78521.1  (AC005312)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]  | 1E-127      |            | transferase        |
| 13565_at<br>(AL035601.21_AT)  | emb CAB38206.1 <br>(AL035601) auxin-responsive<br>GH3-like protein [Arabidopsis<br>thaliana]   | 0           |            |                    |
| 13584_at<br>(AC007127.23_AT)  | gb AAD25137.1 AC007127_3<br>(AC007127) putative<br>ubiquitin-like protein<br>[Arabidopsis thaliana]                                    | 0           |            |                    |
| 13588_at<br>(AL021961.24_AT)  | emb CAA17552.1 <br>(AL021961) Phosphoglycerate<br>dehydrogenase - like protein<br>[Arabidopsis thaliana]                               | 0           |            | dehydro-<br>genase |
| 13589_at<br>(AC000132.24_AT)  | gb AAB60745.1  (AC000132)<br>ESTs<br>gb ATTS1236,gb T43334,gb N<br>97019,gb AA395203 come<br>from this gene. [Arabidopsis<br>thaliana] | 2E-91       |            |                    |
| 13604_at<br>(AC000104.20_AT)  | gb AAB70431.1  (AC000104)<br>F19P19.10 [Arabidopsis<br>thaliana]   | 0           | EC_2.7.1.- |                    |
| 13605_at<br>(AL078470.75_AT)  | emb CAB43918.1 <br>(AL078470) 26S proteasome<br>subunit 4-like protein<br>[Arabidopsis thaliana]                                       | 0           |            | ATPase             |
| 13617_at<br>(AC006592.64_AT)  | gb AAD22351.1 AC006592_8<br>(AC006592) putative<br>mitochondrial dicarboxylate<br>carrier protein [Arabidopsis<br>thaliana]            | 1E-152      |            |                    |
| 13627_at<br>(AL035394.196_AT) | emb CAA23036.1 <br>(AL035394) putative Na <sup>+</sup> /H <sup>+</sup> -<br>exchanging protein<br>[Arabidopsis thaliana]               | 0           |            |                    |

| ProbeSet                            | Description   | Blast Score | EC # | Family |
|-------------------------------------|---|-------------|------|--------|
| 13645_at<br>(AC000098.8_AT)         | gb AAB71443.1  (AC000098)<br>EST gb ATT50295 comes<br>from this gene. [Arabidopsis<br>thaliana]   | 6E-19       |      |        |
| 13647_at<br>(AF000657.22_AT)        | gb AAB72161.1  (AF000657)<br>unknown protein [Arabidopsis<br>thaliana]  | 0           |      |        |
| 13656_at<br>(AC007138.31_AT)        | gb AAD22649.1 AC007138_1<br>3 (AC007138) predicted<br>protein of unknown function<br>[Arabidopsis thaliana]   | 0           |      |        |
| 13659_at<br>(AL022347.46_AT)        | emb CAA18462.1 <br>(AL022347) serine/threonine<br>kinase-like protein<br>[Arabidopsis thaliana]   | 0           |      |        |
| 13666_s_at<br>(INDOLE3GPS_S_A<br>T) | gb AAD25838.1 AC006951_1<br>7 (AC006951) putative<br>indole-3-glycerol phosphate<br>synthase [Arabidopsis<br>thaliana]  | 0           |      |        |
| 13680_s_at<br>(LOX1_S_AT)           | gb AAA17036.1  (U01843)<br>lipoxygenase 1 [Arabidopsis<br>thaliana]   | 0           |      |        |
| 13685_s_at<br>(MLOLIKE2_S_AT)       | gb AAD25552.1 AC005850_9<br>(AC005850) Highly Similar<br>to Mlo proteins [Arabidopsis<br>thaliana]  | 0           |      |        |
| 13688_s_at<br>(MONOPTEROS_S_<br>AT) | gb AAC60794.1  (AF037229)<br>transcription factor<br>[Arabidopsis thaliana]   | 0           |      |        |
| 13697_at (NI16_AT)                  | No hits found.  |             |      |        |
| 13705_s_at<br>(AC003671X_S_AT)      | gb AAC18810.1  (AC003671)<br>Strong similarity to trehalose-<br>6-phosphate synthase homolog<br>from A. thaliana chromosome<br>4 contig gb Z97344. ESTs<br>gb H37594, gb R65023,<br>gb H37578 and gb R64855<br>come from this gene.<br>[Arabidopsis thaliana] | 0           |      |        |
| 13706_s_at<br>(AC005724X_S_AT)      | gb AAD08939.1  (AC005724)<br>putative trehalose-6-phosphate<br>synthase [Arabidopsis<br>thaliana]   | 0           |      |        |

| ProbeSet                          | Description   | Blast Score | EC # | Family                 |
|-----------------------------------|---|-------------|------|------------------------|
| 13716_at<br>(NOVARTIS103_RC_AT)   | emb CAA62665.1  (X91259)<br>lectin like protein<br>[Arabidopsis thaliana]                                       | 6E-70       |      |                        |
| 13718_at<br>(NOVARTIS105_RC_AT)   | emb CAA96522.1  (Z72152)<br>AMP-binding protein<br>[Brassica napus]   | 6E-19       |      |                        |
| 13746_at<br>(NOVARTIS121_RC_AT)   | gb AAF18699.1 AC010795_1<br>4 (AC010795) hypothetical<br>protein [Arabidopsis thaliana]                         | 8E-80       |      |                        |
| 13751_at<br>(NOVARTIS127_AT)      | gb AAF16751.1 AC010155_4<br>(AC010155) F3M18.8<br>[Arabidopsis thaliana]  | 2E-27       |      |                        |
| 13755_at<br>(NOVARTIS15_AT)       | emb CAA16797.1 <br>(AL021713) receptor<br>serine/threonine kinase-like<br>protein [Arabidopsis thaliana]        | 1E-114      |      |                        |
| 13763_at<br>(NOVARTIS21_AT)       | gb AAF24849.1 AC012679_2<br>0 (AC012679) putative<br>calmodulin-binding protein<br>[Arabidopsis thaliana]       | 1E-105      |      |                        |
| 13764_at<br>(NOVARTIS22_AT)       | gb AAD39641.1 AC007591_6<br>(AC007591) F9L1.6<br>[Arabidopsis thaliana]   | 2E-30       |      |                        |
| 13789_at<br>(AJ132436.2_AT)       | emb CAB41008.1 <br>(AJ132436) GA 2-oxidase<br>[Arabidopsis thaliana]  | 0           |      | oxidase                |
| 13803_at<br>(Z97341.376_AT)       | emb CAB10444.1  (Z97341)<br>cyanohydrin lyase like protein<br>[Arabidopsis thaliana]                            | 1E-154      |      |                        |
| 13818_s_at<br>(AC006218.175_S_AT) | gb AAD15433.1  (AC006218)<br>putative aspartate<br>aminotransferase [Arabidopsis<br>thaliana]                   | 0           |      |                        |
| 13825_s_at<br>(AF104919.22_S_AT)  | gb AAC72875.1  (AF104919)<br>Arabidopsis thaliana ABC1<br>protein (GB:AJ001158)                                 | 0           |      |                        |
| 13834_at<br>(AL080237.29_AT)      | emb CAB45784.1 <br>(AL080237) cyclic nucleotide<br>gated channel (CNGC4) like<br>protein [Arabidopsis thaliana] | 0           |      |                        |
| 13842_at<br>(AC002396.12_AT)      | gb AAC00572.1  (AC002396)<br>similar to zinc<br>metalloproteinases<br>[Arabidopsis thaliana]                    | 0           |      | metallo-<br>proteinase |

| ProbeSet                              | Description  | Blast Score | EC # | Family |
|---------------------------------------|--|-------------|------|--------|
| 13848_at<br>(AC003981.31_AT)          | gb AAC14057.1  (AC003981)<br>F22O13.31 [Arabidopsis<br>thaliana]   | 0           |      |        |
| 13880_s_at<br>(AL049480.183_S_A<br>T) | emb CAB39611.1 <br>(AL049480) possible<br>apospory-associated like<br>protein [Arabidopsis thaliana]   | 0           |      |        |
| 13896_at<br>(AC004473.8_AT)           | gb AAC24048.1  (AC004473)<br>Strong similarity to trehalose-<br>6-phosphate synthase homolog<br>gb 2245136 from A. thaliana<br>chromosome 4 contig<br>gb Z97344. [Arabidopsis<br>thaliana] | 0           |      |        |
| 13908_s_at<br>(A71590.1_S_AT)         | emb CAB42588.1  (A71590)<br>unnamed protein product<br>[Arabidopsis thaliana]  | 9E-62       |      |        |
| 13918_at<br>(AC005388.29_AT)          | gb AAC64884.1  (AC005388)<br>Strong similarity to F21B7.33<br>gi 2809264 from A. thaliana<br>BAC gb AC002560. EST<br>gb N65119 comes from this<br>gene. [Arabidopsis thaliana]             | 0           |      |        |
| 13920_at<br>(AC005990.53_AT)          | gb AAC98026.1  (AC005990)<br>EST gb AA597511 comes<br>from this gene. [Arabidopsis<br>thaliana]  | 1E-131      |      |        |
| 13944_at<br>(U89959.24_AT)            | gb AAC24380.1  (U89959)<br>Unknown protein<br>[Arabidopsis thaliana]   | 0           |      |        |
| 13949_s_at<br>(Z97343.352_S_AT)       | emb CAB10528.1  (Z97343)<br>thioesterase like protein<br>[Arabidopsis thaliana]  | 1E-167      |      |        |
| 13963_at<br>(AL021711.26_AT)          | emb CAA16746.1 <br>(AL021711) putative protein<br>[Arabidopsis thaliana]   | 1E-137      |      |        |
| 13964_at<br>(AL021889.3_AT)           | emb CAA17126.2 <br>(AL021889) N-acetylornithine<br>deacetylase-like protein,<br>fragment [Arabidopsis<br>thaliana]   | 0           |      |        |
| 13966_at<br>(AL022023.172_AT)         | emb CAA17775.1 <br>(AL022023) putative protein<br>[Arabidopsis thaliana]   | 1E-173      |      |        |

| ProbeSet                              | Description  | Blast Score | EC # | Family             |
|---------------------------------------|--|-------------|------|--------------------|
| 13999_at<br>(AF071527.56_AT)          | gb AAD11584.1 AAD11584<br>(AF071527) hypothetical<br>protein [Arabidopsis thaliana]  | 1E-173      |      |                    |
| 14015_s_at<br>(A71588.1_S_AT)         | emb CAB42612.1  (A71639)<br>unnamed protein product<br>[Arabidopsis thaliana]  | 1E-63       |      |                    |
| 14016_s_at<br>(A71596.1_S_AT)         | emb CAB42592.1  (A71596)<br>unnamed protein product<br>[Arabidopsis thaliana]  | 3E-58       |      |                    |
| 14025_s_at<br>(AC007293.3_S_AT)       | gb AAD12260.1  (AF098632)<br>subtilisin-like protease<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 14026_at<br>(AC000106.5_AT)           | gb AAB70397.1  (AC000106)<br>Similar to probable Mg-<br>dependent ATPase<br>(pir S56671). ESTs<br>gb T46782,gb AA04798 come<br>from this gene. [Arabidopsis<br>thaliana] | 0           |      |                    |
| 14030_at<br>(AC005970.225_AT)         | gb AAC95171.1  (AC005970)<br>putative protein kinase<br>[Arabidopsis thaliana]   | 0           |      | kinase             |
| 14032_at<br>(AL035601.11_AT)          | emb CAB38204.1 <br>(AL035601) cytochrome<br>P450-like protein<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 14041_at<br>(AC003970.28_AT)          | gb AAC33208.1  (AC003970)<br>Highly similar to cinnamyl<br>alcohol dehydrogenase,<br>gi 1143445 [Arabidopsis<br>thaliana]  | 0           |      | dehydro-<br>genase |
| 14052_at<br>(AC004122.24_AT)          | gb AAC34333.1  (AC004122)<br>Highly Similar to branched-<br>chain amino acid<br>aminotransferase [Arabidopsis<br>thaliana]   | 1E-174      |      |                    |
| 14068_s_at<br>(AC006922.197_S_A<br>T) | gb AAD31580.1 AC006922_1<br>2 (AC006922) putative<br>farnesylated protein<br>[Arabidopsis thaliana]  | 1E-132      |      |                    |
| 14070_at<br>(AL049658.217_AT)         | emb CAB41143.1 <br>(AL049658) putative peptide<br>transporter [Arabidopsis<br>thaliana]  | 0           |      |                    |

| ProbeSet                              | Description  | Blast Score | EC # | Family  |
|---------------------------------------|--|-------------|------|---------|
| 14083_at<br>(AC005662.56_AT)          | gb AAC78535.1  (AC005662)<br>putative embryo-abundant<br>protein [Arabidopsis thaliana]  | 1E-155      |      |         |
| 14089_at<br>(AC006223.65_AT)          | gb AAD15390.1  (AC006223)<br>putative hydrolase<br>[Arabidopsis thaliana]  | 1E-135      |      |         |
| 14100_at<br>(AF002109.108_AT)         | gb AAB95282.1  (AF002109)<br>putative peroxisomal<br>membrane carrier protein<br>[Arabidopsis thaliana]                                      | 1E-166      |      |         |
| 14110_i_at<br>(AL035528.279_I_A<br>T) | emb CAB36854.1 <br>(AL035528) putative disease<br>resistance protein<br>[Arabidopsis thaliana]   | 0           |      | disease |
| 14116_at<br>(AF077407.30_AT)          | gb AAC26243.1  (AF077407)<br>contains similarity to sugar<br>transporters (Pfam:<br>sugar_tr.hmm, score: 395.39)<br>[Arabidopsis thaliana]   | 0           |      |         |
| 14122_at<br>(AF058826.23_AT)          | gb AAC13608.1  (AF058826)<br>similar to eukaryotic protein<br>kinase domains (Pfam:<br>pkinase.hmm, score: 189.74)<br>[Arabidopsis thaliana] | 0           |      | kinase  |
| 14139_at<br>(NOVARTIS30_AT)           | gb AAD09343.1  (AF026538)<br>ABA-responsive protein<br>[Hordeum vulgare]   | 7E-36       |      |         |
| 14141_at<br>(NOVARTIS31_AT)           |  |             |      |         |
| 14148_at<br>(NOVARTIS38_AT)           | gb AAF34713.1 AF224762_1<br>(AF224762) SigA binding<br>protein [Arabidopsis thaliana]  | 4E-53       |      |         |
| 14170_at<br>(NOVARTIS51_AT)           | gb AAF29406.1 AC022354_5<br>(AC022354) unknown protein<br>[Arabidopsis thaliana]   | 9E-26       |      |         |
| 14197_at<br>(NOVARTIS71_AT)           |  |             |      |         |
| 14214_at<br>(NOVARTIS83_AT)           | gb AAF24849.1 AC012679_2<br>0 (AC012679) putative<br>calmodulin-binding protein<br>[Arabidopsis thaliana]                                    | 2E-92       |      |         |
| 14223_at<br>(NOVARTIS9_AT)            | emb CAA19683.1 <br>(AL024486) putative protein<br>[Arabidopsis thaliana]   | 0           |      |         |



| ProbeSet                          | Description   | Blast Score | EC # | Family |
|-----------------------------------|---|-------------|------|--------|
| 14240_s_at<br>(NR1_S_AT)          | gb AAF08556.1 AC012193_5<br>(AC012193) nitrate reductase<br>1 (NR1) [Arabidopsis<br>thaliana]   | 1E-148      |      |        |
| 14242_s_at<br>(NRA_S_AT)          | gb AAF19225.1 AC007505_1<br>(AC007505) nitrate reductase<br>[Arabidopsis thaliana]  | 0           |      |        |
| 14248_at<br>(PAD3_AT)             | gb AAD31062.1 AC007357_1<br>1 (AC007357) Strong<br>similarity to gb X97864<br>cytochrome P450 from<br>Arabidopsis thaliana and is a<br>member of the PF 00067<br>Cytochrome P450 family.<br>ESTs gb N65665, gb T14112,<br>gb T76255, gb T20906 and<br>gb AI100027 come from this<br>gene. | 0           |      |        |
| 14249_i_at<br>(PAD4_I_AT)         | emb CAB43438.1 <br>(AL050300) putative protein<br>[Arabidopsis thaliana]  | 0           |      |        |
| 14254_s_at (PAL1-<br>MRNA_S_AT)   | gb AAD18156.2  (AC006260)<br>phenylalanine ammonia lyase<br>(PAL1) [Arabidopsis   | 1E-134      |      |        |
| 14256_f_at (PAL1-<br>INTRON_F_AT) | gb AAD18156.2  (AC006260)<br>phenylalanine ammonia lyase<br>(PAL1) [Arabidopsis thaliana]   | 7E-77       |      |        |
| 14257_s_at (PAL2-<br>MRNA_S_AT)   | gb AAC18871.1  (L33678)<br>phenylalanine ammonia lyase<br>[Arabidopsis thaliana]  | 0           |      |        |
| 14320_at<br>(AC005956.54_AT)      | gb AAD23719.1 AC005956_8<br>(AC005956) putative RING<br>zinc finger protein<br>[Arabidopsis thaliana]   | 1E-110      |      |        |
| 14381_at<br>(AC002521.68_AT)      | gb AAC05341.1  (AC002521)<br>unknown protein [Arabidopsis<br>thaliana]  | 0           |      |        |
| 14408_at<br>(AC002291.14_AT)      | gb AAC00635.1  (AC002291)<br>Unknown protein<br>[Arabidopsis thaliana]  | 0           |      |        |
| 14448_at<br>(AC002387.243_AT)     | gb AAB82641.1  (AC002387)<br>putative auxin-regulated<br>protein [Arabidopsis thaliana]   | 2E-94       |      |        |

| ProbeSet                      | Description   | Blast Score | EC # | Family                   |
|-------------------------------|---|-------------|------|--------------------------|
| 14450_at<br>(AC002986.49_AT)  | gb AAC17046.1  (AC002986)<br>EST gb N65759 comes from<br>this gene. [Arabidopsis<br>thaliana]     | 4E-78       |      |                          |
| 14459_at<br>(AC006200.69_AT)  | gb AAD14519.1  (AC006200)<br>unknown protein [Arabidopsis<br>thaliana]                            | 0           |      | kinase                   |
| 14460_at<br>(AC006201.21_AT)  | gb AAD20117.1  (AC006201)<br>unknown protein [Arabidopsis<br>thaliana]                            | 0           |      |                          |
| 14461_at<br>(AC006202.73_AT)  | gb AAD29832.1 AC006202_1<br>0 (AC006202) putative<br>carbonic anhydrase<br>[Arabidopsis thaliana] | 1E-134      |      |                          |
| 14468_at<br>(AC007576.62_AT)  | gb AAD39306.1 AC007576_2<br>9 (AC007576) Unknown<br>protein [Arabidopsis thaliana]                | 2E-89       |      |                          |
| 14475_at<br>(AL021811.121_AT) | emb CAA16965.1 <br>(AL021811) putative protein<br>[Arabidopsis thaliana]                          | 0           |      |                          |
| 14487_at<br>(Z97341.343_AT)   | emb CAB46039.1  (Z97341)<br>HSP like protein [Arabidopsis<br>thaliana]                            | 1E-160      |      |                          |
| 14498_at<br>(AC004261.51_AT)  | gb AAD11996.1  (AC004261)<br>unknown protein [Arabidopsis<br>thaliana]                            | 2E-43       |      |                          |
| 14584_at<br>(AC007658.25_AT)  | gb AAD32844.1 AC007658_3<br>(AC007658) unknown protein<br>[Arabidopsis thaliana]                  | 2E-74       |      |                          |
| 14591_at<br>(AL035440.107_AT) | emb CAB36521.1 <br>(AL035440) putative protein<br>[Arabidopsis thaliana]                          | 1E-178      |      |                          |
| 14609_at<br>(AC002340.147_AT) | gb AAC02748.1  (AC002340)<br>putative cytochrome P450<br>[Arabidopsis thaliana]                   | 0           |      |                          |
| 14614_at<br>(AC004165.66_AT)  | gb AAC16958.1  (AC004165)<br>putative glucosyltransferase<br>[Arabidopsis thaliana]               | 0           |      | glucosy-<br>ltransferase |
| 14620_s_at<br>(PAT1_S_AT)     | gb AAA32835.1  (M96073)<br>phosphoribosylanthranilate<br>transferase [Arabidopsis<br>thaliana]    | 0           |      |                          |

| ProbeSet                                      | Description   | Blast Score | EC # | Family |
|---|---|-------------|------|--------|
| 14635_s_at<br>(PR1_S_AT)                      | gb AAC69381.1  (AC005398)<br>pathogenesis-related PR-1-like<br>protein [Arabidopsis thaliana]   | 1E-94       |      |        |
| 14638_s_at<br>(PRXCB_S_AT)                    | emb CAA50677.1  (X71794)<br>peroxidase [Arabidopsis<br>thaliana]  | 0           |      |        |
| 14640_s_at<br>(PUTATIVEMLOHI<br>_S_AT)        | gb AAC28997.1  (AC004697)<br>similar to Mlo proteins from<br>H. vulgare [Arabidopsis<br>thaliana]   | 0           |      |        |
| 14643_s_at<br>(RAR047_S_AT)                   | gb AAD20078.1  (AC006836)<br>putative steroid<br>sulfotransferase [Arabidopsis<br>thaliana]   | 1E-162      |      |        |
| 14660_s_at<br>(THIOREDOXL_S_<br>AT)           | gb AAB86519.1  (AC002329)<br>putative thioredoxin reductase<br>[Arabidopsis thaliana]   | 2E-34       |      |        |
| 14663_s_at<br>(TREHALASEPREC<br>USOR_RC_S_AT) | gb AAB63620.1  (AC002343)<br>trehalase precursor isolog<br>[Arabidopsis thaliana]   | 0           |      |        |
| 14667_s_at<br>(TRPB_S_AT)                     | gb AAA32878.1  (M23872)<br>tryptophan synthase beta<br>subunit [Arabidopsis thaliana]   | 0           |      |        |
| 14672_s_at<br>(TSA1_S_AT)                     | gb AAC49117.1  (U18993)<br>tryptophan synthase alpha<br>chain [Arabidopsis thaliana]  | 1E-158      |      |        |
| 14673_s_at<br>(TSB2_S_AT)                     | gb AAA32879.1  (M81620)<br>tryptophan synthase beta-<br>subunit [Arabidopsis thaliana]<br>thaliana]   | 0           |      |        |
| 14675_s_at<br>(VSP_S_AT)                      | dbj BAA22096.1  (D85191)<br>vegetative storage protein<br>[Arabidopsis thaliana]  | 1E-146      |      |        |
| 14682_i_at<br>(WT1012A_RC_I_A<br>T)           |   |             |      |        |
| 14686_s_at<br>(WT1073_S_AT)                   | gb AAB60752.1  (AC000132)<br>Similar to A. thaliana<br>receptor-like protein kinase<br>(gb RLK5_ARATH). ESTs<br>gb ATTS0475,gb ATTS4362<br>come from this gene.<br>[Arabidopsis thaliana] | 1E-116      |      |        |

| ProbeSet                             | Description   | Blast Score | EC # | Family |
|--------------------------------------|---|-------------|------|--------|
| 14696_at<br>(WT740_RC_AT)            | gb AAD23643.1 AC007119_9<br>(AC007119) unknown protein<br>[Arabidopsis thaliana]  | 5E-45       |      |        |
| 14697_g_at<br>(WT740_RC_G_AT)        | gb AAD23643.1 AC007119_9<br>(AC007119) unknown protein<br>[Arabidopsis thaliana]  | 5E-45       |      |        |
| 14705_i_at<br>(WT77_RC_I_AT)         | emb CAB69849.1 <br>(AL137189) anthranilate N-<br>benzoyltransferase-like<br>protein [Arabidopsis thaliana]                  | 1E-38       |      |        |
| 14706_r_at<br>(WT77_RC_R_AT)         | emb CAB69849.1 <br>(AL137189) anthranilate N-<br>benzoyltransferase-like<br>protein [Arabidopsis thaliana]                  | 1E-38       |      |        |
| 14711_s_at<br>(ZFPL_S_AT)            | gb AAD25930.1 AF085279_3<br>(AF085279) hypothetical Cys-<br>3-His zinc finger protein<br>[Arabidopsis thaliana]             | 3E-37       |      |        |
| 14735_s_at<br>(AF008124_S_AT)        | gb AAB71832.1  (AF008125)<br>multidrug resistance-<br>associated protein homolog<br>[Arabidopsis thaliana]                  | 0           |      |        |
| 14750_s_at<br>(AF096370.12_S_AT<br>) | gb AAC62777.1  (AF096370)<br>contains similarity to NAM<br>(no apical meristem) -like<br>proteins [Arabidopsis<br>thaliana] | 1E-175      |      |        |
| 14763_at<br>(X86958.1_AT)            | emb CAA60521.1  (X86958)<br>protein kinase catalytic<br>domain (fragment)<br>[Arabidopsis thaliana]                         | 1E-25       |      | kinase |
| 14779_at<br>(AC004680.71_AT)         | gb AAC31851.1  (AC004680)<br>hypothetical protein<br>[Arabidopsis thaliana]   | 0           |      |        |
| 14780_at<br>(AC004683.103_AT)        | gb AAC28770.1  (AC004683)<br>DREB-like AP2 domain<br>transcription factor<br>[Arabidopsis thaliana]                         | 1E-126      |      |        |
| 14786_at<br>(AC005397.115_AT)        | gb AAC62908.1  (AC005397)<br>putative desiccation related<br>protein [Arabidopsis thaliana]                                 | 2E-90       |      |        |
| 14793_at<br>(AC006202.10_AT)         | emb CAB67652.1 <br>(AL132966) putative protein<br>[Arabidopsis thaliana]  | 2E-79       |      |        |

| ProbeSet                        | Description  | Blast Score | EC # | Family                   |
|---------------------------------|--|-------------|------|--------------------------|
| 14838_s_at<br>(M96073.6_S_AT)   | gb AAA32835.1  (M96073)<br>phosphoribosylanthranilate<br>transferase [Arabidopsis<br>thaliana]           | 0           |      |                          |
| 14882_at<br>(AL022605.63_AT)    | emb CAA18753.1 <br>(AL022605) putative protein<br>[Arabidopsis thaliana]                                 | 1E-103      |      |                          |
| 14884_at<br>(AL031032.95_AT)    | emb CAA19873.1 <br>(AL031032) putative protein<br>[Arabidopsis thaliana]                                 | 1E-149      |      |                          |
| 14895_s_at<br>(Z97344.138_S_AT) | emb CAB10562.1  (Z97344)<br>acetylornithine deacetylase<br>[Arabidopsis thaliana]                        | 4E-68       |      |                          |
| 14900_at<br>(AC000348.12_AT)    | gb AAB61488.1  (AC000348)<br>T7N9.12 [Arabidopsis<br>thaliana]   | 0           |      |                          |
| 14923_at<br>(AC006283.158_AT)   | gb AAD20693.1  (AC006283)<br>unknown protein [Arabidopsis<br>thaliana]                                   | 0           |      |                          |
| 14924_at<br>(AC006283.46_AT)    | gb AAD20686.1  (AC006283)<br>hypothetical protein<br>[Arabidopsis thaliana]                              | 7E-85       |      |                          |
| 14928_at<br>(AC006569.88_AT)    | gb AAD21756.1  (AC006569)<br>unknown protein [Arabidopsis<br>thaliana]                                   | 0           |      |                          |
| 14959_at<br>(AC007202.26_AT)    | gb AAD30230.1 AC007202_1<br>2 (AC007202) T8K14.13<br>[Arabidopsis thaliana]                              | 0           |      |                          |
| 14972_at<br>(AC005499.38_AT)    | gb AAC67344.1  (AC005499)<br>unknown protein [Arabidopsis<br>thaliana]                                   | 1E-123      |      |                          |
| 14978_at<br>(AC002333.49_AT)    | gb AAB64024.1  (AC002333)<br>putative glucosyltransferase<br>[Arabidopsis thaliana]                      | 0           |      | glucosyl-<br>transferase |
| 15032_at<br>(AC002294.8_AT)     | gb AAB71471.1  (AC002294)<br>Unknown protein<br>[Arabidopsis thaliana]                                   | 1E-115      |      |                          |
| 15042_at<br>(AL021961.3_AT)     | emb CAA17549.1 <br>(AL021961) cinnamyl alcohol<br>dehydrogenase - like protein<br>[Arabidopsis thaliana] | 1E-167      |      |                          |

| ProbeSet                              | Description  | Blast Score | EC # | Family |
|---------------------------------------|--|-------------|------|--------|
| 15052_at<br>(AC002332.103_AT)         | gb AAB80656.1  (AC002332)<br>putative calcium-binding EF-<br>hand protein [Arabidopsis<br>thaliana]  | 1E-120      |      |        |
| 15073_at<br>(AC007069.93_AT)          | gb AAD21785.1  (AC007069)<br>putative purple acid<br>phosphatase [Arabidopsis<br>thaliana]           | 0           |      |        |
| 15085_s_at<br>(AL031018.274_S_A<br>T) | emb CAA19817.1 <br>(AL031018) putative protein<br>[Arabidopsis thaliana]                             | 0           |      |        |
| 15088_s_at<br>(AC002311.37_S_A<br>T)  | gb AAB72158.1  (AF000657)<br>unknown protein [Arabidopsis<br>thaliana]                               | 9E-48       |      |        |
| 15091_at<br>(AC004683.97_AT)          | gb AAC28768.1  (AC004683)<br>unknown protein [Arabidopsis<br>thaliana]                               | 1E-101      |      |        |
| 15098_s_at<br>(ATU26945_S_AT)         | emb CAA21463.1 <br>(AL031986) senescence-<br>associated protein sen1<br>[Arabidopsis thaliana]       | 1E-103      |      |        |
| 15116_f_at<br>(AF121356_F_AT)         | gb AAD28243.1 AF121356_1<br>(AF121356) peroxiredoxin<br>TPx2 [Arabidopsis thaliana]                  | 6E-81       |      |        |
| 15123_s_at<br>(ATU40857_S_AT)         | gb AAC49283.1  (U40857)<br>AIG2 [Arabidopsis thaliana]   | 3E-97       |      |        |
| 15124_s_at<br>(ATU59508_S_AT)         | gb AAB40615.1  (U59508)<br>osmotic stress-induced proline<br>dehydrogenase [Arabidopsis<br>thaliana] | 0           |      |        |
| 15125_f_at<br>(D85190_F_AT)           | dbj BAA22095.1  (D85190)<br>vegetative storage protein<br>[Arabidopsis thaliana]                     | 1E-142      |      |        |
| 15129_s_at<br>(AF030386_S_AT)         | gb AAB86938.1  (AF030386)<br>NOI protein [Arabidopsis<br>thaliana]                                   | 1E-30       |      |        |
| 15132_s_at<br>(AF121878_S_AT)         | gb AAD30449.1 AF121878_1<br>(AF121878) cytidine<br>deaminase [Arabidopsis<br>thaliana]               | 1E-169      |      |        |
| 15137_s_at<br>(ATU57320_S_AT)         | gb AAB47973.1  (U57320)<br>blue copper-binding protein II<br>[Arabidopsis thaliana]                  | 5E-70       |      |        |

| ProbeSet                       | Description  | Blast Score | EC # | Family |
|--------------------------------|--|-------------|------|--------|
| 15140_s_at<br>(ATU93845_S_AT)  | gb AAB52420.1  (U93845)<br>Arabidopsis thaliana ER-type<br>calcium pump protein,<br>complete sequence                | 0           |      |        |
| 15141_s_at<br>(D85191_S_AT)    | dbj BAA22096.1  (D85191)<br>vegetative storage protein<br>[Arabidopsis thaliana]                                     | 1E-146      |      |        |
| 15154_s_at<br>(ATHMTGDAS_S_AT) | emb CAB51206.1 <br>(AL096860) glutamine-<br>dependent asparagine<br>synthetase [Arabidopsis<br>thaliana]             | 0           |      |        |
| 15161_s_at<br>(ATU90522_S_AT)  | gb AAB53975.1  (U90522)<br>lysine-ketoglutarate<br>reductase/saccharopine<br>dehydrogenase [Arabidopsis<br>thaliana] | 0           |      |        |
| 15162_s_at<br>(U01880_S_AT)    | gb AAA20642.1  (U01880)<br>pre-hevein-like protein<br>[Arabidopsis thaliana]   | 1E-113      |      |        |
| 15188_s_at<br>(AF081202_S_AT)  | gb AAC31606.1  (AF081202)<br>villin 2 [Arabidopsis thaliana]   | 0           |      |        |
| 15192_s_at<br>(ATHERD1_S_AT)   | dbj BAA04506.1  (D17582)<br>ERD1 protein [Arabidopsis<br>thaliana]   | 0           |      |        |
| 15196_s_at<br>(ATU43412_S_AT)  | gb AAC49573.1  (U43412) 3'-<br>phosphoadenosine 5'-<br>phosphosulfate reductase<br>[Arabidopsis thaliana]            | 0           |      |        |
| 15197_s_at<br>(ATU52851_S_AT)  | gb AAB09723.1  (U52851)<br>arginine decarboxylase<br>[Arabidopsis thaliana]  | 0           |      |        |
| 15199_s_at<br>(AB005804_S_AT)  | dbj BAA28624.1  (AB005804)<br>aldehyde oxidase [Arabidopsis<br>thaliana]   | 0           |      |        |
| 15211_s_at<br>(ATH243813_S_AT) | emb CAB51027.1 <br>(AJ243813) glutathione<br>synthetase [Arabidopsis<br>thaliana]                                    | 0           |      |        |
| 15216_s_at<br>(ATU75191_S_AT)  | gb AAB51576.1  (U75198)<br>germin-like protein<br>[Arabidopsis thaliana]   | 8E-97       |      |        |

| ProbeSet                              | Description  | Blast Score | EC # | Family      |
|---------------------------------------|--|-------------|------|-------------|
| 15342_at<br>(AC006593.101_AT)         | gb AAD20671.1  (AC006593)<br>unknown protein [Arabidopsis<br>thaliana]   | 0           |      |             |
| 15379_at<br>(AC002335.182_AT)         | gb AAB64323.1  (AC002335)<br>unknown protein [Arabidopsis<br>thaliana]   | 7E-75       |      |             |
| 15389_at<br>(AC004786.100_AT)         | gb AAC32433.1  (AC004786)<br>unknown protein [Arabidopsis<br>thaliana]   | 2E-37       |      |             |
| 15406_at<br>(AC006931.179_AT)         | gb AAD21731.1  (AC006931)<br>unknown protein [Arabidopsis<br>thaliana]   | 2E-98       |      |             |
| 15431_at<br>(AL030978.64_AT)          | emb CAA19722.1 <br>(AL030978) putative protein<br>[Arabidopsis thaliana]   | 2E-51       |      |             |
| 15463_at<br>(AL031326.226_AT)         |  |             |      |             |
| 15479_at<br>(AL049483.205_AT)         | emb CAB39671.1 <br>(AL049483) putative protein<br>[Arabidopsis thaliana]   | 4E-68       |      |             |
| 15483_s_at<br>(AC005819.20_S_A<br>T)  | gb AAC69922.1  (AC005819)<br>putative cytochrome b5<br>[Arabidopsis thaliana]  | 4E-58       |      |             |
| 15485_at<br>(AC006233.109_AT)         | gb AAD41998.1 AC006233_1<br>0 (AC006233) unknown<br>protein [Arabidopsis thaliana]   | 1E-169      |      |             |
| 15496_at<br>(AC006282.167_AT)         | gb AAD20156.1  (AC006282)<br>putative glucosyl transferase<br>[Arabidopsis thaliana]   | 0           |      | transferase |
| 15518_at<br>(AC005322.28_AT)          | gb AAC97999.1  (AC005322)<br>ESTs gb H36249,<br>gb AA59732 and<br>gb AA651219 come from this<br>gene. [Arabidopsis thaliana] | 1E-125      |      |             |
| 15522_i_at<br>(AL078637.213_I_A<br>T) | emb CAB45071.1 <br>(AL078637) putative protein<br>[Arabidopsis thaliana]   | 8E-48       |      |             |
| 15523_s_at<br>(AL078637.213_S_A<br>T) | emb CAB45071.1 <br>(AL078637) putative protein<br>[Arabidopsis thaliana]   | 8E-48       |      |             |
| 15524_at<br>(AC005508.25_AT)          | gb AAD14499.1  (AC005508)<br>44123 [Arabidopsis thaliana]  | 0           |      |             |



| ProbeSet                              | Description  | Blast Score | EC #        | Family                   |
|---------------------------------------|--|-------------|-------------|--------------------------|
| 15526_at<br>(AC004122.16_AT)          | gb AAC34332.1  (AC004122)<br>Unknown protein<br>[Arabidopsis thaliana]   | 0           |             |                          |
| 15531_i_at<br>(AL078637.191_I_A<br>T) | emb CAB45069.1 <br>(AL078637) putative protein<br>[Arabidopsis thaliana]                                       | 0           |             |                          |
| 15532_r_at<br>(AL078637.191_R_A<br>T) | emb CAB45069.1 <br>(AL078637) putative protein<br>[Arabidopsis thaliana]                                       | 0           |             |                          |
| 15540_at<br>(AC006585.205_AT)         | gb AAD18030.1  (AF118129)<br>Tsi1-interacting protein TSIP1<br>[Nicotiana tabacum]                             | 1E-33       |             |                          |
| 15543_at<br>(AF096371.10_AT)          | gb AAC62794.1  (AF096371)<br>T2L5.6 gene product<br>[Arabidopsis thaliana]                                     | 1E-108      |             |                          |
| 15544_at<br>(AL021633.110_AT)         | emb CAA16532.1 <br>(AL021633) predicted protein<br>[Arabidopsis thaliana]                                      | 1E-147      |             |                          |
| 15547_at<br>(AC005970.122_AT)         | gb AAC95168.1  (AC005970)<br>unknown protein [Arabidopsis<br>thaliana]   | 1E-110      |             |                          |
| 15551_at<br>(AL035440.289_AT)         | emb CAB36537.1 <br>(AL035440) putative<br>dihydrolipoamide<br>succinyltransferase<br>[Arabidopsis thaliana]    | 0           | EC_2.3.1.61 | succinyl-<br>transferase |
| 15580_s_at<br>(AF057043_S_AT)         | gb AAC13497.1  (AF057043)<br>acyl-CoA oxidase<br>[Arabidopsis thaliana]  | 0           |             |                          |
| 15582_s_at<br>(ATH131392_S_AT)        | emb CAA10364.1 <br>(AJ131392) alternative<br>oxidase [Arabidopsis thaliana]                                    | 0           |             |                          |
| 15594_s_at<br>(ATU56635_S_AT)         | gb AAB01222.1  (U56635)<br>glutamate dehydrogenase 2<br>[Arabidopsis thaliana]                                 | 0           |             |                          |
| 15613_s_at<br>(ATHHOMEOA_S_<br>AT)    | emb CAA79670.1  (Z19602)<br>HAT4 [Arabidopsis thaliana]  | 1E-144      |             |                          |
| 15614_s_at<br>(ATHMERI5B_S_A<br>T)    | emb CAB52471.1 <br>(AL109796) xyloglucan endo-<br>1, 4-beta-D-glucanase<br>precursor [Arabidopsis<br>thaliana] | 1E-162      |             |                          |

| ProbeSet                        | Description  | Blast Score | EC # | Family |
|---------------------------------|--|-------------|------|--------|
| 15617_s_at<br>(ATHSAR1_S_AT)    | gb AAA56991.1  (M90418)<br>formerly called HAT24;<br>synaptobrevin-related protein<br>[Arabidopsis thaliana]           | 1E-112      |      |        |
| 15622_s_at<br>(ATU43945_S_AT)   | gb AAB40594.1  (U43945)<br>strictosidine synthase<br>[Arabidopsis thaliana]  | 0           |      |        |
| 15625_s_at<br>(ATU74610_S_AT)   | gb AAB17995.1  (U74610)<br>glyoxalase II [Arabidopsis<br>thaliana]   | 1E-141      |      |        |
| 15629_s_at<br>(AB003280_S_AT)   | dbj BAA24440.1  (AB010407)<br>phosphoglycerate<br>dehydrogenase [Arabidopsis<br>thaliana]                              | 0           |      |        |
| 15632_s_at<br>(AB012570_S_AT)   | dbj BAA37112.1  (AB012570)<br>ATHP3 [Arabidopsis thaliana]   | 7E-77       |      |        |
| 15641_s_at<br>(AF117063_S_AT)   | gb AAD10829.1  (AF117063)<br>putative inositol<br>polyphosphate 5-phosphatase<br>At5P2 [Arabidopsis thaliana]          | 0           |      |        |
| 15646_s_at<br>(ATHSAT1G_S_AT)   | gb AAC37474.1  (L42212)<br>serine acetyltransferase<br>[Arabidopsis thaliana]  | 0           |      |        |
| 15665_s_at<br>(AF022658_S_AT)   | gb AAB80922.1  (AF022658)<br>putative c2h2 zinc finger<br>transcription factor<br>[Arabidopsis thaliana]               | 3E-94       |      |        |
| 15669_s_at<br>(AF047834_S_AT)   | gb AAF24813.1 AC007592_6<br>(AC007592) F12K11.9<br>[Arabidopsis thaliana]  | 0           |      |        |
| 15670_s_at<br>(AF061638_S_AT)   | gb AAC64005.1  (AF061638)<br>branched-chain alpha-keto<br>acid decarboxylase E1 beta<br>subunit [Arabidopsis thaliana] | 0           |      |        |
| 15672_s_at<br>(AF082299_S_AT)   |  |             |      |        |
| 15674_s_at<br>(AF091844_S_AT)   | gb AAC61769.1  (AF091844)<br>aminoalcoholphosphotransferase<br>[Arabidopsis thaliana]                                  | 0           |      |        |
| 15680_s_at<br>(ATHATPK19B_S_AT) | dbj BAA07661.1  (D42061)<br>ribosomal-protein S6 kinase<br>homolog [Arabidopsis<br>thaliana]                           | 0           |      |        |

| ProbeSet                             | Description   | Blast Score | EC # | Family            |
|--------------------------------------|---|-------------|------|-------------------|
| 15778_at<br>(X98676.2_AT)            | emb CAA67234.1  (X98676)<br>zinc finger protein<br>[Arabidopsis thaliana]                                   | 8E-87       |      |                   |
| 15779_g_at<br>(X98676.2_G_AT)        | emb CAA67234.1  (X98676)<br>zinc finger protein<br>[Arabidopsis thaliana]                                   | 8E-87       |      |                   |
| 15792_at<br>(AC002341.106_AT)        | gb AAB67625.1  (AC002341)<br>hypothetical protein<br>[Arabidopsis thaliana]                                 | 1E-174      |      |                   |
| 15798_at<br>(AC002521.173_AT)        | gb AAC05351.1  (AC002521)<br>putative receptor-like protein<br>kinase [Arabidopsis thaliana]                | 0           |      |                   |
| 15815_s_at<br>(Z97342.366_S_AT)      | emb CAB10487.1  (Z97342)<br>hypothetical protein<br>[Arabidopsis thaliana]                                  | 1E-175      |      |                   |
| 15839_at<br>(AC005662.203_AT)        | gb AAC78548.1  (AC005662)<br>unknown protein [Arabidopsis<br>thaliana]                                      | 8E-22       |      |                   |
| 15859_at<br>(AC006587.164_AT)        | gb AAD21491.1  (AC006587)<br>unknown protein [Arabidopsis<br>thaliana]                                      | 4E-17       |      |                   |
| 15866_s_at<br>(AC007133.59_S_A<br>T) | gb AAC79625.1  (AC005770)<br>unknown protein [Arabidopsis<br>thaliana]                                      | 5E-27       |      |                   |
| 15874_at<br>(AL022223.106_AT)        | emb CAA18223.1 <br>(AL022223) putative protein<br>[Arabidopsis thaliana]                                    | 1E-133      |      |                   |
| 15886_at<br>(AL078637.204_AT)        | emb CAB45070.1 <br>(AL078637) putative protein<br>[Arabidopsis thaliana]                                    | 1E-126      |      | phosphorylas<br>e |
| 15900_at<br>(AC005311.74_AT)         |   |             |      |                   |
| 15919_at<br>(AC007060.42_AT)         | gb AAD25764.1 AC007060_2<br>2 (AC007060) EST<br>gb AA721821 comes from this<br>gene. [Arabidopsis thaliana] | 0           |      |                   |
| 15921_s_at<br>(AC007067.1_S_AT)      | gb AAD39561.1 AC007067_1<br>(AC007067) T10O24.1<br>[Arabidopsis thaliana]                                   | 1E-24       |      |                   |
| 15924_at<br>(AC007138.61_AT)         | gb AAD22658.1 AC007138_2<br>2 (AC007138) predicted<br>protein of unknown function<br>[Arabidopsis thaliana] | 8E-94       |      |                   |

| ProbeSet                              | Description   | Blast Score | EC # | Family      |
|---------------------------------------|---|-------------|------|-------------|
| 15970_s_at<br>(X71794.2_S_AT)         | emb CAA50677.1  (X71794)<br>peroxidase [Arabidopsis<br>thaliana]  | 0           |      |             |
| 15978_at<br>(X68592.6_AT)             | emb CAA48579.1  (X68592)<br>adenosine nucleotide<br>translocator [Arabidopsis<br>thaliana]              | 0           |      | translocase |
| 15982_s_at<br>(AC006260.78_S_A<br>T)  | emb CAA66863.1  (X98190)<br>peroxidase ATP2a<br>[Arabidopsis thaliana]                                  | 0           |      |             |
| 16001_at<br>(AF035385.2_AT)           | gb AAC39468.1  (AF035385)<br>unknown [Arabidopsis<br>thaliana]  | 4E-72       |      |             |
| 16003_s_at<br>(AL021749.64_S_AT<br>)  | emb CAA16877.1 <br>(AL021749) ADP, ATP<br>carrier-like protein<br>[Arabidopsis thaliana]                | 0           |      |             |
| 16021_s_at<br>(AL022224.182_S_A<br>T) | emb CAA18251.1 <br>(AL022224) endomembrane-<br>associated protein<br>[Arabidopsis thaliana]             | 3E-63       |      |             |
| 16031_at<br>(X94248.1_AT)             | emb CAA63932.1  (X94248)<br>ferritin [Arabidopsis thaliana]   | 1E-136      |      |             |
| 16043_at<br>(AC005489.17_AT)          | gb AAD32879.1 AC005489_1<br>7 (AC005489) F14N23.17<br>[Arabidopsis thaliana]                            | 0           |      |             |
| 16053_i_at<br>(Y14251.4_I_AT)         | emb CAA74639.1  (Y14251)<br>glutathione S-transferase<br>[Arabidopsis thaliana]                         | 1E-110      |      | transferase |
| 16054_s_at<br>(Y14251.4_S_AT)         | emb CAA74639.1  (Y14251)<br>glutathione S-transferase<br>[Arabidopsis thaliana]                         | 1E-110      |      |             |
| 16058_s_at<br>(ATU94495_S_AT)         | gb AAD24836.1 AC007071_8<br>(AC007071) putative<br>glutathione peroxidase<br>[Arabidopsis thaliana]     | 5E-95       |      |             |
| 16059_s_at<br>(D88206_S_AT)           | dbj BAA24694.1  (D88206)<br>protein kinase [Arabidopsis<br>thaliana]                                    | 0           |      |             |
| 16063_s_at<br>(AB008103_S_AT)         | dbj BAA32418.1  (AB008103)<br>ethylene responsive element<br>binding factor 1 [Arabidopsis<br>thaliana] | 1E-126      |      |             |

| ProbeSet                           | Description  | Blast Score | EC # | Family |
|------------------------------------|--|-------------|------|--------|
| 16073_f_at<br>(AF062908_F_AT)      | gb AAC83630.1  (AF062908)<br>putative transcription factor<br>[Arabidopsis thaliana]                                 | 1E-122      |      |        |
| 16080_f_at<br>(AF118822_F_AT)      | gb AAD20612.1  (AF118822)<br>senescence-associated protein<br>[Arabidopsis thaliana]                                 | 3E-26       |      |        |
| 16083_s_at<br>(AF153283_S_AT)      | gb AAD34615.1 AF153283_1<br>(AF153283) putative<br>progesterone-binding protein<br>homolog [Arabidopsis<br>thaliana] | 1E-116      |      |        |
| 16087_s_at<br>(ATHATPK6A_S_A<br>T) | dbj BAA07656.1  (D42056)<br>ribosomal-protein S6 kinase<br>homolog [Arabidopsis<br>thaliana]                         | 0           |      |        |
| 16091_s_at<br>(ATHHSP83_S_AT)      | gb AAA32822.1  (M62984)<br>heat shock protein 83<br>[Arabidopsis thaliana]   | 0           |      |        |
| 16092_s_at<br>(ATHKAT1_S_AT)       | gb AAA32824.1  (M86990)<br>potassium channel protein<br>[Arabidopsis thaliana]                                       | 0           |      |        |
| 16103_s_at<br>(ATU60445_S_AT)      | gb AAD51782.1 AF145299_1<br>(AF145299) 14-3-3 protein<br>GF14 nu [Arabidopsis<br>thaliana]                           | 1E-148      |      |        |
| 16105_s_at<br>(ATU68017_S_AT)      | gb AAC31756.1  (U68017)<br>heat shock transcription factor<br>4 [Arabidopsis thaliana]                               | 1E-113      |      |        |
| 16108_s_at<br>(D78604_S_AT)        | gb AAD03379.1  (AC005967)<br>putative cytochrome P450<br>[Arabidopsis thaliana]                                      | 0           |      |        |
| 16130_s_at<br>(AF078683_S_AT)      | gb AAC68664.1  (AF078683)<br>RING-H2 finger protein<br>RHA1a [Arabidopsis thaliana]                                  | 3E-96       |      |        |
| 16133_s_at<br>(AF089810_S_AT)      | gb AAF26045.1 AC015986_8<br>(AC015986) ARG1 protein<br>(Altered Response to Gravity)<br>[Arabidopsis thaliana]       | 0           |      |        |
| 16134_s_at<br>(AF132016_S_AT)      | gb AAD33584.1 AF132016_1<br>(AF132016) RING-H2 zinc<br>finger protein ATL6<br>[Arabidopsis thaliana]                 | 0           |      |        |
| 16159_s_at<br>(ATU37697_S_AT)      | gb AAB67841.1  (U37697)<br>glutathione reductase<br>[Arabidopsis thaliana]   | 0           |      |        |

| ProbeSet                             | Description   | Blast Score | EC # | Family      |
|--------------------------------------|---|-------------|------|-------------|
| 16161_s_at<br>(ATU39072_S_AT)        | gb AAA91165.1  (U39072)<br>AtGRP2b [Arabidopsis<br>thaliana]  | 8E-57       |      |             |
| 16173_s_at<br>(D78607_S_AT)          | dbj BAA28539.1  (D78607)<br>cytochrome P450<br>monooxygenase [Arabidopsis<br>thaliana]  | 0           |      |             |
| 16203_at<br>(AC007519.53_AT)         | gb AAD46036.1 AC007519_2<br>1 (AC007519) Contains<br>similarity to gb M74161<br>inositol polyphosphate 5-<br>phosphatase from Homo<br>sapiens and contains a<br>PF 00783 inositol<br>polyphosphate phosphatase<br>catalytic domain. [Arabidopsis<br>thaliana] | 0           |      |             |
| 16230_at<br>(AL049655.78_AT)         | emb CAB41089.1 <br>(AL049655) putative protein<br>[Arabidopsis thaliana]  | 0           |      |             |
| 16232_s_at<br>(AL080252.77_S_AT<br>) | emb CAB45796.1 <br>(AL080252) putative protein<br>[Arabidopsis thaliana]  | 1E-119      |      |             |
| 16233_at<br>(AL080254.83_AT)         | emb CAB45846.1 <br>(AL080254) putative protein<br>[Arabidopsis thaliana]  | 0           |      |             |
| 16236_g_at<br>(X92657.3_G_AT)        | emb CAA63346.1  (X92657)<br>cationic amino acid<br>transporter [Arabidopsis<br>thaliana]  | 1E-37       |      |             |
| 16272_at<br>(AC006304.136_AT)        | gb AAD20108.1  (AC006304)<br>hypothetical protein<br>[Arabidopsis thaliana]   | 0           |      |             |
| 16288_at<br>(AF024504.17_AT)         | gb AAB80790.2  (AF024504)<br>similar to prolyl 4-hydroxylase<br>alpha subunit [Arabidopsis<br>thaliana]   | 1E-143      |      | hydroxylase |
| 16298_at<br>(AL021890.71_AT)         | emb CAA17152.1 <br>(AL021890) putative protein<br>[Arabidopsis thaliana]  | 1E-68       |      |             |
| 16299_at<br>(AL024486.185_AT)        | emb CAA19705.1 <br>(AL024486) putative protein<br>[Arabidopsis thaliana]  | 1E-170      |      |             |

| ProbeSet                          | Description   | Blast Score | EC # | Family             |
|-----------------------------------|---|-------------|------|--------------------|
| 16301_s_at<br>(AL031018.105_S_AT) | emb CAA19807.1 <br>(AL031018) hypothetical<br>protein [Arabidopsis thaliana]  | 1E-149      |      |                    |
| 16306_at<br>(AL049751.112_AT)     | emb CAB41935.1 <br>(AL049751) putative protein<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 16329_s_at<br>(AF013294.17_S_AT)  | emb CAA10659.1 <br>(AJ132387) Ca <sup>2+</sup> -ATPase<br>[Arabidopsis thaliana]<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 16335_at<br>(AL079347.105_AT)     | emb CAB45450.1 <br>(AL079347) xanthine<br>dehydrogenase-like protein<br>[Arabidopsis thaliana]  | 0           |      | dehydro-<br>genase |
| 16340_at<br>(AC004255.15_AT)      | gb AAC13905.1  (AC004255)<br>T1F9.15 [Arabidopsis<br>thaliana]  | 0           |      |                    |
| 16357_at<br>(AF149413.38_AT)      | gb AAD40144.1 AF149413_2<br>5 (AF149413) contains<br>similarity to protein kinase<br>domains (Pfam F00069,<br>Score=162.6, E=6.8e-45,<br>N=1) and leucien rich repeats<br>(Pfam PF00560, Score=210.7,<br>E=2.2e-59, N=10)<br>[Arabidopsis thaliana] | 0           |      | kinase             |
| 16363_at<br>(AC004255.14_AT)      | gb AAC13904.1  (AC004255)<br>T1F9.14 [Arabidopsis<br>thaliana]  | 0           |      |                    |
| 16383_at<br>(AC006300.64_AT)      | gb AAD20719.1  (AC006300)<br>putative disease resistance<br>protein [Arabidopsis thaliana]  | 0           |      | disease            |
| 16391_at<br>(AL050351.194_AT)     | emb CAB43642.1 <br>(AL050351) receptor protein<br>kinase-like protein<br>[Arabidopsis thaliana]   | 0           |      |                    |
| 16398_s_at<br>(AL022603.3_S_AT)   | emb CAA20204.1 <br>(AL031187) serine/threonine<br>kinase-like protein<br>[Arabidopsis thaliana]   | 0           |      |                    |
| 16405_at<br>(AC005850.9_AT)       | gb AAD25549.1 AC005850_6<br>(AC005850) Putative<br>serine/threonine kinase<br>[Arabidopsis thaliana]  | 0           |      | kinase             |

| ProbeSet                              | Description   | Blast Score | EC # | Family             |
|---------------------------------------|---|-------------|------|--------------------|
| 16409_at<br>(AC004393.2_AT)           | gb AAC18783.1  (AC004393)<br>Strong similarity to receptor<br>kinase gb M80238 from A.  | 0           |      | kinase             |
| 16440_s_at<br>(AF002109.137_S_A<br>T) | gb AAB95285.1  (AF002109)<br>putative nematode-resistance<br>protein [Arabidopsis thaliana]   | 0           |      |                    |
| 16457_s_at<br>(AC005397.17_S_A<br>T)  | gb AAD23036.1 AC006526_2<br>(AC006526) unknown protein<br>[Arabidopsis thaliana]  | 3E-29       |      |                    |
| 16461_i_at<br>(AC004683.79_I_AT<br>)  | gb AAC28766.1  (AC004683)<br>peroxidase [Arabidopsis<br>thaliana]   | 0           |      | peroxidase         |
| 16462_s_at<br>(AC004683.79_S_A<br>T)  | gb AAC28766.1  (AC004683)<br>peroxidase [Arabidopsis<br>thaliana]   | 0           |      |                    |
| 16465_at<br>(Y08892.1_AT)             | emb CAA70105.1  (Y08892)<br>Hsc70-G8 protein<br>[Arabidopsis thaliana]  | 4E-46       |      |                    |
| 16470_s_at<br>(AF068299.4_S_AT)       | gb AAD14544.1  (AF068299)<br>gamma-glutamylcysteine<br>synthetase [Arabidopsis<br>thaliana]   | 0           |      |                    |
| 16483_at<br>(X68053_AT)               | emb CAA48189.1  (X68053)<br>transcription factor<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 16496_s_at<br>(AF030386.1_S_AT)       | gb AAB86938.1  (AF030386)<br>NOI protein [Arabidopsis<br>thaliana]  | 1E-30       |      |                    |
| 16510_at<br>(AL034567.198_AT)         | emb CAA22575.1 <br>(AL034567) putative protein<br>[Arabidopsis thaliana]  | 1E-168      |      |                    |
| 16522_at<br>(X77500.2_AT)             | emb CAA54631.1  (X77500)<br>amino acid transporter<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 16524_at<br>(AC006577.38_AT)          | gb AAD25783.1 AC006577_1<br>9 (AC006577) Strong<br>similarity to gb S77096<br>aldehyde dehydrogenase<br>homolog from Brassica napus<br>and is a member of PF 00171<br>Aldehyde dehydrogenase<br>family. ESTs gb T46213,<br>gb T42164, gb T43682,<br>gb N96380, gb T42973,<br>gb Z34663, gb Z4 | 0           |      | dehydro-<br>genase |



| ProbeSet                           | Description   | Blast Score | EC # | Family      |
|------------------------------------|---|-------------|------|-------------|
| 16526_at<br>(Z49227.1_AT)          | emb CAA89201.2  (Z49227)<br>adenine nucleotide translocase<br>[Arabidopsis thaliana]                            | 0           |      | translocase |
| 16538_s_at<br>(AB010259_S_AT)      | dbj BAA28347.1  (AB010259)<br>DRH1 [Arabidopsis thaliana]   | 0           |      |             |
| 16541_s_at<br>(AB023423_S_AT)      | dbj BAA75015.1  (AB023423)<br>sulfate transporter<br>[Arabidopsis thaliana]                                     | 0           |      |             |
| 16545_s_at<br>(AF037229_S_AT)      | gb AAC60794.1  (AF037229)<br>transcription factor<br>[Arabidopsis thaliana]                                     | 0           |      |             |
| 16553_f_at<br>(AF078821_F_AT)      | gb AAC68670.1  (AF078821)<br>RING-H2 finger protein<br>RHA1b [Arabidopsis thaliana]                             | 2E-93       |      |             |
| 16568_s_at<br>(ATHATCDPK_S_A<br>T) | gb AAB03246.1  (U31835)<br>calmodulin-domain protein<br>kinase CDPK isoform 6<br>[Arabidopsis thaliana]         | 0           |      |             |
| 16570_s_at<br>(ATHCDPKA_S_AT<br>)  | gb AAF27092.1 AC011809_1<br>(AC011809) calcium-<br>dependent protein kinase 1<br>[Arabidopsis thaliana]         | 0           |      |             |
| 16578_s_at<br>(ATHRPRP1B_S_A<br>T) | emb CAB68132.1 <br>(AL137080) beta-1, 3-<br>glucanase 2 (BG2)<br>[Arabidopsis thaliana]                         | 0           |      |             |
| 16589_s_at<br>(ATU26937_S_AT)      | gb AAD24605.1 AC005825_1<br>2 (AC005825) putative MYB<br>family transcription factor<br>[Arabidopsis thaliana]  | 1E-143      |      |             |
| 16594_s_at<br>(ATU39783_S_AT)      | gb AAB82307.1  (U39783)<br>amino acid transport protein<br>[Arabidopsis thaliana]                               | 0           |      |             |
| 16603_s_at<br>(ATU81293_S_AT)      | gb AAB58497.1  (U81293)<br>UDP-glucose:indole-3-acetate<br>beta-D-glucosyltransferase<br>[Arabidopsis thaliana] | 0           |      |             |
| 16609_s_at<br>(AB008104_S_AT)      | dbj BAA32419.1  (AB008104)<br>ethylene responsive element<br>binding factor 2 [Arabidopsis<br>thaliana]         | 1E-112      |      |             |

| ProbeSet                      | Description   | Blast Score | EC # | Family                |
|-------------------------------|---|-------------|------|-----------------------|
| 16611_s_at<br>(AB008782_S_AT) | dbj BAA23424.1  (AB008782)<br>sulfate transporter<br>[Arabidopsis thaliana]   | 0           |      |                       |
| 16635_s_at<br>(AF126057_S_AT) | emb CAB43670.1 <br>(AL050352) putative protein<br>[Arabidopsis thaliana]  | 0           |      |                       |
| 16638_s_at<br>(AF139098_S_AT) | gb AAD37511.1 AF139098_1<br>(AF139098) putative zinc<br>finger protein [Arabidopsis<br>thaliana]  | 1E-103      |      |                       |
| 16646_s_at<br>(ATHDHS1_S_AT)  | gb AAA32784.1  (M74819) 3-<br>deoxy-D-arabino-<br>heptulosonate y-phosphate<br>synthase [Arabidopsis<br>thaliana]   | 0           |      |                       |
| 16649_s_at<br>(ATHORF_S_AT)   | gb AAF16576.1 AC012563_2<br>9 (AC012563) putative S-<br>adenosyl-L-methionine:trans-<br>caffeoyl-Coenzyme A 3-O-<br>methyltransferase<br>[Arabidopsis thaliana] | 1E-121      |      |                       |
| 16701_at<br>(AC005312.61_AT)  | gb AAC78514.1  (AC005312)<br>putative phloem-specific<br>lectin [Arabidopsis thaliana]  | 1E-170      |      |                       |
| 16721_at<br>(AC006533.58_AT)  | gb AAD32285.1 AC006533_9<br>(AC006533) putative<br>poly(ADP-ribose)<br>glycohydrolase [Arabidopsis<br>thaliana]   | 0           |      | kinase                |
| 16747_at<br>(AL021713.3_AT)   | emb CAA16788.1 <br>(AL021713) DNA binding-<br>like protein [Arabidopsis<br>thaliana]  | 0           |      |                       |
| 16753_at<br>(AL031032.110_AT) | emb CAA19874.1 <br>(AL031032) putative protein<br>[Arabidopsis thaliana]  | 0           |      |                       |
| 16781_at<br>(AC002392.100_AT) | gb AAD12030.1  (AC002392)<br>putative receptor-like protein<br>kinase [Arabidopsis thaliana]  | 0           |      | kinase                |
| 16810_at<br>(AC002339.46_AT)  | gb AAC02763.1  (AC002339)<br>putative polygalacturonase<br>[Arabidopsis thaliana]   | 0           |      | Polygalactur<br>onase |

| ProbeSet                              | Description   | Blast Score | EC # | Family                 |
|---------------------------------------|---|-------------|------|------------------------|
| 16817_s_at<br>(AL096882.91_S_AT)      | emb CAB51412.1 <br>(AL096882) ACC synthase<br>(AtACS-6) [Arabidopsis<br>thaliana]                 | 0           |      |                        |
| 16859_at<br>(AL035523.135_AT)         | emb CAB36742.1 <br>(AL035523) alpha-amylase-<br>like protein [Arabidopsis<br>thaliana]            | 0           |      |                        |
| 16864_i_at<br>(AF037367.4_I_AT)       | emb CAB66108.1 <br>(AL133248) endo-<br>polygalacturonase<br>[Arabidopsis thaliana]                | 0           |      | polygalac-<br>turonase |
| 16865_s_at<br>(AF037367.4_S_AT)       | emb CAB66108.1 <br>(AL133248) endo-<br>polygalacturonase<br>[Arabidopsis thaliana]                | 0           |      |                        |
| 16888_s_at<br>(AC004684.174_S_A<br>T) | gb AAC23647.1  (AC004684)<br>putative alcohol<br>dehydrogenase [Arabidopsis<br>thaliana]          | 1E-170      |      |                        |
| 16902_at<br>(AC007119.67_AT)          | gb AAD23643.1 AC007119_9<br>(AC007119) unknown protein<br>[Arabidopsis thaliana]                  | 6E-91       |      |                        |
| 16903_g_at<br>(AC007119.67_G_A<br>T)  | gb AAD23643.1 AC007119_9<br>(AC007119) unknown protein<br>[Arabidopsis thaliana]                  | 6E-91       |      |                        |
| 16908_at<br>(AC002396.22_AT)          | gb AAC00577.1  (AC002396)<br>pyruvate dehydrogenase E1<br>alpha subunit [Arabidopsis<br>thaliana] | 0           |      |                        |
| 16914_s_at<br>(AL049500.57_S_AT<br>)  | emb CAB39936.1 <br>(AL049500) osmotin<br>precursor [Arabidopsis<br>thaliana]                      | 1E-143      |      |                        |
| 16916_s_at<br>(X77199.8_S_AT)         | emb CAA54420.1  (X77199)<br>heat shock cognate 70-2<br>[Arabidopsis thaliana]                     | 3E-52       |      |                        |
| 16927_s_at<br>(AF035384.2_S_AT)       | gb AAC39467.1  (AF035384)<br>endo-xyloglucan transferase<br>[Arabidopsis thaliana]                | 2E-65       |      |                        |
| 16940_g_at<br>(AC002334.110_G_<br>AT) | gb AAC04922.1  (AC002334)<br>putative synaptobrevin<br>[Arabidopsis thaliana]                     | 1E-123      |      |                        |

| ProbeSet                         | Description  | Blast Score | EC # | Family       |
|----------------------------------|--|-------------|------|--------------|
| 16951_i_at<br>(AC005662.30_I_AT) | gb AAC78532.1  (AC005662)<br>calmodulin-like protein<br>[Arabidopsis thaliana]                           | 1E-88       |      |              |
| 16952_s_at<br>(AC005662.30_S_AT) | gb AAC78532.1  (AC005662)<br>calmodulin-like protein<br>[Arabidopsis thaliana]                           | 1E-88       |      |              |
| 16955_at<br>(AL031326.215_AT)    | emb CAA20468.1 <br>(AL031326) putative protein<br>[Arabidopsis thaliana]                                 | 4E-87       |      |              |
| 16968_at<br>(AL021961.93_AT)     | emb CAA17559.1 <br>(AL021961)<br>glucosyltransferase -like<br>protein [Arabidopsis thaliana]             | 0           |      |              |
| 16970_s_at<br>(Y18291.5_S_AT)    | emb CAA77109.1  (Y18291)<br>uncoupling protein<br>[Arabidopsis thaliana]                                 | 1E-175      |      |              |
| 16972_at<br>(AC004261.89_AT)     | gb AAA87347.1  (M88307)<br>calmodulin [Brassica juncea]  | 5E-81       |      |              |
| 16981_s_at<br>(U35829.2_S_AT)    | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]                                       | 4E-64       |      |              |
| 16989_at<br>(AL030978.46_AT)     | emb CAA19720.1 <br>(AL030978) GH3 like protein<br>[Arabidopsis thaliana]                                 | 0           |      |              |
| 16995_at<br>(AC002391.188_AT)    | gb AAB87114.1  (AC002391)<br>unknown protein [Arabidopsis<br>thaliana]                                   | 0           |      |              |
| 17007_at<br>(AC005896.26_AT)     | gb AAC98046.1  (AC005896)<br>putative adenylate kinase<br>[Arabidopsis thaliana]                         | 1E-155      |      | kinase       |
| 17008_at<br>(AC006585.212_AT)    | gb AAD23027.1 AC006585_2<br>2 (AC006585) putative<br>tyrosine aminotransferase<br>[Arabidopsis thaliana] | 0           |      | transaminase |
| 17009_at<br>(AL021633.163_AT)    | emb CAA16537.1 <br>(AL021633) putative protein<br>[Arabidopsis thaliana]                                 | 0           |      |              |
| 17039_s_at<br>(D78602_S_AT)      | dbj BAA28534.1  (D78602)<br>cytochrome P450<br>monooxygenase [Arabidopsis<br>thaliana]                   | 0           |      |              |
| 17041_s_at<br>(D89631_S_AT)      | gb AAC14417.1  (AF049236)<br>unknown [Arabidopsis<br>thaliana]   | 0           |      |              |

| ProbeSet                       | Description   | Blast Score | EC # | Family   |
|--------------------------------|---|-------------|------|----------|
| 17051_s_at<br>(AF098947_S_AT)  | gb AAD09952.1  (AF098947)<br>CTF2B [Arabidopsis thaliana]   | 0           |      |          |
| 17066_s_at<br>(ATHLIPOXY_S_AT) | gb AAA17036.1  (U01843)<br>lipoxygenase 1 [Arabidopsis thaliana]  | 0           |      |          |
| 17073_s_at<br>(ATTS4391_S_AT)  | gb AAD20078.1  (AC006836)<br>putative steroid<br>sulfotransferase [Arabidopsis thaliana]  | 1E-162      |      |          |
| 17075_s_at<br>(ATU09961_S_AT)  | gb AAA19628.1  (U09961)<br>nitrilase [Arabidopsis thaliana]   | 0           |      |          |
| 17083_s_at<br>(ATU18770_S_AT)  | gb AAD25838.1 AC006951_1<br>7 (AC006951) putative<br>indole-3-glycerol phosphate<br>synthase [Arabidopsis thaliana]             | 0           |      |          |
| 17104_s_at<br>(D88541_S_AT)    | dbj BAA13640.1  (D88541)<br>phosphoserine<br>aminotransferase [Arabidopsis thaliana]  | 0           |      |          |
| 17111_s_at<br>(ATHACSC_S_AT)   | emb CAA78260.1  (Z12614)<br>1-aminocyclopropane 1-<br>carboxylate synthase<br>[Arabidopsis]                                     | 0           |      |          |
| 17119_s_at<br>(AF132212_S_AT)  | gb AAD38925.1 AF132212_1<br>(AF132212) OPDA-reductase<br>homolog [Arabidopsis thaliana]   | 0           |      |          |
| 17128_s_at<br>(ATHRPRP1A_S_AT) | gb AAC69381.1  (AC005398)<br>pathogenesis-related PR-1-like<br>protein [Arabidopsis]  | 1E-94       |      |          |
| 17180_at<br>(AF007270.30_AT)   | gb AAB61058.1  (AF007270)<br>contains similarity to GATA-<br>type zinc fingers<br>(PS:PS00344) [Arabidopsis thaliana]           | 0           |      |          |
| 17187_at<br>(AF128396.2_AT)    | gb AAD17371.1  (AF128396)<br>similar to arginases (Pfam:<br>PF00491, Score=353.2;<br>E=1.4e-119, N=1)<br>[Arabidopsis thaliana] | 0           |      | arginase |

| ProbeSet                             | Description  | Blast Score | EC # | Family              |
|--------------------------------------|--|-------------|------|---------------------|
| 17300_at<br>(X66017.2_AT)            | emb CAA46815.1  (X66017)<br>NADPH-ferrihemoprotein<br>reductase [Arabidopsis<br>thaliana]                  | 0           |      | reductase           |
| 17303_s_at<br>(AC004683.25_S_A<br>T) | gb AAC67339.2  (AC005499)<br>putative WRKY-type DNA<br>binding protein [Arabidopsis<br>thaliana]           | 0           |      |                     |
| 17323_at<br>(U95973.69_AT)           | gb AAB65477.1  (U95973)<br>Ser/Thr protein kinase isolog<br>[Arabidopsis thaliana]                         | 0           |      | kinase              |
| 17338_at<br>(AC002535.97_AT)         | gb AAC62855.1  (AC002535)<br>putative pectinesterase<br>[Arabidopsis thaliana]                             | 0           |      | pectin-<br>esterase |
| 17341_at<br>(AL021713.89_AT)         | emb CAA16797.1 <br>(AL021713) receptor<br>serine/threonine kinase-like<br>protein [Arabidopsis thaliana]   | 0           |      |                     |
| 17352_at<br>(AC007127.33_AT)         | gb AAD25140.1 AC007127_6<br>(AC007127) putative protein<br>kinase [Arabidopsis thaliana]                   | 0           |      | kinase              |
| 17356_s_at<br>(Z97338.190_S_AT)      | emb CAB10307.1  (Z97338)<br>UTP-glucose<br>glucosyltransferase<br>[Arabidopsis thaliana]                   | 0           |      |                     |
| 17371_at<br>(AF076243.44_AT)         | gb AAD29762.1 AF076243_9<br>(AF076243) putative receptor-<br>like protein kinase<br>[Arabidopsis thaliana] | 0           |      |                     |
| 17376_at<br>(AL021890.218_AT)        | emb CAA17164.1 <br>(AL021890) putative protein<br>[Arabidopsis thaliana]                                   | 1E-119      |      |                     |
| 17379_at<br>(AF085279.9_AT)          | gb AAF18728.1 AC018721_3<br>(AC018721) putative CCH-<br>type zinc finger protein<br>[Arabidopsis thaliana] | 0           |      |                     |
| 17380_at<br>(AL021961.39_AT)         | emb CAA17554.1 <br>(AL021961) putative protein<br>[Arabidopsis thaliana]                                   | 0           |      |                     |
| 17398_at<br>(AC002535.143_AT)        | gb AAC62863.1  (AC002535)<br>putative protein disulfide-<br>isomerase [Arabidopsis<br>thaliana]            | 0           |      | isomerase           |

| ProbeSet                             | Description   | Blast Score | EC # | Family |
|--------------------------------------|---|-------------|------|--------|
| 17413_s_at<br>(AJ006961.4_S_AT)      | emb CAA67551.1  (X99097)<br>peroxidase [Arabidopsis<br>thaliana]  | 2E-84       |      |        |
| 17451_at<br>(AC002343.47_AT)         | gb AAB63613.1  (AC002343)<br>unknown protein [Arabidopsis<br>thaliana]  | 0           |      |        |
| 17452_g_at<br>(AC002343.47_G_A<br>T) | gb AAB63613.1  (AC002343)<br>unknown protein [Arabidopsis<br>thaliana]  | 0           |      |        |
| 17458_at<br>(AC006260.91_AT)         | gb AAD18148.1  (AC006260)<br>unknown protein [Arabidopsis<br>thaliana]  | 1E-138      |      |        |
| 17464_at<br>(AC000132.72_AT)         | gb AAB60752.1  (AC000132)<br>Similar to A. thaliana<br>receptor-like protein kinase<br>(gb RLK5_ARATH). ESTs<br>gb ATTS0475,gb ATTS4362<br>come from this gene.<br>[Arabidopsis thaliana] | 0           |      | kinase |
| 17482_s_at<br>(Z97343.441_S_AT)      | emb CAB10533.1  (Z97343)<br>GTP-binding RAB1C like<br>protein [Arabidopsis thaliana]  | 1E-112      |      |        |
| 17484_at<br>(X79052.2_AT)            | emb CAA55654.1  (X79052)<br>SRG1 [Arabidopsis thaliana]   | 0           |      |        |
| 17485_s_at<br>(Z97340.345_S_AT)      | emb CAB10405.1  (Z97340)<br>beta-1, 3-glucanase class I<br>precursor [Arabidopsis<br>thaliana]  | 1E-169      |      |        |
| 17487_s_at<br>(U18993.2_S_AT)        | gb AAC49117.1  (U18993)<br>tryptophan synthase alpha<br>chain [Arabidopsis thaliana]  | 1E-158      |      |        |
| 17490_s_at<br>(M90416.2_S_AT)        | gb AAF01532.1 AC009325_2<br>(AC009325) homeobox-<br>leucine zipper protein HAT5<br>(HD-ZIP protein 5) (HD-ZIP<br>protein ATHB-1)<br>[Arabidopsis thaliana]                                | 4E-52       |      |        |
| 17500_s_at<br>(ATHCALLGA_S_A<br>T)   | emb CAB42906.1 <br>(AL049862) calmodulin-like<br>protein [Arabidopsis thaliana]   | 6E-99       |      |        |
| 17511_s_at<br>(AF067605_S_AT)        | gb AAB71482.1  (AC002294)<br>similar to S-linalool synthase<br>gp U58314 1491939<br>[Arabidopsis thaliana]  | 0           |      |        |

| ProbeSet                      | Description  | Blast Score | EC # | Family |
|-------------------------------|--|-------------|------|--------|
| 17514_s_at<br>(AF076277_S_AT) | gb AAD03545.1  (AF076278)<br>ethylene response factor 1<br>[Arabidopsis thaliana]  | 1E-104      |      |        |
| 17516_s_at<br>(AF072536_S_AT) | gb AAC24592.1  (AF072536)<br>H-protein promoter binding<br>factor-1 [Arabidopsis thaliana]   | 0           |      |        |
| 17522_s_at<br>(D78606_S_AT)   | dbj BAA28538.1  (D78606)<br>cytochrome P450<br>monooxygenase [Arabidopsis<br>thaliana]   | 0           |      |        |
| 17533_s_at<br>(ATU43488_S_AT) | gb AAB18367.1  (U43488)<br>xyloglucan<br>endotransglycosylase-related<br>protein [Arabidopsis thaliana]  | 1E-171      |      |        |
| 17544_s_at<br>(ATU40856_S_AT) | gb AAC49282.1  (U40856)<br>AIG1 [Arabidopsis thaliana]   | 1E-150      |      |        |
| 17548_s_at<br>(AF118823_S_AT) | gb AAD20613.1  (AF118823)<br>senescence-associated protein<br>[Arabidopsis thaliana]   | 2E-26       |      |        |
| 17578_at<br>(AF093604_AT)     | gb AAF00071.1 AF093604_1<br>(AF093604) apyrase<br>[Arabidopsis thaliana]   | 0           |      |        |
| 17585_s_at<br>(AF134487_S_AT) | emb CAA06460.1 <br>(AJ005261) cytidine<br>deaminase [Arabidopsis<br>thaliana]  | 1E-168      |      |        |
| 17595_s_at<br>(AF166352_S_AT) | gb AAC28764.1  (AC004683)<br>putative beta-alanine-pyruvate<br>aminotransferase [Arabidopsis<br>thaliana]  | 0           |      |        |
| 17636_at<br>(AF077409.7_AT)   | gb AAC28219.1  (AF077409)<br>contains similarity to C3HC4-<br>type zinc fingers (Pfam: zf-<br>C3HC4.hmm, score: 32.94)<br>[Arabidopsis thaliana] | 1E-138      |      |        |
| 17648_at<br>(AL021684.43_AT)  | emb CAA16674.1 <br>(AL021684) predicted protein<br>[Arabidopsis thaliana]  | 0           |      |        |
| 17653_at<br>(AL035679.144_AT) | emb CAB38823.1 <br>(AL035679) putative protein<br>[Arabidopsis thaliana]   | 0           |      |        |
| 17702_at<br>(AC005700.212_AT) | gb AAC69951.1  (AC005700)<br>Mutator-like transposase<br>[Arabidopsis thaliana]  | 0           |      |        |



| ProbeSet                              | Description  | Blast Score | EC # | Family             |
|---------------------------------------|--|-------------|------|--------------------|
| 17719_at<br>(AC006592.17_AT)          | gb AAD22346.1 AC006592_3<br>(AC006592) hypothetical<br>protein [Arabidopsis thaliana]  | 0           |      |                    |
| 17744_s_at<br>(AC004684.168_S_A<br>T) | gb AAC23646.1  (AC004684)<br>putative alcohol<br>dehydrogenase [Arabidopsis<br>thaliana]   | 1E-170      |      |                    |
| 17752_at<br>(AC003974.37_AT)          | gb AAC04483.1  (AC003974)<br>putative protein kinase<br>[Arabidopsis thaliana]   | 0           |      | kinase             |
| 17758_at<br>(AF076243.41_AT)          | gb AAD29761.1 AF076243_8<br>(AF076243) putative receptor-<br>like protein kinase<br>[Arabidopsis thaliana]   | 0           |      |                    |
| 17775_at<br>(AC004392.2_AT)           | gb AAC28500.1  (AC004392)<br>Similar to glucose-6-<br>phosphate/phosphate-<br>translocator (GPT)<br>gb AF020814 from Pisum<br>sativum. [Arabidopsis<br>thaliana] | 0           |      |                    |
| 17781_at<br>(AL049746.177_AT)         | emb CAB41861.1 <br>(AL049746) ABC transporter-<br>like protein [Arabidopsis<br>thaliana]   | 0           |      |                    |
| 17840_s_at<br>(AC002333.223_S_A<br>T) | gb AAB64049.1  (AC002333)<br>putative endochitinase<br>[Arabidopsis thaliana]  | 1E-161      |      |                    |
| 17854_at<br>(Z99707.366_AT)           | emb CAB16762.1  (Z99707)<br>caltractin-like protein<br>[Arabidopsis thaliana]  | 1E-91       |      |                    |
| 17860_at<br>(AL078467.4_AT)           | emb CAB43873.1 <br>(AL078467) putative protein<br>[Arabidopsis thaliana]   | 1E-177      |      |                    |
| 17876_at<br>(AJ007587.2_AT)           | emb CAA07574.1 <br>(AJ007587) monooxygenase<br>[Arabidopsis thaliana]  | 0           |      | mono-<br>oxygenase |
| 17877_g_at<br>(AJ007587.2_G_AT)       | emb CAA07574.1 <br>(AJ007587) monooxygenase<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 17881_at<br>(AC002391.54_AT)          | gb AAB87100.1  (AC002391)<br>putative WRKY-type DNA-<br>binding protein [Arabidopsis<br>thaliana]  | 1E-133      |      |                    |

| ProbeSet                              | Description   | Blast Score | EC #        | Family              |
|---------------------------------------|---|-------------|-------------|---------------------|
| 17882_at<br>(AL035523.49_AT)          | emb CAB36734.1 <br>(AL035523) PROTEIN<br>TRANSPORT PROTEIN<br>SEC61 GAMMA SUBUNIT-<br>like [Arabidopsis thaliana]   | 1E-24       |             |                     |
| 17893_at<br>(AC004401.135_AT)         | gb AAC17827.1  (AC004401)<br>similar to late embryogenesis<br>abundant proteins<br>[Arabidopsis thaliana]   | 2E-43       |             |                     |
| 17894_at<br>(AC005724.44_AT)          | gb AAD08938.1  (AC005724)<br>unknown protein [Arabidopsis<br>thaliana]  | 1E-147      |             |                     |
| 17899_at<br>(Z97339.197_AT)           | emb CAB10339.1  (Z97339)<br>hypothetical protein<br>[Arabidopsis thaliana]  | 5E-55       |             |                     |
| 17900_s_at<br>(AC000106.13_S_A<br>T)  | gb AAB70401.1  (AC000106)<br>Similar to Glycine SRC2<br>(gb AB000130). ESTs<br>gb H76869,gb T21700,gb ATT<br>S5089 come from this gene.<br>[Arabidopsis thaliana] | 2E-97       |             |                     |
| 17907_s_at<br>(AC004684.165_S_A<br>T) | gb AAC23645.1  (AC004684)<br>unknown protein [Arabidopsis<br>thaliana]  | 4E-24       |             |                     |
| 17930_s_at<br>(AJ006960.4_S_AT)       | emb CAA07352.1 <br>(AJ006960) peroxidase<br>[Arabidopsis thaliana]  | 1E-111      |             |                     |
| 17945_at<br>(Z97341.411_AT)           | emb CAB10448.1  (Z97341)<br>limonene cyclase like protein<br>[Arabidopsis thaliana]   | 0           |             |                     |
| 17955_at<br>(AL021768.242_AT)         | emb CAA16940.1 <br>(AL021768) small GTP-<br>binding protein-like<br>[Arabidopsis thaliana]  | 1E-110      |             |                     |
| 17956_i_at<br>(AC005967.32_I_AT<br>)  | gb AAD03381.1  (AC005967)<br>putative leucine<br>aminopeptidase [Arabidopsis<br>thaliana]   | 0           | EC_3.4.11.1 | amino-<br>peptidase |
| 17963_at<br>(AL049730.88_AT)          | emb CAB41717.1 <br>(AL049730) pEARLI 1-like<br>protein [Arabidopsis thaliana]   | 6E-46       |             |                     |
| 17967_at<br>(AL096859.32_AT)          | emb CAB51172.1 <br>(AL096859) protein kinase 6-<br>like protein [Arabidopsis<br>thaliana]   | 0           |             |                     |

| ProbeSet                              | Description  | Blast Score | EC # | Family                   |
|---------------------------------------|--|-------------|------|--------------------------|
| 18010_s_at<br>(AJ001264_S_AT)         | emb CAA77109.1  (Y18291)<br>uncoupling protein<br>[Arabidopsis thaliana]                               | 1E-175      |      |                          |
| 18012_s_at<br>(AJ002295_S_AT)         | emb CAB59428.1 <br>(AJ002295) inositol-1,4,5-<br>trisphosphate 5-Phosphatase<br>[Arabidopsis thaliana] | 0           |      |                          |
| 18045_at<br>(AJ011976_AT)             | emb CAA71798.1  (Y10845)<br>O-acetylserine(thiol) lyase<br>[Brassica juncea]                           | 1E-148      |      |                          |
| 18054_at<br>(AJ238846_AT)             | emb CAB54517.1 <br>(AJ238846) SGP1 monomeric<br>G-protein [Arabidopsis<br>thaliana]                    | 1E-150      |      |                          |
| 18109_s_at<br>(AC002391.206_S_A<br>T) | gb AAB87118.1  (AC002391)<br>putative metal ion transporter<br>(NRAMP) [Arabidopsis<br>thaliana]       | 0           |      |                          |
| 18121_s_at<br>(AC002337.21_S_A<br>T)  | gb AAB63819.1  (AC002337)<br>MYB transcription factor<br>(Atmyb2) [Arabidopsis<br>thaliana]            | 1E-167      |      |                          |
| 18122_at<br>(AC002338.110_AT)         | gb AAC16938.1  (AC002338)<br>putative protein kinase<br>[Arabidopsis thaliana]                         | 0           |      | kinase                   |
| 18140_at<br>(Z97341.319_AT)           | emb CAB10440.1  (Z97341)<br>growth regulator like protein<br>[Arabidopsis thaliana]                    | 0           |      |                          |
| 18148_at<br>(AC004669.25_AT)          | gb AAC20719.1  (AC004669)<br>putative dioxygenase<br>[Arabidopsis thaliana]                            | 0           |      |                          |
| 18176_at<br>(AL035540.31_AT)          | emb CAB37503.1 <br>(AL035540) protein kinase<br>like protein [Arabidopsis<br>thaliana]                 | 0           |      | kinase                   |
| 18194_i_at<br>(AL096859.227_I_A<br>T) | emb CAB51196.1 <br>(AL096859) glucuronosyl<br>transferase-like protein<br>[Arabidopsis thaliana]       | 0           |      | glucosyl-<br>transferase |
| 18213_at<br>(AL022140.126_AT)         | emb CAA18110.1 <br>(AL022140) putative protein<br>[Arabidopsis thaliana]                               | 1E-174      |      |                          |
| 18216_at<br>(X95573.2_AT)             | gb AAF24959.1 AC012375_2<br>2 (AC012375) T22C5.18<br>[Arabidopsis thaliana]                            | 1E-119      |      |                          |

| ProbeSet                          | Description   | Blast Score | EC # | Family |
|-----------------------------------|---|-------------|------|--------|
| 18217_g_at<br>(X95573.2_G_AT)     | gb AAF24959.1 AC012375_2<br>2 (AC012375) T22C5.18<br>[Arabidopsis thaliana]                               | 1E-119      |      |        |
| 18224_s_at<br>(AL021890.57_S_AT)  | emb CAA17150.1 <br>(AL021890) putative protein<br>[Arabidopsis thaliana]                                  | 4E-81       |      |        |
| 18226_s_at<br>(AC002343.142_S_AT) | emb CAB51645.1 <br>(AL109619) putative protein<br>[Arabidopsis thaliana]                                  | 0           |      |        |
| 18228_at<br>(X91259.1_AT)         | emb CAA62665.1  (X91259)<br>lectin like protein<br>[Arabidopsis thaliana]                                 | 1E-142      |      |        |
| 18234_at<br>(AC000348.3_AT)       | gb AAB61479.1  (AC000348)<br>T7N9.3 [Arabidopsis thaliana]  | 1E-154      |      |        |
| 18236_s_at<br>(AC004683.69_S_AT)  | gb AAC28764.1  (AC004683)<br>putative beta-alanine-pyruvate<br>aminotransferase [Arabidopsis<br>thaliana] | 0           |      |        |
| 18241_at<br>(AC006580.71_AT)      | gb AAD21720.1  (AC006931)<br>putative RNA-binding protein<br>[Arabidopsis thaliana]                       | 0           |      |        |
| 18242_g_at<br>(AC006580.71_G_AT)  | gb AAD21720.1  (AC006931)<br>putative RNA-binding protein<br>[Arabidopsis thaliana]                       | 0           |      |        |
| 18255_at<br>(AC005770.25_AT)      | gb AAC79625.1  (AC005770)<br>unknown protein [Arabidopsis<br>thaliana]                                    | 0           |      |        |
| 18258_s_at<br>(AC006439.222_S_AT) | gb AAD15515.1  (AC006439)<br>unknown protein [Arabidopsis<br>thaliana]                                    | 4E-58       |      |        |
| 18263_at<br>(AC005724.36_AT)      | gb AAD08937.1  (AC005724)<br>unknown protein [Arabidopsis<br>thaliana]                                    | 1E-139      |      |        |
| 18266_at<br>(AC004684.33_AT)      | gb AAC23628.1  (AC004684)<br>unknown protein [Arabidopsis<br>thaliana]                                    | 0           |      |        |
| 18267_at<br>(AC006223.23_AT)      | gb AAD15384.1  (AC006223)<br>unknown protein [Arabidopsis<br>thaliana]                                    | 2E-38       |      |        |
| 18268_s_at<br>(AC006418.38_S_AT)  | gb AAD20161.1  (AC006418)<br>putative ubiquitin<br>[Arabidopsis thaliana]                                 | 0           |      |        |

| ProbeSet                              | Description   | Blast Score | EC #       | Family    |
|---------------------------------------|---|-------------|------------|-----------|
| 18280_at<br>(AC007369.2_AT)           | gb AAD30591.1 AC007369_1<br>(AC007369) Unknown protein<br>[Arabidopsis thaliana]  | 1E-135      |            |           |
| 18284_at<br>(AL021961.67_AT)          | emb CAA17557.1 <br>(AL021961) putative protein<br>[Arabidopsis thaliana]  | 6E-97       |            |           |
| 18287_at<br>(AC007661.142_AT)         | gb AAD32777.1 AC007661_1<br>4 (AC007661) unknown<br>protein [Arabidopsis thaliana]  | 0           |            |           |
| 18299_s_at<br>(M23872.2_S_AT)         | gb AAA32878.1  (M23872)<br>tryptophan synthase beta<br>subunit [Arabidopsis thaliana]   | 0           |            |           |
| 18314_i_at<br>(AL078579.83_I_AT<br>)  | emb CAB43971.1 <br>(AL078579) putative beta-<br>glucosidase [Arabidopsis<br>thaliana]   | 0           |            |           |
| 18348_at<br>(AL022603.104_AT)         | emb CAA18711.1 <br>(AL022603) putative protein<br>[Arabidopsis thaliana]  | 1E-160      |            |           |
| 18456_s_at<br>(AC004697.159_S_A<br>T) | gb AAC28997.1  (AC004697)<br>similar to Mlo proteins from<br>H. vulgare [Arabidopsis<br>thaliana]   | 0           |            |           |
| 18508_s_at<br>(AC006532.89_S_A<br>T)  | gb AAA33709.1  (L16797)<br>glutamate decarboxylase<br>[Petunia x hybrida]   | 0           |            |           |
| 18544_at<br>(AC007060.14_AT)          | gb AAD25749.1 AC007060_7<br>(AC007060) Strong similarity<br>to F19I3.8 gi 3033381 putative<br>UDP-galactose-4-epimerase<br>from Arabidopsis thaliana<br>BAC gb AC004238 and is a<br>member of PF 01370 the NAD<br>dependent<br>epimerase/dehydratase family.<br>EST gb AA597338 comes fro | 0           | EC_5.1.3.2 | epimerase |
| 18582_s_at<br>(AC003671.36_S_A<br>T)  | gb AAC18810.1  (AC003671)<br>Strong similarity to trehalose-<br>6-phosphate synthase homolog<br>from A. thaliana chromosome<br>4 contig gb Z97344. ESTs<br>gb H37594, gb R65023,<br>gb H37578 and gb R64855<br>come from this gene.<br>[Arabidopsis thaliana]                             | 0           |            |           |

| ProbeSet                          | Description   | Blast Score | EC # | Family |
|-----------------------------------|---|-------------|------|--------|
| 18587_s_at<br>(AC007166.53_S_AT)  | gb AAF18667.1 AC007166_9<br>(AC007166) unknown protein<br>[Arabidopsis thaliana]  | 0           |      |        |
| 18590_at<br>(AJ222713.4_AT)       | emb CAA10955.1 <br>(AJ222713) unnamed protein<br>product [Arabidopsis thaliana]   | 1E-151      |      |        |
| 18591_at<br>(X74756.2_AT)         | emb CAA52772.1  (X74756)<br>ATAF2 [Arabidopsis thaliana]  | 2E-99       |      |        |
| 18596_at<br>(AC005698.13_AT)      | gb AAD43614.1 AC005698_1<br>3 (AC005698) T3P18.13<br>[Arabidopsis thaliana]   | 0           |      |        |
| 18597_at<br>(AL080282.74_AT)      | emb CAB45881.1 <br>(AL080282) berberine bridge<br>enzyme-like protein<br>[Arabidopsis thaliana]   | 0           |      |        |
| 18601_s_at<br>(AC002387.279_S_AT) | gb AAF18602.1 AC002387_1<br>(AC002387) putative<br>microtubule-associated protein<br>[Arabidopsis thaliana]   | 8E-45       |      |        |
| 18604_at<br>(AF069298.31_AT)      | gb AAC19273.1  (AF069298)<br>similar to several small<br>proteins (~100 aa) that are<br>induced by heat, auxin,<br>ethylene and wounding such<br>as Phaseolus aureus indole-3-<br>acetic acid induced protein<br>ARG (SW:32292)<br>[Arabidopsis thaliana] | 3E-82       |      |        |
| 18622_g_at<br>(AJ005902.2_G_AT)   | emb CAA06759.1 <br>(AJ005902) vag2<br>[Arabidopsis thaliana]  | 3E-47       |      |        |
| 18625_at<br>(AC005278.22_AT)      | gb AAC72125.1  (AC005278)<br>ESTs gb H36966, gb R65511,<br>gb T42324 and gb T20569<br>come from this gene.<br>[Arabidopsis thaliana]  | 0           |      |        |
| 18631_at<br>(AC002510.112_AT)     | gb AAB84346.1  (AC002510)<br>unknown protein [Arabidopsis<br>thaliana]  | 0           |      |        |

| ProbeSet                             | Description  | Blast Score | EC # | Family |
|--------------------------------------|--|-------------|------|--------|
| 18636_at<br>(AC006577.22_AT)         | gb AAD25775.1 AC006577_1<br>(AC006577) Belongs to the<br>PF 00657<br>Lipase/Acylhydrolase with<br>GDSE-motif family. ESTs<br>gb T75865, gb R30449,<br>gb AI239373, gb F19931 and<br>gb F19930 come from this<br>gene. [Arabidopsis thaliana] | 0           |      |        |
| 18650_s_at<br>(AF013294.25_S_AT<br>) | gb AAB62867.1  (AF013294)<br>AT0Z11 gene product<br>[Arabidopsis thaliana]   | 3E-41       |      |        |
| 18662_s_at<br>(AC002343.20_S_A<br>T) | gb AAB63608.1  (AC002343)<br>unknown protein [Arabidopsis<br>thaliana]   | 0           |      |        |
| 18668_at<br>(AJ249794_AT)            | emb CAB56692.1 <br>(AJ249794).lipoxygenase<br>[Arabidopsis thaliana]   | 0           |      |        |
| 18681_at<br>(L23573_AT)              | gb AAD02548.1  (AF049922)<br>PGP169-12 [Petunia x<br>hybrida]  | 8E-94       |      |        |
| 18686_s_at<br>(U18126_S_AT)          | gb AAA57314.1  (U18126)<br>inner mitochondrial<br>membrane protein<br>[Arabidopsis thaliana]   | 2E-78       |      |        |
| 18698_s_at<br>(X17528_S_AT)          | emb CAA35570.1  (X17528)<br>citrate synthetase [Arabidopsis<br>thaliana]   | 0           |      |        |
| 18720_s_at<br>(X92419_S_AT)          | emb CAB52583.1  (X92420)<br>SNAP25AB protein<br>[Arabidopsis thaliana]   | 1E-157      |      |        |
| 18735_s_at<br>(Z29490_S_AT)          | emb CAA82626.1  (Z29490)<br>gamma-glutamylcysteine<br>synthetase [Arabidopsis<br>thaliana]   | 0           |      |        |
| 18753_s_at<br>(AF118222.28_S_AT<br>) | gb AAD03425.1  (AF118222)<br>contains similarity to<br>Iron/Ascorbate family of<br>oxidoreductases (Pfam:<br>PF00671, Score=297.8,<br>E=1.3e-85, N=1) [Arabidopsis<br>thaliana]  | 0           |      |        |
| 18782_at<br>(AC003040.90_AT)         | gb AAC23760.1  (AC003040)<br>putative protein kinase<br>[Arabidopsis thaliana]   | 0           |      | kinase |

| ProbeSet                        | Description   | Blast Score | EC #        | Family             |
|---------------------------------|---|-------------|-------------|--------------------|
| 18803_at<br>(AC005315.94_AT)    | gb AAC33232.1  (AC005315)<br>putative SCARECROW gene<br>regulator [Arabidopsis<br>thaliana]   | 0           |             |                    |
| 18885_at<br>(AC006921.147_AT)   | gb AAD21443.1  (AC006921)<br>unknown protein [Arabidopsis<br>thaliana]  | 1E-126      |             |                    |
| 18888_at<br>(AC007591.68_AT)    | gb AAD39666.1 AC007591_3<br>1 (AC007591) Is a member of<br>the PF 00903 glyoxalase<br>family. ESTs gb T44721,<br>gb T21844 and gb AA395404<br>come from this gene.<br>[Arabidopsis thaliana]  | 1E-101      |             |                    |
| 18899_s_at<br>(X13434.1_S_AT)   | emb CAA79494.1  (Z19050)<br>nitrate reductase [Arabidopsis<br>thaliana]   | 0           |             |                    |
| 18908_i_at<br>(AF055848.2_I_AT) | gb AAC62611.1  (AF055848)<br>subtilisin-like protease<br>[Arabidopsis thaliana]   | 0           |             | protease           |
| 18909_s_at<br>(AF055848.2_S_AT) | gb AAC62611.1  (AF055848)<br>subtilisin-like protease<br>[Arabidopsis thaliana]   | 0           |             |                    |
| 18928_at<br>(AC002333.181_AT)   | gb AAB64044.1  (AC002333)<br>putative endochitinase<br>[Arabidopsis thaliana]   | 1E-157      |             | endo-<br>chitinase |
| 18930_at<br>(AC005990.57_AT)    | gb AAC98028.1  (AC005990)<br>Similar to gb L19255 carbonic<br>anhydrase from Nicotiana<br>tabacum and a member of the<br>prokaryotic-type carbonic<br>anhydrase family PF 00484.<br>EST gb Z235745 comes from<br>this gene. [Arabidopsis<br>thaliana] | 1E-149      |             | anhydrase          |
| 18933_at<br>(AC007020.48_AT)    | gb AAD25665.1 AC007020_7<br>(AC007020) putative ferritin<br>[Arabidopsis thaliana]  | 1E-129      |             |                    |
| 18936_at<br>(AJ003119.4_AT)     | emb CAA05875.1 <br>(AJ003119) protein<br>phosphatase 2C [Arabidopsis<br>thaliana]   | 0           | EC_3.1.3.16 | phosphatase        |
| 18949_at<br>(Z54136.1_AT)       | emb CAA90809.1  (Z54136)<br>MYB-related protein<br>[Arabidopsis thaliana]   | 1E-145      |             |                    |



| ProbeSet                              | Description  | Blast Score | EC #        | Family             |
|---------------------------------------|--|-------------|-------------|--------------------|
| 18953_at<br>(AF077955.1_AT)           | gb AAC69851.1  (AF077955)<br>branched-chain alpha keto-<br>acid dehydrogenase E1 alpha<br>subunit [Arabidopsis thaliana]   | 0           |             | dehydro-<br>genase |
| 18963_at<br>(AC004561.99_AT)          | gb AAC95194.1  (AC004561)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]  | 1E-127      |             | transferase        |
| 18966_at<br>(AC004561.106_AT)         | gb AAC95196.1  (AC004561)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]  | 1E-128      |             | transferase        |
| 18980_at<br>(U78721.20_AT)            | gb AAC69126.1  (U78721)<br>putative protein phosphatase<br>2C [Arabidopsis thaliana]   | 0           | EC_3.1.3.16 | phosphatase        |
| 19019_i_at<br>(X82623.2_I_AT)         | emb CAA57943.1  (X82623)<br>SRG2At [Arabidopsis<br>thaliana]   | 7E-33       |             |                    |
| 19060_at<br>(AC003671.34_AT)          | gb AAC18809.1  (AC003671)<br>Similar to high affinity<br>potassium transporter, HAK1<br>protein gb U22945 from<br>Schwanniomyces occidentalis.<br>[Arabidopsis thaliana] | 0           |             |                    |
| 19092_at<br>(AL078606.188_AT)         | emb CAB44327.1 <br>(AL078606) protein kinase-<br>like protein [Arabidopsis<br>thaliana]  | 0           |             | kinase             |
| 19110_s_at<br>(X86947.2_S_AT)         | emb CAA60510.1  (X86947)<br>Protein Kinase catalytic<br>domain (fragment)<br>[Arabidopsis thaliana]  | 4E-27       |             |                    |
| 19132_s_at<br>(AL022603.298_S_A<br>T) | emb CAA18722.1 <br>(AL022603) putative NADPH<br>quinone oxidoreductase<br>[Arabidopsis thaliana]   | 0           |             |                    |
| 19137_at<br>(X74755.2_AT)             | emb CAA52771.1  (X74755)<br>ATAF1 [Arabidopsis thaliana]   | 1E-138      |             |                    |
| 19140_at<br>(AC005170.24_AT)          | gb AAC63657.1  (AC005170)<br>unknown protein [Arabidopsis<br>thaliana]   | 9E-83       |             |                    |

| ProbeSet                             | Description   | Blast Score | EC # | Family |
|--------------------------------------|---|-------------|------|--------|
| 19161_at<br>(AL078579.9_AT)          | emb CAB43966.1 <br>(AL078579) putative acyl-<br>CoA binding protein<br>[Arabidopsis thaliana]         | 0           |      |        |
| 19171_at<br>(AC002335.160_AT)        | gb AAB64325.1  (AC002335)<br>putative trypsin inhibitor<br>[Arabidopsis thaliana]                     | 7E-40       |      |        |
| 19178_at<br>(Y18227.2_AT)            | dbj BAA86999.1  (AB035137)<br>blue copper binding protein<br>[Arabidopsis thaliana]                   | 2E-57       |      |        |
| 19181_s_at<br>(AF053065.2_S_AT)      | gb AAC39464.1  (AF053065)<br>late embryogenesis abundant<br>protein homolog [Arabidopsis<br>thaliana] | 3E-34       |      |        |
| 19182_at<br>(AL031804.245_AT)        | emb CAA21214.1 <br>(AL031804) putative protein<br>[Arabidopsis thaliana]                              | 0           |      |        |
| 19207_at<br>(AC006069.117_AT)        | gb AAD12704.1  (AC006069)<br>unknown protein [Arabidopsis<br>thaliana]                                | 1E-104      |      |        |
| 19230_at<br>(AC003113.15_AT)         | gb AAB96860.1  (AC003113)<br>F25O1.15 [Arabidopsis<br>thaliana]                                       | 1E-134      |      |        |
| 19247_at<br>(AF071527.44_AT)         | gb AAD11587.1  AAD11587<br>(AF071527) hypothetical<br>protein [Arabidopsis thaliana]                  | 0           |      |        |
| 19257_s_at<br>(AC000104.57_S_A<br>T) | emb CAA44318.1  (X62461)<br>H1flk [Arabidopsis thaliana]<br>[Arabidopsis thaliana]                    | 0           |      |        |
| 19284_at<br>(AC003028.196_AT)        | gb AAC27173.1  (AC003028)<br>putative anthocyanidin<br>synthase [Arabidopsis<br>thaliana]             | 0           |      |        |
| 19288_at<br>(AC005824.130_AT)        | gb AAC73031.1  (AC005824)<br>putative cytochrome P450<br>[Arabidopsis thaliana]                       | 0           |      |        |
| 19325_at<br>(AL022604.42_AT)         | emb CAA18731.1 <br>(AL022604) putative protein<br>[Arabidopsis thaliana]                              | 0           |      |        |
| 19364_at<br>(AL022023.142_AT)        | emb CAA17779.1 <br>(AL022023) putative protein<br>[Arabidopsis thaliana]                              | 1E-126      |      |        |

| ProbeSet                             | Description  | Blast Score | EC #        | Family                   |
|--------------------------------------|--|-------------|-------------|--------------------------|
| 19376_at<br>(AF024504.11_AT)         | gb AAB80784.2  (AF024504)<br>putative chloroplast nucleoid<br>DNA binding protein<br>[Arabidopsis thaliana]  | 0           |             | peptidase                |
| 19383_at<br>(AC006200.203_AT)        | gb AAD14534.1  (AC006200)<br>unknown protein [Arabidopsis<br>thaliana]   | 1E-101      |             |                          |
| 19395_at<br>(AF007270.32_AT)         | gb AAB61059.1  (AF007270)<br>contains similarity to DNA<br>polymerase III, alpha chain<br>(SP:P47277) [Arabidopsis<br>thaliana]  | 1E-179      |             | polymerase               |
| 19405_at<br>(AJ223803.1_AT)          | emb CAA11553.1 <br>(AJ223803) 2-oxoglutarate<br>dehydrogenase E2 subunit<br>[Arabidopsis thaliana]   | 0           | EC_2.3.1.61 | succinyl-<br>transferase |
| 19407_at<br>(AC004697.81_AT)         | gb AAC28981.1  (AC004697)<br>putative adenylate kinase<br>[Arabidopsis thaliana]   | 1E-142      | EC_2.7.4.3  | kinase                   |
| 19409_at<br>(AC007357.56_AT)         | gb AAD31077.1 AC007357_2<br>6 (AC007357) EST<br>gb T21221 comes from this<br>gene. [Arabidopsis thaliana]  | 2E-17       |             |                          |
| 19411_at<br>(AC007661.104_AT)        | gb AAD32774.1 AC007661_1<br>1 (AC007661) unknown<br>protein [Arabidopsis thaliana]   | 1E-110      |             |                          |
| 19421_at<br>(X70990.4_AT)            | emb CAA50317.1  (X70990)<br>sucrose synthase [Arabidopsis<br>thaliana]   | 0           |             | synthase                 |
| 19432_s_at<br>(AL035680.11_S_AT<br>) | gb AAA32879.1  (M81620)<br>tryptophan synthase beta-<br>subunit [Arabidopsis thaliana]   | 0           |             |                          |
| 19451_at<br>(AC004392.6_AT)          | gb AAC28502.1  (AC004392)<br>Similar to F4I1.26 putative<br>beta-glucosidase gi 3128187<br>from A. thaliana BAC<br>gb AC004521. ESTs<br>gb N97083, gb F19868 and<br>gb F15482 come from this<br>gene. [Arabidopsis thaliana] | 0           |             | glucosidase              |
| 19460_s_at<br>(AC000132.66_S_A<br>T) | gb AAB60749.1  (AC000132)<br>Identical to A. thaliana<br>HEMA2 (gb U27118).<br>[Arabidopsis thaliana]  | 0           |             |                          |

| ProbeSet                         | Description   | Blast Score | EC #        | Family          |
|----------------------------------|---|-------------|-------------|-----------------|
| 19462_s_at<br>(AF001168.2_S_AT)  | emb CAB75467.1 <br>(AL138659) serine/threonine-specific kinase lecRK1 precursor, lectin                       | 0           |             |                 |
| 19464_at<br>(AC005560.51_AT)     | gb AAC67338.1  (AC005560) putative MAP kinase [Arabidopsis thaliana]  | 0           | EC_2.7.1.37 | kinase          |
| 19465_at<br>(AL021768.96_AT)     | emb CAA16929.1 <br>(AL021768) resistance protein RPP5-like [Arabidopsis thaliana]                             | 0           |             |                 |
| 19531_at<br>(AL021960.91_AT)     | emb CAA17531.1 <br>(AL021960) amino acid transport protein AAT1 [Arabidopsis thaliana]                        | 0           |             |                 |
| 19546_at<br>(AC005398.172_AT)    | gb AAC69380.1  (AC005398) putative endoxyloglucan glycosyltransferase [Arabidopsis thaliana]                  | 0           |             | transferase     |
| 19555_at<br>(AF058919.48_AT)     | gb AAC13630.1  (AF058919) F6N23.8 gene product [Arabidopsis thaliana]   | 0           |             |                 |
| 19591_at<br>(AJ010735.4_AT)      | emb CAA09330.1 <br>(AJ010735) gr1-protein [Arabidopsis thaliana]  | 0           | EC_3.5.-.-  | amido-hydrolase |
| 19614_at<br>(AC003970.32_AT)     | gb AAC33210.1  (AC003970) Highly similar to cinnamyl alcohol dehydrogenase, gi 1143445 [Arabidopsis thaliana] | 0           |             | dehydro-genase  |
| 19623_at<br>(AF000657.40_AT)     | gb AAB72175.1  (AF000657) cytochrome C [Arabidopsis thaliana]   | 2E-63       |             |                 |
| 19624_at<br>(AL049481.196_AT)    | emb CAB39628.1 <br>(AL049481) cytochrome c [Arabidopsis thaliana]   | 4E-63       |             |                 |
| 19625_s_at<br>(AC002311.26_S_AT) | gb AAC00610.1  (AC002311) Putative sulphate transporter protein#protein [Arabidopsis thaliana]                | 0           |             |                 |
| 19635_at<br>(AL049746.38_AT)     | emb CAB41856.1 <br>(AL049746) ABC-type transport-like protein [Arabidopsis thaliana]                          | 0           |             |                 |

| ProbeSet                              | Description   | Blast Score | EC #       | Family             |
|---------------------------------------|---|-------------|------------|--------------------|
| 19639_at<br>(AL080252.22_AT)          | emb CAB45788.1 <br>(AL080252) putative protein<br>[Arabidopsis thaliana]                            | 0           |            |                    |
| 19640_at<br>(AC004561.78_AT)          | gb AAC95192.1  (AC004561)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]       | 1E-118      |            | transferase        |
| 19641_at<br>(AC004561.66_AT)          | gb AAC95189.1  (AC004561)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]       | 1E-128      |            | transferase        |
| 19645_at<br>(AC004561.70_AT)          | gb AAC95190.1  (AC004561)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]       | 1E-130      |            | transferase        |
| 19646_s_at<br>(AC005819.55_S_A<br>T)  | gb AAC69925.1  (AC005819)<br>homeodomain transcription<br>factor (ATHB-7)<br>[Arabidopsis thaliana] | 9E-57       |            |                    |
| 19655_at<br>(Y14199.1_AT)             | emb CAA74591.1  (Y14199)<br>MAP3K delta-1 protein kinase<br>[Arabidopsis thaliana]                  | 0           |            | kinase             |
| 19667_at<br>(AL021710.5_AT)           | emb CAA16716.1 <br>(AL021710) glycolate oxidase<br>- like protein [Arabidopsis<br>thaliana]         | 0           | EC_1.1.2.3 | dehydro-<br>genase |
| 19672_at<br>(AC005687.19_AT)          | gb AAC36019.1  (AC005687)<br>RAP2.6 [Arabidopsis<br>thaliana]                                       | 1E-105      |            |                    |
| 19673_g_at<br>(AC005687.19_G_A<br>T)  | gb AAC36019.1  (AC005687)<br>RAP2.6 [Arabidopsis<br>thaliana]                                       | 1E-105      |            |                    |
| 19700_s_at<br>(AL031326.154_S_A<br>T) | emb CAA20463.1 <br>(AL031326) putative protein<br>[Arabidopsis thaliana]                            | 8E-85       |            |                    |
| 19701_s_at<br>(AC005724.67_S_A<br>T)  | gb AAD08939.1  (AC005724)<br>putative trehalose-6-phosphate<br>synthase [Arabidopsis<br>thaliana]   | 0           |            |                    |
| 19704_i_at<br>(AJ005927.2_I_AT)       | emb CAA06769.1 <br>(AJ005927) squalene<br>epoxidase homologue<br>[Arabidopsis thaliana]             | 0           |            | epoxidase          |

| ProbeSet                             | Description  | Blast Score | EC #       | Family              |
|--------------------------------------|--|-------------|------------|---------------------|
| 19707_s_at<br>(Z95768.3_S_AT)        | emb CAB09200.1  (Z95768)<br>R2R3-MYB transcription<br>factor [Arabidopsis thaliana]                                    | 4E-21       |            |                     |
| 19741_at<br>(AL049171.72_AT)         | emb CAB38956.1 <br>(AL049171) pyrophosphate-<br>dependent phosphofructo-1-<br>kinase [Arabidopsis thaliana]            | 0           |            | kinase              |
| 19755_at<br>(AC006593.64_AT)         | gb AAD20668.1  (AC006593)<br>ethylene reponse factor-like<br>AP2 domain transcription<br>factor [Arabidopsis thaliana] | 8E-97       |            |                     |
| 19762_at<br>(AL035527.204_AT)        | emb CAB36812.1 <br>(AL035527) peptide<br>transporter-like protein<br>[Arabidopsis thaliana]                            | 0           |            |                     |
| 19818_i_at<br>(AL021749.33_I_AT<br>) | emb CAA16876.2 <br>(AL021749) hypothetical<br>protein [Arabidopsis thaliana]   | 0           |            |                     |
| 19819_s_at<br>(AL021749.33_S_AT<br>) | emb CAA16876.2 <br>(AL021749) hypothetical<br>protein [Arabidopsis thaliana]   | 0           |            |                     |
| 19844_at<br>(AJ007588.2_AT)          | emb CAB37510.1 <br>(AL035540) monooxygenase<br>2 (MO2) [Arabidopsis<br>thaliana]                                       | 0           |            | mono-<br>oxygenase  |
| 19845_g_at<br>(AJ007588.2_G_AT)      | emb CAB37510.1 <br>(AL035540) monooxygenase<br>2 (MO2) [Arabidopsis<br>thaliana]                                       | 0           |            |                     |
| 19848_s_at<br>(AC004261.94_S_A<br>T) | dbj BAA08282.1  (D45848)<br>calmodulin-related protein<br>[Arabidopsis thaliana]                                       | 0           |            |                     |
| 19851_at<br>(U23794.3_AT)            | gb AAB60293.1  (U23794)<br>ILR1 [Arabidopsis thaliana]   | 0           | EC_3.5.-.- | amido-<br>hydrolase |
| 19878_at<br>(AL080252.102_AT)        | emb CAB45799.1 <br>(AL080252) nodulin-like<br>protein [Arabidopsis thaliana]   | 0           |            |                     |
| 19879_s_at<br>(Z97338.342_S_AT)      | emb CAB10321.1  (Z97338)<br>UFD1 like protein<br>[Arabidopsis thaliana]  | 0           |            |                     |
| 19881_at<br>(AC004077.49_AT)         | gb AAC26705.1  (AC004077)<br>putative trans-<br>prenyltransferase [Arabidopsis<br>thaliana]                            | 1E-154      |            | synthetase          |

| ProbeSet                             | Description  | Blast Score | EC # | Family             |
|--------------------------------------|--|-------------|------|--------------------|
| 19892_at<br>(AC005770.30_AT)         | gb AAC79626.1  (AC005770)<br>putative protease inhibitor<br>[Arabidopsis thaliana]   | 2E-36       |      | protease           |
| 19894_at<br>(AJ001809.1_AT)          | emb CAA05025.1 <br>(AJ001809) succinate<br>dehydrogenase flavoprotein<br>alpha subunit [Arabidopsis<br>thaliana]   | 0           |      | dehydro-<br>genase |
| 19895_s_at<br>(U77347.4_S_AT)        | gb AAC49679.1  (U77347)<br>lethal leaf-spot 1 homolog<br>[Arabidopsis thaliana]  | 0           |      |                    |
| 19916_at<br>(AC006577.34_AT)         | gb AAD25781.1 AC006577_1<br>7 (AC006577) EST<br>gb R64848 comes from this<br>gene. [Arabidopsis thaliana]  | 0           |      |                    |
| 19944_at<br>(AC002130.4_AT)          | gb AAB95233.1  (AC002130)<br>F1N21.4 [Arabidopsis<br>thaliana]   | 1E-128      |      |                    |
| 19946_at<br>(AC004482.14_AT)         | gb AAF18611.1 AC005170_1<br>(AC005170) similar to<br>senescence-associated protein<br>[Arabidopsis thaliana]   | 4E-92       |      |                    |
| 19956_at<br>(AC006282.11_AT)         | gb AAD20139.1  (AC006282)<br>unknown protein [Arabidopsis<br>thaliana]   | 1E-123      |      |                    |
| 19960_at<br>(AL035527.360_AT)        | emb CAB36823.1 <br>(AL035527) putative protein<br>[Arabidopsis thaliana]   | 1E-128      |      |                    |
| 19970_s_at<br>(AC003674.10_S_A<br>T) | gb AAB97120.1  (AC003674)<br>unknown protein [Arabidopsis<br>thaliana]   | 1E-154      |      |                    |
| 19982_at<br>(AC002986.28_AT)         | gb AAC17040.1  (AC002986)<br>Similarity to A. thaliana gene<br>product F21M12.20,<br>gb AC000132. EST<br>gb Z25651 comes from this<br>gene. [Arabidopsis thaliana] | 0           |      |                    |
| 19991_at<br>(AC007017.124_AT)        | gb AAD21459.1  (AC007017)<br>similar to harpin-induced<br>protein hin1 from tobacco<br>[Arabidopsis thaliana]  | 1E-112      |      |                    |
| 20017_at<br>(AC004521.66_AT)         | gb AAC16079.1  (AC004521)<br>unknown protein [Arabidopsis<br>thaliana]   | 1E-101      |      |                    |

| ProbeSet                             | Description  | Blast Score | EC #       | Family                      |
|--------------------------------------|--|-------------|------------|-----------------------------|
| 20023_at<br>(AC006577.46_AT)         | gb AAD25787.1 AC006577_2<br>3 (AC006577) Similar to<br>gi 1653162 (p)ppGpp 3-<br>pyrophosphohydrolase from<br>Synechocystis sp genome<br>gb D90911. EST gb W43807<br>comes from this gene.<br>[Arabidopsis thaliana] | 0           | EC_2.7.6.5 | pyro-<br>phospho-<br>kinase |
| 20030_at<br>(AL078637.51_AT)         | emb CAB45058.1 <br>(AL078637) putative protein<br>[Arabidopsis thaliana]   | 1E-168      |            |                             |
| 20051_at<br>(AC000106.38_AT)         | gb AAB70412.1  (AC000106)<br>Similar to Saccharomyces<br>hypothetical protein YDR051c<br>(gb Z49209). ESTs<br>gb T44436,gb 42252 come<br>from this gene. [Arabidopsis<br>thaliana]                                   | 1E-166      |            |                             |
| 20053_at<br>(AC002292.27_AT)         | gb AAB71973.1  (AC002292)<br>Unknown protein<br>[Arabidopsis thaliana]   | 0           |            |                             |
| 20096_at<br>(AC004238.31_AT)         | gb AAC12821.1  (AC004238)<br>putative berberine bridge<br>enzyme [Arabidopsis thaliana]  | 0           |            |                             |
| 20098_at<br>(AC004697.123_AT)        | gb AAC28986.1  (AC004697)<br>similar to latex allergen from<br>Hevea brasiliensis<br>[Arabidopsis thaliana]  | 0           |            |                             |
| 20133_i_at<br>(AC007178.71_I_AT<br>) | gb AAD28682.1 AC007178_2<br>(AC007178) hypothetical<br>protein [Arabidopsis thaliana]  | 0           |            |                             |
| 20134_s_at<br>(AC007178.71_S_A<br>T) | gb AAD28682.1 AC007178_2<br>(AC007178) hypothetical<br>protein [Arabidopsis thaliana]  | 0           |            |                             |
| 20142_at<br>(AL035521.155_AT)        | emb CAB36716.1 <br>(AL035521) extra-large G-<br>protein-like [Arabidopsis<br>thaliana]   | 0           |            |                             |
| 20144_at<br>(AL079350.68_AT)         | emb CAB45516.1 <br>(AL079350) receptor kinase-<br>like protein [Arabidopsis<br>thaliana]   | 0           |            | kinase                      |
| 20165_at<br>(AC002311.16_AT)         | gb AAC00605.1  (AC002311)<br>Unknown protein<br>[Arabidopsis thaliana]   | 6E-75       |            |                             |



| ProbeSet                             | Description  | Blast Score | EC #        | Family      |
|--------------------------------------|--|-------------|-------------|-------------|
| 20179_at<br>(AL035538.229_AT)        | emb CAB37546.1 <br>(AL035538) putative protei<br>[Arabidopsis thaliana]  | 2E-63       |             |             |
| 20189_at<br>(AC005489.2_AT)          | gb AAD32864.1 AC005489_2<br>(AC005489) F14N23.2<br>[Arabidopsis thaliana]  | 2E-57       |             |             |
| 20194_at<br>(AC007584.48_AT)         | gb AAD32907.1 AC007584_5<br>(AC007584) unknown protein<br>[Arabidopsis thaliana]   | 0           |             |             |
| 20200_at<br>(AL050400.67_AT)         | emb CAB43700.1 <br>(AL050400) hypothetical<br>protein [Arabidopsis thaliana]   | 4E-95       |             |             |
| 20215_s_at<br>(AF117125.2_S_AT)      | gb AAD29957.1  (AF117125)<br>endoplasmic reticulum-type<br>calcium-transporting ATPase<br>4 [Arabidopsis thaliana]                       | 0           |             |             |
| 20223_at<br>(AL022347.145_AT)        | emb CAA18469.1 <br>(AL022347) serine/threonine<br>kinase-like protein<br>[Arabidopsis thaliana]  | 0           |             | kinase      |
| 20238_at<br>(X74514.2_AT)            | emb CAA52619.1  (X74514)<br>beta-fructofuranosidase<br>[Arabidopsis thaliana]  | 0           | EC_3.2.1.26 | hydrolase   |
| 20239_g_at<br>(X74514.2_G_AT)        | emb CAA52619.1  (X74514)<br>beta-fructofuranosidase<br>[Arabidopsis thaliana]  | 0           |             |             |
| 20245_s_at<br>(AC005309.97_S_A<br>T) | emb CAA05625.1 <br>(AJ002584) AtMRP4<br>[Arabidopsis thaliana]<br>thaliana]  | 0           |             |             |
| 20246_s_at<br>(AF084037.3_S_AT)      | gb AAC95354.1  (AF084037)<br>receptor-like protein kinase<br>[Arabidopsis thaliana]  | 0           |             |             |
| 20247_at<br>(AC004392.4_AT)          | gb AAC28501.1  (AC004392)<br>Similar to beta-glucosidase<br>BGQ60 precursor gb L41869<br>from Hordeum vulgare.<br>[Arabidopsis thaliana] | 0           | EC_3.2.1.21 | glucosidase |
| 20258_at<br>(AF130252.1_AT)          | gb AAD28759.1 AF130252_1<br>(AF130252) calcium<br>dependent protein kinase CP4<br>[Arabidopsis thaliana]                                 | 0           |             | kinase      |

| ProbeSet                             | Description  | Blast Score | EC # | Family                   |
|--------------------------------------|--|-------------|------|--------------------------|
| 20262_at<br>(AC002294.26_AT)         | gb AAB71480.1  (AC002294)<br>Similar to transcription factor<br>gb Z46606 1658307 and others<br>[Arabidopsis thaliana]         | 0           |      |                          |
| 20263_at<br>(AB004798.1_AT)          | dbj BAA20519.1  (AB004798)<br>ascorbate oxidase<br>[Arabidopsis thaliana]  | 0           |      | oxidase                  |
| 20269_at<br>(AC002387.237_AT)        | gb AAB82640.1  (AC002387)<br>putative pectinesterase<br>[Arabidopsis thaliana]   | 0           |      | pectin-<br>esterase      |
| 20271_at<br>(Z99707.27_AT)           | emb CAB16771.1  (Z99707)<br>cytochrome P450-like protein<br>[Arabidopsis thaliana]   | 0           |      |                          |
| 20285_s_at<br>(AC003674.18_S_A<br>T) | gb AAB97121.1  (AC003674)<br>putative protein kinase<br>[Arabidopsis thaliana]   | 0           |      |                          |
| 20287_at<br>(Y14590.5_AT)            | emb CAA74930.1  (Y14590)<br>class IV chitinase<br>[Arabidopsis thaliana]   | 1E-166      |      | chitinase                |
| 20288_g_at<br>(Y14590.5_G_AT)        | emb CAA74930.1  (Y14590)<br>class IV chitinase<br>[Arabidopsis thaliana]   | 1E-166      |      |                          |
| 20291_s_at<br>(M92353.4_S_AT)        | gb AAA32738.1  (M92353)<br>anthranilate synthase alpha<br>subunit [Arabidopsis thaliana]                                       | 0           |      |                          |
| 20297_at<br>(AC007153.27_AT)         | gb AAD30627.1 AC007153_1<br>9 (AC007153) Similar to<br>indole-3-acetate beta-<br>glucosyltransferase<br>[Arabidopsis thaliana] | 0           |      | glucosyl-<br>transferase |
| 20323_at<br>(AC004561.62_AT)         | gb AAC95188.1  (AC004561)<br>putative small heat shock<br>protein [Arabidopsis thaliana]                                       | 1E-67       |      |                          |
| 20346_at<br>(AL031135.156_AT)        | emb CAA20030.1 <br>(AL031135) protein kinase -<br>like protein [Arabidopsis<br>thaliana]                                       | 0           |      |                          |
| 20348_at<br>(AC005967.35_AT)         | gb AAD03382.1  (AC005967)<br>putative limonene cyclase<br>[Arabidopsis thaliana]   | 0           |      | cyclase                  |
| 20356_at<br>(AC004561.74_AT)         | gb AAC95191.1  (AC004561)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]                                  | 1E-128      |      | transferase              |

| ProbeSet                         | Description   | Blast Score | EC # | Family                |
|----------------------------------|---|-------------|------|-----------------------|
| 20365_s_at<br>(AC005850.19_S_AT) | gb AAD25552.1 AC005850_9<br>(AC005850) Highly Similar<br>to Mlo proteins [Arabidopsis<br>thaliana]      | 0           |      |                       |
| 20370_at<br>(AC004561.263_AT)    | gb AAC95219.1  (AC004561)<br>putative tropinone reductase<br>[Arabidopsis thaliana]                     | 1E-151      |      | reductase             |
| 20382_s_at<br>(AC002338.35_S_AT) | gb AAC16930.1  (AC002338)<br>putative WRKY-type DNA<br>binding protein [Arabidopsis<br>thaliana]        | 0           |      |                       |
| 20420_at<br>(AL024486.131_AT)    | emb CAA19698.1 <br>(AL024486) putative chitinase<br>[Arabidopsis thaliana]                              | 0           |      | chitinase             |
| 20421_at<br>(U81294.2_AT)        | emb CAB10242.1  (Z97336)<br>germin precursor oxalate<br>oxidase [Arabidopsis thaliana]                  | 8E-98       |      |                       |
| 20422_g_at<br>(U81294.2_G_AT)    | emb CAB10242.1  (Z97336)<br>germin precursor oxalate<br>oxidase [Arabidopsis thaliana]                  | 8E-98       |      |                       |
| 20432_at<br>(U43486.2_AT)        | gb AAB18365.1  (U43486)<br>xyloglucan<br>endotransglycosylase-related<br>protein [Arabidopsis thaliana] | 1E-174      |      | trans-<br>glycosylase |
| 20433_at<br>(AC006232.147_AT)    | gb AAD15611.1  (AC006232)<br>putative beta-1,3-glucanase<br>[Arabidopsis thaliana]                      | 0           |      | glucanase             |
| 20450_at<br>(AJ005930.2_AT)      | emb CAA06772.1 <br>(AJ005930) squalene<br>epoxidase homologue<br>[Arabidopsis thaliana]                 | 0           |      | epoxidase             |
| 20461_at<br>(AL049480.157_AT)    | emb CAB39609.1 <br>(AL049480) pumilio-like<br>protein [Arabidopsis thaliana]                            | 0           |      |                       |
| 20462_at<br>(U82399.2_AT)        | gb AAB40725.1  (U82399)<br>putative protein kinase PK1<br>[Arabidopsis thaliana]                        | 3E-38       |      | kinase                |
| 20479_i_at<br>(AF069495.2_I_AT)  | emb CAB38908.1 <br>(AL035708) cytochrome<br>P450-like protein<br>[Arabidopsis thaliana]                 | 0           |      |                       |
| 20480_s_at<br>(AF069495.2_S_AT)  | emb CAB38908.1 <br>(AL035708) cytochrome<br>P450-like protein<br>[Arabidopsis thaliana]                 | 0           |      |                       |

| ProbeSet                              | Description  | Blast Score | EC #             | Family    |
|---------------------------------------|--|-------------|------------------|-----------|
| 20485_at<br>(AC007660.131_AT)         | gb AAD32811.1 AC007660_1<br>2 (AC007660) putative two-<br>component response regulator<br>protein [Arabidopsis thaliana] | 1E-80       |                  |           |
| 20491_at<br>(AC004561.146_AT)         | gb AAC95203.1  (AC004561)<br>putative tropinone reductase<br>[Arabidopsis thaliana]                                      | 1E-153      | EC_1.1.1.10<br>0 | reductase |
| 20511_at<br>(AC007290.24_AT)          | gb AAD26884.1 AC007290_3<br>(AC007290) putative<br>nucleotide-binding protein<br>[Arabidopsis thaliana]                  | 0           |                  |           |
| 20517_at<br>(Y17722.7_AT)             | emb CAB50690.1  (Y17722)<br>telomere repeat-binding<br>protein TRP1 [Arabidopsis<br>thaliana]                            | 0           |                  |           |
| 20529_at<br>(Z97341.125_AT)           | emb CAB10426.1  (Z97341)<br>cysteine proteinase inhibitor<br>like protein [Arabidopsis<br>thaliana]                      | 7E-35       |                  |           |
| 20551_at<br>(AC006081.211_AT)         | gb AAD24395.1 AC006081_7<br>(AC006081) unknown protein<br>[Arabidopsis thaliana]   | 0           |                  |           |
| 20572_s_at<br>(AC005560.229_S_A<br>T) | gb AAD12710.1  (AC006069)<br>unknown protein [Arabidopsis<br>thaliana]   | 1E-114      |                  |           |
| 20577_at<br>(AL078464.72_AT)          | emb CAB43841.1 <br>(AL078464) putative protein<br>[Arabidopsis thaliana]   | 0           |                  |           |
| 20584_at<br>(AC004450.75_AT)          | gb AAC64305.1  (AC004450)<br>putative clathrin binding<br>protein (epsin) [Arabidopsis<br>thaliana]                      | 2E-92       |                  |           |
| 20586_i_at<br>(AC005824.195_I_A<br>T) | gb AAC73042.1  (AC005824)<br>unknown protein [Arabidopsis<br>thaliana]   | 2E-67       |                  |           |
| 20587_s_at<br>(AC005824.195_S_A<br>T) | gb AAC73042.1  (AC005824)<br>unknown protein [Arabidopsis<br>thaliana]   | 2E-67       |                  |           |
| 20589_at<br>(AF081066.3_AT)           | gb AAC31939.1  (AF081066)<br>IAA-amino acid hydrolase<br>homolog ILL3 [Arabidopsis<br>thaliana]                          | 0           |                  | hydrolase |
| 20591_at<br>(AL080252.115_AT)         | emb CAB45800.1 <br>(AL080252) nodulin-like<br>protein [Arabidopsis thaliana]   | 0           |                  |           |

| ProbeSet                             | Description   | Blast Score | EC #       | Family             |
|--------------------------------------|---|-------------|------------|--------------------|
| 20646_at<br>(AC002291.20_AT)         | gb AAC00619.1  (AC002291)<br>Unknown protein<br>[Arabidopsis thaliana]                                    | 0           |            |                    |
| 20656_at<br>(AL035396.46_AT)         | emb CAA23064.1 <br>(AL035396) putative protein<br>[Arabidopsis thaliana]                                  | 0           |            |                    |
| 20658_s_at<br>(AL050400.70_S_AT<br>) | emb CAB43701.1 <br>(AL050400) beta-carotene<br>hydroxylase [Arabidopsis<br>thaliana]                      | 1E-165      |            |                    |
| 20669_s_at<br>(AC002388.6_S_AT)      | gb AAD32838.1 AC007659_2<br>0 (AC007659) unknown<br>protein [Arabidopsis thaliana]                        | 0           |            |                    |
| 20685_at<br>(AL049751.46_AT)         | emb CAB41928.1 <br>(AL049751) short-chain<br>alcohol dehydrogenase like<br>protein [Arabidopsis thaliana] | 1E-139      | EC_1.1.1.- | dehydro-<br>genase |
| 20686_at<br>(Y14424.2_AT)            | emb CAA74766.1  (Y14424)<br>hypothetical protein<br>[Arabidopsis thaliana]                                | 0           |            |                    |
| 20689_s_at<br>(AC002335.19_S_A<br>T) | gb AAB64310.1  (AC002335)<br>putative calcium binding<br>protein [Arabidopsis thaliana]                   | 6E-95       |            |                    |
| 20715_at<br>(AF079183.1_AT)          | gb AAC69857.1  (AF079183)<br>RING-H2 finger protein<br>RHG1a [Arabidopsis thaliana]                       | 4E-99       |            |                    |

**Table 4a** Probe Sets in addition to those in Table 4b corresponding to genes, the expression of which is induced by *Pseudomonas* infection in wild-type *Arabidopsis* plants and perturbed in at least one mutant plant

| ProbeSet                        | Description  | Blast Score |
|---------------------------------|--|-------------|
| 13716_at<br>(NOVARTIS103_RC_AT) | emb CAA62665.1  (X91259) lectin like protein [ <i>Arabidopsis thaliana</i> ]                             | 6E-70       |
| 13751_at<br>(NOVARTIS127_AT)    | gb AAF16751.1 AC010155_4 (AC010155) F3M18.8 [ <i>Arabidopsis thaliana</i> ]                              | 2E-27       |
| 13755_at<br>(NOVARTIS15_AT)     | emb CAA16797.1  (AL021713) receptor serine/threonine kinase-like protein [ <i>Arabidopsis thaliana</i> ] | 1E-114      |
| 13763_at<br>(NOVARTIS21_AT)     | gb AAF24849.1 AC012679_20 (AC012679) putative calmodulin-binding protein [ <i>Arabidopsis thaliana</i> ] | 1E-105      |
| 13764_at<br>(NOVARTIS22_AT)     | gb AAD39641.1 AC007591_6 (AC007591) F9L1.6 [ <i>Arabidopsis thaliana</i> ]                               | 2E-30       |
| 14139_at<br>(NOVARTIS30_AT)     | gb AAD09343.1  (AF026538) ABA-responsive protein [ <i>Hordeum vulgare</i> ]                              | 7E-36       |
| 14148_at<br>(NOVARTIS38_AT)     | gb AAF34713.1 AF224762_1 (AF224762) SigA binding protein [ <i>Arabidopsis thaliana</i> ]                 | 4E-53       |
| 14170_at<br>(NOVARTIS51_AT)     | gb AAF29406.1 AC022354_5 (AC022354) unknown protein [ <i>Arabidopsis thaliana</i> ]                      | 9E-26       |
| 14197_at<br>(NOVARTIS71_AT)     | No hits found.   |             |
| 14214_at<br>(NOVARTIS83_AT)     | gb AAF24849.1 AC012679_20 (AC012679) putative calmodulin-binding protein [ <i>Arabidopsis thaliana</i> ] | 2E-92       |
| 14223_at (NOVARTIS9_AT)         | emb CAA19683.1  (AL024486) putative protein [ <i>Arabidopsis thaliana</i> ]                              | 0           |

**Table 4b** Probe Sets as referred to in Table 3 corresponding to genes, the expression of which is induced by *Pseudomonas* infection in wild-type *Arabidopsis* plants and perturbed in at least one mutant plant

|                            |                            |
|----------------------------|----------------------------|
| 11997_at (AC005967.4_AT)   | 12004_at (AL022023.132_AT) |
| 12007_at (Z99708.249_AT)   | 12037_at (AC004005.174_AT) |
| 12072_at (AL035396.4_AT)   | 12079_s_at (A71597.1_S_AT) |
| 12081_at (AC001645.140_AT) | 12115_at (AL033545.26_AT)  |
| 12136_at (AC007591.60_AT)  | 12150_at (AC004005.151_AT) |
| 12198_at (AC006954.90_AT)  | 12216_at (AC007119.56_AT)  |
| 12233_at (AJ001807.1_AT)   | 12307_at (AC002392.162_AT) |

|   |                                |
|---|--------------------------------|
| 12314_at (AC001229.28_AT)                 | 12317_at (AC004138.27_AT)      |
| 12323_at (AC002333.18_AT)                 | 12332_s_at (AB023448.2_S_AT)   |
| 12341_s_at (AL021637.176_S_AT)            | 12349_s_at (X84728.6_S_AT)     |
| 12369_at (AC002535.59_AT)                 | 12400_at (X98453.1_AT)         |
| 12438_at (AL021710.83_AT)                 | 12449_s_at (AC002343.179_S_AT) |
| 12454_at (AC006232.164_AT)                | 12475_at (Y11794.1_AT)         |
| 12487_at (AC004411.126_AT)                | 12497_at (AC006533.51_AT)      |
| 12500_s_at (AF081067.3_S_AT)              | 12525_at (AC006587.85_AT)      |
| 12530_at (Z99707.184_AT)                  | 12535_at (AL035538.156_AT)     |
| 12538_at (AF033205.2_AT)                  | 12574_at (X82624.2_AT)         |
| 12584_at (AC004521.233_AT)                | 12626_at (AC006234.95_AT)      |
| 12645_at (AL021712.56_AT)                 | 12712_f_at (Z95774_F_AT)       |
| 12736_f_at (Z97048_F_AT)                  | 12744_at (AC001645.15_AT)      |
| 12760_g_at (AC005278.32_G_AT)             | 12764_f_at (AC004138.69_F_AT)  |
| 12772_at (AC005278.34_AT)                 | 12797_s_at (AC007138.25_S_AT)  |
| 12851_s_at (ACCSYN1_S_AT)                 | 12879_s_at (AIG1_S_AT)         |
| 12880_s_at (AIG2_S_AT)                    | 12883_s_at (APX_S_AT)          |
| 12889_s_at (ASA1_S_AT)                    | 12891_at (ATACS6_AT)           |
| 12892_g_at (ATACS6_G_AT)                  | 12904_s_at (ATERF1_S_AT)       |
| 12905_s_at (ATERF2_S_AT)                  | 12911_s_at (ATG6PDHE5_S_AT)    |
| 12921_s_at (ATHHMGCOAR_S_AT)              | 12951_at (AC005489.5_AT)       |
| 12958_at (AC002332.249_AT)                | 12965_at (AL021711.118_AT)     |
| 12966_s_at (AL023094.197_S_AT)            | 12989_s_at (AC004077.149_S_AT) |
| 13003_s_at (AB021936.1_S_AT)              | 13014_at (U93215.87_AT)        |
| 13040_at (AC002392.134_AT)                | 13070_at (AC006919.171_AT)     |
| 13094_at (AL035523.163_AT)                | 13134_s_at (AC002337.9_S_AT)   |
| 13152_s_at (AC005322.24_S_AT)             | 13154_s_at (AC002333.210_S_AT) |
| 13157_at (AC002409.35_AT)                 | 13176_at (AL031394.56_AT)      |
| 13177_at (AL049640.42_AT)                 | 13190_s_at (ATTHIREDA_S_AT)    |
| 13211_s_at (BCHI_S_AT)                    | 13212_s_at (BGL2_S_AT)         |
| 13215_s_at (cafferoylcoamethyltrans_S_AT) | 13219_s_at (CHI4_S_AT)         |
| 13243_r_at (ELI32_R_AT)                   | 13244_s_at (ELI32_S_AT)        |
| 13255_i_at (gammaglutamyltranspepti_I_AT) | 13266_s_at (GST4_S_AT)         |
| 13273_s_at (HSF4_S_AT)                    | 13275_f_at (HSP174_F_AT)       |
| 13277_i_at (HSP176A_I_AT)                 | 13285_s_at (HSP83_S_AT)        |
| 13312_at (AC006223.75_AT)                 | 13370_at (AC005322.4_AT)       |
| 13381_at (AC006580.8_AT)                  | 13395_at (AL035528.202_AT)     |
| 13435_at (AF003102.3_AT)                  | 13437_at (AF096371.8_AT)       |

|   |                                |
|---|--------------------------------|
| 13467_at (AL096860.198_AT)              | 13536_at (AL021636.47_AT)      |
| 13538_at (AL080254.75_AT)               | 13565_at (AL035601.21_AT)      |
| 13588_at (AL021961.24_AT)               | 13589_at (AC000132.24_AT)      |
| 13617_at (AC006592.64_AT)               | 13627_at (AL035394.196_AT)     |
| 13645_at (AC000098.8_AT)                | 13656_at (AC007138.31_AT)      |
| 13659_at (AL022347.46_AT)               | 13666_s_at (INDOLE3GPS_S_AT)   |
| 13680_s_at (LOX1_S_AT)                  | 13685_s_at (MLOLIKE2_S_AT)     |
| 13688_s_at (MONOPTEROS_S_AT)            | 13697_at (NI16_AT)             |
| 13705_s_at (AC003671X_S_AT)             | 13789_at (AJ132436.2_AT)       |
| 13803_at (Z97341.376_AT)                | 13818_s_at (AC006218.175_S_AT) |
| 13842_at (AC002396.12_AT)               | 13848_at (AC003981.31_AT)      |
| 13880_s_at (AL049480.183_S_AT)          | 13908_s_at (A71590.1_S_AT)     |
| 13918_at (AC005388.29_AT)               | 13920_at (AC005990.53_AT)      |
| 13949_s_at (Z97343.352_S_AT)            | 13963_at (AL021711.26_AT)      |
| 13999_at (AF071527.56_AT)               | 14015_s_at (A71588.1_S_AT)     |
| 14016_s_at (A71596.1_S_AT)              | 14025_s_at (AC007293.3_S_AT)   |
| 14030_at (AC005970.225_AT)              | 14032_at (AL035601.11_AT)      |
| 14041_at (AC003970.28_AT)               | 14052_at (AC004122.24_AT)      |
| 14068_s_at (AC006922.197_S_AT)          | 14070_at (AL049658.217_AT)     |
| 14089_at (AC006223.65_AT)               | 14100_at (AF002109.108_AT)     |
| 14110_i_at (AL035528.279_I_AT)          | 14240_s_at (NR1_S_AT)          |
| 14242_s_at (NRA_S_AT)                   | 14248_at (PAD3_AT)             |
| 14249_i_at (PAD4_I_AT)                  | 14254_s_at (PAL1-MRNA_S_AT)    |
| 14256_f_at (PAL1-INTRON_F_AT)           | 14320_at (AC005956.54_AT)      |
| 14408_at (AC002291.14_AT)               | 14448_at (AC002387.243_AT)     |
| 14450_at (AC002986.49_AT)               | 14461_at (AC006202.73_AT)      |
| 14468_at (AC007576.62_AT)               | 14475_at (AL021811.121_AT)     |
| 14487_at (Z97341.343_AT)                | 14584_at (AC007658.25_AT)      |
| 14609_at (AC002340.147_AT)              | 14620_s_at (PAT1_S_AT)         |
| 14635_s_at (PR.1_S_AT)                  | 14638_s_at (PRXCB_S_AT)        |
| 14640_s_at (putativemloHI_S_AT)         | 14643_s_at (RAR047_S_AT)       |
| 14663_s_at (trehalaseprecursor_RC_S_AT) | 14667_s_at (TRPB_S_AT)         |
| 14672_s_at (TSA1_S_AT)                  | 14673_s_at (TSB2_S_AT)         |
| 14675_s_at (VSP_S_AT)                   | 14705_i_at (WT77_RC_I_AT)      |
| 14706_r_at (WT77_RC_R_AT)               | 14711_s_at (ZFPL_S_AT)         |
| 14735_s_at (AF008124_S_AT)              | 14750_s_at (AF096370.12_S_AT)  |
| 14763_at (X86958.1_AT)                  | 14779_at (AC004680.71_AT)      |
| 14780_at (AC004683.103_AT)              | 14786_at (AC005397.115_AT)     |



|                                |                                |
|--------------------------------|--------------------------------|
| 14793_at (AC006202.10_AT)      | 14882_at (AL022605.63_AT)      |
| 14900_at (AC000348.12_AT)      | 14923_at (AC006283.158_AT)     |
| 14924_at (AC006283.46_AT)      | 14928_at (AC006569.88_AT)      |
| 14959_at (AC007202.26_AT)      | 14978_at (AC002333.49_AT)      |
| 15052_at (AC002332.103_AT)     | 15073_at (AC007069.93_AT)      |
| 15088_s_at (AC002311.37_S_AT)  | 15091_at (AC004683.97_AT)      |
| 15098_s_at (ATU26945_S_AT)     | 15116_f_at (AF121356_F_AT)     |
| 15123_s_at (ATU40857_S_AT)     | 15124_s_at (ATU59508_S_AT)     |
| 15125_f_at (D85190_F_AT)       | 15132_s_at (AF121878_S_AT)     |
| 15137_s_at (ATU57320_S_AT)     | 15141_s_at (D85191_S_AT)       |
| 15154_s_at (ATHMTGDAS_S_AT)    | 15161_s_at (ATU90522_S_AT)     |
| 15162_s_at (U01880_S_AT)       | 15188_s_at (AF081202_S_AT)     |
| 15196_s_at (ATU43412_S_AT)     | 15199_s_at (AB005804_S_AT)     |
| 15211_s_at (ATH243813_S_AT)    | 15216_s_at (ATU75191_S_AT)     |
| 15342_at (AC006593.101_AT)     | 15406_at (AC006931.179_AT)     |
| 15431_at (AL030978.64_AT)      | 15479_at (AL049483.205_AT)     |
| 15483_s_at (AC005819.20_S_AT)  | 15485_at (AC006233.109_AT)     |
| 15496_at (AC006282.167_AT)     | 15518_at (AC005322.28_AT)      |
| 15522_i_at (AL078637.213_I_AT) | 15523_s_at (AL078637.213_S_AT) |
| 15524_at (AC005508.25_AT)      | 15526_at (AC004122.16_AT)      |
| 15531_i_at (AL078637.191_I_AT) | 15532_r_at (AL078637.191_R_AT) |
| 15540_at (AC006585.205_AT)     | 15543_at (AF096371.10_AT)      |
| 15544_at (AL021633.110_AT)     | 15547_at (AC005970.122_AT)     |
| 15551_at (AL035440.289_AT)     | 15594_s_at (ATU56635_S_AT)     |
| 15613_s_at (ATHHOMEOA_S_AT)    | 15622_s_at (ATU43945_S_AT)     |
| 15625_s_at (ATU74610_S_AT)     | 15632_s_at (AB012570_S_AT)     |
| 15641_s_at (AF117063_S_AT)     | 15665_s_at (AF022658_S_AT)     |
| 15669_s_at (AF047834_S_AT)     | 15670_s_at (AF061638_S_AT)     |
| 15672_s_at (AF082299_S_AT)     | 15778_at (X98676.2_AT)         |
| 15779_g_at (X98676.2_G_AT)     | 15792_at (AC002341.106_AT)     |
| 15798_at (AC002521.173_AT)     | 15815_s_at (Z97342.366_S_AT)   |
| 15839_at (AC005662.203_AT)     | 15859_at (AC006587.164_AT)     |
| 15866_s_at (AC007133.59_S_AT)  | 15874_at (AL022223.106_AT)     |
| 15886_at (AL078637.204_AT)     | 15919_at (AC007060.42_AT)      |
| 15921_s_at (AC007067.1_S_AT)   | 15924_at (AC007138.61_AT)      |
| 15970_s_at (X71794.2_S_AT)     | 15978_at (X68592.6_AT)         |
| 15982_s_at (AC006260.78_S_AT)  | 16001_at (AF035385.2_AT)       |
| 16003_s_at (AL021749.64_S_AT)  | 16021_s_at (AL022224.182_S_AT) |

|                                |                                |
|--------------------------------|--------------------------------|
| 16031_at (X94248.1_AT)         | 16043_at (AC005489.17_AT)      |
| 16053_i_at (Y14251.4_I_AT)     | 16063_s_at (AB008103_S_AT)     |
| 16073_f_at (AF062908_F_AT)     | 16083_s_at (AF153283_S_AT)     |
| 16091_s_at (ATHHSP83_S_AT)     | 16103_s_at (ATU60445_S_AT)     |
| 16105_s_at (ATU68017_S_AT)     | 16108_s_at (D78604_S_AT)       |
| 16130_s_at (AF078683_S_AT)     | 16173_s_at (D78607_S_AT)       |
| 16203_at (AC007519.53_AT)      | 16232_s_at (AL080252.77_S_AT)  |
| 16233_at (AL080254.83_AT)      | 16272_at (AC006304.136_AT)     |
| 16298_at (AL021890.71_AT)      | 16299_at (AL024486.185_AT)     |
| 16301_s_at (AL031018.105_S_AT) | 16306_at (AL049751.112_AT)     |
| 16335_at (AL079347.105_AT)     | 16340_at (AC004255.15_AT)      |
| 16357_at (AF149413.38_AT)      | 16363_at (AC004255.14_AT)      |
| 16383_at (AC006300.64_AT)      | 16398_s_at (AL022603.3_S_AT)   |
| 16405_at (AC005850.9_AT)       | 16409_at (AC004393.2_AT)       |
| 16440_s_at (AF002109.137_S_AT) | 16461_i_at (AC004683.79_I_AT)  |
| 16462_s_at (AC004683.79_S_AT)  | 16465_at (Y08892.1_AT)         |
| 16470_s_at (AF068299.4_S_AT)   | 16510_at (AL034567.198_AT)     |
| 16522_at (X77500.2_AT)         | 16526_at (Z49227.1_AT)         |
| 16541_s_at (AB023423_S_AT)     | 16545_s_at (AF037229_S_AT)     |
| 16553_f_at (AF078821_F_AT)     | 16570_s_at (ATHCDPKA_S_AT)     |
| 16578_s_at (ATHRPRP1B_S_AT)    | 16589_s_at (ATU26937_S_AT)     |
| 16594_s_at (ATU39783_S_AT)     | 16603_s_at (ATU81293_S_AT)     |
| 16609_s_at (AB008104_S_AT)     | 16611_s_at (AB008782_S_AT)     |
| 16638_s_at (AF139098_S_AT)     | 16646_s_at (ATHDHS1_S_AT)      |
| 16649_s_at (ATHORF_S_AT)       | 16701_at (AC005312.61_AT)      |
| 16721_at (AC006533.58_AT)      | 16747_at (AL021713.3_AT)       |
| 16781_at (AC002392.100_AT)     | 16810_at (AC002339.46_AT)      |
| 16859_at (AL035523.135_AT)     | 16864_i_at (AF037367.4_I_AT)   |
| 16865_s_at (AF037367.4_S_AT)   | 16888_s_at (AC004684.174_S_AT) |
| 16914_s_at (AL049500.57_S_AT)  | 16916_s_at (X77199.8_S_AT)     |
| 16968_at (AL021961.93_AT)      | 16989_at (AL030978.46_AT)      |
| 16995_at (AC002391.188_AT)     | 17007_at (AC005896.26_AT)      |
| 17008_at (AC006585.212_AT)     | 17039_s_at (D78602_S_AT)       |
| 17041_s_at (D89631_S_AT)       | 17051_s_at (AF098947_S_AT)     |
| 17066_s_at (ATHLIPOXY_S_AT)    | 17073_s_at (ATT54391_S_AT)     |
| 17075_s_at (ATU09961_S_AT)     | 17104_s_at (D88541_S_AT)       |
| 17111_s_at (ATHACSC_S_AT)      | 17119_s_at (AF132212_S_AT)     |
| 17128_s_at (ATHRPRP1A_S_AT)    | 17180_at (AF007270.30_AT)      |

|                                |                                |
|--------------------------------|--------------------------------|
| 17187_at (AF128396.2_AT)       | 17300_at (X66017.2_AT)         |
| 17303_s_at (AC004683.25_S_AT)  | 17323_at (U95973.69_AT)        |
| 17338_at (AC002535.97_AT)      | 17341_at (AL021713.89_AT)      |
| 17356_s_at (Z97338.190_S_AT)   | 17371_at (AF076243.44_AT)      |
| 17379_at (AF085279.9_AT)       | 17413_s_at (AJ006961.4_S_AT)   |
| 17451_at (AC002343.47_AT)      | 17452_g_at (AC002343.47_G_AT)  |
| 17484_at (X79052.2_AT)         | 17485_s_at (Z97340.345_S_AT)   |
| 17487_s_at (U18993.2_S_AT)     | 17511_s_at (AF067605_S_AT)     |
| 17514_s_at (AF076277_S_AT)     | 17522_s_at (D78606_S_AT)       |
| 17533_s_at (ATU43488_S_AT)     | 17544_s_at (ATU40856_S_AT)     |
| 17548_s_at (AF118823_S_AT)     | 17585_s_at (AF134487_S_AT)     |
| 17595_s_at (AF166352_S_AT)     | 17648_at (AL021684.43_AT)      |
| 17653_at (AL035679.144_AT)     | 17702_at (AC005700.212_AT)     |
| 17719_at (AC006592.17_AT)      | 17752_at (AC003974.37_AT)      |
| 17758_at (AF076243.41_AT)      | 17775_at (AC004392.2_AT)       |
| 17781_at (AL049746.177_AT)     | 17840_s_at (AC002333.223_S_AT) |
| 17860_at (AL078467.4_AT)       | 17876_at (AJ007587.2_AT)       |
| 17877_g_at (AJ007587.2_G_AT)   | 17893_at (AC004401.135_AT)     |
| 17899_at (Z97339.197_AT)       | 17930_s_at (AJ006960.4_S_AT)   |
| 17945_at (Z97341.411_AT)       | 17963_at (AL049730.88_AT)      |
| 18012_s_at (AJ002295_S_AT)     | 18054_at (AJ238846_AT)         |
| 18109_s_at (AC002391.206_S_AT) | 18121_s_at (AC002337.21_S_AT)  |
| 18122_at (AC002338.110_AT)     | 18148_at (AC004669.25_AT)      |
| 18176_at (AL035540.31_AT)      | 18194_i_at (AL096859.227_I_AT) |
| 18213_at (AL022140.126_AT)     | 18224_s_at (AL021890.57_S_AT)  |
| 18228_at (X91259.1_AT)         | 18236_s_at (AC004683.69_S_AT)  |
| 18242_g_at (AC006580.71_G_AT)  | 18255_at (AC005770.25_AT)      |
| 18258_s_at (AC006439.222_S_AT) | 18263_at (AC005724.36_AT)      |
| 18266_at (AC004684.33_AT)      | 18267_at (AC006223.23_AT)      |
| 18299_s_at (M23872.2_S_AT)     | 18456_s_at (AC004697.159_S_AT) |
| 18508_s_at (AC006532.89_S_AT)  | 18544_at (AC007060.14_AT)      |
| 18582_s_at (AC003671.36_S_AT)  | 18587_s_at (AC007166.53_S_AT)  |
| 18590_at (AJ222713.4_AT)       | 18591_at (X74756.2_AT)         |
| 18596_at (AC005698.13_AT)      | 18597_at (AL080282.74_AT)      |
| 18604_at (AF069298.31_AT)      | 18622_g_at (AJ005902.2_G_AT)   |
| 18625_at (AC005278.22_AT)      | 18631_at (AC002510.112_AT)     |
| 18636_at (AC006577.22_AT)      | 18668_at (AJ249794_AT)         |
| 18686_s_at (U18126_S_AT)       | 18698_s_at (X17528_S_AT)       |

|                               |                                |
|-------------------------------|--------------------------------|
| 18735_s_at (Z29490_S_AT)      | 18753_s_at (AF118222.28_S_AT)  |
| 18782_at (AC003040.90_AT)     | 18803_at (AC005315.94_AT)      |
| 18885_at (AC006921.147_AT)    | 18888_at (AC007591.68_AT)      |
| 18899_s_at (X13434.1_S_AT)    | 18908_i_at (AF055848.2_I_AT)   |
| 18909_s_at (AF055848.2_S_AT)  | 18928_at (AC002333.181_AT)     |
| 18930_at (AC005990.57_AT)     | 18933_at (AC007020.48_AT)      |
| 18949_at (Z54136.1_AT)        | 18953_at (AF077955.1_AT)       |
| 18963_at (AC004561.99_AT)     | 18966_at (AC004561.106_AT)     |
| 19019_i_at (X82623.2_I_AT)    | 19060_at (AC003671.34_AT)      |
| 19092_at (AL078606.188_AT)    | 19110_s_at (X86947.2_S_AT)     |
| 19137_at (X74755.2_AT)        | 19140_at (AC005170.24_AT)      |
| 19181_s_at (AF053065.2_S_AT)  | 19207_at (AC006069.117_AT)     |
| 19247_at (AF071527.44_AT)     | 19257_s_at (AC000104.57_S_AT)  |
| 19284_at (AC003028.196_AT)    | 19288_at (AC005824.130_AT)     |
| 19325_at (AL022604.42_AT)     | 19364_at (AL022023.142_AT)     |
| 19383_at (AC006200.203_AT)    | 19395_at (AF007270.32_AT)      |
| 19405_at (AJ223803.1_AT)      | 19407_at (AC004697.81_AT)      |
| 19409_at (AC007357.56_AT)     | 19411_at (AC007661.104_AT)     |
| 19421_at (X70990.4_AT)        | 19432_s_at (AL035680.11_S_AT)  |
| 19451_at (AC004392.6_AT)      | 19460_s_at (AC000132.66_S_AT)  |
| 19465_at (AL021768.96_AT)     | 19546_at (AC005398.172_AT)     |
| 19555_at (AF058919.48_AT)     | 19591_at (AJ010735.4_AT)       |
| 19614_at (AC003970.32_AT)     | 19624_at (AL049481.196_AT)     |
| 19625_s_at (AC002311.26_S_AT) | 19635_at (AL049746.38_AT)      |
| 19640_at (AC004561.78_AT)     | 19641_at (AC004561.66_AT)      |
| 19645_at (AC004561.70_AT)     | 19655_at (Y14199.1_AT)         |
| 19667_at (AL021710.5_AT)      | 19672_at (AC005687.19_AT)      |
| 19673_g_at (AC005687.19_G_AT) | 19700_s_at (AL031326.154_S_AT) |
| 19704_i_at (AJ005927.2_I_AT)  | 19707_s_at (Z95768.3_S_AT)     |
| 19741_at (AL049171.72_AT)     | 19755_at (AC006593.64_AT)      |
| 19762_at (AL035527.204_AT)    | 19818_i_at (AL021749.33_I_AT)  |
| 19844_at (AJ007588.2_AT)      | 19848_s_at (AC004261.94_S_AT)  |
| 19851_at (U23794.3_AT)        | 19881_at (AC004077.49_AT)      |
| 19892_at (AC005770.30_AT)     | 19895_s_at (U77347.4_S_AT)     |
| 19944_at (AC002130.4_AT)      | 19956_at (AC006282.11_AT)      |
| 19960_at (AL035527.360_AT)    | 19970_s_at (AC003674.10_S_AT)  |
| 19982_at (AC002986.28_AT)     | 19991_at (AC007017.124_AT)     |
| 20017_at (AC004521.66_AT)     | 20023_at (AC006577.46_AT)      |

|                                |                               |
|--------------------------------|-------------------------------|
| 20030_at (AL078637.51_AT)      | 20051_at (AC000106.38_AT)     |
| 20053_at (AC002292.27_AT)      | 20096_at (AC004238.31_AT)     |
| 20098_at (AC004697.123_AT)     | 20133_i_at (AC007178.71_I_AT) |
| 20134_s_at (AC007178.71_S_AT)  | 20144_at (AL079350.68_AT)     |
| 20179_at (AL035538.229_AT)     | 20189_at (AC005489.2_AT)      |
| 20194_at (AC007584.48_AT)      | 20223_at (AL022347.145_AT)    |
| 20238_at (X74514.2_AT)         | 20239_g_at (X74514.2_G_AT)    |
| 20245_s_at (AC005309.97_S_AT)  | 20247_at (AC004392.4_AT)      |
| 20263_at (AB004798.1_AT)       | 20269_at (AC002387.237_AT)    |
| 20271_at (Z99707.27_AT)        | 20287_at (Y14590.5_AT)        |
| 20288_g_at (Y14590.5_G_AT)     | 20291_s_at (M92353.4_S_AT)    |
| 20297_at (AC007153.27_AT)      | 20323_at (AC004561.62_AT)     |
| 20348_at (AC005967.35_AT)      | 20356_at (AC004561.74_AT)     |
| 20365_s_at (AC005850.19_S_AT)  | 20370_at (AC004561.263_AT)    |
| 20420_at (AL024486.131_AT)     | 20421_at (U81294.2_AT)        |
| 20422_g_at (U81294.2_G_AT)     | 20432_at (U43486.2_AT)        |
| 20433_at (AC006232.147_AT)     | 20450_at (AJ005930.2_AT)      |
| 20461_at (AL049480.157_AT)     | 20462_at (U82399.2_AT)        |
| 20479_i_at (AF069495.2_I_AT)   | 20480_s_at (AF069495.2_S_AT)  |
| 20485_at (AC007660.131_AT)     | 20529_at (Z97341.125_AT)      |
| 20572_s_at (AC005560.229_S_AT) | 20589_at (AF081066.3_AT)      |
| 20646_at (AC002291.20_AT)      | 20656_at (AL035396.46_AT)     |
| 20658_s_at (AL050400.70_S_AT)  | 20669_s_at (AC002388.6_S_AT)  |
| 20685_at (AL049751.46_AT)      | 20686_at (Y14424.2_AT)        |
| 20689_s_at (AC002335.19_S_AT)  |                               |

**Table 5** Probe Sets as referred to in Tables 4a and 4b corresponding to genes encoding regulatory proteins, the expression of which is increased after infection of wild-type *Arabidopsis*, and altered after infection of at least one mutant *Arabidopsis*, with *Pseudomonas syringae*

| ProbeSet                     | Regulatory factor                             |
|------------------------------|---|
| 12497_at (AC006533.51_AT)    | putative receptor-like protein kinase         |
| 16409_at (AC004393.2_AT)     | putative receptor-like kinase                 |
| 12307_at (AC002392.162_AT)   | putative receptor-like protein kinase         |
| 13659_at (AL022347.46_AT)    | serine/threonine kinase-like protein          |
| 17341_at (AL021713.89_AT)    | receptor serine/threonine kinase-like protein |
| 12958_at (AC002332.249_AT)   | putative protein kinase                       |
| 17758_at (AF076243.41_AT)    | putative receptor-like protein kinase         |
| 16570_s_at (ATHCDPKA_S_AT)   | CDPK1 calcium dependent protein kinase 1      |
| 19092_at (AL078606.188_AT)   | protein kinase-like protein                   |
| 17371_at (AF076243.44_AT)    | putative receptor-like protein kinase         |
| 20223_at (AL022347.145_AT)   | putative ser/thr protein kinase               |
| 19655_at (Y14199.1_AT)       | MAP3K delta-1 protein kinase                  |
| 14030_at (AC005970.225_AT)   | putative protein kinase                       |
| 20462_at (U82399.2_AT)       | putative protein kinase PK1                   |
| 16781_at (AC002392.100_AT)   | putative receptor-like protein kinase         |
| 14763_at (X86958.1_AT)       | protein kinase catalytic domain (fragment)    |
| 17752_at (AC003974.37_AT)    | putative protein kinase                       |
| 13755_at (NOVARTIS15_AT)     | putative ser/thr kinase                       |
| 13370_at (AC005322.4_AT)     | similar to Dsor1 protein kinase               |
| 20144_at (AL079350.68_AT)    | receptor kinase-like protein                  |
| 16357_at (AF149413.38_AT)    | putative protein kinase                       |
| 18176_at (AL035540.31_AT)    | putative protein kinase                       |
| 15798_at (AC002521.173_AT)   | putative receptor-like protein kinase         |
| 12965_at (AL021711.118_AT)   | protein kinase-like protein                   |
| 16398_s_at (AL022603.3_S_AT) | putative ser/thr protein kinase               |
| 18122_at (AC002338.110_AT)   | putative protein kinase                       |
| 17323_at (U95973.69_AT)      | putative ser/thr protein kinase               |
| 18782_at (AC003040.90_AT)    | putative protein kinase                       |
| 16405_at (AC005850.9_AT)     | putative serine/threonine protein kinase      |

| ProbeSet                        | Regulatory factor   |
|---------------------------------|---|
| 14110_i_at (AL035528.279_I_AT)  | putative R protein. Like Hcr9-9A, Lycopersicon pimpinellifolium F18A5_290 chr.4 |
| 14214_at (NOVARTIS83_AT)        | putative calmodulin-binding protein (duplicate)                                 |
| 13763_at (NOVARTIS21_AT)        | putative calmodulin-binding protein (duplicate)                                 |
| 12438_at (AL021710.83_AT)       | membrane-bound small GTP-binding - like protein                                 |
| 19848_s_at (AC004261.94_S_AT)   | calmodulin-related protein  |
| 16103_s_at (ATU60445_S_AT)      | GRF7 general regulatory factor encoding 14-3-3 protein                          |
| 14249_i_at (PAD4_I_AT)          | PAD4 phytoalexin deficient 4  |
| 19465_at (AL021768.96_AT)       | RPP5-like NBS-LRR resistance protein  |
| 14640_s_at (putativemloHI_S_AT) | Mlo-like (duplicate)  |
| 18456_s_at (AC004697.159_S_AT)  | Mlo-like (duplicate)  |
| 14320_at (AC005956.54_AT)       | putative RING zinc finger protein   |
| 18054_at (AJ238846_AT)          | SGP1 monomeric G-protein  |
| 16130_s_at (AF078683_S_AT)      | RHA1a RING-H2 finger protein  |
| 20485_at (AC007660.131_AT)      | putative two-component response regulator protein                               |
| 15052_at (AC002332.103_AT)      | putative calcium-binding EF-hand protein  |
| 15632_s_at (AB012570_S_AT)      | ATHP3 two-compoent phosphorelay mediator with a single HPt domain               |
| 16553_f_at (AF078821_F_AT)      | RHA1b RING-H2 finger protein  |
| 13685_s_at (MLOLIKE2_S_AT)      | Mlo-like 2 (duplicate)  |
| 20365_s_at (AC005850.19_S_AT)   | Mlo-like 2 (duplicate)  |
| 13312_at (AC006223.75_AT)       | putative disease resistance protein   |
| 17180_at (AF007270.30_AT)       | similar to GATA-type zinc fingers   |
| 15779_g_at (X98676.2_G_AT)      | zinc finger protein (duplicate)   |
| 15778_at (X98676.2_AT)          | zinc finger protein (duplicate)   |
| 14711_s_at (ZFPL_S_AT)          | hypothetical Cys-3-His zinc finger protein                                      |
| 17379_at (AF085279.9_AT)        | putative CCCH-type zinc finger protein  |
| 12525_at (AC006587.85_AT)       | putative DOF zinc finger protein  |
| 16589_s_at (ATU26937_S_AT)      | AtMYB7 transcription factor   |
| 12712_f_at (Z95774_F_AT)        | AtMYB51 R2R3 myb transcription factor   |
| 13273_s_at (HSF4_S_AT)          | HSF4 heat shock transcription factor 4 (duplicate)                              |
| 16105_s_at (ATU68017_S_AT)      | HSF4 heat shock transcription factor 4 (duplicate)                              |
| 18803_at (AC005315.94_AT)       | putative SCARECROW homeobox gene regulator                                      |

| ProbeSet                       | Regulatory factor  |
|--------------------------------|--|
| 12905_s_at (ATERF2_S_AT)       | EREBP2 ethylene responsive element binding factor 2 (duplicate)        |
| 16609_s_at (AB008104_S_AT)     | EREBP2 ethylene responsive element binding factor 2 (duplicate)        |
| 19755_at (AC006593.64_AT)      | ethylene reponse factor-like AP2 domain transcription factor           |
| 17514_s_at (AF076277_S_AT)     | ERF1 ethylene response factor 1 transcription factor                   |
| 18121_s_at (AC002337.21_S_AT)  | AtMYB2 MYB transcription factor  |
| 13688_s_at (monopteros_S_AT)   | transcription factor   |
| 12904_s_at (ATERF1_S_AT)       | EREBP1 ethylene responsive element binding factor 1 (duplicate)        |
| 16063_s_at (AB008103_S_AT)     | EREBP1 ethylene responsive element binding factor 1 (duplicate)        |
| 16638_s_at (AF139098_S_AT)     | putative zinc finger protein   |
| 16545_s_at (AF037229_S_AT)     | transcription factor   |
| 13435_at (AF003102.3_AT)       | RAP2.9 AP2 domain containing putative transcription factor             |
| 15665_s_at (AF022658_S_AT)     | putative c2h2 zinc finger transcription factor                         |
| 19673_g_at (AC005687.19_G_AT)  | RAP2.6 AP2 domain containing putative transcription factor (duplicate) |
| 19672_at (AC005687.19_AT)      | RAP2.6 AP2 domain containing putative transcription factor (duplicate) |
| 14780_at (AC004683.103_AT)     | DREB-like AP2 domain transcription factor                              |
| 17303_s_at (AC004683.25_S_AT)  | WRKY33 transcription factor  |
| 18949_at (Z54136.1_AT)         | MYB-related protein  |
| 19707_s_at (Z95768.3_S_AT)     | AtMYB44 transcription factor   |
| 16073_f_at (AF062908_F_AT)     | putative transcription factor  |
| 12966_s_at (AL023094.197_S_AT) | ATB2 bZIP transcription factor   |
| 12736_f_at (Z97048_F_AT)       | AtMYB13 transcription factor   |



**Table 6** Probe Sets as referred to in Tables 4a and 4b corresponding to genes, the expression of which is increased after infection of wild-type *Arabidopsis*, and altered after infection of at least one mutant *Arabidopsis* having a mutation that results in enhanced susceptibility to *Pseudomonas*

|                                |                               |
|--------------------------------|-------------------------------|
| 12079_s_at (A71597.1_S_AT)     | 12081_at (AC001645.140_AT)    |
| 12115_at (AL033545.26_AT)      | 12150_at (AC004005.151_AT)    |
| 12233_at (AJ001807.1_AT)       | 12307_at (AC002392.162_AT)    |
| 12314_at (AC001229.28_AT)      | 12332_s_at (AB023448.2_S_AT)  |
| 12341_s_at (AL021637.176_S_AT) | 12400_at (X98453.1_AT)        |
| 12438_at (AL021710.83_AT)      | 12475_at (Y11794.1_AT)        |
| 12487_at (AC004411.126_AT)     | 12497_at (AC006533.51_AT)     |
| 12500_s_at (AF081067.3_S_AT)   | 12530_at (Z99707.184_AT)      |
| 12574_at (X82624.2_AT)         | 12584_at (AC004521.233_AT)    |
| 12712_f_at (Z95774_F_AT)       | 12760_g_at (AC005278.32_G_AT) |
| 12764_f_at (AC004138.69_F_AT)  | 12772_at (AC005278.34_AT)     |
| 12851_s_at (ACCSYN1_S_AT)      | 12879_s_at (AIG1_S_AT)        |
| 12883_s_at (APX_S_AT)          | 12889_s_at (ASA1_S_AT)        |
| 12891_at (ATACS6_AT)           | 12892_g_at (ATACS6_G_AT)      |
| 12905_s_at (ATERF2_S_AT)       | 12911_s_at (ATG6PDHE5_S_AT)   |
| 12921_s_at (ATHHMGCOAR_S_AT)   | 12958_at (AC002332.249_AT)    |
| 12989_s_at (AC004077.149_S_AT) | 13003_s_at (AB021936.1_S_AT)  |
| 13014_at (U93215.87_AT)        | 13040_at (AC002392.134_AT)    |
| 13070_at (AC006919.171_AT)     | 13094_at (AL035523.163_AT)    |
| 13157_at (AC002409.35_AT)      | 13177_at (AL049640.42_AT)     |
| 13190_s_at (ATTHIREDA_S_AT)    | 13211_s_at (BCHI_S_AT)        |
| 13212_s_at (BGL2_S_AT)         | 13219_s_at (CHI4_S_AT)        |
| 13266_s_at (GST4_S_AT)         | 13273_s_at (HSF4_S_AT)        |
| 13275_f_at (HSP174_F_AT)       | 13370_at (AC005322.4_AT)      |
| 13381_at (AC006580.8_AT)       | 13435_at (AF003102.3_AT)      |
| 13467_at (AL096860.198_AT)     | 13538_at (AL080254.75_AT)     |
| 13565_at (AL035601.21_AT)      | 13588_at (AL021961.24_AT)     |
| 13589_at (AC000132.24_AT)      | 13617_at (AC006592.64_AT)     |
| 13627_at (AL035394.196_AT)     | 13656_at (AC007138.31_AT)     |
| 13659_at (AL022347.46_AT)      | 13666_s_at (INDOLE3GPS_S_AT)  |
| 13680_s_at (LOX1_S_AT)         | 13688_s_at (MONOPTEROS_S_AT)  |
| 13697_at (NI16_AT)             | 13716_at (NOVARTIS103_RC_AT)  |
| 13751_at (NOVARTIS127_AT)      | 13755_at (NOVARTIS15_AT)      |
| 13763_at (NOVARTIS21_AT)       | 13764_at (NOVARTIS22_AT)      |

|                                |   |
|--------------------------------|---|
| 13818_s_at (AC006218.175_S_AT) | 13842_at (AC002396.12_AT)               |
| 13880_s_at (AL049480.183_S_AT) | 13920_at (AC005990.53_AT)               |
| 14015_s_at (A71588.1_S_AT)     | 14016_s_at (A71596.1_S_AT)              |
| 14025_s_at (AC007293.3_S_AT)   | 14030_at (AC005970.225_AT)              |
| 14032_at (AL035601.11_AT)      | 14041_at (AC003970.28_AT)               |
| 14052_at (AC004122.24_AT)      | 14068_s_at (AC006922.197_S_AT)          |
| 14110_i_at (AL035528.279_I_AT) | 14139_at (NOVARTIS30_AT)                |
| 14148_at (NOVARTIS38_AT)       | 14170_at (NOVARTIS51_AT)                |
| 14214_at (NOVARTIS83_AT)       | 14242_s_at (NRA_S_AT)                   |
| 14248_at (PAD3_AT)             | 14249_i_at (PAD4_I_AT)                  |
| 14408_at (AC002291.14_AT)      | 14450_at (AC002986.49_AT)               |
| 14461_at (AC006202.73_AT)      | 14487_at (Z97341.343_AT)                |
| 14584_at (AC007658.25_AT)      | 14609_at (AC002340.147_AT)              |
| 14620_s_at (PAT1_S_AT)         | 14635_s_at (PR.1_S_AT)                  |
| 14638_s_at (PRXCB_S_AT)        | 14663_s_at (trehalaseprecursor_RC_S_AT) |
| 14667_s_at (TRPB_S_AT)         | 14672_s_at (TSA1_S_AT)                  |
| 14673_s_at (TSB2_S_AT)         | 14705_i_at (WT77_RC_I_AT)               |
| 14706_r_at (WT77_RC_R_AT)      | 14735_s_at (AF008124_S_AT)              |
| 14763_at (X86958.1_AT)         | 14786_at (AC005397.115_AT)              |
| 14882_at (AL022605.63_AT)      | 14924_at (AC006283.46_AT)               |
| 14959_at (AC007202.26_AT)      | 14978_at (AC002333.49_AT)               |
| 15073_at (AC007069.93_AT)      | 15091_at (AC004683.97_AT)               |
| 15098_s_at (ATU26945_S_AT)     | 15116_f_at (AF121356_F_AT)              |
| 15124_s_at (ATU59508_S_AT)     | 15125_f_at (D85190_F_AT)                |
| 15137_s_at (ATU57320_S_AT)     | 15154_s_at (ATHMTGDAS_S_AT)             |
| 15162_s_at (U01880_S_AT)       | 15196_s_at (ATU43412_S_AT)              |
| 15199_s_at (AB005804_S_AT)     | 15216_s_at (ATU75191_S_AT)              |
| 15406_at (AC006931.179_AT)     | 15431_at (AL030978.64_AT)               |
| 15483_s_at (AC005819.20_S_AT)  | 15496_at (AC006282.167_AT)              |
| 15518_at (AC005322.28_AT)      | 15522_i_at (AL078637.213_I_AT)          |
| 15526_at (AC004122.16_AT)      | 15531_i_at (AL078637.191_I_AT)          |
| 15540_at (AC006585.205_AT)     | 15543_at (AF096371.10_AT)               |
| 15544_at (AL021633.110_AT)     | 15551_at (AL035440.289_AT)              |
| 15594_s_at (ATU56635_S_AT)     | 15622_s_at (ATU43945_S_AT)              |
| 15632_s_at (AB012570_S_AT)     | 15665_s_at (AF022658_S_AT)              |
| 15672_s_at (AF082299_S_AT)     | 15778_at (X98676.2_AT)                  |
| 15779_g_at (X98676.2_G_AT)     | 15815_s_at (Z97342.366_S_AT)            |
| 15839_at (AC005662.203_AT)     | 15859_at (AC006587.164_AT)              |

|                                |                                |
|--------------------------------|--------------------------------|
| 15866_s_at (AC007133.59_S_AT)  | 15874_at (AL022223.106_AT)     |
| 15886_at (AL078637.204_AT)     | 15924_at (AC007138.61_AT)      |
| 15970_s_at (X71794.2_S_AT)     | 15978_at (X68592.6_AT)         |
| 15982_s_at (AC006260.78_S_AT)  | 16001_at (AF035385.2_AT)       |
| 16003_s_at (AL021749.64_S_AT)  | 16021_s_at (AL022224.182_S_AT) |
| 16031_at (X94248.1_AT)         | 16043_at (AC005489.17_AT)      |
| 16053_i_at (Y14251.4_I_AT)     | 16083_s_at (AF153283_S_AT)     |
| 16091_s_at (ATHHSP83_S_AT)     | 16103_s_at (ATU60445_S_AT)     |
| 16105_s_at (ATU68017_S_AT)     | 16108_s_at (D78604_S_AT)       |
| 16130_s_at (AF078683_S_AT)     | 16233_at (AL080254.83_AT)      |
| 16272_at (AC006304.136_AT)     | 16298_at (AL021890.71_AT)      |
| 16299_at (AL024486.185_AT)     | 16306_at (AL049751.112_AT)     |
| 16340_at (AC004255.15_AT)      | 16357_at (AF149413.38_AT)      |
| 16363_at (AC004255.14_AT)      | 16383_at (AC006300.64_AT)      |
| 16409_at (AC004393.2_AT)       | 16461_i_at (AC004683.79_I_AT)  |
| 16462_s_at (AC004683.79_S_AT)  | 16470_s_at (AF068299.4_S_AT)   |
| 16545_s_at (AF037229_S_AT)     | 16570_s_at (ATHCDPKA_S_AT)     |
| 16578_s_at (ATHRPRP1B_S_AT)    | 16589_s_at (ATU26937_S_AT)     |
| 16594_s_at (ATU39783_S_AT)     | 16603_s_at (ATU81293_S_AT)     |
| 16609_s_at (AB008104_S_AT)     | 16701_at (AC005312.61_AT)      |
| 16721_at (AC006533.58_AT)      | 16781_at (AC002392.100_AT)     |
| 16810_at (AC002339.46_AT)      | 16859_at (AL035523.135_AT)     |
| 16864_i_at (AF037367.4_I_AT)   | 16865_s_at (AF037367.4_S_AT)   |
| 16914_s_at (AL049500.57_S_AT)  | 16968_at (AL021961.93_AT)      |
| 16989_at (AL030978.46_AT)      | 16995_at (AC002391.188_AT)     |
| 17007_at (AC005896.26_AT)      | 17051_s_at (AF098947_S_AT)     |
| 17066_s_at (ATHLIPOXY_S_AT)    | 17075_s_at (ATU09961_S_AT)     |
| 17128_s_at (ATHRPRP1A_S_AT)    | 17180_at (AF007270.30_AT)      |
| 17338_at (AC002535.97_AT)      | 17341_at (AL021713.89_AT)      |
| 17356_s_at (Z97338.190_S_AT)   | 17371_at (AF076243.44_AT)      |
| 17413_s_at (AJ006961.4_S_AT)   | 17451_at (AC002343.47_AT)      |
| 17452_g_at (AC002343.47_G_AT)  | 17484_at (X79052.2_AT)         |
| 17485_s_at (Z97340.345_S_AT)   | 17487_s_at (U18993.2_S_AT)     |
| 17533_s_at (ATU43488_S_AT)     | 17544_s_at (ATU40856_S_AT)     |
| 17585_s_at (AF134487_S_AT)     | 17653_at (AL035679.144_AT)     |
| 17752_at (AC003974.37_AT)      | 17758_at (AF076243.41_AT)      |
| 17775_at (AC004392.2_AT)       | 17781_at (AL049746.177_AT)     |
| 17840_s_at (AC002333.223_S_AT) | 17877_g_at (AJ007587.2_G_AT)   |

|                                |                                |
|--------------------------------|--------------------------------|
| 17893_at (AC004401.135_AT)     | 17899_at (Z97339.197_AT)       |
| 17930_s_at (AJ006960.4_S_AT)   | 17945_at (Z97341.411_AT)       |
| 17963_at (AL049730.88_AT)      | 18054_at (AJ238846_AT)         |
| 18109_s_at (AC002391.206_S_AT) | 18148_at (AC004669.25_AT)      |
| 18176_at (AL035540.31_AT)      | 18194_i_at (AL096859.227_I_AT) |
| 18213_at (AL022140.126_AT)     | 18228_at (X91259.1_AT)         |
| 18242_g_at (AC006580.71_G_AT)  | 18255_at (AC005770.25_AT)      |
| 18258_s_at (AC006439.222_S_AT) | 18267_at (AC006223.23_AT)      |
| 18508_s_at (AC006532.89_S_AT)  | 18544_at (AC007060.14_AT)      |
| 18590_at (AJ222713.4_AT)       | 18591_at (X74756.2_AT)         |
| 18596_at (AC005698.13_AT)      | 18604_at (AF069298.31_AT)      |
| 18622_g_at (AJ005902.2_G_AT)   | 18625_at (AC005278.22_AT)      |
| 18686_s_at (U18126_S_AT)       | 18698_s_at (X17528_S_AT)       |
| 18735_s_at (Z29490_S_AT)       | 18753_s_at (AF118222.28_S_AT)  |
| 18803_at (AC005315.94_AT)      | 18885_at (AC006921.147_AT)     |
| 18888_at (AC007591.68_AT)      | 18899_s_at (X13434.1_S_AT)     |
| 18909_s_at (AF055848.2_S_AT)   | 18928_at (AC002333.181_AT)     |
| 18930_at (AC005990.57_AT)      | 18933_at (AC007020.48_AT)      |
| 18953_at (AF077955.1_AT)       | 18966_at (AC004561.106_AT)     |
| 19019_i_at (X82623.2_I_AT)     | 19137_at (X74755.2_AT)         |
| 19140_at (AC005170.24_AT)      | 19181_s_at (AF053065.2_S_AT)   |
| 19207_at (AC006069.117_AT)     | 19247_at (AF071527.44_AT)      |
| 19257_s_at (AC000104.57_S_AT)  | 19288_at (AC005824.130_AT)     |
| 19325_at (AL022604.42_AT)      | 19364_at (AL022023.142_AT)     |
| 19383_at (AC006200.203_AT)     | 19405_at (AJ223803.1_AT)       |
| 19407_at (AC004697.81_AT)      | 19409_at (AC007357.56_AT)      |
| 19411_at (AC007661.104_AT)     | 19421_at (X70990.4_AT)         |
| 19432_s_at (AL035680.11_S_AT)  | 19451_at (AC004392.6_AT)       |
| 19460_s_at (AC000132.66_S_AT)  | 19465_at (AL021768.96_AT)      |
| 19546_at (AC005398.172_AT)     | 19555_at (AF058919.48_AT)      |
| 19624_at (AL049481.196_AT)     | 19641_at (AC004561.66_AT)      |
| 19645_at (AC004561.70_AT)      | 19667_at (AL021710.5_AT)       |
| 19741_at (AL049171.72_AT)      | 19755_at (AC006593.64_AT)      |
| 19818_i_at (AL021749.33_I_AT)  | 19848_s_at (AC004261.94_S_AT)  |
| 19881_at (AC004077.49_AT)      | 19892_at (AC005770.30_AT)      |
| 19944_at (AC002130.4_AT)       | 19956_at (AC006282.11_AT)      |
| 19970_s_at (AC003674.10_S_AT)  | 19982_at (AC002986.28_AT)      |
| 19991_at (AC007017.124_AT)     | 20017_at (AC004521.66_AT)      |

|                                |                               |
|--------------------------------|-------------------------------|
| 20030_at (AL078637.51_AT)      | 20051_at (AC000106.38_AT)     |
| 20098_at (AC004697.123_AT)     | 20133_i_at (AC007178.71_I_AT) |
| 20134_s_at (AC007178.71_S_AT)  | 20144_at (AL079350.68_AT)     |
| 20238_at (X74514.2_AT)         | 20239_g_at (X74514.2_G_AT)    |
| 20245_s_at (AC005309.97_S_AT)  | 20247_at (AC004392.4_AT)      |
| 20263_at (AB004798.1_AT)       | 20269_at (AC002387.237_AT)    |
| 20271_at (Z99707.27_AT)        | 20287_at (Y14590.5_AT)        |
| 20288_g_at (Y14590.5_G_AT)     | 20291_s_at (M92353.4_S_AT)    |
| 20297_at (AC007153.27_AT)      | 20323_at (AC004561.62_AT)     |
| 20348_at (AC005967.35_AT)      | 20356_at (AC004561.74_AT)     |
| 20370_at (AC004561.263_AT)     | 20420_at (AL024486.131_AT)    |
| 20421_at (U81294.2_AT)         | 20422_g_at (U81294.2_G_AT)    |
| 20432_at (U43486.2_AT)         | 20450_at (AJ005930.2_AT)      |
| 20461_at (AL049480.157_AT)     | 20462_at (U82399.2_AT)        |
| 20479_i_at (AF069495.2_I_AT)   | 20480_s_at (AF069495.2_S_AT)  |
| 20485_at (AC007660.131_AT)     | 20529_at (Z97341.125_AT)      |
| 20572_s_at (AC005560.229_S_AT) | 20589_at (AF081066.3_AT)      |
| 20669_s_at (AC002388.6_S_AT)   | 20685_at (AL049751.46_AT)     |
| 20689_s_at (AC002335.19_S_AT)  |                               |

**Table 7** Probe Sets as referred to in Tables 4a and 4b corresponding to genes, the expression of which is increased after infection of wild-type *Arabidopsis*, and altered after infection of at least one mutant *Arabidopsis* having a mutation in a gene whose expression is important for resistance to necrotrophic fungi

|   |                                |
|---|--------------------------------|
| 11997_at (AC005967.4_AT)                  | 12004_at (AL022023.132_AT)     |
| 12037_at (AC004005.174_AT)                | 12072_at (AL035396.4_AT)       |
| 12079_s_at (A71597.1_S_AT)                | 12081_at (AC001645.140_AT)     |
| 12115_at (AL033545.26_AT)                 | 12136_at (AC007591.60_AT)      |
| 12150_at (AC004005.151_AT)                | 12216_at (AC007119.56_AT)      |
| 12317_at (AC004138.27_AT)                 | 12323_at (AC002333.18_AT)      |
| 12332_s_at (AB023448.2_S_AT)              | 12369_at (AC002535.59_AT)      |
| 12400_at (X98453.1_AT)                    | 12449_s_at (AC002343.179_S_AT) |
| 12475_at (Y11794.1_AT)                    | 12487_at (AC004411.126_AT)     |
| 12500_s_at (AF081067.3_S_AT)              | 12530_at (Z99707.184_AT)       |
| 12535_at (AL035538.156_AT)                | 12538_at (AF033205.2_AT)       |
| 12574_at (X82624.2_AT)                    | 12584_at (AC004521.233_AT)     |
| 12626_at (AC006234.95_AT)                 | 12645_at (AL021712.56_AT)      |
| 12744_at (AC001645.15_AT)                 | 12760_g_at (AC005278.32_G_AT)  |
| 12772_at (AC005278.34_AT)                 | 12797_s_at (AC007138.25_S_AT)  |
| 12851_s_at (ACCSYN1_S_AT)                 | 12880_s_at (AIG2_S_AT)         |
| 12889_s_at (ASA1_S_AT)                    | 12904_s_at (ATERF1_S_AT)       |
| 12905_s_at (ATERF2_S_AT)                  | 12921_s_at (ATHHMGCOAR_S_AT)   |
| 12951_at (AC005489.5_AT)                  | 12965_at (AL021711.118_AT)     |
| 12966_s_at (AL023094.197_S_AT)            | 12989_s_at (AC004077.149_S_AT) |
| 13040_at (AC002392.134_AT)                | 13094_at (AL035523.163_AT)     |
| 13134_s_at (AC002337.9_S_AT)              | 13152_s_at (AC005322.24_S_AT)  |
| 13176_at (AL031394.56_AT)                 | 13211_s_at (BCHI_S_AT)         |
| 13215_s_at (cafferoylcoAmethyltrans_S_AT) | 13243_r_at (ELI32_R_AT)        |
| 13244_s_at (ELI32_S_AT)                   | 13266_s_at (GST4_S_AT)         |
| 13275_f_at (HSP174_F_AT)                  | 13277_i_at (HSP176A_I_AT)      |
| 13312_at (AC006223.75_AT)                 | 13370_at (AC005322.4_AT)       |
| 13381_at (AC006580.8_AT)                  | 13395_at (AL035528.202_AT)     |
| 13435_at (AF003102.3_AT)                  | 13437_at (AF096371.8_AT)       |
| 13536_at (AL021636.47_AT)                 | 13538_at (AL080254.75_AT)      |
| 13565_at (AL035601.21_AT)                 | 13588_at (AL021961.24_AT)      |
| 13680_s_at (LOX1_S_AT)                    | 13688_s_at (monopteros_S_AT)   |
| 13751_at (NOVARTIS127_AT)                 | 13755_at (NOVARTIS15_AT)       |
| 13764_at (NOVARTIS22_AT)                  | 13803_at (Z97341.376_AT)       |

|                                |   |
|--------------------------------|---|
| 13848_at (AC003981.31_AT)      | 13918_at (AC005388.29_AT)               |
| 13949_s_at (Z97343.352_S_AT)   | 13999_at (AF071527.56_AT)               |
| 14015_s_at (A71588.1_S_AT)     | 14016_s_at (A71596.1_S_AT)              |
| 14041_at (AC003970.28_AT)      | 14052_at (AC004122.24_AT)               |
| 14068_s_at (AC006922.197_S_AT) | 14070_at (AL049658.217_AT)              |
| 14089_at (AC006223.65_AT)      | 14100_at (AF002109.108_AT)              |
| 14197_at (NOVARTIS71_AT)       | 14240_s_at (NR1_S_AT)                   |
| 14450_at (AC002986.49_AT)      | 14461_at (AC006202.73_AT)               |
| 14475_at (AL021811.121_AT)     | 14584_at (AC007658.25_AT)               |
| 14609_at (AC002340.147_AT)     | 14663_s_at (trehalaseprecursor_RC_S_AT) |
| 14667_s_at (TRPB_S_AT)         | 14673_s_at (TSB2_S_AT)                  |
| 14675_s_at (VSP_S_AT)          | 14705_i_at (WT77_RC_I_AT)               |
| 14706_r_at (WT77_RC_R_AT)      | 14735_s_at (AF008124_S_AT)              |
| 14750_s_at (AF096370.12_S_AT)  | 14779_at (AC004680.71_AT)               |
| 14786_at (AC005397.115_AT)     | 14978_at (AC002333.49_AT)               |
| 15052_at (AC002332.103_AT)     | 15073_at (AC007069.93_AT)               |
| 15088_s_at (AC002311.37_S_AT)  | 15091_at (AC004683.97_AT)               |
| 15098_s_at (ATU26945_S_AT)     | 15124_s_at (ATU59508_S_AT)              |
| 15125_f_at (D85190_F_AT)       | 15137_s_at (ATU57320_S_AT)              |
| 15141_s_at (D85191_S_AT)       | 15154_s_at (ATHMTGDAS_S_AT)             |
| 15161_s_at (ATU90522_S_AT)     | 15162_s_at (U01880_S_AT)                |
| 15188_s_at (AF081202_S_AT)     | 15196_s_at (ATU43412_S_AT)              |
| 15211_s_at (ATH243813_S_AT)    | 15342_at (AC006593.101_AT)              |
| 15406_at (AC006931.179_AT)     | 15479_at (AL049483.205_AT)              |
| 15483_s_at (AC005819.20_S_AT)  | 15485_at (AC006233.109_AT)              |
| 15496_at (AC006282.167_AT)     | 15522_i_at (AL078637.213_I_AT)          |
| 15523_s_at (AL078637.213_S_AT) | 15524_at (AC005508.25_AT)               |
| 15526_at (AC004122.16_AT)      | 15531_i_at (AL078637.191_I_AT)          |
| 15532_r_at (AL078637.191_R_AT) | 15547_at (AC005970.122_AT)              |
| 15594_s_at (ATU56635_S_AT)     | 15622_s_at (ATU43945_S_AT)              |
| 15625_s_at (ATU74610_S_AT)     | 15632_s_at (AB012570_S_AT)              |
| 15665_s_at (AF022658_S_AT)     | 15670_s_at (AF061638_S_AT)              |
| 15792_at (AC002341.106_AT)     | 15798_at (AC002521.173_AT)              |
| 15839_at (AC005662.203_AT)     | 15859_at (AC006587.164_AT)              |
| 15874_at (AL022223.106_AT)     | 15886_at (AL078637.204_AT)              |
| 15982_s_at (AC006260.78_S_AT)  | 16063_s_at (AB008103_S_AT)              |
| 16073_f_at (AF062908_F_AT)     | 16108_s_at (D78604_S_AT)                |
| 16130_s_at (AF078683_S_AT)     | 16173_s_at (D78607_S_AT)                |

|                                |                                |
|--------------------------------|--------------------------------|
| 16203_at (AC007519.53_AT)      | 16272_at (AC006304.136_AT)     |
| 16298_at (AL021890.71_AT)      | 16301_s_at (AL031018.105_S_AT) |
| 16306_at (AL049751.112_AT)     | 16357_at (AF149413.38_AT)      |
| 16363_at (AC004255.14_AT)      | 16398_s_at (AL022603.3_S_AT)   |
| 16405_at (AC005850.9_AT)       | 16461_i_at (AC004683.79_I_AT)  |
| 16462_s_at (AC004683.79_S_AT)  | 16470_s_at (AF068299.4_S_AT)   |
| 16510_at (AL034567.198_AT)     | 16522_at (X77500.2_AT)         |
| 16526_at (Z49227.1_AT)         | 16541_s_at (AB023423_S_AT)     |
| 16545_s_at (AF037229_S_AT)     | 16553_f_at (AF078821_F_AT)     |
| 16589_s_at (ATU26937_S_AT)     | 16603_s_at (ATU81293_S_AT)     |
| 16609_s_at (AB008104_S_AT)     | 16611_s_at (AB008782_S_AT)     |
| 16638_s_at (AF139098_S_AT)     | 16646_s_at (ATHDHS1_S_AT)      |
| 16649_s_at (ATHORF_S_AT)       | 16701_at (AC005312.61_AT)      |
| 16747_at (AL021713.3_AT)       | 16781_at (AC002392.100_AT)     |
| 16810_at (AC002339.46_AT)      | 16864_i_at (AF037367.4_I_AT)   |
| 16865_s_at (AF037367.4_S_AT)   | 16914_s_at (AL049500.57_S_AT)  |
| 16968_at (AL021961.93_AT)      | 16989_at (AL030978.46_AT)      |
| 16995_at (AC002391.188_AT)     | 17008_at (AC006585.212_AT)     |
| 17041_s_at (D89631_S_AT)       | 17066_s_at (ATHLIPOXY_S_AT)    |
| 17075_s_at (ATU09961_S_AT)     | 17104_s_at (D88541_S_AT)       |
| 17111_s_at (ATHACSC_S_AT)      | 17119_s_at (AF132212_S_AT)     |
| 17187_at (AF128396.2_AT)       | 17300_at (X66017.2_AT)         |
| 17323_at (U95973.69_AT)        | 17338_at (AC002535.97_AT)      |
| 17356_s_at (Z97338.190_S_AT)   | 17485_s_at (Z97340.345_S_AT)   |
| 17511_s_at (AF067605_S_AT)     | 17514_s_at (AF076277_S_AT)     |
| 17522_s_at (D78606_S_AT)       | 17533_s_at (ATU43488_S_AT)     |
| 17548_s_at (AF118823_S_AT)     | 17595_s_at (AF166352_S_AT)     |
| 17648_at (AL021684.43_AT)      | 17702_at (AC005700.212_AT)     |
| 17719_at (AC006592.17_AT)      | 17775_at (AC004392.2_AT)       |
| 17860_at (AL078467.4_AT)       | 17877_g_at (AJ007587.2_G_AT)   |
| 17893_at (AC004401.135_AT)     | 17945_at (Z97341.411_AT)       |
| 18109_s_at (AC002391.206_S_AT) | 18121_s_at (AC002337.21_S_AT)  |
| 18122_at (AC002338.110_AT)     | 18148_at (AC004669.25_AT)      |
| 18176_at (AL035540.31_AT)      | 18213_at (AL022140.126_AT)     |
| 18224_s_at (AL021890.57_S_AT)  | 18228_at (X91259.1_AT)         |
| 18236_s_at (AC004683.69_S_AT)  | 18258_s_at (AC006439.222_S_AT) |
| 18266_at (AC004684.33_AT)      | 18299_s_at (M23872.2_S_AT)     |
| 18587_s_at (AC007166.53_S_AT)  | 18590_at (AJ222713.4_AT)       |



|                                |                               |
|--------------------------------|-------------------------------|
| 18596_at (AC005698.13_AT)      | 18636_at (AC006577.22_AT)     |
| 18668_at (AJ249794_AT)         | 18735_s_at (Z29490_S_AT)      |
| 18782_at (AC003040.90_AT)      | 18803_at (AC005315.94_AT)     |
| 18888_at (AC007591.68_AT)      | 18899_s_at (X13434.1_S_AT)    |
| 18908_i_at (AF055848.2_I_AT)   | 18933_at (AC007020.48_AT)     |
| 18949_at (Z54136.1_AT)         | 18953_at (AF077955.1_AT)      |
| 18963_at (AC004561.99_AT)      | 19019_i_at (X82623.2_I_AT)    |
| 19060_at (AC003671.34_AT)      | 19110_s_at (X86947.2_S_AT)    |
| 19137_at (X74755.2_AT)         | 19257_s_at (AC000104.57_S_AT) |
| 19284_at (AC003028.196_AT)     | 19288_at (AC005824.130_AT)    |
| 19325_at (AL022604.42_AT)      | 19364_at (AL022023.142_AT)    |
| 19383_at (AC006200.203_AT)     | 19395_at (AF007270.32_AT)     |
| 19409_at (AC007357.56_AT)      | 19432_s_at (AL035680.11_S_AT) |
| 19451_at (AC004392.6_AT)       | 19460_s_at (AC000132.66_S_AT) |
| 19546_at (AC005398.172_AT)     | 19591_at (AJ010735.4_AT)      |
| 19614_at (AC003970.32_AT)      | 19635_at (AL049746.38_AT)     |
| 19640_at (AC004561.78_AT)      | 19641_at (AC004561.66_AT)     |
| 19645_at (AC004561.70_AT)      | 19667_at (AL021710.5_AT)      |
| 19672_at (AC005687.19_AT)      | 19673_g_at (AC005687.19_G_AT) |
| 19700_s_at (AL031326.154_S_AT) | 19704_i_at (AJ005927.2_I_AT)  |
| 19707_s_at (Z95768.3_S_AT)     | 19755_at (AC006593.64_AT)     |
| 19762_at (AL035527.204_AT)     | 19818_i_at (AL021749.33_I_AT) |
| 19851_at (U23794.3_AT)         | 19895_s_at (U77347.4_S_AT)    |
| 19956_at (AC006282.11_AT)      | 19960_at (AL035527.360_AT)    |
| 20023_at (AC006577.46_AT)      | 20051_at (AC000106.38_AT)     |
| 20053_at (AC002292.27_AT)      | 20096_at (AC004238.31_AT)     |
| 20133_i_at (AC007178.71_I_AT)  | 20134_s_at (AC007178.71_S_AT) |
| 20144_at (AL079350.68_AT)      | 20179_at (AL035538.229_AT)    |
| 20189_at (AC005489.2_AT)       | 20194_at (AC007584.48_AT)     |
| 20247_at (AC004392.4_AT)       | 20263_at (AB004798.1_AT)      |
| 20291_s_at (M92353.4_S_AT)     | 20297_at (AC007153.27_AT)     |
| 20323_at (AC004561.62_AT)      | 20348_at (AC005967.35_AT)     |
| 20356_at (AC004561.74_AT)      | 20370_at (AC004561.263_AT)    |
| 20433_at (AC006232.147_AT)     | 20450_at (AJ005930.2_AT)      |
| 20479_i_at (AF069495.2_I_AT)   | 20485_at (AC007660.131_AT)    |
| 20572_s_at (AC005560.229_S_AT) | 20646_at (AC002291.20_AT)     |
| 20656_at (AL035396.46_AT)      | 20658_s_at (AL050400.70_S_AT) |
| 20669_s_at (AC002388.6_S_AT)   | 20686_at (Y14424.2_AT)        |



**Table 8** Probe Sets as referred to in Table 6 corresponding to genes, the expression of which is increased after infection of wild-type *Arabidopsis* and altered after infection of at least one mutant *Arabidopsis* having a mutation in a gene that interferes with salicylic acid dependent signaling

|                                |                                |
|--------------------------------|--------------------------------|
| 12079_s_at (A71597.1_S_AT)     | 12081_at (AC001645.140_AT)     |
| 12115_at (AL033545.26_AT)      | 12150_at (AC004005.151_AT)     |
| 12233_at (AJ001807.1_AT)       | 12307_at (AC002392.162_AT)     |
| 12314_at (AC001229.28_AT)      | 12332_s_at (AB023448.2_S_AT)   |
| 12341_s_at (AL021637.176_S_AT) | 12400_at (X98453.1_AT)         |
| 12438_at (AL021710.83_AT)      | 12475_at (Y11794.1_AT)         |
| 12487_at (AC004411.126_AT)     | 12497_at (AC006533.51_AT)      |
| 12500_s_at (AF081067.3_S_AT)   | 12530_at (Z99707.184_AT)       |
| 12574_at (X82624.2_AT)         | 12584_at (AC004521.233_AT)     |
| 12712_f_at (Z95774_F_AT)       | 12760_g_at (AC005278.32_G_AT)  |
| 12764_f_at (AC004138.69_F_AT)  | 12879_s_at (AIG1_S_AT)         |
| 12883_s_at (APX_S_AT)          | 12889_s_at (ASA1_S_AT)         |
| 12891_at (ATACS6_AT)           | 12892_g_at (ATACS6_G_AT)       |
| 12905_s_at (ATERF2_S_AT)       | 12911_s_at (ATG6PDHE5_S_AT)    |
| 12921_s_at (ATHHMGCOAR_S_AT)   | 12958_at (AC002332.249_AT)     |
| 12989_s_at (AC004077.149_S_AT) | 13003_s_at (AB021936.1_S_AT)   |
| 13014_at (U93215.87_AT)        | 13040_at (AC002392.134_AT)     |
| 13070_at (AC006919.171_AT)     | 13094_at (AL035523.163_AT)     |
| 13157_at (AC002409.35_AT)      | 13190_s_at (ATTHIREDA_S_AT)    |
| 13211_s_at (BCHI_S_AT)         | 13212_s_at (BGL2_S_AT)         |
| 13219_s_at (CHI4_S_AT)         | 13273_s_at (HSF4_S_AT)         |
| 13275_f_at (HSP174_F_AT)       | 13370_at (AC005322.4_AT)       |
| 13381_at (AC006580.8_AT)       | 13538_at (AL080254.75_AT)      |
| 13565_at (AL035601.21_AT)      | 13588_at (AL021961.24_AT)      |
| 13589_at (AC000132.24_AT)      | 13617_at (AC006592.64_AT)      |
| 13627_at (AL035394.196_AT)     | 13656_at (AC007138.31_AT)      |
| 13659_at (AL022347.46_AT)      | 13666_s_at (INDOLE3GPS_S_AT)   |
| 13680_s_at (LOX1_S_AT)         | 13697_at (NI16_AT)             |
| 13751_at (NOVARTIS127_AT)      | 13755_at (NOVARTIS15_AT)       |
| 13763_at (NOVARTIS21_AT)       | 13764_at (NOVARTIS22_AT)       |
| 13818_s_at (AC006218.175_S_AT) | 13880_s_at (AL049480.183_S_AT) |
| 13920_at (AC005990.53_AT)      | 14015_s_at (A71588.1_S_AT)     |
| 14016_s_at (A71596.1_S_AT)     | 14025_s_at (AC007293.3_S_AT)   |
| 14030_at (AC005970.225_AT)     | 14032_at (AL035601.11_AT)      |

|   |                                |
|---|--------------------------------|
| 14041_at (AC003970.28_AT)               | 14052_at (AC004122.24_AT)      |
| 14068_s_at (AC006922.197_S_AT)          | 14110_i_at (AL035528.279_I_AT) |
| 14139_at (NOVARTIS30_AT)                | 14148_at (NOVARTIS38_AT)       |
| 14170_at (NOVARTIS51_AT)                | 14214_at (NOVARTIS83_AT)       |
| 14242_s_at (NRA_S_AT)                   | 14248_at (PAD3_AT)             |
| 14249_i_at (PAD4_I_AT)                  | 14408_at (AC002291.14_AT)      |
| 14450_at (AC002986.49_AT)               | 14461_at (AC006202.73_AT)      |
| 14487_at (Z97341.343_AT)                | 14584_at (AC007658.25_AT)      |
| 14609_at (AC002340.147_AT)              | 14620_s_at (PAT1_S_AT)         |
| 14635_s_at (PR.1_S_AT)                  | 14638_s_at (PRXCB_S_AT)        |
| 14663_s_at (trehalaseprecursor_RC_S_AT) | 14667_s_at (TRPB_S_AT)         |
| 14672_s_at (TSA1_S_AT)                  | 14673_s_at (TSB2_S_AT)         |
| 14735_s_at (AF008124_S_AT)              | 14763_at (X86958.1_AT)         |
| 14786_at (AC005397.115_AT)              | 14882_at (AL022605.63_AT)      |
| 14959_at (AC007202.26_AT)               | 14978_at (AC002333.49_AT)      |
| 15073_at (AC007069.93_AT)               | 15091_at (AC004683.97_AT)      |
| 15116_f_at (AF121356_F_AT)              | 15137_s_at (ATU57320_S_AT)     |
| 15162_s_at (U01880_S_AT)                | 15196_s_at (ATU43412_S_AT)     |
| 15199_s_at (AB005804_S_AT)              | 15216_s_at (ATU75191_S_AT)     |
| 15431_at (AL030978.64_AT)               | 15483_s_at (AC005819.20_S_AT)  |
| 15496_at (AC006282.167_AT)              | 15518_at (AC005322.28_AT)      |
| 15526_at (AC004122.16_AT)               | 15540_at (AC006585.205_AT)     |
| 15543_at (AF096371.10_AT)               | 15544_at (AL021633.110_AT)     |
| 15551_at (AL035440.289_AT)              | 15594_s_at (ATU56635_S_AT)     |
| 15622_s_at (ATU43945_S_AT)              | 15632_s_at (AB012570_S_AT)     |
| 15672_s_at (AF082299_S_AT)              | 15778_at (X98676.2_AT)         |
| 15779_g_at (X98676.2_G_AT)              | 15815_s_at (Z97342.366_S_AT)   |
| 15839_at (AC005662.203_AT)              | 15859_at (AC006587.164_AT)     |
| 15866_s_at (AC007133.59_S_AT)           | 15874_at (AL022223.106_AT)     |
| 15924_at (AC007138.61_AT)               | 15970_s_at (X71794.2_S_AT)     |
| 15978_at (X68592.6_AT)                  | 15982_s_at (AC006260.78_S_AT)  |
| 16003_s_at (AL021749.64_S_AT)           | 16021_s_at (AL022224.182_S_AT) |
| 16031_at (X94248.1_AT)                  | 16043_at (AC005489.17_AT)      |
| 16053_i_at (Y14251.4_I_AT)              | 16083_s_at (AF153283_S_AT)     |
| 16091_s_at (ATHHSP83_S_AT)              | 16103_s_at (ATU60445_S_AT)     |
| 16105_s_at (ATU68017_S_AT)              | 16108_s_at (D78604_S_AT)       |
| 16130_s_at (AF078683_S_AT)              | 16233_at (AL080254.83_AT)      |
| 16272_at (AC006304.136_AT)              | 16299_at (AL024486.185_AT)     |

|                                |                                |
|--------------------------------|--------------------------------|
| 16306_at (AL049751.112_AT)     | 16340_at (AC004255.15_AT)      |
| 16357_at (AF149413.38_AT)      | 16363_at (AC004255.14_AT)      |
| 16383_at (AC006300.64_AT)      | 16409_at (AC004393.2_AT)       |
| 16461_i_at (AC004683.79_I_AT)  | 16462_s_at (AC004683.79_S_AT)  |
| 16470_s_at (AF068299.4_S_AT)   | 16570_s_at (ATHCDPKA_S_AT)     |
| 16578_s_at (ATHRPRP1B_S_AT)    | 16589_s_at (ATU26937_S_AT)     |
| 16594_s_at (ATU39783_S_AT)     | 16603_s_at (ATU81293_S_AT)     |
| 16609_s_at (AB008104_S_AT)     | 16701_at (AC005312.61_AT)      |
| 16721_at (AC006533.58_AT)      | 16781_at (AC002392.100_AT)     |
| 16810_at (AC002339.46_AT)      | 16864_i_at (AF037367.4_I_AT)   |
| 16865_s_at (AF037367.4_S_AT)   | 16914_s_at (AL049500.57_S_AT)  |
| 16968_at (AL021961.93_AT)      | 16995_at (AC002391.188_AT)     |
| 17007_at (AC005896.26_AT)      | 17051_s_at (AF098947_S_AT)     |
| 17066_s_at (ATHLIPOXY_S_AT)    | 17075_s_at (ATU09961_S_AT)     |
| 17128_s_at (ATHRPRP1A_S_AT)    | 17180_at (AF007270.30_AT)      |
| 17338_at (AC002535.97_AT)      | 17341_at (AL021713.89_AT)      |
| 17356_s_at (Z97338.190_S_AT)   | 17413_s_at (AJ006961.4_S_AT)   |
| 17451_at (AC002343.47_AT)      | 17452_g_at (AC002343.47_G_AT)  |
| 17484_at (X79052.2_AT)         | 17485_s_at (Z97340.345_S_AT)   |
| 17487_s_at (U18993.2_S_AT)     | 17533_s_at (ATU43488_S_AT)     |
| 17544_s_at (ATU40856_S_AT)     | 17585_s_at (AF134487_S_AT)     |
| 17653_at (AL035679.144_AT)     | 17752_at (AC003974.37_AT)      |
| 17758_at (AF076243.41_AT)      | 17775_at (AC004392.2_AT)       |
| 17781_at (AL049746.177_AT)     | 17840_s_at (AC002333.223_S_AT) |
| 17893_at (AC004401.135_AT)     | 17899_at (Z97339.197_AT)       |
| 17930_s_at (AJ006960.4_S_AT)   | 17963_at (AL049730.88_AT)      |
| 18054_at (AJ238846_AT)         | 18148_at (AC004669.25_AT)      |
| 18194_i_at (AL096859.227_I_AT) | 18213_at (AL022140.126_AT)     |
| 18242_g_at (AC006580.71_G_AT)  | 18255_at (AC005770.25_AT)      |
| 18258_s_at (AC006439.222_S_AT) | 18508_s_at (AC006532.89_S_AT)  |
| 18544_at (AC007060.14_AT)      | 18590_at (AJ222713.4_AT)       |
| 18591_at (X74756.2_AT)         | 18596_at (AC005698.13_AT)      |
| 18604_at (AF069298.31_AT)      | 18622_g_at (AJ005902.2_G_AT)   |
| 18625_at (AC005278.22_AT)      | 18686_s_at (U18126_S_AT)       |
| 18698_s_at (X17528_S_AT)       | 18735_s_at (Z29490_S_AT)       |
| 18753_s_at (AF118222.28_S_AT)  | 18803_at (AC005315.94_AT)      |
| 18885_at (AC006921.147_AT)     | 18888_at (AC007591.68_AT)      |
| 18909_s_at (AF055848.2_S_AT)   | 18928_at (AC002333.181_AT)     |

|                               |                                |
|-------------------------------|--------------------------------|
| 18930_at (AC005990.57_AT)     | 18933_at (AC007020.48_AT)      |
| 18953_at (AF077955.1_AT)      | 18966_at (AC004561.106_AT)     |
| 19019_i_at (X82623.2_I_AT)    | 19137_at (X74755.2_AT)         |
| 19140_at (AC005170.24_AT)     | 19181_s_at (AF053065.2_S_AT)   |
| 19207_at (AC006069.117_AT)    | 19247_at (AF071527.44_AT)      |
| 19257_s_at (AC000104.57_S_AT) | 19325_at (AL022604.42_AT)      |
| 19364_at (AL022023.142_AT)    | 19407_at (AC004697.81_AT)      |
| 19409_at (AC007357.56_AT)     | 19411_at (AC007661.104_AT)     |
| 19421_at (X70990.4_AT)        | 19432_s_at (AL035680.11_S_AT)  |
| 19451_at (AC004392.6_AT)      | 19460_s_at (AC000132.66_S_AT)  |
| 19465_at (AL021768.96_AT)     | 19546_at (AC005398.172_AT)     |
| 19555_at (AF058919.48_AT)     | 19624_at (AL049481.196_AT)     |
| 19641_at (AC004561.66_AT)     | 19667_at (AL021710.5_AT)       |
| 19741_at (AL049171.72_AT)     | 19755_at (AC006593.64_AT)      |
| 19818_i_at (AL021749.33_I_AT) | 19881_at (AC004077.49_AT)      |
| 19892_at (AC005770.30_AT)     | 19944_at (AC002130.4_AT)       |
| 19956_at (AC006282.11_AT)     | 19970_s_at (AC003674.10_S_AT)  |
| 19982_at (AC002986.28_AT)     | 19991_at (AC007017.124_AT)     |
| 20017_at (AC004521.66_AT)     | 20030_at (AL078637.51_AT)      |
| 20051_at (AC000106.38_AT)     | 20098_at (AC004697.123_AT)     |
| 20133_i_at (AC007178.71_I_AT) | 20134_s_at (AC007178.71_S_AT)  |
| 20144_at (AL079350.68_AT)     | 20245_s_at (AC005309.97_S_AT)  |
| 20247_at (AC004392.4_AT)      | 20269_at (AC002387.237_AT)     |
| 20271_at (Z99707.27_AT)       | 20287_at (Y14590.5_AT)         |
| 20288_g_at (Y14590.5_G_AT)    | 20291_s_at (M92353.4_S_AT)     |
| 20323_at (AC004561.62_AT)     | 20356_at (AC004561.74_AT)      |
| 20370_at (AC004561.263_AT)    | 20420_at (AL024486.131_AT)     |
| 20421_at (U81294.2_AT)        | 20422_g_at (U81294.2_G_AT)     |
| 20432_at (U43486.2_AT)        | 20450_at (AJ005930.2_AT)       |
| 20461_at (AL049480.157_AT)    | 20462_at (U82399.2_AT)         |
| 20479_i_at (AF069495.2_I_AT)  | 20480_s_at (AF069495.2_S_AT)   |
| 20529_at (Z97341.125_AT)      | 20572_s_at (AC005560.229_S_AT) |
| 20589_at (AF081066.3_AT)      | 20669_s_at (AC002388.6_S_AT)   |
| 20685_at (AL049751.46_AT)     | 20689_s_at (AC002335.19_S_AT)  |

**Table 9** Probe Sets corresponding to genes, the expression of which is downregulated (repressed) in an avrRpt-2-dependent manner in *Arabidopsis*

| ProbeSet                       | Description  |
|--------------------------------|--|
| 11991_g_at (AC002387.210_G_AT) | 11991_g_at<br>(AC002387.210_G_AT)gb AAB82645.1 <br>(AC002387) unknown protein [Arabidopsis thaliana]   |
| 12091_at (AC004450.116_AT)     | 12091_at (AC004450.116_AT)gb AAC64313.1 <br>(AC004450) unknown protein [Arabidopsis thaliana]  |
| 12124_s_at (Z97337.149_S_AT)   | 12124_s_at (Z97337.149_S_AT)emb CAB10270.1 <br>(Z97337) imidazoleglycerol-phosphate dehydratase [Arabidopsis thaliana]                               |
| 12125_at (Z97341.99_AT)        | 12125_at (Z97341.99_AT)emb CAB10421.1 <br>(Z97341) hypothetical protein [Arabidopsis thaliana]   |
| 12160_at (AC006284.117_AT)     | 12160_at (AC006284.117_AT)gb AAD17436.1 <br>(AC006284) unknown protein [Arabidopsis thaliana]  |
| 12191_at (AC006068.35_AT)      | 12191_at (AC006068.35_AT)gb AAD15440.1 <br>(AC006068) unknown protein [Arabidopsis thaliana]   |
| 12193_at (AC006072.132_AT)     | 12193_at (AC006072.132_AT)gb AAD13708.1 <br>(AC006072) unknown protein [Arabidopsis thaliana]  |
| 12223_s_at (AC007168.178_S_AT) | 12223_s_at<br>(AC007168.178_S_AT)gb AAF18668.1 AC007168_1<br>(AC007168) unknown protein [Arabidopsis thaliana]                                       |
| 12290_at (Y09418.2_AT)         | 12290_at (Y09418.2_AT)emb CAA70572.1 <br>(Y09418) CDPK-related protein kinase [Arabidopsis thaliana]   |
| 12421_at (AJ002414.1_AT)       | 12421_at (AJ002414.1_AT)emb CAA05398.1 <br>(AJ002414) hnRNP-like protein [Arabidopsis thaliana]  |
| 12460_s_at (AC006920.129_S_AT) | 12460_s_at<br>(AC006920.129_S_AT)gb AAD22284.1 AC006920_8<br>(AC006920) DNA-directed RNA polymerase II, third largest subunit [Arabidopsis thaliana] |
| 12493_g_at (Y09095.1_G_AT)     | 12493_g_at (Y09095.1_G_AT)emb CAA70310.1 <br>(Y09095) chloride channel [Arabidopsis thaliana]  |

| ProbeSet                       | Description  |
|--------------------------------|--|
| 12559_at (AC005727.83_AT)      | 12559_at (AC005727.83_AT)gb AAC79586.1  (AC005727) putative DOF zinc finger protein [Arabidopsis thaliana]   |
| 12560_at (AC005825.57_AT)      | 12560_at (AC005825.57_AT)gb AAD24598.1 AC005825_5 (AC005825) putative chloroplast outer membrane protein [Arabidopsis thaliana]  |
| 12561_at (AL021687.107_AT)     | 12561_at (AL021687.107_AT)emb CAA16701.1  (AL021687) putative protein [Arabidopsis thaliana]   |
| 12855_f_at (ADH_F_AT)          | 12855_f_at (ADH_F_AT)gb AAC00625.1  (AC002291) Alcohol Dehydrogenase [Arabidopsis thaliana]  |
| 12962_at (AC004697.165_AT)     | 12962_at (AC004697.165_AT)gb AAC28988.1  (AC004697) putative ABC transporter [Arabidopsis thaliana]  |
| 13221_at (CHS-WHOLE-SEQ_AT)    | 13221_at (CHS-WHOLE-SEQ_AT)gb AAF23561.1 AF112086_1 (AF112086) chalcone synthase [Arabidopsis thaliana]  |
| 13253_f_at (FPS1_F_AT)         | 13253_f_at (FPS1_F_AT)gb AAB49290.1  (U80605) farnesyl diphosphate synthase precursor [Arabidopsis thaliana]   |
| 13459_at (AF013294.21_AT)      | 13459_at (AF013294.21_AT)gb AAB62852.1  (AF013294) similar to auxin-induced protein [Arabidopsis thaliana]   |
| 13708_s_at (AC007259X_S_AT)    | 13708_s_at (AC007259X_S_AT)gb AAD50000.1 AC007259_1 3 (AC007259) Similar to protein kinases [Arabidopsis thaliana]   |
| 14428_s_at (AL023094.342_S_AT) | 14428_s_at (AL023094.342_S_AT)emb CAA18852.1  (AL023094) putative protein [Arabidopsis thaliana]   |
| 15175_s_at (ATU28215_S_AT)     | 15175_s_at (ATU28215_S_AT)gb AAC62130.1  (AC005169) hexokinase (ATHXK2) [Arabidopsis thaliana]   |
| 15943_at (AC007202.16_AT)      | 15943_at (AC007202.16_AT)gb AAD30227.1 AC007202_9 (AC007202) Strong similarity to gb Y14272 3-deoxy-D-manno-2-octulosonate-8-phosphate synthase from Pisum sativum. ESTs gb AA067485 and gb AI100551 come from this gene. [Arabidopsis thaliana] |



| ProbeSet                   | Description  |
|----------------------------|--|
| 19219_at (AC007019.185_AT) | 19219_at (AC007019.185_AT)gb AAD20413.1 <br>(AC007019) unknown protein [Arabidopsis<br>thaliana] |

**Table 10A:** Expression data for 9 probe sets corresponding to genes that are specifically induced during incompatible interaction within 3 hours

| ProbeSet   | Col  |         |      |      |         |      |
|------------|------|---------|------|------|---------|------|
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12094_at   | ud   | 303     | ud   | 19   | 28      | ud   |
| 12128_at   | 84   | 255     | ud   | 0    | 181     | ud   |
| 12712_f_at | 10   | 278     | ud   | ud   | 90      | 20   |
| 13763_at   | 18   | 176     | 16   | 64   | 274     | 40   |
| 14882_at   | 24   | 385     | 11   | 62   | 457     | 46   |
| 16398_s_at | 16   | 104     | 5    | 17   | 65      | 18   |
| 16536_s_at | 24   | 346     | 19   | ud   | 19      | 30   |
| 17180_at   | 50   | 361     | 26   | 93   | 367     | 76   |
| 19970_s_at | 18   | 175     | 8    | 14   | 65      | 11   |
| ProbeSet   | Ws   |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12094_at   | ud   | 104     | ud   | ud   | 29      | ud   |
| 12128_at   | ud   | 77      | ud   | ud   | 29      | ud   |
| 12712_f_at | ud   | 146     | ud   | ud   | 46      | ud   |
| 13763_at   | ud   | 90      | ud   | 15   | 378     | 25   |
| 14882_at   | 19   | 255     | 7    | 23   | 291     | 32   |
| 16398_s_at | 9    | 93      | 7    | 4    | 49      | ud   |
| 16536_s_at | ud   | 409     | ud   | ud   | 4       | 8    |
| 17180_at   | ud   | 200     | ud   | ud   | 259     | ud   |
| 19970_s_at | 11   | 146     | 5    | 4    | 46      | 6    |
| ProbeSet   | Ler  |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12094_at   | ud   | 121     | ud   | 24   | 41      | ud   |
| 12128_at   | 71   | 160     | ud   | 33   | 145     | ud   |
| 12712_f_at | 19   | 113     | 11   | 24   | 46      | 14   |
| 13763_at   | 22   | 85      | 19   | 139  | 248     | 43   |
| 14882_at   | 50   | 227     | 25   | 121  | 321     | 41   |
| 16398_s_at | 20   | 85      | 20   | 23   | 57      | 10   |

|            |      |         |      |      |         |      |
|------------|------|---------|------|------|---------|------|
| 16536_s_at | 13   | 110     | 29   | ud   | ud      | 78   |
| 17180_at   | 46   | 181     | 18   | 123  | 214     | 37   |
| 19970_s_at | 23   | 140     | 13   | 32   | 55      | 22   |
| ProbeSet   | Cvi  |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12094_at   | 18   | 166     | ud   | 32   | 54      | ud   |
| 12128_at   | 75   | 204     | 37   | 52   | 168     | 28   |
| 12712_f_at | 11   | 92      | ud   | 14   | 39      | 13   |
| 13763_at   | 42   | 166     | 21   | 133  | 344     | 25   |
| 14882_at   | 45   | 213     | 34   | 86   | 417     | 26   |
| 16398_s_at | 12   | 52      | 16   | 25   | 67      | 14   |
| 16536_s_at | 15   | 164     | 30   | ud   | 8       | 23   |
| 17180_at   | 45   | 203     | ud   | 152  | 263     | ud   |
| 19970_s_at | 25   | 142     | 11   | 19   | 68      | 13   |

**Table 10B:** Expression data for 18 probe sets corresponding to genes that are specifically induced during incompatible interaction in 6 hours (but not within 3 hours)

| ProbeSet   | Col  |         |      |      |         |      |
|------------|------|---------|------|------|---------|------|
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12711_f_at | ud   | ud      | ud   | ud   | 100     | ud   |
| 13312_at   | ud   | 118     | ud   | 36   | 176     | 11   |
| 13370_at   | ud   | ud      | ud   | ud   | 144     | ud   |
| 13818_s_at | ud   | 9       | ud   | 65   | 476     | 21   |
| 14609_at   | ud   | 11      | 4    | 23   | 233     | 21   |
| 14635_s_at | ud   | ud      | ud   | 42   | 360     | 13   |
| 14931_at   | ud   | ud      | ud   | 10   | 284     | 16   |
| 15120_s_at | ud   | ud      | ud   | ud   | 214     | 12   |
| 16357_at   | 8    | 70      | 6    | 14   | 102     | 26   |
| 16968_at   | 19   | 28      | 7    | 33   | 257     | 19   |
| 17134_at   | ud   | ud      | ud   | ud   | 199     | ud   |
| 17371_at   | ud   | 16      | ud   | 31   | 191     | 19   |
| 17485_s_at | ud   | ud      | ud   | ud   | 86      | 7    |
| 18631_at   | 10   | 105     | ud   | 32   | 92      | 17   |
| 19451_at   | ud   | ud      | ud   | ud   | 116     | ud   |
| 20323_at   | 20   | 51      | 19   | 12   | 280     | 14   |
| 20356_at   | ud   | ud      | ud   | ud   | 82      | ud   |
| 20421_at   | ud   | ud      | ud   | 26   | 685     | 20   |
| ProbeSet   | Ws   |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12711_f_at | ud   | ud      | ud   | ud   | 86      | ud   |
| 13312_at   | ud   | 34      | ud   | 11   | 114     | 2    |
| 13370_at   | ud   | ud      | ud   | 20   | 70      | ud   |
| 13818_s_at | ud   | 12      | 8    | 10   | 422     | 10   |
| 14609_at   | ud   | 12      | ud   | ud   | 184     | 3    |
| 14635_s_at | ud   | ud      | ud   | ud   | 264     | ud   |
| 14931_at   | ud   | ud      | ud   | ud   | 36      | ud   |
| 15120_s_at | ud   | 13      | ud   | ud   | 89      | ud   |
| 16357_at   | ud   | 31      | ud   | ud   | 80      | 6    |
| 16968_at   | ud   | 5       | ud   | 12   | 54      | ud   |

|            |      |         |      |      |         |      |
|------------|------|---------|------|------|---------|------|
| 17134_at   | ud   | ud      | ud   | ud   | 27      | ud   |
| 17371_at   | ud   | ud      | ud   | ud   | 133     | ud   |
| 17485_s_at | ud   | ud      | ud   | 8    | 85      | ud   |
| 18631_at   | 10   | 103     | ud   | 17   | 81      | 10   |
| 19451_at   | ud   | 6       | ud   | ud   | 52      | ud   |
| 20323_at   | ud   | 60      | 11   | ud   | 119     | ud   |
| 20356_at   | ud   | ud      | ud   | ud   | 96      | ud   |
| 20421_at   | ud   | 5       | ud   | ud   | 67      | ud   |
| ProbeSet   | Ler  |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12711_f_at | ud   | ud      | 5    | 15   | 32      | ud   |
| 13312_at   | ud   | 48      | ud   | 133  | 158     | 12   |
| 13370_at   | ud   | ud      | ud   | 36   | 122     | ud   |
| 13818_s_at | ud   | ud      | 4    | 26   | 211     | ud   |
| 14609_at   | ud   | ud      | 5    | 30   | 359     | 15   |
| 14635_s_at | ud   | ud      | ud   | 22   | 310     | 17   |
| 14931_at   | ud   | ud      | ud   | ud   | 149     | ud   |
| 15120_s_at | ud   | ud      | ud   | 17   | 252     | ud   |
| 16357_at   | 14   | 47      | 10   | 23   | 117     | 11   |
| 16968_at   | 14   | 18      | 5    | ud   | 105     | 5    |
| 17134_at   | ud   | ud      | ud   | ud   | 112     | ud   |
| 17371_at   | ud   | ud      | ud   | ud   | ud      | ud   |
| 17485_s_at | ud   | ud      | ud   | ud   | 49      | ud   |
| 18631_at   | 16   | 63      | 27   | 34   | 68      | ud   |
| 19451_at   | 5    | 7       | 4    | 14   | 123     | ud   |
| 20323_at   | 11   | 6       | 8    | ud   | 11      | ud   |
| 20356_at   | ud   | 5       | ud   | 17   | 111     | ud   |
| 20421_at   | ud   | ud      | ud   | 7    | 329     | ud   |
| ProbeSet   | Cvi  |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 12711_f_at | ud   | ud      | ud   | 15   | 46      | 7    |
| 13312_at   | 10   | 64      | 5    | 60   | 116     | ud   |
| 13370_at   | ud   | ud      | 18   | 40   | 88      | ud   |
| 13818_s_at | 4    | ud      | 7    | 47   | 274     | 10   |

|            |    |    |    |     |     |    |
|------------|----|----|----|-----|-----|----|
| 14609_at   | 8  | 23 | 6  | 127 | 778 | ud |
| 14635_s_at | 6  | ud | ud | 179 | 519 | 44 |
| 14931_at   | ud | ud | ud | 23  | 182 | ud |
| 15120_s_at | ud | 17 | ud | 31  | 150 | ud |
| 16357_at   | 17 | 52 | 16 | 32  | 132 | 16 |
| 16968_at   | 16 | 20 | 18 | ud  | 56  | 3  |
| 17134_at   | ud | ud | ud | ud  | 121 | ud |
| 17371_at   | ud | ud | ud | ud  | ud  | ud |
| 17485_s_at | ud | ud | ud | ud  | 55  | ud |
| 18631_at   | 11 | 41 | 10 | 24  | 85  | ud |
| 19451_at   | 4  | 6  | ud | 19  | 413 | ud |
| 20323_at   | 7  | ud | ud | ud  | 11  | ud |
| 20356_at   | ud | ud | ud | 13  | 107 | ud |
| 20421_at   | ud | 20 | ud | ud  | 293 | ud |

**Table 10C:** Expression data for 6 probe sets corresponding to genes that are activated by *P. syringae* in 6 hours. Most of them are compatible interaction-specific/preferential

| ProbeSet   | Col  |         |      |      |         |      |
|------------|------|---------|------|------|---------|------|
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 13243_r_at | 44   | ud      | 24   | 123  | 91      | 14   |
| 14573_at   | ud   | ud      | ud   | 234  | 32      | 23   |
| 14613_at   | ud   | ud      | 9    | 138  | ud      | 10   |
| 18122_at   | 34   | 38      | 25   | 169  | 96      | 28   |
| 19150_at   | ud   | ud      | 3    | 97   | 4       | 6    |
| 19673_g_at | 69   | 34      | 27   | 596  | 290     | ud   |
| ProbeSet   | Ws   |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 13243_r_at | 27   | 38      | 16   | 132  | 96      | 17   |
| 14573_at   | ud   | ud      | ud   | 377  | 63      | 12   |
| 14613_at   | ud   | 10      | ud   | 136  | ud      | ud   |
| 18122_at   | 33   | 56      | 30   | 235  | 76      | 24   |
| 19150_at   | ud   | ud      | ud   | 218  | 9       | ud   |
| 19673_g_at | 21   | 6       | ud   | 496  | 312     | ud   |
| ProbeSet   | Ler  |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 13243_r_at | 66   | 52      | 45   | 149  | 59      | 7    |
| 14573_at   | ud   | ud      | ud   | 22   | 17      | 9    |
| 14613_at   | ud   | ud      | 14   | 102  | 13      | 13   |
| 18122_at   | 42   | 58      | 42   | 136  | 60      | 20   |
| 19150_at   | ud   | ud      | ud   | 24   | 7       | ud   |
| 19673_g_at | 30   | 2       | 10   | 426  | 181     | ud   |
| ProbeSet   | Cvi  |         |      |      |         |      |
|            | 3 hr |         |      | 6 hr |         |      |
|            | Pst  | Pst/aR2 | cont | Pst  | Pst/aR2 | cont |
| 13243_r_at | 62   | 42      | 55   | 142  | 116     | ud   |
| 14573_at   | ud   | ud      | ud   | 25   | 22      | ud   |
| 14613_at   | 13   | ud      | 13   | 63   | ud      | ud   |

|            |    |    |    |            |     |    |
|------------|----|----|----|------------|-----|----|
| 18122_at   | 36 | 43 | 41 | <b>151</b> | 81  | 28 |
| 19150_at   | ud | 6  | ud | <b>11</b>  | 3   | ud |
| 19673_g_at | ud | ud | 26 | <b>416</b> | 184 | ud |

Pst/aR2 ... represents the incompatible interaction

Pst ... *P. syringae* pv. tomato

aR2 ... avrRpt2

ud ... undetectable



**Table 11** Probe Sets corresponding to genes induced early after infection of different *Arabidopsis* ecotypes with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121 (at 3 or 6 hours) or by estradiol inducible

| ProbeSet                                  | Description  |
|---|--|
| 12737_f_at (Z97049_F_AT)                  | 12737_f_at (Z97049_F_AT)emb CAA74603.1  (Y14207) R2R3-MYB transcription factor [Arabidopsis thaliana]                  |
| 12908_s_at (ATERF5_S_AT)                  | 12908_s_at (ATERF5_S_AT)dbj BAA32422.1  (AB008107) ethylene responsive element binding factor 5 [Arabidopsis thaliana] |
| 16398_s_at (AL022603.3_S_AT)              | 16398_s_at (AL022603.3_S_AT)emb CAA20204.1  (AL031187) serine/threonine kinase-like protein [Arabidopsis thaliana]     |
| 17105_s_at (AF055357_S_AT)                | 17105_s_at (AF055357_S_AT)gb AAC39479.1  (AF055357) respiratory burst oxidase protein D [Arabidopsis thaliana]         |
| 19970_s_at (AC003674.10_S_AT)             | 19970_s_at (AC003674.10_S_AT)gb AAB97120.1  (AC003674) unknown protein [Arabidopsis thaliana]                          |
| 20620_g_at (AC005896.161_G_AT)            | 20620_g_at (AC005896.161_G_AT)gb AAC98070.1  (AC005896) putative C2H2-type zinc finger protein [Arabidopsis thaliana]  |
| 12094_at (AC006223.143_AT)                | 12094_at (AC006223.143_AT)gb AAD15401.1  (AC006223) putative alanine acetyl transferase [Arabidopsis thaliana]         |
| 13255_i_at (gammaglutamyltranspepti_I_AT) | 13255_i_at (GAMMAGLUTAMYLTRANSPEPTI_I_AT)emb CAA89206.1  (Z49240) gamma-glutamyl transpeptidase [Arabidopsis thaliana] |
| 12712_f_at (Z95774_F_AT)                  | 12712_f_at (Z95774_F_AT)emb CAB09206.1  (Z95774) R2R3-MYB transcription factor [Arabidopsis thaliana]                  |
| 13015_s_at (X98673.2_S_AT)                | 13015_s_at (X98673.2_S_AT)emb CAA67232.1  (X98674) zinc finger protein [Arabidopsis thaliana]                          |
| 13842_at (AC002396.12_AT)                 | 13842_at (AC002396.12_AT)gb AAC00572.1  (AC002396) similar to zinc metalloproteinases [Arabidopsis thaliana]           |

| ProbeSet                      | Description  |
|-------------------------------|--|
| 14235_at (NOVARTIS97_AT)      | 14235_at (NOVARTIS97_AT)gb AAF14671.1 AC011713_19 (AC011713) Similar to gb Z48431 DNA-binding protein from Avena fatua. [Arabidopsis thaliana]           |
| 15779_g_at (X98676.2_G_AT)    | 15779_g_at (X98676.2_G_AT)emb CAA67234.1  (X98676) zinc finger protein [Arabidopsis thaliana]  |
| 16059_s_at (D88206_S_AT)      | 16059_s_at (D88206_S_AT)dbj BAA24694.1  (D88206) protein kinase [Arabidopsis thaliana]   |
| 17180_at (AF007270.30_AT)     | 17180_at (AF007270.30_AT)gb AAB61058.1  (AF007270) contains similarity to GATA-type zinc fingers (PS:PS00344) [Arabidopsis thaliana]                     |
| 17303_s_at (AC004683.25_S_AT) | 17303_s_at (AC004683.25_S_AT)gb AAC67339.2  (AC005499) putative WRKY-type DNA binding protein [Arabidopsis thaliana]                                     |
| 12571_s_at (AF149413.18_S_AT) | 12571_s_at (AF149413.18_S_AT)gb AAD40138.1 AF149413_19 (AF149413) Arabidopsis thaliana ferrochelatase-I (SW:P42043); Pfam                                |
| 12642_at (AC006920.138_AT)    | 12642_at (AC006920.138_AT)gb AAD22285.1 AC006920_9 (AC006920) unknown protein [Arabidopsis thaliana]   |
| 13177_at (AL049640.42_AT)     | 13177_at (AL049640.42_AT)emb CAB40989.1  (AL049640) growth factor like protein [Arabidopsis thaliana]  |
| 14162_at (NOVARTIS46_AT)      | 14162_at (NOVARTIS46_AT)No hits found less than or equal to 1e-15.   |
| 14214_at (NOVARTIS83_AT)      | 14214_at (NOVARTIS83_AT)gb AAF24849.1 AC012679_20 (AC012679) putative calmodulin-binding protein [Arabidopsis thaliana]                                  |
| 14217_at (NOVARTIS85_RC_AT)   | 14217_at (NOVARTIS85_RC_AT)gb AAF07816.1 AC011020_16 (AC011020) putative receptor protein kinase [Arabidopsis thaliana] thaliana. [Arabidopsis thaliana] |
| 14711_s_at (ZFPL_S_AT)        | 14711_s_at (ZFPL_S_AT)gb AAD25930.1 AF085279_3 (AF085279) hypothetical Cys-3-His zinc finger protein [Arabidopsis thaliana]                              |

| ProbeSet                      | Description   |
|-------------------------------|---|
| 14882_at (AL022605.63_AT)     | 14882_at (AL022605.63_AT)emb CAA18753.1 <br>(AL022605) putative protein [Arabidopsis thaliana]  |
| 15431_at (AL030978.64_AT)     | 15431_at (AL030978.64_AT)emb CAA19722.1 <br>(AL030978) putative protein [Arabidopsis thaliana]  |
| 15680_s_at (ATHATPK19B_S_AT)  | 15680_s_at<br>(ATHATPK19B_S_AT)dbj BAA07661.1 <br>(D42061) ribosomal-protein S6 kinase homolog<br>[Arabidopsis thaliana]  |
| 17379_at (AF085279.9_AT)      | 17379_at<br>(AF085279.9_AT)gb AAF18728.1 AC018721_3<br>(AC018721) putative CCH-type zinc finger<br>protein [Arabidopsis thaliana]   |
| 18267_at (AC006223.23_AT)     | 18267_at (AC006223.23_AT)gb AAD15384.1 <br>(AC006223) unknown protein [Arabidopsis thaliana]  |
| 18604_at (AF069298.31_AT)     | 18604_at (AF069298.31_AT)gb AAC19273.1 <br>(AF069298) similar to several small proteins<br>(~100 aa) that are induced by heat, auxin, ethylene<br>and wounding such as Phaseolus aureus indole-3-<br>acetic acid induced protein ARG (SW:32292)<br>[Arabidopsis thaliana] |
| 19460_s_at (AC000132.66_S_AT) | 19460_s_at<br>(AC000132.66_S_AT)gb AAB60749.1 <br>(AC000132) Identical to A. thaliana HEMA2<br>(gb U27118). [Arabidopsis thaliana]  |
| 15775_at (AL079344.196_AT)    | 15775_at (AL079344.196_AT)emb CAB45334.1 <br>(AL079344) cytokinin oxidase-like protein<br>[Arabidopsis thaliana]  |
| 16939_at (AC002334.110_AT)    | 16939_at (AC002334.110_AT)gb AAC04922.1 <br>(AC002334) putative synaptobrevin [Arabidopsis thaliana]  |
| 18672_s_at (D13983_S_AT)      | 18672_s_at (D13983_S_AT)dbj BAA03090.1 <br>(D13983) chloroplast envelope Ca <sup>2+</sup> -ATPase<br>precursor [Arabidopsis thaliana]   |

**Table 12** Probe Sets corresponding to genes repressed early after infection of *Arabidopsis* with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121

| ProbeSet                      | Description   |
|-------------------------------|---|
| 12212_at (AL049711.168_AT)    | 12212_at (AL049711.168_AT)emb CAB41327.1  (AL049711) putative protein [Arabidopsis thaliana]  |
| 17555_s_at (ATU89296_S_AT)    | 17555_s_at (ATU89296_S_AT)gb AAB62404.1  (U89296) auxin response transcription factor 3; ARF3 [Arabidopsis thaliana]  |
| 17975_at (AF175998_AT)        | 17975_at (AF175998_AT)gb AAD53103.1 AF175998_1 (AF175998) putative transcription factor [Arabidopsis thaliana]  |
| 14386_at (AC005309.177_AT)    | 14386_at (AC005309.177_AT)gb AAC63645.1  (AC005309) unknown protein [Arabidopsis thaliana]  |
| 16613_s_at (AF012657_S_AT)    | 16613_s_at (AF012657_S_AT)gb AAC49845.1  (AF012657) putative potassium transporter AtKT2p [Arabidopsis thaliana]  |
| 18916_s_at (X92393.1_S_AT)    | 18916_s_at (X92393.1_S_AT)emb CAA63131.1  (X92393) KNAT4 homeobox protein [Arabidopsis thaliana]  |
| 20443_s_at (AC006341.42_S_AT) | 20443_s_at (AC006341.42_S_AT)gb AAD34693.1 AC006341_21 (AC006341) Similar to gb AF069494 cytochrome P450 from Sinapis alba and is a member of the PF 00067 Cytochrome P450 family. EST gb F14190 comes from this gene. [Arabidopsis thaliana] |
| 20524_at (AC005698.12_AT)     | 20524_at (AC005698.12_AT)gb AAD43613.1 AC005698_12 (AC005698) T3P18.12 [Arabidopsis thaliana]   |
| 12609_at (X92975.2_AT)        | 12609_at (X92975.2_AT)emb CAA63553.1  (X92975) xyloglucan endo-transglycosylase [Arabidopsis thaliana]  |
| 20061_at (AC005508.23_AT)     | 20061_at (AC005508.23_AT)gb AAD14498.1  (AC005508) 37496 [Arabidopsis thaliana]   |

**Table 13a** Probe Sets corresponding to genes, the expression of which is induced in an incompatible interaction at 3 and/or 6 hours after infection of four *Arabidopsis* ecotypes with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121

| ProbeSet                       | Description  |
|--------------------------------|--|
| 12128_at (AC004261.157_AT)     | 12128_at (AC004261.157_AT)gb AAD12010.1  (AC004261) unknown protein [Arabidopsis thaliana]   |
| 12335_at (AC004411.73_AT)      | 12335_at (AC004411.73_AT)gb AAC34243.1  (AC004411) putative protein kinase [Arabidopsis thaliana]  |
| 12347_at (AC007258.28_AT)      | 12347_at (AC007258.28_AT)gb AAD39325.1 AC007258_14 (AC007258) Putative ATPase [Arabidopsis thaliana]   |
| 12497_at (AC006533.51_AT)      | 12497_at (AC006533.51_AT)gb AAD32284.1 AC006533_8 (AC006533) putative receptor-like protein kinase [Arabidopsis thaliana]  |
| 12759_at (AC005278.32_AT)      | 12759_at (AC005278.32_AT)gb AAC72120.1  (AC005278) Strong similarity to gb D14550 extracellular dermal glycoprotein (EDGP) precursor from <i>Daucus carota</i> . ESTs gb 84105 and gb AI100071 come from this gene. [Arabidopsis thaliana] |
| 12790_s_at (AL021635.58_S_AT)  | 12790_s_at (AL021635.58_S_AT)emb CAA16554.1  (AL021635) cytochrome P450 like protein [Arabidopsis thaliana]  |
| 12801_at (AC005223.34_AT)      | 12801_at (AC005223.34_AT)gb AAD10652.1  (AC005223) Unknown protein [Arabidopsis thaliana]  |
| 12904_s_at (ATERF1_S_AT)       | 12904_s_at (ATERF1_S_AT)dbj BAA32418.1  (AB008103) ethylene responsive element binding factor 1 [Arabidopsis thaliana]   |
| 12909_s_at (ATERF6_S_AT)       | 12909_s_at (ATERF6_S_AT)emb CAB10530.1  (Z97343) EREBP-4 like protein [Arabidopsis thaliana]   |
| 12989_s_at (AC004077.149_S_AT) | 12989_s_at (AC004077.149_S_AT)gb AAC26690.1  (AC004077) putative cytochrome P450 [Arabidopsis thaliana]  |
| 13014_at (U93215.87_AT)        | 13014_at (U93215.87_AT)gb AAB63082.1  (U93215) putative lipase [Arabidopsis thaliana]  |

| ProbeSet                                  | Description  |
|---|--|
| 13110_at (AF074021.34_AT)                 | 13110_at (AF074021.34_AT)gb AAD29776.1 AF074021_8 (AF074021) putative symbiosis-related protein [Arabidopsis thaliana]                                     |
| 13115_at (AC000375.44_AT)                 | 13115_at (AC000375.44_AT)gb AAB60774.1  (AC000375) ESTs gb U75592,gb T13956,gb T43869 come from from this gene. [Arabidopsis thaliana]                     |
| 13256_s_at (gammaglutamyltranspepti_S_AT) | 13256_s_at (gammaglutamyltranspepti_S_AT)emb CAA89206.1  (Z49240) gamma-glutamyl transpeptidase [Arabidopsis thaliana]                                     |
| 13270_at (HSF21_AT)                       | 13270_at (HSF21_AT)gb AAC31792.1  (U68561) heat shock transcription factor 21 [Arabidopsis thaliana]   |
| 13271_g_at (HSF21_G_AT)                   | 13271_g_at (HSF21_G_AT)gb AAC31792.1  (U68561) heat shock transcription factor 21 [Arabidopsis thaliana]   |
| 13273_s_at (HSF4_S_AT)                    | 13273_s_at (HSF4_S_AT)gb AAC31756.1  (U68017) heat shock transcription factor 4 [Arabidopsis thaliana]   |
| 13279_s_at (HSP176II_S_AT)                | 13279_s_at (HSP176II_S_AT)emb CAA45039.1  (X63443) heat shock protein 17.6-II [Arabidopsis thaliana]   |
| 13370_at (AC005322.4_AT)                  | 13370_at (AC005322.4_AT)gb AAC97990.1  (AC005322) Strong similarity to Dsor1 protein kinase gb D13782 from Drosophila melanogaster. [Arabidopsis thaliana] |
| 13645_at (AC000098.8_AT)                  | 13645_at (AC000098.8_AT)gb AAB71443.1  (AC000098) EST gb ATTS0295 comes from this gene. [Arabidopsis thaliana]   |
| 13656_at (AC007138.31_AT)                 | 13656_at (AC007138.31_AT)gb AAD22649.1 AC007138_13 (AC007138) predicted protein of unknown function [Arabidopsis thaliana]                                 |
| 13714_at (NOVARTIS101_RC_AT)              | 13714_at gb AAF18728.1 AC018721_3 (AC018721) putative CCH-type zinc finger protein [Arabidopsis thaliana]  |
| 13806_at (AC002354.15_AT)                 | 13806_at (AC002354.15_AT)gb AAB81668.1  (AC002354) NAM (no apical meristem)-like protein [Arabidopsis thaliana]  |

| ProbeSet                       | Description  |
|--------------------------------|--|
| 13818_s_at (AC006218.175_S_AT) | 13818_s_at<br>(AC006218.175_S_AT)gb AAD15433.1 <br>(AC006218) putative aspartate aminotransferase<br>[Arabidopsis thaliana]  |
| 14030_at (AC005970.225_AT)     | 14030_at (AC005970.225_AT)gb AAC95171.1 <br>(AC005970) putative protein kinase [Arabidopsis<br>thaliana]   |
| 14139_at (NOVARTIS30_AT)       | 14139_at gb AAD09343.1  (AF026538) ABA-<br>responsive protein [Hordeum vulgare]  |
| 14141_at (NOVARTIS31_AT)       | 14141_at No hits found less than or equal to 1e-15.  |
| 14170_at (NOVARTIS51_AT)       | 14170_at gb AAF29406.1 AC022354_5 (AC022354)<br>unknown protein [Arabidopsis thaliana]   |
| 14248_at (PAD3_AT)             | 14248_at<br>(PAD3_AT)gb AAD31062.1 AC007357_11<br>(AC007357) Strong similarity to gb X97864<br>cytochrome P450 from Arabidopsis thaliana and is a<br>member of the PF 00067 Cytochrome P450 family.<br>ESTs gb N65665, gb T14112, gb T76255, gb T20906<br>and gb AI100027 come from this gene. |
| 14431_at (AL035394.279_AT)     | 14431_at (AL035394.279_AT)emb CAA23047.1 <br>(AL035394) putative protein [Arabidopsis thaliana]  |
| 14605_at (AC006193.6_AT)       | 14605_at<br>(AC006193.6_AT)gb AAD38247.1 AC006193_3<br>(AC006193) very similar to alcohol dehydrogenase<br>[Arabidopsis thaliana]  |
| 14608_at (AC007357.49_AT)      | 14608_at<br>(AC007357.49_AT)gb AAD31074.1 AC007357_23<br>(AC007357) Similar to gb AF038007 FIC1 gene<br>from Homo sapiens and is a member of the PF 00122<br>E1-E2 ATPase family. ESTs gb T45045 and<br>gb AA394473 come from this gene. [Arabidopsis<br>thaliana]                             |
| 14609_at (AC002340.147_AT)     | 14609_at (AC002340.147_AT)gb AAC02748.1 <br>(AC002340) putative cytochrome P450 [Arabidopsis<br>thaliana]  |
| 14697_g_at (WT740_RC_G_AT)     | 14697_g_at<br>(WT740_RC_G_AT)gb AAD23643.1 AC007119_9<br>(AC007119) unknown protein [Arabidopsis thaliana]   |
| 14763_at (X86958.1_AT)         | 14763_at (X86958.1_AT)emb CAA60521.1 <br>(X86958) protein kinase catalytic domain (fragment)<br>[Arabidopsis thaliana]   |

| ProbeSet                      | Description  |
|-------------------------------|--|
| 14959_at (AC007202.26_AT)     | 14959_at (AC007202.26_AT)gb AAD30230.1 AC007202_12 (AC007202) T8K14.13 [Arabidopsis thaliana]                              |
| 14978_at (AC002333.49_AT)     | 14978_at (AC002333.49_AT)gb AAB64024.1  (AC002333) putative glucosyltransferase [Arabidopsis thaliana]                     |
| 15039_at (AF001308.67_AT)     | 15039_at (AF001308.67_AT)gb AAC78710.1  (AF001308) drought-induced-19-like 1 [Arabidopsis thaliana]                        |
| 15040_g_at (AF001308.67_G_AT) | 15040_g_at (AF001308.67_G_AT)gb AAC78710.1  (AF001308) drought-induced-19-like 1 [Arabidopsis thaliana]                    |
| 15120_s_at (ATU10034_S_AT)    | 15120_s_at (ATU10034_S_AT)gb AAA93132.1  (U10034) glutamate decarboxylase [Arabidopsis thaliana]                           |
| 15137_s_at (ATU57320_S_AT)    | 15137_s_at (ATU57320_S_AT)gb AAB47973.1  (U57320) blue copper-binding protein II [Arabidopsis thaliana]                    |
| 15203_s_at (AB013887_S_AT)    | 15203_s_at (AB013887_S_AT)dbj BAA34251.1  (AB013887) RAV2 [Arabidopsis thaliana]   |
| 17376_at (AL021890.218_AT)    | 17376_at (AL021890.218_AT)emb CAA17164.1  (AL021890) putative protein [Arabidopsis thaliana]                               |
| 17381_at (Z99708.402_AT)      | 17381_at (Z99708.402_AT)emb CAB16811.1  (Z99708) putative protein [Arabidopsis thaliana]                                   |
| 15551_at (AL035440.289_AT)    | 15551_at (AL035440.289_AT)emb CAB36537.1  (AL035440) putative dihydrolipoamide succinyltransferase [Arabidopsis thaliana]  |
| 15582_s_at (ATH131392_S_AT)   | 15582_s_at (ATH131392_S_AT)emb CAA10364.1  (AJ131392) alternative oxidase [Arabidopsis thaliana]                           |
| 15665_s_at (AF022658_S_AT)    | 15665_s_at (AF022658_S_AT)gb AAB80922.1  (AF022658) putative c2h2 zinc finger transcription factor [Arabidopsis thaliana]  |
| 15672_s_at (AF082299_S_AT)    | 15672_s_at (AF082299_S_AT)No hits found less than or equal to 1e-15.   |
| 15919_at (AC007060.42_AT)     | 15919_at (AC007060.42_AT)gb AAD25764.1 AC007060_22 (AC007060) EST gb AA721821 comes from this gene. [Arabidopsis thaliana] |



| ProbeSet                      | Description   |
|-------------------------------|---|
| 15924_at (AC007138.61_AT)     | 15924_at<br>(AC007138.61_AT)gb AAD22658.1 AC007138_22<br>(AC007138) predicted protein of unknown function<br>[Arabidopsis thaliana]   |
| 15978_at (X68592.6_AT)        | 15978_at (X68592.6_AT)emb CAA48579.1 <br>(X68592) adenosine nucleotide translocator<br>[Arabidopsis thaliana]   |
| 16003_s_at (AL021749.64_S_AT) | 16003_s_at<br>(AL021749.64_S_AT)emb CAA16877.1 <br>(AL021749) ADP, ATP carrier-like protein<br>[Arabidopsis thaliana]   |
| 16053_i_at (Y14251.4_I_AT)    | 16053_i_at (Y14251.4_I_AT)emb CAA74639.1 <br>(Y14251) glutathione S-transferase [Arabidopsis<br>thaliana]   |
| 16077_s_at (AF085230_S_AT)    | 16077_s_at (AF085230_S_AT)gb AAD16046.1 <br>(AF085230) phytochelatin synthase 1 [Arabidopsis<br>thaliana]   |
| 16134_s_at (AF132016_S_AT)    | 16134_s_at<br>(AF132016_S_AT)gb AAD33584.1 AF132016_1<br>(AF132016) RING-H2 zinc finger protein ATL6<br>[Arabidopsis thaliana]  |
| 16151_s_at (ATHPRKINA_S_AT)   | 16151_s_at (ATHPRKINA_S_AT)gb AAA18853.1 <br>(L07248) protein kinase [Arabidopsis thaliana]   |
| 16232_s_at (AL080252.77_S_AT) | 16232_s_at<br>(AL080252.77_S_AT)emb CAB45796.1 <br>(AL080252) putative protein [Arabidopsis thaliana]   |
| 16357_at (AF149413.38_AT)     | 16357_at<br>(AF149413.38_AT)gb AAD40144.1 AF149413_25<br>(AF149413) contains similarity to protein kinase<br>domains (Pfam F00069, Score=162.6, E=6.8e-45,<br>N=1) and leucien rich repeats (Pfam PF00560,<br>Score=210.7, E=2.2e-59, N=10) [Arabidopsis<br>thaliana] |
| 16442_s_at (AJ002551.2_S_AT)  | 16442_s_at (AJ002551.2_S_AT)emb CAA05547.1 <br>(AJ002551) heat shock protein 70 [Arabidopsis<br>thaliana]   |
| 16457_s_at (AC005397.17_S_AT) | 16457_s_at<br>(AC005397.17_S_AT)gb AAD23036.1 AC006526_2<br>(AC006526) unknown protein [Arabidopsis<br>thaliana]  |
| 16539_s_at (AB013301_S_AT)    | 16539_s_at (AB013301_S_AT)emb CAB10530.1 <br>(Z97343) EREBP-4 like protein [Arabidopsis<br>thaliana]  |

| ProbeSet                   | Description   |
|----------------------------|---|
| 16570_s_at (ATHCDPKA_S_AT) | 16570_s_at<br>(ATHCDPKA_S_AT)gb AAF27092.1 AC011809_1<br>(AC011809) calcium-dependent protein kinase 1<br>[Arabidopsis thaliana]      |
| 16620_s_at (AF051338_S_AT) | 16620_s_at (AF051338_S_AT)gb AAC05572.1 <br>(AF051338) xyloglucan endotransglycosylase related<br>protein [Arabidopsis thaliana]      |
| 16638_s_at (AF139098_S_AT) | 16638_s_at<br>(AF139098_S_AT)gb AAD37511.1 AF139098_1<br>(AF139098) putative zinc finger protein<br>[Arabidopsis thaliana]            |
| 16712_at (AC006068.67_AT)  | 16712_at (AC006068.67_AT)gb AAD15444.1 <br>(AC006068) putative glycogenin [Arabidopsis<br>thaliana]                                   |
| 16721_at (AC006533.58_AT)  | 16721_at<br>(AC006533.58_AT)gb AAD32285.1 AC006533_9<br>(AC006533) putative poly(ADP-ribose)<br>glycohydrolase [Arabidopsis thaliana] |
| 16753_at (AL031032.110_AT) | 16753_at (AL031032.110_AT)emb CAA19874.1 <br>(AL031032) putative protein [Arabidopsis thaliana]                                       |
| 16916_s_at (X77199.8_S_AT) | 16916_s_at (X77199.8_S_AT)emb CAA54420.1 <br>(X77199) heat shock cognate 70-2 [Arabidopsis<br>thaliana]                               |
| 16968_at (AL021961.93_AT)  | 16968_at (AL021961.93_AT)emb CAA17559.1 <br>(AL021961) glucosyltransferase -like protein<br>[Arabidopsis thaliana]                    |
| 17051_s_at (AF098947_S_AT) | 17051_s_at (AF098947_S_AT)gb AAD09952.1 <br>(AF098947) CTF2B [Arabidopsis thaliana]   |
| 17323_at (U95973.69_AT)    | 17323_at (U95973.69_AT)gb AAB65477.1 <br>(U95973) Ser/Thr protein kinase isolog [Arabidopsis<br>thaliana]                             |
| 17352_at (AC007127.33_AT)  | 17352_at<br>(AC007127.33_AT)gb AAD25140.1 AC007127_6<br>(AC007127) putative protein kinase [Arabidopsis<br>thaliana]                  |
| 17477_s_at (X63443.2_S_AT) | 17477_s_at (X63443.2_S_AT)emb CAA45039.1 <br>(X63443) heat shock protein 17.6-II [Arabidopsis<br>thaliana]                            |
| 17579_s_at (AF093753_S_AT) | 17579_s_at<br>(AF093753_S_AT)gb AAD50593.1 AF093753_1<br>(AF093753) phytochelatase synthase [Arabidopsis<br>thaliana]                 |

| ProbeSet                       | Description  |
|--------------------------------|--|
| 17775_at (AC004392.2_AT)       | 17775_at (AC004392.2_AT)gb AAC28500.1  (AC004392) Similar to glucose-6-phosphate/phosphate-translocator (GPT) gb AF020814 from Pisum sativum. [Arabidopsis thaliana]   |
| 17840_s_at (AC002333.223_S_AT) | 17840_s_at (AC002333.223_S_AT)gb AAB64049.1  (AC002333) putative endochitinase [Arabidopsis thaliana]  |
| 17886_at (AC004484.151_AT)     | 17886_at (AC004484.151_AT)gb AAC14530.1  (AC004484) unknown protein [Arabidopsis thaliana]   |
| 17894_at (AC005724.44_AT)      | 17894_at (AC005724.44_AT)gb AAD08938.1  (AC005724) unknown protein [Arabidopsis thaliana]  |
| 17907_s_at (AC004684.165_S_AT) | 17907_s_at (AC004684.165_S_AT)gb AAC23645.1  (AC004684) unknown protein [Arabidopsis thaliana]   |
| 17990_at (AF178075_AT)         | 17990_at (AF178075_AT)emb CAB41312.1  (AL049711) putative calmodulin [Arabidopsis thaliana]  |
| 18167_s_at (AL021711.23_S_AT)  | 18167_s_at (AL021711.23_S_AT)emb CAA16745.1  (AL021711) heat shock transcription factor-like protein [Arabidopsis thaliana]  |
| 18255_at (AC005770.25_AT)      | 18255_at (AC005770.25_AT)gb AAC79625.1  (AC005770) unknown protein [Arabidopsis thaliana]  |
| 18263_at (AC005724.36_AT)      | 18263_at (AC005724.36_AT)gb AAD08937.1  (AC005724) unknown protein [Arabidopsis thaliana]  |
| 18284_at (AL021961.67_AT)      | 18284_at (AL021961.67_AT)emb CAA17557.1  (AL021961) putative protein [Arabidopsis thaliana]  |
| 18544_at (AC007060.14_AT)      | 18544_at (AC007060.14_AT)gb AAD25749.1 AC007060_7 (AC007060) Strong similarity to F19I3.8 gi 3033381 putative UDP-galactose-4-epimerase from Arabidopsis thaliana BAC gb AC004238 and is a member of PF01370 the NAD dependent epimerase/dehydratase family. EST gb AA597338 comes from this gene. |
| 18600_at (L76926.1_AT)         | 18600_at (L76926.1_AT)gb AAC77829.1  (L76926) putative zinc finger protein [Arabidopsis thaliana]  |
| 18662_s_at (AC002343.20_S_AT)  | 18662_s_at (AC002343.20_S_AT)gb AAB63608.1  (AC002343) unknown protein [Arabidopsis thaliana]  |
| 18896_at (AC002329.51_AT)      | 18896_at (AC002329.51_AT)gb AAA67317.1  (L19262) 3-hydroxy-3-methylglutaryl-CoA reductase [Arabidopsis thaliana]   |

| ProbeSet                     | Description   |
|------------------------------|---|
| 19247_at (AF071527.44_AT)    | 19247_at (AF071527.44_AT)gb AAD11587.1 AAD11587 (AF071527) hypothetical protein [Arabidopsis thaliana]  |
| 19411_at (AC007661.104_AT)   | 19411_at (AC007661.104_AT)gb AAD32774.1 AC007661_11 (AC007661) unknown protein [Arabidopsis thaliana]   |
| 19451_at (AC004392.6_AT)     | 19451_at (AC004392.6_AT)gb AAC28502.1  (AC004392) Similar to F4I1.26 putative beta-glucosidase gi 3128187 from A. thaliana BAC gb AC004521. ESTs gb N97083, gb F19868 and gb F15482 come from this gene. [Arabidopsis thaliana] |
| 19462_s_at (AF001168.2_S_AT) | 19462_s_at (AF001168.2_S_AT)emb CAB75467.1  (AL138659) serine/threonine-specific kinase lecRK1 precursor, lectin  |
| 19640_at (AC004561.78_AT)    | 19640_at (AC004561.78_AT)gb AAC95192.1  (AC004561) putative glutathione S-transferase [Arabidopsis thaliana]  |
| 19916_at (AC006577.34_AT)    | 19916_at (AC006577.34_AT)gb AAD25781.1 AC006577_17 (AC006577) EST gb R64848 comes from this gene. [Arabidopsis thaliana]  |
| 19991_at (AC007017.124_AT)   | 19991_at (AC007017.124_AT)gb AAD21459.1  (AC007017) similar to harpin-induced protein hin1 from tobacco [Arabidopsis thaliana]  |
| 20051_at (AC000106.38_AT)    | 20051_at (AC000106.38_AT)gb AAB70412.1  (AC000106) Similar to Saccharomyces hypothetical protein YDR051c (gb Z49209). ESTs gb T44436,gb 42252 come from this gene. [Arabidopsis thaliana]                                       |
| 20142_at (AL035521.155_AT)   | 20142_at (AL035521.155_AT)emb CAB36716.1  (AL035521) extra-large G-protein-like [Arabidopsis thaliana]  |
| 20199_at (AL050300.89_AT)    | 20199_at (AL050300.89_AT)emb CAB43428.1  (AL050300) putative protein [Arabidopsis thaliana]   |
| 20269_at (AC002387.237_AT)   | 20269_at (AC002387.237_AT)gb AAB82640.1  (AC002387) putative pectinesterase [Arabidopsis thaliana]  |
| 20335_s_at (Y14208.2_S_AT)   | 20335_s_at (Y14208.2_S_AT)emb CAA74604.1  (Y14208) R2R3-MYB transcription factor [Arabidopsis thaliana]   |

| ProbeSet                      | Description   |
|-------------------------------|---|
| 20356_at (AC004561.74_AT)     | 20356_at (AC004561.74_AT)gb AAC95191.1 <br>(AC004561) putative glutathione S-transferase<br>[Arabidopsis thaliana]      |
| 20372_at (AL021713.24_AT)     | 20372_at (AL021713.24_AT)emb CAA16792.1 <br>(AL021713) putative protein [Arabidopsis thaliana]                          |
| 20421_at (U81294.2_AT)        | 20421_at (U81294.2_AT)emb CAB10242.1 <br>(Z97336) germin precursor oxalate oxidase<br>[Arabidopsis thaliana]            |
| 20590_at (AL035540.159_AT)    | 20590_at (AL035540.159_AT)emb CAB37511.1 <br>(AL035540) Phospholipase like protein [Arabidopsis<br>thaliana]            |
| 20619_at (AC005896.161_AT)    | 20619_at (AC005896.161_AT)gb AAC98070.1 <br>(AC005896) putative C2H2-type zinc finger protein<br>[Arabidopsis thaliana] |
| 20686_at (Y14424.2_AT)        | 20686_at (Y14424.2_AT)emb CAA74766.1 <br>(Y14424) hypothetical protein [Arabidopsis thaliana]                           |
| 20689_s_at (AC002335.19_S_AT) | 20689_s_at (AC002335.19_S_AT)gb AAB64310.1 <br>(AC002335) putative calcium binding protein<br>[Arabidopsis thaliana]    |

**Table 13b** Probe Sets corresponding to genes, the expression of which is decreased in an incompatible interaction at 3 and/or 6 hours after infection of four *Arabidopsis* ecotypes with *Pseudomonas syringae* pv tomato DC3000, *P. maculicola* ES4326 and *P. phaseolica* NPS3121

| ProbeSet                   | Description   |
|----------------------------|---|
| 12048_at (AF001308.46_AT)  | 12048_at (AF001308.46_AT)gb AAC78704.1 <br>(AF001308) predicted glycosyl transferase<br>[Arabidopsis thaliana]                  |
| 12218_at (AJ242588.2_AT)   | 12218_at (AJ242588.2_AT)emb CAB43344.1 <br>(AJ242588) 1-deoxy-d-xylulose-5-phosphate<br>reductoisomerase [Arabidopsis thaliana] |
| 12727_f_at (Z95799_F_AT)   | 12727_f_at<br>(Z95799_F_AT)gb AAD53097.1 AF175992_1<br>(AF175992) putative transcription factor<br>[Arabidopsis thaliana]       |
| 13144_at (AC007017.246_AT) | 13144_at (AC007017.246_AT)gb AAD21471.1 <br>(AC007017) unknown protein [Arabidopsis thaliana]                                   |

| ProbeSet                      | Description  |
|-------------------------------|--|
| 13161_at (AF002109.89_AT)     | 13161_at (AF002109.89_AT)gb AAB95278.1  (AF002109) putative isoamylase [Arabidopsis thaliana]                                    |
| 13220_s_at (CHS-EXON1_S_AT)   | 13220_s_at (CHS-EXON1_S_AT)gb AAA32771.1  (M20308) chalcone synthase [Arabidopsis thaliana]                                      |
| 13482_at (AC005896.195_AT)    | 13482_at (AC005896.195_AT)gb AAC98071.1  (AC005896) nodulin-like protein [Arabidopsis thaliana]                                  |
| 13547_s_at (AC004450.15_S_AT) | 13547_s_at (AC004450.15_S_AT)gb AAC64299.1  (AC004450) 3-isopropylmalate dehydratase, small subunit [Arabidopsis thaliana]       |
| 13637_at (AL049482.9_AT)      | 13637_at (AL049482.9_AT)emb CAB39634.1  (AL049482) AX110P-like protein [Arabidopsis thaliana]                                    |
| 13651_at (AL035538.320_AT)    | 13651_at (AL035538.320_AT)emb CAB37556.1  (AL035538) putative protein [Arabidopsis thaliana]                                     |
| 15832_at (AC004561.255_AT)    | 15832_at (AC004561.255_AT)gb AAC95217.1  (AC004561) unknown protein [Arabidopsis thaliana]                                       |
| 16110_s_at (AB004822_S_AT)    | 16110_s_at (AB004822_S_AT)dbj BAA22215.1  (AB004822) plastid RNA polymerase sigma-subunit [Arabidopsis thaliana]                 |
| 16229_at (AL049638.148_AT)    | 16229_at (AL049638.148_AT)emb CAB40944.1  (AL049638) putative transport protein [Arabidopsis thaliana]                           |
| 16351_at (AL021684.194_AT)    | 16351_at (AL021684.194_AT)emb CAA16688.1  (AL021684) receptor protein kinase - like protein [Arabidopsis thaliana]               |
| 16468_at (AF049870.5_AT)      | 16468_at (AF049870.5_AT)gb AAD02499.1  (AF049870) thaumatin-like protein [Arabidopsis thaliana]                                  |
| 16548_s_at (AF054617_S_AT)    | 16548_s_at (AF054617_S_AT)gb AAC25108.1  (AF054617) one helix protein [Arabidopsis thaliana]                                     |
| 16978_g_at (AF076641.2_G_AT)  | 16978_g_at (AF076641.2_G_AT)gb AAD46064.1 AF076641_1 (AF076641) homeodomain leucine-zipper protein ATHB16 [Arabidopsis thaliana] |
| 17018_s_at (ATU18929_S_AT)    | 17018_s_at (ATU18929_S_AT)gb AAA79982.1  (U18929) cytochrome p450 dependent monooxygenase [Arabidopsis thaliana]                 |
| 17494_s_at (ATU30478_S_AT)    | 17494_s_at (ATU30478_S_AT)gb AAB38071.1  (U30478) expansin At-EXP5 [Arabidopsis thaliana]  |

| ProbeSet                      | Description  |
|-------------------------------|--|
| 17823_s_at (AC006555.10_S_AT) | 17823_s_at<br>(AC006555.10_S_AT)gb AAD26909.1 AC007233_1 (AC007233) putative beta-1,3-glucanase [Arabidopsis thaliana]   |
| 18215_at (Z97335.114_AT)      | 18215_at (Z97335.114_AT)emb CAB46000.1  (Z97335) selenium-binding protein like [Arabidopsis thaliana]  |
| 18301_s_at (AL022223.48_S_AT) | 18301_s_at<br>(AL022223.48_S_AT)emb CAA18218.1  (AL022223) fructose-bisphosphate aldolase [Arabidopsis thaliana]   |
| 18471_at (AC006533.103_AT)    | 18471_at<br>(AC006533.103_AT)gb AAD32293.1 AC006533_17 (AC006533) putative glucosyltransferase [Arabidopsis thaliana]  |
| 18634_s_at (Z97343.468_S_AT)  | 18634_s_at (Z97343.468_S_AT)emb CAB10536.1  (Z97343) hypothetical protein [Arabidopsis thaliana]   |
| 18976_at (AC000106.31_AT)     | 18976_at (AC000106.31_AT)gb AAB70409.1  (AC000106) Contains similarity to Rhodococcus amidase (gb D16207). ESTs gb T20504,gb H36650,gb N97423,gb H36595 come from this gene. [Arabidopsis thaliana]  |
| 18984_at (AC003096.100_AT)    | 18984_at (AC003096.100_AT)gb AAC16266.1  (AC003096) unknown protein [Arabidopsis thaliana]   |
| 19494_at (AC007296.26_AT)     | 19494_at<br>(AC007296.26_AT)gb AAD30251.1 AC007296_12 (AC007296) Strong similarity to gb X95759 soluble-starch-synthase precursor (SSIII) from Solanum tuberosum. [Arabidopsis thaliana]   |
| 20442_i_at (AC006341.42_I_AT) | 20442_i_at<br>(AC006341.42_I_AT)gb AAD34693.1 AC006341_21 (AC006341) Similar to gb AF069494 cytochrome P450 from Sinapis alba and is a member of the PF 00067 Cytochrome P450 family. EST gb F14190 comes from this gene. [Arabidopsis thaliana] |

**Table 14** Probe Sets as referred to in Table 3 corresponding to genes with promoters conferring low basal expression in all ecotypes (Col, Ler, Ws, and Cvi) and high inducibility in Col

| ProbeSet                       | Description  |
|--------------------------------|--------------|
| 20421_at (U81294.2_AT)         | see Table 3  |
| 20142_at (AL035521.155_AT)     | see Table 3  |
| 14635_s_at (PR.1_S_AT)         | see Table 3  |
| 12908_s_at (ATERF5_S_AT)       | see Table 11 |
| 20620_g_at (AC005896.161_G_AT) | see Table 11 |



**Table 15** Probe Sets as referred to in Table 3 corresponding to genes with promoters inducing expression in *Pseudomonas syringae* pv. *maculiola*-infected *Arabidopsis*

|   |                                |
|---|--------------------------------|
| 12891_at (ATACS6_AT)                      | 13015_s_at (X98673.2_S_AT)     |
| 13100_at (AC003680.50_AT)                 | 13115_at (AC000375.44_AT)      |
| 13217_s_at (CALMODULINLIKE_S_AT)          | 13467_at (AL096860.198_AT)     |
| 13645_at (AC000098.8_AT)                  | 13818_s_at (AC006218.175_S_AT) |
| 14032_at (AL035601.11_AT)                 | 14248_at (PAD3_AT)             |
| 14609_at (AC002340.147_AT)                | 15116_f_at (AF121356_F_AT)     |
| 15622_s_at (ATU43945_S_AT)                | 16173_s_at (D78607_S_AT)       |
| 17485_s_at (Z97340.345_S_AT)              | 17511_s_at (AF067605_S_AT)     |
| 17548_s_at (AF118823_S_AT)                | 17775_at (AC004392.2_AT)       |
| 17930_s_at (AJ006960.4_S_AT)              | 19284_at (AC003028.196_AT)     |
| 19546_at (AC005398.172_AT)                | 19640_at (AC004561.78_AT)      |
| 20134_s_at (AC007178.71_S_AT)             | 20194_at (AC007584.48_AT)      |
| 20348_at (AC005967.35_AT)                 | 12892_g_at (ATACS6_G_AT)       |
| 12904_s_at (ATERF1_S_AT)                  | 12989_s_at (AC004077.149_S_AT) |
| 13215_s_at (cafferoylcoAmethyltrans_S_AT) | 13565_at (AL035601.21_AT)      |
| 13627_at (AL035394.196_AT)                | 16649_s_at (ATHORF_S_AT)       |
| 16914_s_at (AL049500.57_S_AT)             | 16995_at (AC002391.188_AT)     |
| 19991_at (AC007017.124_AT)                | 20356_at (AC004561.74_AT)      |

**Table 16** Probe Sets corresponding to genes with promoters inducing expression in *Botrytis cinerea*-infected *Arabidopsis*

| Probe Set                                    | Description  |
|--|--|
| 12115_at (AL033545.26_AT)                    | emb CAA22152.1  (AL033545) extensin-like protein [Arabidopsis thaliana]  |
| 12574_at (X82624.2_AT)                       | emb CAA57944.1  (X82624) SRG2At [Arabidopsis thaliana]   |
| 12989_s_at (AC004077.149_S_AT)               | gb AAC26690.1  (AC004077) putative cytochrome P450 [Arabidopsis thaliana]  |
| 13015_s_at (X98673.2_S_AT)                   | emb CAA67232.1  (X98674) zinc finger protein [Arabidopsis thaliana]  |
| 13100_at (AC003680.50_AT)                    | gb AAC06158.1  (AC003680) putative cytochrome P450 [Arabidopsis thaliana]  |
| 13215_s_at<br>(cafferoylcoAmethyltrans_S_AT) | gb AAF16576.1 AC012563_29 (AC012563) putative S-adenosyl-L-methionine:trans-caffeoyl-Coenzyme A 3-O-methyltransferase [Arabidopsis thaliana] |
| 13565_at (AL035601.21_AT)                    | emb CAB38206.1  (AL035601) auxin-responsive GH3-like protein [Arabidopsis thaliana]  |
| 14015_s_at (A71588.1_S_AT)                   | emb CAB42612.1  (A71639) unnamed protein product [Arabidopsis thaliana]  |
| 14016_s_at (A71596.1_S_AT)                   | emb CAB42592.1  (A71596) unnamed protein product [Arabidopsis thaliana]  |
| 14609_at (AC002340.147_AT)                   | gb AAC02748.1  (AC002340) putative cytochrome P450 [Arabidopsis thaliana]  |
| 16638_s_at (AF139098_S_AT)                   | gb AAD37511.1 AF139098_1 (AF139098) putative zinc finger protein [Arabidopsis thaliana]  |
| 16649_s_at (ATHORF_S_AT)                     | gb AAF16576.1 AC012563_29 (AC012563) putative S-adenosyl-L-methionine:trans-caffeoyl-Coenzyme A 3-O-methyltransferase [Arabidopsis thaliana] |
| 16892_at (U37336.3_AT)                       | gb AAC49135.1  (U37336) senescence-specific cysteine protease  |
| 16914_s_at (AL049500.57_S_AT)                | emb CAB39936.1  (AL049500) osmotin precursor [Arabidopsis thaliana]  |
| 17533_s_at (ATU43488_S_AT)                   | gb AAB18367.1  (U43488) xyloglucan endotransglycosylase-related protein [Arabidopsis thaliana]   |
| 17744_s_at (AC004684.168_S_AT)               | gb AAC23646.1  (AC004684) putative alcohol dehydrogenase [Arabidopsis thaliana]  |

| Probe Set                  | Description   |
|----------------------------|---|
| 18716_at (X91916_AT)       |   |
| 18888_at (AC007591.68_AT)  | gb AAD39666.1 AC007591_31 (AC007591) Is a member of the PF 00903 gyloxalase family. ESTs gb T44721, gb T21844 and gb AA395404 come from this gene. [Arabidopsis thaliana] |
| 19019_i_at (X82623.2_I_AT) | emb CAA57943.1  (X82623) SRG2At [Arabidopsis thaliana]  |
| 19284_at (AC003028.196_AT) | gb AAC27173.1  (AC003028) putative anthocyanidin synthase [Arabidopsis thaliana]  |
| 19762_at (AL035527.204_AT) | emb CAB36812.1  (AL035527) peptide transporter-like protein [Arabidopsis thaliana]  |
| 19991_at (AC007017.124_AT) | gb AAD21459.1  (AC007017) similar to harpin-induced protein hin1 from tobacco [Arabidopsis thaliana]  |
| 20356_at (AC004561.74_AT)  | gb AAC95191.1  (AC004561) putative glutathione S-transferase [Arabidopsis thaliana]   |

**Table 17** Probe sets which correspond to genes downregulated in response to infection by 5 different viruses

| Probe Set                         | Description  | Function            | Blast   |
|-----------------------------------|--|---------------------|---------|
| 17191_i_at<br>(AL021961.178_I_AT) | emb CAA17564.1  (AL021961)<br>putative protein [Arabidopsis thaliana]  | DNA binding protein | MyB TF  |
| 18866_at<br>(AC005917.178_AT)     | gb AAD10163.1  (AC005917)<br>putative Ta11-like non-LTR<br>retroelement protein<br>[Arabidopsis thaliana]              | transposon          |         |
| 20678_at<br>(AC007296.30_AT)      | gb AAD30253.1 AC007296_14<br>(AC007296) ESTs gb R65381<br>and gb T44635 come from this<br>gene. [Arabidopsis thaliana] | unknown             | unknown |
| 12356_at<br>(X99952.1_AT)         | emb CAA68212.1  (X99952)<br>peroxidase [Arabidopsis thaliana]  | metabolic protein   |         |
| 12448_at<br>(AC002337.58_AT)      | gb AAB63824.1  (AC002337)<br>putative acyl-CoA synthetase<br>[Arabidopsis thaliana]                                    | metabolic protein   |         |
| 12854_s_at<br>(ACS1_S_AT)         | gb AAA96006.1  (U26542) 1-<br>aminocyclopropane-1-<br>carboxylate synthase-like protein<br>[Arabidopsis thaliana]      | hormone response    |         |
| 13812_s_at<br>(AC005275.104_S_AT) | gb AAD14468.1  (AC005275)<br>putative GH3-like protein<br>[Arabidopsis thaliana]                                       | hormone response    |         |
| 14530_at<br>(AL021889.231_AT)     | emb CAA17145.1  (AL021889)<br>putative protein [Arabidopsis thaliana]  | unknown             | unknown |
| 16461_i_at<br>(AC004683.79_I_AT)  | gb AAC28766.1  (AC004683)<br>peroxidase [Arabidopsis thaliana]   | metabolic protein   |         |
| 16462_s_at<br>(AC004683.79_S_AT)  | gb AAC28766.1  (AC004683)<br>peroxidase [Arabidopsis thaliana]   | metabolic protein   |         |
| 17305_at<br>(U25649.3_AT)         | gb AAC50023.1  (U25649)<br>ATPME2 precursor [Arabidopsis thaliana]   | cell wall polymer   |         |
| 14965_at<br>(AC002329.22_AT)      | gb AAB86507.1  (AC002329)<br>unknown protein [Arabidopsis thaliana]  | unknown             | unknown |

| Probe Set                        | Description  | Function              | Blast   |
|----------------------------------|--|-----------------------|---|
| 15578_s_at<br>(AF004213_S_AT)    | gb AAC49746.1  (AF004213)<br>ethylene-insensitive3-like1<br>[Arabidopsis thaliana]   | hormone<br>response   |   |
| 16062_s_at<br>(AB007789_S_AT)    | gb AAD15976.1  (AF074601)<br>CRT/DRE binding factor 2<br>[Arabidopsis thaliana]  | stress response       |   |
| 16111_f_at<br>(AB007788_F_AT)    | dbj BAA33435.1  (AB013816)<br>DREB1B [Arabidopsis thaliana]<br>[Arabidopsis thaliana]  | stress response       |   |
| 16434_at<br>(AL021711.157_AT)    | emb CAA16754.1  (AL021711)<br>putative protein [Arabidopsis<br>thaliana]   | metabolic<br>protein  | similar to APG<br>(non proline-rich<br>region)<br>[Arabidopsis<br>thaliana]<br>/putative GDSL-<br>motif<br>lipase/hydrolase |
| 16891_s_at<br>(AF080120.33_S_AT) | gb AAC35539.1  (AF080120)<br>contains similarity to glycosyl<br>hydrolases family 9 (Pfam:<br>glycosyl_hydro5.hmm, score:<br>88.03) [Arabidopsis thaliana] | cell wall<br>polymer  |   |
| 16990_at<br>(AC004684.91_AT)     | gb AAC23634.1  (AC004684)<br>putative expansin [Arabidopsis<br>thaliana]   | cell wall<br>polymer  |   |
| 17577_g_at<br>(AF087820_G_AT)    | gb AAD52697.1 AF087820_1<br>(AF087820) auxin transport<br>protein [Arabidopsis thaliana]   | hormone<br>response   |   |
| 17743_at<br>(AC002341.99_AT)     | gb AAB67624.1  (AC002341)<br>putative peroxidase [Arabidopsis<br>thaliana]   | metabolic<br>protein  |   |
| 19017_at<br>(AL035709.69_AT)     | emb CAB38928.1  (AL035709)<br>endo-xyloglucan transferase-like<br>protein [Arabidopsis thaliana]   | cell wall<br>polymer  |   |
| 19396_at<br>(AJ001855.2_AT)      | emb CAA05054.1  (AJ001855)<br>alpha subunit of F-actin capping<br>protein [Arabidopsis thaliana]   | structural<br>protein |   |
| 19660_at<br>(AC002336.29_AT)     | gb AAB87577.1  (AC002336)<br>putative expansin [Arabidopsis<br>thaliana]   | cell wall<br>polymer  |   |

| Probe Set                         | Description   | Function               | Blast                              |
|-----------------------------------|---|------------------------|------------------------------------|
| 20675_at<br>(AC006234.204_AT)     | gb AAD20920.1  (AC006234)<br>beta-expansin [Arabidopsis<br>thaliana]                              | cell wall<br>polymer   |                                    |
| 12086_s_at<br>(AC002409.88_S_AT)  | gb AAB86456.1  (AC002409)<br>unknown protein [Arabidopsis<br>thaliana]                            | unknown                |                                    |
| 13728_at<br>(NOVARTIS111_AT)      | gb AAF27057.1 AC008262_6<br>(AC008262) F4N2.12<br>[Arabidopsis thaliana]                          | unknown                | unknown                            |
| 14770_s_at<br>(AC002338.167_S_AT) | gb AAB63092.1  (U93215)<br>putative MYB family<br>transcription factor [Arabidopsis<br>thaliana]  | DNA binding<br>protein |                                    |
| 15067_at<br>(AC004683.36_AT)      | gb AAC28758.1  (AC004683)<br>unknown protein [Arabidopsis<br>thaliana]                            | membrane<br>protein    | transporter or<br>ferroportin      |
| 15154_s_at<br>(ATHMTGDAS_S_AT)    | emb CAB51206.1  (AL096860)<br>glutamine-dependent asparagine<br>synthetase [Arabidopsis thaliana] | metabolic<br>protein   |                                    |
| 15544_at<br>(AL021633.110_AT)     | emb CAA16532.1  (AL021633)<br>predicted protein [Arabidopsis<br>thaliana]                         | membrane<br>protein    | E. coli cation<br>transporter ChaC |
| 15631_s_at<br>(AB005805_S_AT)     | dbj BAA28625.1  (AB005805)<br>aldehyde oxidase [Arabidopsis<br>thaliana]                          | metabolic<br>protein   |                                    |
| 16048_at<br>(X78586.2_AT)         | emb CAA55323.1  (X78586)<br>Dr4 [Arabidopsis thaliana]  | stress response        |                                    |
| 16090_s_at<br>(ATHFAD8A_S_AT)     | gb AAB60302.1  (U08216)<br>chloroplast linoleate desaturase<br>[Arabidopsis thaliana]             | metabolic<br>protein   |                                    |
| 16164_s_at<br>(ATU47029_S_AT)     | gb AAC49302.1  (U47029)<br>ERECTA [Arabidopsis thaliana]  | receptor/kinase        |                                    |
| 16241_at<br>(AL022604.84_AT)      | emb CAA18733.1  (AL022604)<br>putative protein [Arabidopsis<br>thaliana]                          | unknown                | unknown                            |
| 16327_at<br>(AC002334.12_AT)      | gb AAF18589.1 AC002332_1<br>(AC002332) putative myosin<br>heavy chain [Arabidopsis<br>thaliana]   | structural<br>protein  |                                    |

| Probe Set                        | Description  | Function             | Blast   |
|----------------------------------|--|----------------------|---|
| 16406_at<br>(AC006921.33_AT)     | gb AAD21431.1  (AC006921)<br>putative protein kinase<br>[Arabidopsis thaliana]               | receptor/kinase      |   |
| 16868_at<br>(AL035679.123_AT)    | emb CAB38821.1  (AL035679)<br>putative endo-1, 4-beta-<br>glucanase [Arabidopsis thaliana]   | cell wall<br>polymer |   |
| 17362_s_at<br>(Z97338.181_S_AT)  | emb CAB10305.1  (Z97338)<br>glucosyltransferase [Arabidopsis<br>thaliana]                    | metabolic<br>protein |   |
| 17916_at<br>(U22428.2_AT)        | gb AAB03100.1  (U22428)<br>starch branching enzyme class II<br>[Arabidopsis thaliana]        | metabolic<br>protein |   |
| 18515_at<br>(AC007063.215_AT)    | gb AAF24822.1 AC007592_15<br>(AC007592) F12K11.17<br>[Arabidopsis thaliana]                  | metabolic<br>protein | putative 1,3-<br>beta-D-glucan<br>synthase<br>[Arabidopsis<br>thaliana] |
| 18635_at<br>(AC004005.44_AT)     | gb AAC23401.1  (AC004005)<br>unknown protein [Arabidopsis<br>thaliana]                       | metabolic<br>protein | methyl Cl<br>transferase  |
| 18667_at<br>(AJ249442_AT)        | emb CAB55758.1  (AJ249442)<br>putative AUX1-like permease<br>[Arabidopsis thaliana]          | hormone<br>response  |   |
| 18683_s_at<br>(L27158_S_AT)      | gb AAB60302.1  (U08216)<br>chloroplast linoleate desaturase<br>[Arabidopsis thaliana]        | metabolic<br>protein |   |
| 18694_s_at<br>(U89272_S_AT)      | gb AAB61458.1  (U89272)<br>chloroplast membrane protein<br>ALBINO3 [Arabidopsis<br>thaliana] | membrane<br>protein  |   |
| 19870_s_at<br>(AL021811.48_S_AT) | emb CAA16958.1  (AL021811)<br>putative protein [Arabidopsis<br>thaliana]                     | unknown              | unknown   |

**Table 18** Probe sets which correspond to genes upregulated in response to infection by 5 different viruses

| Probeset                          | Description  | Time   | Function                  | Blast   |
|-----------------------------------|--|--------|---------------------------|---|
| 12904_s_at<br>(ATERF1_S_AT)       | dbj BAA32418.1  (AB008103)<br>ethylene responsive element<br>binding factor 1 [Arabidopsis<br>thaliana]                    | 5dpi   | hormone<br>response       |   |
| 16063_s_at<br>(AB008103_S_AT)     | dbj BAA32418.1  (AB008103)<br>ethylene responsive element<br>binding factor 1 [Arabidopsis<br>thaliana]                    | 5dpi   | hormone<br>response       |   |
| 12906_s_at<br>(ATERF3_S_AT)       | dbj BAA32420.1  (AB008105)<br>ethylene responsive element<br>binding factor 3 [Arabidopsis<br>thaliana]                    | 1dpi   | stress<br>response        |   |
| 16537_s_at<br>(AB008111_S_AT)     | dbj BAA28953.1  (AB008111)<br>Atrboh F [Arabidopsis thaliana]  | 4dpi   | stress<br>response        |   |
| 16610_s_at<br>(AB008490_S_AT)     | dbj BAA34729.1  (AB008490)<br>response regulator 7<br>[Arabidopsis thaliana]   | 1dpi   | signaling not<br>kinase   |   |
| 13645_at<br>(AC000098.8_AT)       | gb AAB71443.1  (AC000098)<br>EST gb ATTS0295 comes from<br>this gene. [Arabidopsis thaliana]                               | 4-5dpi | unknown                   | unknown   |
| 19388_at<br>(AC000104.61_AT)      | gb AAB70450.1  (AC000104)<br>ESTs gb N65789,gb T04628<br>come from this gene.<br>[Arabidopsis thaliana]                    | 4dpi   | kinase/recep<br>tor       | prenylated<br>Rab receptor                                      |
| 19368_at<br>(AC000348.22_AT)      | gb AAB61497.1  (AC000348)<br>T7N9.21 [Arabidopsis thaliana]  | 5dpi   | unknown                   | unknown   |
| 13115_at<br>(AC000375.44_AT)      | gb AAB60774.1  (AC000375)<br>ESTs<br>gb U75592,gb T13956,gb T4386<br>9 come from from this gene.<br>[Arabidopsis thaliana] | 4dpi   | DNA<br>binding<br>protein | putative<br>WRKY DNA<br>binding<br>protein<br>[Oryza<br>sativa] |
| 14964_at<br>(AC001229.8_AT)       | gb AAB60905.1  (AC001229)<br>F5I14.4 gene product<br>[Arabidopsis thaliana]  | 5dpi   | unknown                   | unknown   |
| 17840_s_at<br>(AC002333.223_S_AT) | gb AAB64049.1  (AC002333)<br>putative endochitinase<br>[Arabidopsis thaliana]  | 5dpi   | stress<br>response        |   |



| Probeset                          | Description  | Time   | Function                | Blast   |
|-----------------------------------|--|--------|-------------------------|---------|
| 15379_at<br>(AC002335.182_AT)     | gb AAB64323.1  (AC002335)<br>unknown protein [Arabidopsis<br>thaliana]                               | 1dpi   | unknown                 | unknown |
| 20689_s_at<br>(AC002335.19_S_AT)  | gb AAB64310.1  (AC002335)<br>putative calcium binding protein<br>[Arabidopsis thaliana]              | 5dpi   | stress<br>response      |         |
| 12449_s_at<br>(AC002343.179_S_AT) | gb AAB63623.1  (AC002343)<br>cellulose synthase isolog<br>[Arabidopsis thaliana]                     | 4-5dpi | cell wall<br>polymer    |         |
| 13842_at<br>(AC002396.12_AT)      | gb AAC00572.1  (AC002396)<br>similar to zinc<br>metalloproteinases [Arabidopsis<br>thaliana]         | 5dpi   | protein<br>processing   |         |
| 19424_at<br>(AC002396.44_AT)      | gb AAC00588.1  (AC002396)<br>glucose-6-phosphate 1-<br>dehydrogenase [Arabidopsis<br>thaliana]       | 5dpi   | metabolic<br>protein    |         |
| 13100_at<br>(AC003680.50_AT)      | gb AAC06158.1  (AC003680)<br>putative cytochrome P450<br>[Arabidopsis thaliana]                      | 5dpi   | metabolic<br>protein    |         |
| 16365_at<br>(AC003974.136_AT)     | gb AAC04495.1  (AC003974)<br>putative disease resistance<br>protein [Arabidopsis thaliana]           | 2-5dpi | stress<br>response      |         |
| 17752_at<br>(AC003974.37_AT)      | gb AAC04483.1  (AC003974)<br>putative protein kinase<br>[Arabidopsis thaliana]                       | 4dpi   | kinase/recep<br>tor     |         |
| 20409_g_at<br>(AC004077.132_G_AT) | emb CAB40383.1  (AJ131180)<br>ribosomal protein S14<br>[Arabidopsis thaliana]                        | 5dpi   | protein<br>processing   |         |
| 13048_s_at<br>(AC004138.22_S_AT)  | gb AAC32906.1  (AC004138)<br>putative basic blue protein<br>(plantacyanin) [Arabidopsis<br>thaliana] | 4dpi   | stress<br>response      |         |
| 12764_f_at<br>(AC004138.69_F_AT)  | gb AAC32912.1  (AC004138)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]        | 5dpi   | stress<br>response      |         |
| 13895_at<br>(AC004218.63_AT)      | gb AAC27833.1  (AC004218)<br>putative phospholipase<br>[Arabidopsis thaliana]                        | 4dpi   | signaling not<br>kinase |         |

| Probeset                          | Description  | Time | Function                | Blast   |
|-----------------------------------|--|------|-------------------------|---------|
| 20477_at<br>(AC004238.154_AT)     | gb AAC12841.1  (AC004238)<br>putative UDP-N-<br>acetylglucosamine<br>pyrophosphorylase [Arabidopsis<br>thaliana]                                     | 5dpi | metabolic<br>protein    |         |
| 16972_at<br>(AC004261.89_AT)      | gb AAA87347.1  (M88307)<br>calmodulin [Brassica juncea]  | 5dpi | signaling not<br>kinase |         |
| 19848_s_at<br>(AC004261.94_S_AT)  | dbj BAA08282.1  (D45848)<br>calmodulin-related protein<br>[Arabidopsis thaliana]   | 5dpi | signaling not<br>kinase |         |
| 19044_at<br>(AC004392.38_AT)      | gb AAC28514.1  (AC004392)<br>Strong similarity to gi 2160138<br>F19K23.6 gene product from A.<br>thaliana BAC gb AC000375.<br>[Arabidopsis thaliana] | 5dpi | unknown                 | unknown |
| 19640_at<br>(AC004561.78_AT)      | gb AAC95192.1  (AC004561)<br>putative glutathione S-<br>transferase [Arabidopsis<br>thaliana]  | 5dpi | stress<br>response      |         |
| 20491_at<br>(AC004561.146_AT)     | gb AAC95203.1  (AC004561)<br>putative tropinone reductase<br>[Arabidopsis thaliana]  | 5dpi | metabolic<br>protein    |         |
| 16888_s_at<br>(AC004684.174_S_AT) | gb AAC23647.1  (AC004684)<br>putative alcohol dehydrogenase<br>[Arabidopsis thaliana]  | 4dpi | stress<br>response      |         |
| 15389_at<br>(AC004786.100_AT)     | gb AAC32433.1  (AC004786)<br>unknown protein [Arabidopsis<br>thaliana]   | 4dpi | unknown                 | unknown |
| 18844_at<br>(AC005315.131_AT)     | gb AAC33239.1  (AC005315)<br>putative ligand-gated ion<br>channel protein [Arabidopsis<br>thaliana]  | 4dpi | signaling not<br>kinase |         |
| 17128_s_at<br>(ATHRPRP1A_S_AT)    | gb AAC69381.1  (AC005398)<br>pathogenesis-related PR-1-like<br>protein [Arabidopsis]   | 5dpi | stress<br>response      |         |
| 17894_at<br>(AC005724.44_AT)      | gb AAD08938.1  (AC005724)<br>unknown protein [Arabidopsis<br>thaliana]   | 5dpi | unknown                 | unknown |
| 18263_at<br>(AC005724.36_AT)      | gb AAD08937.1  (AC005724)<br>unknown protein [Arabidopsis<br>thaliana]   | 5dpi | unknown                 | unknown |

| Probeset                        | Description   | Time         | Function             | Blast   |
|---------------------------------|---|--------------|----------------------|---|
| 15539_at<br>(AC005770.21_AT)    | gb AAC79601.1  (AC005770)<br>unknown protein [Arabidopsis<br>thaliana]  | 2dpi<br>4dpi | unknown              | unknown   |
| 13920_at<br>(AC005990.53_AT)    | gb AAC98026.1  (AC005990)<br>EST gb AA597511 comes from<br>this gene. [Arabidopsis thaliana]                                | 5dpi         | unknown              | unknown   |
| 18267_at<br>(AC006223.23_AT)    | gb AAD15384.1  (AC006223)<br>unknown protein [Arabidopsis<br>thaliana]  | 4dpi         | unknown              | unknown   |
| 12627_at<br>(AC006533.99_AT)    | gb AAD32292.1 AC006533_16<br>(AC006533) putative protein<br>kinase [Arabidopsis thaliana]                                   | 1dpi         | kinase/recep<br>tor  |   |
| 12497_at<br>(AC006533.51_AT)    | gb AAD32284.1 AC006533_8<br>(AC006533) putative receptor-<br>like protein kinase [Arabidopsis<br>thaliana]                  | 5dpi         | kinase/recep<br>tor  |   |
| 12656_at<br>(AC006569.43_AT)    | gb AAD21751.1  (AC006569)<br>unknown protein [Arabidopsis<br>thaliana]  | 5dpi         | unknown              | unknown   |
| 17008_at<br>(AC006585.212_AT)   | gb AAD23027.1 AC006585_22<br>(AC006585) putative tyrosine<br>aminotransferase [Arabidopsis<br>thaliana]                     | 4-5dpi       | metabolic<br>protein |   |
| 19386_at<br>(AC006592.51_AT)    | No hits found less than or equal<br>to 1e-15.   | 2dpi         | metabolic<br>protein |   |
| 13617_at<br>(AC006592.64_AT)    | gb AAD22351.1 AC006592_8<br>(AC006592) putative<br>mitochondrial dicarboxylate<br>carrier protein [Arabidopsis<br>thaliana] | 5dpi         | unknown              | unknown   |
| 15473_at<br>(AC006836.125_AT)   | gb AAD20072.1  (AC006836)<br>putative ubiquinone biosynthesis<br>protein [Arabidopsis thaliana]                             | 5dpi         | metabolic<br>protein |   |
| 12642_at<br>(AC006920.138_AT)   | gb AAD22285.1 AC006920_9<br>(AC006920) unknown protein<br>[Arabidopsis thaliana]  | 5dpi         | metabolic<br>protein | putative<br>xyloglucan<br>fucosyltransf<br>erase<br>[Arabidopsis<br>thaliana] |
| 15921_s_at<br>(AC007067.1_S_AT) | gb AAD39561.1 AC007067_1<br>(AC007067) T10O24.1<br>[Arabidopsis thaliana]   | 4dpi         | unknown              | unknown   |

| Probeset                     | Description  | Time   | Function          | Blast   |
|------------------------------|--|--------|-------------------|---------|
| 14248_at (PAD3_AT)           | gb AAD31062.1 AC007357_11 (AC007357) Strong similarity to gb X97864 cytochrome P450 from <i>Arabidopsis thaliana</i> and is a member of the PF00067 Cytochrome P450 family. ESTs gb N65665, gb T14112, gb T76255, gb T20906 and gb AI100027 come from this gene. | 5dpi   | metabolic protein |         |
| 19903_at (AC007660.40_AT)    | gb AAD32803.1 AC007660_4 (AC007660) unknown protein [ <i>Arabidopsis thaliana</i> ]  | 4dpi   | unknown           | unknown |
| 15487_at (AC007661.87_AT)    | gb AAD32771.1 AC007661_8 (AC007661) unknown protein [ <i>Arabidopsis thaliana</i> ]  | 5dpi   | unknown           | unknown |
| 13751_at (NOVARTIS127_AT)    | gb AAF16751.1 AC010155_4 (AC010155) F3M18.8 [ <i>Arabidopsis thaliana</i> ]  | 4dpi   | unknown           | unknown |
| 13246_at (ERECTAL_AT)        | gb AAF26067.1 AC012562_22 (AC012562) putative protein kinase [ <i>Arabidopsis thaliana</i> ]   | 4dpi   | kinase/receptor   |         |
| 14636_s_at (PR5_S_AT)        | gb AAF21072.1 AC013258_10 (AC013258) thaumatin-like protein [ <i>Arabidopsis thaliana</i> ]  | 2-5dpi | stress response   |         |
| 16153_s_at (ATHRPRP1C_S_AT)  | gb AAF21072.1 AC013258_10 (AC013258) thaumatin-like protein [ <i>Arabidopsis thaliana</i> ]  | 2-5dpi | stress response   |         |
| 16617_s_at (AF029980_S_AT)   | gb AAD01897.1  (AF029980) A37 [ <i>Arabidopsis thaliana</i> ]  | 5dpi   | unknown           | unknown |
| 12538_at (AF033205.2_AT)     | gb AAC02973.1  (AF033205) putative pectin methylesterase [ <i>Arabidopsis thaliana</i> ]   | 4dpi   | cell wall polymer |         |
| 18681_at (L23573_AT)         | gb AAD02548.1  (AF049922) PGP169-12 [ <i>Petunia x hybrida</i> ]   | 4dpi   | membrane protein  |         |
| 19181_s_at (AF053065.2_S_AT) | gb AAC39464.1  (AF053065) late embryogenesis abundant protein homolog [ <i>Arabidopsis thaliana</i> ]  | 5dpi   | stress response   |         |

| Probeset                         | Description  | Time   | Function              | Blast  |
|----------------------------------|--|--------|-----------------------|--|
| 19555_at<br>(AF058919.48_AT)     | gb AAC13630.1  (AF058919)<br>F6N23.8 gene product<br>[Arabidopsis thaliana]  | 4dpi   | metabolic<br>protein  | probable<br>methylene-<br>tetrahydrofol<br>ate<br>dehydrogenas<br>e (NADP+)            |
| 15515_r_at<br>(AF058919.32_R_AT) | gb AAC13627.1  (AF058919)<br>F6N23.26 gene product<br>[Arabidopsis thaliana]   | 5dpi   | metabolic<br>protein  | putative<br>phosphoribos<br>ylanthranilate<br>transferase<br>[Arabidopsis<br>thaliana] |
| 15606_s_at<br>(AF061517_S_AT)    | gb AAC15807.1  (AF061517)<br>putative copper/zinc superoxide<br>dismutase copper chaperone<br>[Arabidopsis thaliana]   | 5dpi   | stress<br>response    |  |
| 18604_at<br>(AF069298.31_AT)     | gb AAC19273.1  (AF069298)<br>similar to several small proteins<br>(~100 aa) that are induced by<br>heat, auxin, ethylene and<br>wounding such as Phaseolus<br>aureus indole-3-acetic acid<br>induced protein ARG<br>(SW:32292) [Arabidopsis<br>thaliana] | 5dpi   | stress<br>response    |  |
| 12002_at<br>(AF069442.47_AT)     | gb AAC79112.1  (AF069442)<br>hypothetical protein<br>[Arabidopsis thaliana]  | 4-5dpi | protein<br>processing | ubiquitin-like   |
| 20480_s_at<br>(AF069495.2_S_AT)  | emb CAB38908.1  (AL035708)<br>cytochrome P450-like protein<br>[Arabidopsis thaliana]   | 4-5dpi | metabolic<br>protein  |  |
| 20479_i_at<br>(AF069495.2_I_AT)  | emb CAB38908.1  (AL035708)<br>cytochrome P450-like protein<br>[Arabidopsis thaliana]   | 5dpi   | metabolic<br>protein  |  |
| 19247_at<br>(AF071527.44_AT)     | gb AAD11587.1  AAD11587<br>(AF071527) hypothetical protein<br>[Arabidopsis thaliana]   | 4-5dpi | unknown               | Ankyrin<br>repeats   |
| 12500_s_at<br>(AF081067.3_S_AT)  | gb AAC32192.1  (AF081067)<br>IAA-Ala hydrolase; IAA-amino<br>acid hydrolase [Arabidopsis<br>thaliana]  | 4-5dpi | hormone<br>response   |  |

| Probeset                        | Description  | Time         | Function                  | Blast                                       |
|---------------------------------|--|--------------|---------------------------|---|
| 14711_s_at<br>(ZFPL_S_AT)       | gb AAD25930.1 AF085279_3<br>(AF085279) hypothetical Cys-3-<br>His zinc finger protein<br>[Arabidopsis thaliana]          | 5dpi         | DNA<br>binding<br>protein |   |
| 20345_at<br>(AF104919.16_AT)    | gb AAC72865.1  (AF104919)<br>similar to class I chitinases<br>(Pfam: PF00182, E=1.2e-142,<br>N=1) [Arabidopsis thaliana] | 4dpi         | stress<br>response        |   |
| 17589_at<br>(AF156783_AT)       | gb AAF00612.1 AF156783_1<br>(AF156783) apyrase<br>[Arabidopsis thaliana]   | 5dpi         | membrane<br>protein       |   |
| 18010_s_at<br>(AJ001264_S_AT)   | emb CAA77109.1  (Y18291)<br>uncoupling protein [Arabidopsis<br>thaliana]   | 5dpi         | stress<br>response        |   |
| 12421_at<br>(AJ002414.1_AT)     | emb CAA05398.1  (AJ002414)<br>hnRNP-like protein<br>[Arabidopsis thaliana]   | 2dpi         | RNA<br>processing         |   |
| 13284_s_at<br>(HSP70_S_AT)      | emb CAA05547.1  (AJ002551)<br>heat shock protein 70<br>[Arabidopsis thaliana]  | 2dpi<br>5dpi | heat shock                |   |
| 16442_s_at<br>(AJ002551.2_S_AT) | emb CAA05547.1  (AJ002551)<br>heat shock protein 70<br>[Arabidopsis thaliana]  | 2dpi         | heat shock                |   |
| 19591_at<br>(AJ010735.4_AT)     | emb CAA09330.1  (AJ010735)<br>gr1-protein [Arabidopsis<br>thaliana]  | 4dpi         | stress<br>response        |   |
| 20507_at<br>(AL021635.67_AT)    | emb CAA16557.1  (AL021635)<br>glycoprotein endopeptidase -<br>like protein [Arabidopsis<br>thaliana]                     | 5dpi         | metabolic<br>protein      |   |
| 19531_at<br>(AL021960.91_AT)    | emb CAA17531.1  (AL021960)<br>amino acid transport protein<br>AAT1 [Arabidopsis thaliana]                                | 5dpi         | membrane<br>protein       |   |
| 14595_at<br>(AL022580.163_AT)   | emb CAA18626.1  (AL022580)<br>putative protein [Arabidopsis<br>thaliana]   | 4dpi         | membrane<br>protein       | putative<br>integral<br>membrane<br>protein |
| 14882_at<br>(AL022605.63_AT)    | emb CAA18753.1  (AL022605)<br>putative protein [Arabidopsis<br>thaliana]   | 2dpi         | protein<br>processing     | vacuole<br>processing                       |
| 20142_at<br>(AL035521.155_AT)   | emb CAB36716.1  (AL035521)<br>extra-large G-protein-like<br>[Arabidopsis thaliana]                                       | 5dpi         | signaling not<br>kinase   |   |

| Probeset                         | Description  | Time         | Function                | Blast  |
|----------------------------------|--|--------------|-------------------------|--|
| 20516_at<br>(AL035523.64_AT)     | emb CAB36736.1  (AL035523)<br>ubiquitin activating enzyme-like<br>protein [Arabidopsis thaliana] | 4dpi         | unknown                 | unknown  |
| 13094_at<br>(AL035523.163_AT)    | emb CAB36745.1  (AL035523)<br>putative protein [Arabidopsis<br>thaliana]                         | 5dpi         | protein<br>processing   |  |
| 14032_at<br>(AL035601.11_AT)     | emb CAB38204.1  (AL035601)<br>cytochrome P450-like protein<br>[Arabidopsis thaliana]             | 2dpi<br>5dpi | metabolic<br>protein    |  |
| 13147_at<br>(AL035678.99_AT)     | emb CAB38794.1  (AL035678)<br>putative protein [Arabidopsis<br>thaliana]                         | 5dpi         | metabolic<br>protein    | HpnA<br>protein,<br>oxidoreducta<br>se/cinnamyl-<br>alcohol<br>dehydrogenas<br>e |
| 19624_at<br>(AL049481.196_AT)    | emb CAB39628.1  (AL049481)<br>cytochrome c [Arabidopsis<br>thaliana]                             | 4dpi         | metabolic<br>protein    |  |
| 14036_at<br>(AL049655.54_AT)     | emb CAB41088.1  (AL049655)<br>protein disulfide-isomerase-like<br>protein [Arabidopsis thaliana] | 4dpi         | protein<br>processing   |  |
| 14249_i_at<br>(PAD4_I_AT)        | emb CAB43438.1  (AL050300)<br>putative protein [Arabidopsis<br>thaliana]                         | 5dpi         | stress<br>response      | stress   |
| 14250_r_at<br>(PAD4_R_AT)        | emb CAB43438.1  (AL050300)<br>putative protein [Arabidopsis<br>thaliana]                         | 5dpi         | stress<br>response      | stress   |
| 16232_s_at<br>(AL080252.77_S_AT) | emb CAB45796.1  (AL080252)<br>putative protein [Arabidopsis<br>thaliana]                         | 5dpi         | signaling not<br>kinase | protein<br>phosphatase<br>2C, putative<br>Ser/The<br>phosphatase<br>2C           |
| 13212_s_at<br>(BGL2_S_AT)        | emb CAB68132.1  (AL137080)<br>beta-1, 3-glucanase 2 (BG2)<br>[Arabidopsis thaliana]              | 2-5dpi       | stress<br>response      |  |
| 16578_s_at<br>(ATHERPRP1B_S_AT)  | emb CAB68132.1  (AL137080)<br>beta-1, 3-glucanase 2 (BG2)<br>[Arabidopsis thaliana]              | 2-5dpi       | stress<br>response      |  |
| 15622_s_at<br>(ATU43945_S_AT)    | gb AAB40594.1  (U43945)<br>strictosidine synthase<br>[Arabidopsis thaliana]                      | 4dpi         | stress<br>response      |  |

| Probeset                        | Description   | Time | Function                  | Blast                   |
|---------------------------------|---|------|---------------------------|-------------------------|
| 14145_at<br>(NOVARTIS35_AT)     | dbj BAA22813.1  (D26015)<br>CND41, chloroplast nucleoid<br>DNA binding protein [Nicotiana<br>tabacum] | 4dpi | DNA<br>binding<br>protein |                         |
| 15680_s_at<br>(ATHATPK19B_S_AT) | dbj BAA07661.1  (D42061)<br>ribosomal-protein S6 kinase<br>homolog [Arabidopsis thaliana]             | 5dpi | signaling not<br>kinase   |                         |
| 16173_s_at<br>(D78607_S_AT)     | dbj BAA28539.1  (D78607)<br>cytochrome P450<br>monooxygenase [Arabidopsis<br>thaliana]                | 4dpi | metabolic<br>protein      |                         |
| 12832_f_at<br>(U33014.2_M_F_AT) | gb AAD03342.1  (L81140)<br>ubiquitin [Pisum sativum]  | 1dpi | protein<br>degradation    |                         |
| 13285_s_at<br>(HSP83_S_AT)      | gb AAA32822.1  (M62984)<br>heat shock protein 83<br>[Arabidopsis thaliana]                            | 5dpi | heat shock                |                         |
| 16091_s_at<br>(ATHHSP83_S_AT)   | gb AAA32822.1  (M62984)<br>heat shock protein 83<br>[Arabidopsis thaliana]                            | 5dpi | heat shock                |                         |
| 13722_at<br>(NOVARTIS108_AT)    | No hits found less than or equal<br>to 1e-15.   | 5dpi | DNA<br>binding<br>protein | transcription<br>factor |
| 13187_i_at<br>(ATTHIRE4_I_AT)   | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]                                    | 5dpi | stress<br>response        |                         |
| 13188_r_at<br>(ATTHIRE4_R_AT)   | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]                                    | 5dpi | stress<br>response        |                         |
| 13189_s_at<br>(ATTHIRE4_S_AT)   | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]                                    | 5dpi | stress<br>response        |                         |
| 16981_s_at<br>(U35829.2_S_AT)   | gb AAC49356.1  (U35829)<br>thioredoxin h [Arabidopsis<br>thaliana]                                    | 5dpi | stress<br>response        |                         |
| 17544_s_at<br>(ATU40856_S_AT)   | gb AAC49282.1  (U40856)<br>AIG1 [Arabidopsis thaliana]  | 5dpi | stress<br>response        |                         |
| 12880_s_at<br>(AIG2_S_AT)       | gb AAC49283.1  (U40857)<br>AIG2 [Arabidopsis thaliana]  | 1dpi | stress<br>response        |                         |
| 17097_s_at<br>(ATU66345_S_AT)   | gb AAC49697.1  (U66345)<br>calreticulin [Arabidopsis<br>thaliana]                                     | 5dpi | protein<br>processing     |                         |



| Probeset                        | Description  | Time   | Function            | Blast  |
|---------------------------------|--|--------|---------------------|--|
| 17027_s_at<br>(ATU72958_S_AT)   | gb AAB38795.1  (U72958)<br>AtHSP23.6-mito [Arabidopsis thaliana] protein (AtHSP23.6-mito) [Arabidopsis thaliana] | 5dpi   | heat shock          |  |
| 14998_at<br>(U93215.42_AT)      | gb AAB63078.1  (U93215)<br>unknown protein [Arabidopsis thaliana]  | 5dpi   | DNA binding protein | Petroselinum crispum transcription factor (WRKY3) WRKY3 DNA-binding protein, TMV response-related gene product |
| 13275_f_at<br>(HSP174_F_AT)     | emb CAA35182.1  (X17293)<br>heat shock protein (AA 1 - 156) [Arabidopsis thaliana]                               | 4-5dpi | heat shock          |  |
| 12831_f_at<br>(U33014.2_5_F_AT) | emb CAA40323.1  (X57003)<br>polyubiquitin protein [Helianthus annuus]  | 1dpi   | protein degradation |  |
| 20238_at<br>(X74514.2_AT)       | emb CAA52619.1  (X74514)<br>beta-fructofuranosidase [Arabidopsis thaliana]                                       | 4dpi   | metabolic protein   |  |
| 15118_s_at<br>(ATHGLUGRFS_S_AT) | emb CAA53051.1  (X75303)<br>glutathione S-transferase [Arabidopsis thaliana]                                     | 5dpi   | stress response     |  |
| 17484_at<br>(X79052.2_AT)       | emb CAA55654.1  (X79052)<br>SRG1 [Arabidopsis thaliana]  | 5dpi   | stress response     |  |
| 16236_g_at<br>(X92657.3_G_AT)   | emb CAA63346.1  (X92657)<br>cationic amino acid transporter [Arabidopsis thaliana]                               | 5dpi   | membrane protein    |  |
| 18216_at<br>(X95573.2_AT)       | gb AAF24959.1 AC012375_22 (AC012375) T22C5.18 [Arabidopsis thaliana]   | 5dpi   | DNA binding protein | salt-tolerance zinc finger protein   |
| 18217_g_at<br>(X95573.2_G_AT)   | gb AAF24959.1 AC012375_22 (AC012375) T22C5.18 [Arabidopsis thaliana]   | 5dpi   | DNA binding protein | A.thaliana mRNA for salt-tolerance zinc finger protein   |

| Probeset                        | Description  | Time   | Function                  | Blast |
|---------------------------------|--|--------|---------------------------|-------|
| 20660_s_at<br>(X97488.2_S_AT)   | emb CAA66120.1  (X97488)<br>beta-transducin like protein<br>[Arabidopsis thaliana]   | 4dpi   | signaling not<br>kinase   |       |
| 13015_s_at<br>(X98673.2_S_AT)   | emb CAA67232.1  (X98674)<br>zinc finger protein [Arabidopsis<br>thaliana]            | 5dpi   | DNA<br>binding<br>protein |       |
| 16465_at<br>(Y08892.1_AT)       | emb CAA70105.1  (Y08892)<br>Hsc70-G8 protein [Arabidopsis<br>thaliana]               | 4-5dpi | heat shock                |       |
| 13277_i_at<br>(HSP176A_I_AT)    | emb CAA74399.1  (Y14070)<br>Heat Shock Protein 17.6A<br>[Arabidopsis thaliana]       | 2-5dpi | heat shock                |       |
| 16054_s_at<br>(Y14251.4_S_AT)   | emb CAA74639.1  (Y14251)<br>glutathione S-transferase<br>[Arabidopsis thaliana]      | 5dpi   | stress<br>response        |       |
| 20287_at<br>(Y14590.5_AT)       | emb CAA74930.1  (Y14590)<br>class IV chitinase [Arabidopsis<br>thaliana]             | 4-5dpi | stress<br>response        |       |
| 19178_at<br>(Y18227.2_AT)       | dbj BAA86999.1  (AB035137)<br>blue copper binding protein<br>[Arabidopsis thaliana]  | 5dpi   | metabolic<br>protein      |       |
| 16943_s_at<br>(Z97339.466_S_AT) | emb CAB10370.1  (Z97339)<br>drought-induced protein like<br>[Arabidopsis thaliana]   | 5dpi   | stress<br>response        |       |
| 13803_at<br>(Z97341.376_AT)     | emb CAB10444.1  (Z97341)<br>cyanohydrin lyase like protein<br>[Arabidopsis thaliana] | 4dpi   | metabolic<br>protein      |       |
| 18140_at<br>(Z97341.319_AT)     | emb CAB10440.1  (Z97341)<br>growth regulator like protein<br>[Arabidopsis thaliana]  | 5dpi   | hormone<br>response       |       |

**Table 19** Correlation of SEQ ID NOs:1-953 (#) representing genes, the expression of which is altered after pathogen infection of *Arabidopsis* with ProbeSet designations

| #  | ProbeSet                          | #  | ProbeSet                     |
|----|-----------------------------------|----|------------------------------|
| 1  | 11991_g_at<br>(AC002387.210_G_AT) | 33 | 12218_at (AJ242588.2_AT)     |
| 2  | 11997_at (AC005967.4_AT)          |    | 12223_s_at                   |
| 3  | 12002_at (AF069442.47_AT)         | 34 | (AC007168.178_S_AT)          |
| 4  | 12004_at (AL022023.132_AT)        | 35 | 12227_at (AC007576.18_AT)    |
| 5  | 12007_at (Z99708.249_AT)          | 36 | 12233_at (AJ001807.1_AT)     |
| 6  | 12037_at (AC004005.174_AT)        | 37 | 12278_at (AJ011674.2_AT)     |
| 7  | 12048_at (AF001308.46_AT)         | 38 | 12290_at (Y09418.2_AT)       |
| 8  | 12051_at (AL021889.94_AT)         | 39 | 12307_at (AC002392.162_AT)   |
| 9  | 12062_at (AC006069.147_AT)        | 40 | 12314_at (AC001229.28_AT)    |
| 10 | 12068_at (AF118223.24_AT)         | 41 | 12317_at (AC004138.27_AT)    |
| 11 | 12072_at (AL035396.4_AT)          | 42 | 12323_at (AC002333.18_AT)    |
| 12 | 12079_s_at (A71597.1_S_AT)        | 43 | 12332_s_at (AB023448.2_S_AT) |
| 13 | 12081_at (AC001645.140_AT)        | 44 | 12335_at (AC004411.73_AT)    |
| 14 | 12086_s_at (AC002409.88_S_AT)     |    | 12341_s_at                   |
| 15 | 12091_at (AC004450.116_AT)        | 45 | (AL021637.176_S_AT)          |
| 16 | 12092_at (AC004793.13_AT)         | 46 | 12347_at (AC007258.28_AT)    |
| 17 | 12094_at (AC006223.143_AT)        | 47 | 12349_s_at (X84728.6_S_AT)   |
| 18 | 12115_at (AL033545.26_AT)         | 48 | 12356_at (X99952.1_AT)       |
| 19 | 12124_s_at (Z97337.149_S_AT)      | 49 | 12369_at (AC002535.59_AT)    |
| 20 | 12125_at (Z97341.99_AT)           | 50 | 12400_at (X98453.1_AT)       |
| 21 | 12128_at (AC004261.157_AT)        | 51 | 12421_at (AJ002414.1_AT)     |
| 22 | 12136_at (AC007591.60_AT)         | 52 | 12438_at (AL021710.83_AT)    |
| 23 | 12150_at (AC004005.151_AT)        | 53 | 12448_at (AC002337.58_AT)    |
| 24 | 12160_at (AC006284.117_AT)        |    | 12449_s_at                   |
| 25 | 12187_at (AC005489.31_AT)         | 54 | (AC002343.179_S_AT)          |
| 26 | 12191_at (AC006068.35_AT)         | 55 | 12454_at (AC006232.164_AT)   |
| 27 | 12193_at (AC006072.132_AT)        |    | 12460_s_at                   |
| 28 | 12198_at (AC006954.90_AT)         | 56 | (AC006920.129_S_AT)          |
| 29 | 12203_at (AL021710.268_AT)        | 57 | 12475_at (Y11794.1_AT)       |
| 30 | 12212_at (AL049711.168_AT)        | 58 | 12487_at (AC004411.126_AT)   |
| 31 | 12216_at (AC007119.56_AT)         | 59 | 12493_g_at (Y09095.1_G_AT)   |
| 32 | 12217_at (AJ223804.1_AT)          | 60 | 12497_at (AC006533.51_AT)    |
|    |                                   | 61 | 12500_s_at (AF081067.3_S_AT) |
|    |                                   | 62 | 12521_at (AF049236.28_AT)    |

| #  | ProbeSet                      | #   | ProbeSet                     |
|----|-------------------------------|-----|------------------------------|
| 63 | 12525_at (AC006587.85_AT)     | 96  | 12832_f_at (U33014.2_M_F_AT) |
| 64 | 12530_at (Z99707.184_AT)      | 97  | 12851_s_at (ACCSYN1_S_AT)    |
| 65 | 12535_at (AL035538.156_AT)    | 98  | 12854_s_at (ACS1_S_AT)       |
| 66 | 12538_at (AF033205.2_AT)      | 99  | 12855_f_at (ADH_F_AT)        |
| 67 | 12559_at (AC005727.83_AT)     | 100 | 12879_s_at (AIG1_S_AT)       |
| 68 | 12560_at (AC005825.57_AT)     | 101 | 12880_s_at (AIG2_S_AT)       |
| 69 | 12561_at (AL021687.107_AT)    | 102 | 12883_s_at (APX_S_AT)        |
|    | 12571_s_at                    | 103 | 12889_s_at (ASA1_S_AT)       |
| 70 | (AF149413.18_S_AT)            | 104 | 12891_at (ATACS6_AT)         |
| 71 | 12574_at (X82624.2_AT)        | 105 | 12892_g_at (ATACS6_G_AT)     |
| 72 | 12584_at (AC004521.233_AT)    | 106 | 12904_s_at (ATERF1_S_AT)     |
| 73 | 12609_at (X92975.2_AT)        | 107 | 12905_s_at (ATERF2_S_AT)     |
| 74 | 12626_at (AC006234.95_AT)     | 108 | 12906_s_at (ATERF3_S_AT)     |
| 75 | 12627_at (AC006533.99_AT)     | 109 | 12908_s_at (ATERF5_S_AT)     |
| 76 | 12642_at (AC006920.138_AT)    | 110 | 12909_s_at (ATERF6_S_AT)     |
| 77 | 12645_at (AL021712.56_AT)     |     | 12911_s_at                   |
| 78 | 12656_at (AC006569.43_AT)     | 111 | (ATG6PDHE5_S_AT)             |
| 79 | 12698_at (AC000106.42_AT)     |     | 12921_s_at                   |
| 80 | 12711_f_at (Z95773_F_AT)      | 112 | (ATHHMGCOAR_S_AT)            |
| 81 | 12712_f_at (Z95774_F_AT)      | 113 | 12930_s_at (ATLLS1_S_AT)     |
| 82 | 12727_f_at (Z95799_F_AT)      | 114 | 12951_at (AC005489.5_AT)     |
| 83 | 12736_f_at (Z97048_F_AT)      | 115 | 12958_at (AC002332.249_AT)   |
| 84 | 12737_f_at (Z97049_F_AT)      | 116 | 12962_at (AC004697.165_AT)   |
| 85 | 12744_at (AC001645.15_AT)     | 117 | 12965_at (AL021711.118_AT)   |
| 86 | 12759_at (AC005278.32_AT)     |     | 12966_s_at                   |
|    | 12760_g_at                    | 118 | (AL023094.197_S_AT)          |
| 87 | (AC005278.32_G_AT)            |     | 12989_s_at                   |
| 88 | 12764_f_at (AC004138.69_F_AT) | 119 | (AC004077.149_S_AT)          |
| 89 | 12772_at (AC005278.34_AT)     | 120 | 13003_s_at (AB021936.1_S_AT) |
| 90 | 12776_at (AL021811.156_AT)    | 121 | 13005_at (AC004683.61_AT)    |
| 91 | 12790_s_at (AL021635.58_S_AT) | 122 | 13014_at (U93215.87_AT)      |
|    | 12797_s_at                    | 123 | 13015_s_at (X98673.2_S_AT)   |
| 92 | (AC007138.25_S_AT)            | 124 | 13025_at (AL050400.20_AT)    |
| 93 | 12801_at (AC005223.34_AT)     | 125 | 13040_at (AC002392.134_AT)   |
| 94 | 12802_at (AL022373.153_AT)    |     | 13048_s_at                   |
| 95 | 12831_f_at (U33014.2_5_F_AT)  | 126 | (AC004138.22_S_AT)           |
|    |                               | 127 | 13070_at (AC006919.171_AT)   |

| #   | ProbeSet                           | #   | ProbeSet                       |
|-----|------------------------------------|-----|--------------------------------|
| 128 | 13094_at (AL035523.163_AT)         | 156 | 13243_r_at (ELI32_R_AT)        |
| 129 | 13100_at (AC003680.50_AT)          | 157 | 13244_s_at (ELI32_S_AT)        |
| 130 | 13110_at (AF074021.34_AT)          | 158 | 13246_at (ERECTAL_AT)          |
| 131 | 13115_at (AC000375.44_AT)          | 159 | 13253_f_at (FPS1_F_AT)         |
| 132 | 13119_at (AC007260.23_AT)          |     | 13255_i_at                     |
| 133 | 13128_at (AL049607.47_AT)          | 160 | (gammaglutamyltranspepti_I_AT) |
| 134 | 13134_s_at (AC002337.9_S_AT)       |     | 13256_s_at                     |
| 135 | 13137_at (AC007169.86_AT)          | 161 | (gammaglutamyltranspepti_S_AT) |
| 136 | 13144_at (AC007017.246_AT)         |     | 13259_s_at                     |
| 137 | 13147_at (AL035678.99_AT)          | 162 | (glutathioneperoxidase1_S_AT)  |
|     | 13152_s_at                         |     | 13261_s_at                     |
| 138 | (AC005322.24_S_AT)                 | 163 | (glutathionereductase1_S_AT)   |
|     | 13154_s_at                         | 164 | 13263_s_at (GST1_RC_S_AT)      |
| 139 | (AC002333.210_S_AT)                | 165 | 13266_s_at (GST4_S_AT)         |
| 140 | 13157_at (AC002409.35_AT)          | 166 | 13270_at (HSF21_AT)            |
| 141 | 13161_at (AF002109.89_AT)          | 167 | 13271_g_at (HSF21_G_AT)        |
|     | 13163_s_at                         | 168 | 13273_s_at (HSF4_S_AT)         |
| 142 | (AC005560.223_S_AT)                | 169 | 13275_f_at (HSP174_F_AT)       |
| 143 | 13176_at (AL031394.56_AT)          | 170 | 13277_i_at (HSP176A_I_AT)      |
| 144 | 13177_at (AL049640.42_AT)          | 171 | 13279_s_at (HSP176II_S_AT)     |
| 145 | 13187_i_at (ATTHIRED4_I_AT)        | 172 | 13284_s_at (HSP70_S_AT)        |
|     | 13188_r_at                         | 173 | 13285_s_at (HSP83_S_AT)        |
| 146 | (ATTHIRED4_R_AT)                   | 174 | 13312_at (AC006223.75_AT)      |
|     | 13189_s_at                         | 175 | 13367_at (AC004680.97_AT)      |
| 147 | (ATTHIRED4_S_AT)                   | 176 | 13370_at (AC005322.4_AT)       |
| 148 | 13190_s_at (ATTHIREDA_S_AT)        | 177 | 13381_at (AC006580.8_AT)       |
| 149 | 13211_s_at (BCHI_S_AT)             | 178 | 13395_at (AL035528.202_AT)     |
| 150 | 13212_s_at (BGL2_S_AT)             | 179 | 13435_at (AF003102.3_AT)       |
|     | 13215_s_at                         | 180 | 13437_at (AF096371.8_AT)       |
| 151 | (CAFFEROYLCOAMETHYLTR<br>ANS_S_AT) | 181 | 13450_at (AL049657.33_AT)      |
|     | 13217_s_at                         | 182 | 13459_at (AF013294.21_AT)      |
| 152 | (CALMODULINLIKE_S_AT)              | 183 | 13467_at (AL096860.198_AT)     |
| 153 | 13219_s_at (CHI4_S_AT)             | 184 | 13480_at (AC005223.15_AT)      |
|     | 13220_s_at (CHS-                   | 185 | 13482_at (AC005896.195_AT)     |
| 154 | EXON1_S_AT)                        | 186 | 13534_at (AF149413.36_AT)      |
|     | 13221_at (CHS-WHOLE-               | 187 | 13536_at (AL021636.47_AT)      |
| 155 | SEQ_AT)                            | 188 | 13538_at (AL080254.75_AT)      |

| #   | ProbeSet                      | #   | ProbeSet                      |
|-----|-------------------------------|-----|-------------------------------|
| 189 | 13547_s_at (AC004450.15_S_AT) | 220 | 13755_at (NOVARTIS15_AT)      |
| 190 | 13564_at (AC005312.113_AT)    | 221 | 13763_at (NOVARTIS21_AT)      |
| 191 | 13565_at (AL035601.21_AT)     | 222 | 13764_at (NOVARTIS22_AT)      |
| 192 | 13584_at (AC007127.23_AT)     | 223 | 13789_at (AJ132436.2_AT)      |
| 193 | 13588_at (AL021961.24_AT)     | 224 | 13803_at (Z97341.376_AT)      |
| 194 | 13589_at (AC000132.24_AT)     | 225 | 13806_at (AC002354.15_AT)     |
| 195 | 13604_at (AC000104.20_AT)     |     | 13812_s_at                    |
| 196 | 13605_at (AL078470.75_AT)     | 226 | (AC005275.104_S_AT)           |
| 197 | 13617_at (AC006592.64_AT)     |     | 13818_s_at                    |
| 198 | 13627_at (AL035394.196_AT)    | 227 | (AC006218.175_S_AT)           |
| 199 | 13637_at (AL049482.9_AT)      | 228 | 13825_s_at (AF104919.22_S_AT) |
| 200 | 13645_at (AC000098.8_AT)      | 229 | 13834_at (AL080237.29_AT)     |
| 201 | 13647_at (AF000657.22_AT)     | 230 | 13842_at (AC002396.12_AT)     |
| 202 | 13651_at (AL035538.320_AT)    | 231 | 13848_at (AC003981.31_AT)     |
| 203 | 13656_at (AC007138.31_AT)     |     | 13880_s_at                    |
| 204 | 13659_at (AL022347.46_AT)     | 232 | (AL049480.183_S_AT)           |
|     | 13666_s_at                    | 233 | 13895_at (AC004218.63_AT)     |
| 205 | (INDOLE3GPS_S_AT)             | 234 | 13896_at (AC004473.8_AT)      |
| 206 | 13680_s_at (LOX1_S_AT)        | 235 | 13908_s_at (A71590.1_S_AT)    |
| 207 | 13685_s_at (MLOLIKE2_S_AT)    | 236 | 13918_at (AC005388.29_AT)     |
|     | 13688_s_at                    | 237 | 13920_at (AC005990.53_AT)     |
| 208 | (MONOPTEROS_S_AT)             | 238 | 13944_at (U89959.24_AT)       |
| 209 | 13697_at (NI16_AT)            | 239 | 13949_s_at (Z97343.352_S_AT)  |
| 210 | 13705_s_at (AC003671X_S_AT)   | 240 | 13963_at (AL021711.26_AT)     |
| 211 | 13706_s_at (AC005724X_S_AT)   | 241 | 13964_at (AL021889.3_AT)      |
| 212 | 13708_s_at (AC007259X_S_AT)   | 242 | 13966_at (AL022023.172_AT)    |
|     | 13714_at                      | 243 | 13999_at (AF071527.56_AT)     |
| 213 | (NOVARTIS101_RC_AT)           | 244 | 14015_s_at (A71588.1_S_AT)    |
|     | 13716_at                      | 245 | 14016_s_at (A71596.1_S_AT)    |
| 214 | (NOVARTIS103_RC_AT)           | 246 | 14025_s_at (AC007293.3_S_AT)  |
|     | 13718_at                      | 247 | 14026_at (AC000106.5_AT)      |
| 215 | (NOVARTIS105_RC_AT)           | 248 | 14030_at (AC005970.225_AT)    |
| 216 | 13722_at (NOVARTIS108_AT)     | 249 | 14032_at (AL035601.11_AT)     |
| 217 | 13728_at (NOVARTIS111_AT)     | 250 | 14036_at (AL049655.54_AT)     |
|     | 13746_at                      | 251 | 14041_at (AC003970.28_AT)     |
| 218 | (NOVARTIS121_RC_AT)           | 252 | 14052_at (AC004122.24_AT)     |
| 219 | 13751_at (NOVARTIS127_AT)     |     |                               |

| #   | ProbeSet                          | #   | ProbeSet                                       |
|-----|-----------------------------------|-----|--|
| 253 | 14068_s_at<br>(AC006922.197_S_AT) | 286 | 14448_at (AC002387.243_AT)                     |
| 254 | 14070_at (AL049658.217_AT)        | 287 | 14450_at (AC002986.49_AT)                      |
| 255 | 14083_at (AC005662.56_AT)         | 288 | 14459_at (AC006200.69_AT)                      |
| 256 | 14089_at (AC006223.65_AT)         | 289 | 14460_at (AC006201.21_AT)                      |
| 257 | 14100_at (AF002109.108_AT)        | 290 | 14461_at (AC006202.73_AT)                      |
| 258 | 14110_i_at (AL035528.279_I_AT)    | 291 | 14468_at (AC007576.62_AT)                      |
| 259 | 14116_at (AF077407.30_AT)         | 292 | 14475_at (AL021811.121_AT)                     |
| 260 | 14122_at (AF058826.23_AT)         | 293 | 14487_at (Z97341.343_AT)                       |
| 261 | 14139_at (NOVARTIS30_AT)          | 294 | 14498_at (AC004261.51_AT)                      |
| 262 | 14141_at (NOVARTIS31_AT)          | 295 | 14530_at (AL021889.231_AT)                     |
| 263 | 14145_at (NOVARTIS35_AT)          | 296 | 14573_at (AF069298.35_AT)                      |
| 264 | 14148_at (NOVARTIS38_AT)          | 297 | 14584_at (AC007658.25_AT)                      |
| 265 | 14162_at (NOVARTIS46_AT)          | 298 | 14591_at (AL035440.107_AT)                     |
| 266 | 14170_at (NOVARTIS51_AT)          | 299 | 14595_at (AL022580.163_AT)                     |
| 267 | 14197_at (NOVARTIS71_AT)          | 300 | 14605_at (AC006193.6_AT)                       |
| 268 | 14214_at (NOVARTIS83_AT)          | 301 | 14608_at (AC007357.49_AT)                      |
| 269 | 14217_at<br>(NOVARTIS85_RC_AT)    | 302 | 14609_at (AC002340.147_AT)                     |
| 270 | 14223_at (NOVARTIS9_AT)           | 303 | 14613_at (AC004669.20_AT)                      |
| 271 | 14235_at (NOVARTIS97_AT)          | 304 | 14614_at (AC004165.66_AT)                      |
| 272 | 14240_s_at (NR1_S_AT)             | 305 | 14620_s_at (PAT1_S_AT)                         |
| 273 | 14242_s_at (NRA_S_AT)             | 306 | 14635_s_at (PR1_S_AT)                          |
| 274 | 14248_at (PAD3_AT)                | 307 | 14636_s_at (PR5_S_AT)                          |
| 275 | 14249_i_at (PAD4_I_AT)            | 308 | 14638_s_at (PRXCB_S_AT)                        |
| 276 | 14250_r_at (PAD4_R_AT)            | 309 | 14640_s_at<br>(PUTATIVEMLOHI_S_AT)             |
| 277 | 14254_s_at (PAL1-MRNA_S_AT)       | 310 | 14643_s_at (RAR047_S_AT)                       |
| 278 | 14256_f_at (PAL1-<br>INTRON_F_AT) | 311 | 14660_s_at<br>(THIOREDOXL_S_AT)                |
| 279 | 14257_s_at (PAL2-MRNA_S_AT)       | 312 | 14663_s_at<br>(TREHALASEPRECURSOR_RC_<br>S_AT) |
| 280 | 14320_at (AC005956.54_AT)         | 313 | 14667_s_at (TRPB_S_AT)                         |
| 281 | 14381_at (AC002521.68_AT)         | 314 | 14672_s_at (TSA1_S_AT)                         |
| 282 | 14386_at (AC005309.177_AT)        | 315 | 14673_s_at (TSB2_S_AT)                         |
| 283 | 14408_at (AC002291.14_AT)         | 316 | 14675_s_at (VSP_S_AT)                          |
| 284 | 14428_s_at<br>(AL023094.342_S_AT) | 317 | 14682_i_at<br>(WT1012A_RC_I_AT)                |
| 285 | 14431_at (AL035394.279_AT)        |     |  |

| #   | ProbeSet                      | #   | ProbeSet                      |
|-----|-------------------------------|-----|-------------------------------|
| 318 | 14686_s_at (WT1073_S_AT)      | 352 | 15067_at (AC004683.36_AT)     |
| 319 | 14696_at (WT740_RC_AT)        | 353 | 15073_at (AC007069.93_AT)     |
| 320 | 14697_g_at (WT740_RC_G_AT)    |     | 15085_s_at                    |
| 321 | 14705_i_at (WT77_RC_I_AT)     | 354 | (AL031018.274_S_AT)           |
| 322 | 14706_r_at (WT77_RC_R_AT)     | 355 | 15088_s_at (AC002311.37_S_AT) |
| 323 | 14711_s_at (ZFPL_S_AT)        | 356 | 15091_at (AC004683.97_AT)     |
| 324 | 14735_s_at (AF008124_S_AT)    | 357 | 15098_s_at (ATU26945_S_AT)    |
| 325 | 14750_s_at (AF096370.12_S_AT) | 358 | 15116_f_at (AF121356_F_AT)    |
| 326 | 14763_at (X86958.1_AT)        |     | 15118_s_at                    |
|     | 14770_s_at                    | 359 | (ATHGLUGRFS_S_AT)             |
| 327 | (AC002338.167_S_AT)           | 360 | 15120_s_at (ATU10034_S_AT)    |
| 328 | 14779_at (AC004680.71_AT)     | 361 | 15123_s_at (ATU40857_S_AT)    |
| 329 | 14780_at (AC004683.103_AT)    | 362 | 15124_s_at (ATU59508_S_AT)    |
| 330 | 14786_at (AC005397.115_AT)    | 363 | 15125_f_at (D85190_F_AT)      |
| 331 | 14793_at (AC006202.10_AT)     | 364 | 15129_s_at (AF030386_S_AT)    |
| 332 | 14838_s_at (M96073.6_S_AT)    | 365 | 15132_s_at (AF121878_S_AT)    |
| 333 | 14882_at (AL022605.63_AT)     | 366 | 15137_s_at (ATU57320_S_AT)    |
| 334 | 14884_at (AL031032.95_AT)     | 367 | 15140_s_at (ATU93845_S_AT)    |
| 335 | 14895_s_at (Z97344.138_S_AT)  | 368 | 15141_s_at (D85191_S_AT)      |
| 336 | 14900_at (AC000348.12_AT)     |     | 15154_s_at                    |
| 337 | 14923_at (AC006283.158_AT)    | 369 | (ATHMTGDAS_S_AT)              |
| 338 | 14924_at (AC006283.46_AT)     | 370 | 15161_s_at (ATU90522_S_AT)    |
| 339 | 14928_at (AC006569.88_AT)     | 371 | 15162_s_at (U01880_S_AT)      |
| 340 | 14931_at (AC006951.173_AT)    | 372 | 15175_s_at (ATU28215_S_AT)    |
| 341 | 14959_at (AC007202.26_AT)     | 373 | 15188_s_at (AF081202_S_AT)    |
| 342 | 14964_at (AC001229.8_AT)      | 374 | 15192_s_at (ATHERD1_S_AT)     |
| 343 | 14965_at (AC002329.22_AT)     | 375 | 15196_s_at (ATU43412_S_AT)    |
| 344 | 14972_at (AC005499.38_AT)     | 376 | 15197_s_at (ATU52851_S_AT)    |
| 345 | 14978_at (AC002333.49_AT)     | 377 | 15199_s_at (AB005804_S_AT)    |
| 346 | 14998_at (U93215.42_AT)       | 378 | 15203_s_at (AB013887_S_AT)    |
| 347 | 15032_at (AC002294.8_AT)      | 379 | 15211_s_at (ATH243813_S_AT)   |
| 348 | 15039_at (AF001308.67_AT)     | 380 | 15216_s_at (ATU75191_S_AT)    |
|     | 15040_g_at                    | 381 | 15342_at (AC006593.101_AT)    |
| 349 | (AF001308.67_G_AT)            | 382 | 15379_at (AC002335.182_AT)    |
| 350 | 15042_at (AL021961.3_AT)      | 383 | 15389_at (AC004786.100_AT)    |
| 351 | 15052_at (AC002332.103_AT)    | 384 | 15406_at (AC006931.179_AT)    |



| #   | ProbeSet                       | #   | ProbeSet                      |
|-----|--------------------------------|-----|-------------------------------|
| 385 | 15431_at (AL030978.64_AT)      | 415 | 15622_s_at (ATU43945_S_AT)    |
| 386 | 15463_at (AL031326.226_AT)     | 416 | 15625_s_at (ATU74610_S_AT)    |
| 387 | 15473_at (AC006836.125_AT)     | 417 | 15629_s_at (AB003280_S_AT)    |
| 388 | 15479_at (AL049483.205_AT)     | 418 | 15631_s_at (AB005805_S_AT)    |
|     | 15483_s_at                     | 419 | 15632_s_at (AB012570_S_AT)    |
| 389 | (AC005819.20_S_AT)             | 420 | 15641_s_at (AF117063_S_AT)    |
| 390 | 15485_at (AC006233.109_AT)     | 421 | 15646_s_at (ATHSAT1G_S_AT)    |
| 391 | 15487_at (AC007661.87_AT)      | 422 | 15665_s_at (AF022658_S_AT)    |
| 392 | 15496_at (AC006282.167_AT)     | 423 | 15669_s_at (AF047834_S_AT)    |
|     | 15515_r_at                     | 424 | 15670_s_at (AF061638_S_AT)    |
| 393 | (AF058919.32_R_AT)             | 425 | 15672_s_at (AF082299_S_AT)    |
| 394 | 15518_at (AC005322.28_AT)      | 426 | 15674_s_at (AF091844_S_AT)    |
| 395 | 15522_i_at (AL078637.213_I_AT) |     | 15680_s_at                    |
|     | 15523_s_at                     | 427 | (ATHATPK19B_S_AT)             |
| 396 | (AL078637.213_S_AT)            | 428 | 15775_at (AL079344.196_AT)    |
| 397 | 15524_at (AC005508.25_AT)      | 429 | 15778_at (X98676.2_AT)        |
| 398 | 15526_at (AC004122.16_AT)      | 430 | 15779_g_at (X98676.2_G_AT)    |
|     | 15531_i_at                     | 431 | 15792_at (AC002341.106_AT)    |
| 399 | (AL078637.191_I_AT)            | 432 | 15798_at (AC002521.173_AT)    |
|     | 15532_r_at                     | 433 | 15815_s_at (Z97342.366_S_AT)  |
| 400 | (AL078637.191_R_AT)            | 434 | 15832_at (AC004561.255_AT)    |
| 401 | 15539_at (AC005770.21_AT)      | 435 | 15839_at (AC005662.203_AT)    |
| 402 | 15540_at (AC006585.205_AT)     | 436 | 15859_at (AC006587.164_AT)    |
| 403 | 15543_at (AF096371.10_AT)      | 437 | 15866_s_at (AC007133.59_S_AT) |
| 404 | 15544_at (AL021633.110_AT)     | 438 | 15874_at (AL022223.106_AT)    |
| 405 | 15547_at (AC005970.122_AT)     | 439 | 15886_at (AL078637.204_AT)    |
| 406 | 15551_at (AL035440.289_AT)     | 440 | 15900_at (AC005311.74_AT)     |
| 407 | 15578_s_at (AF004213_S_AT)     | 441 | 15919_at (AC007060.42_AT)     |
| 408 | 15580_s_at (AF057043_S_AT)     | 442 | 15921_s_at (AC007067.1_S_AT)  |
| 409 | 15582_s_at (ATH131392_S_AT)    | 443 | 15924_at (AC007138.61_AT)     |
| 410 | 15594_s_at (ATU56635_S_AT)     | 444 | 15943_at (AC007202.16_AT)     |
| 411 | 15606_s_at (AF061517_S_AT)     | 445 | 15970_s_at (X71794.2_S_AT)    |
|     | 15613_s_at                     | 446 | 15978_at (X68592.6_AT)        |
| 412 | (ATHHOMEOA_S_AT)               |     | 15982_s_at                    |
|     | 15614_s_at                     | 447 | (AC006260.78_S_AT)            |
| 413 | (ATHMERI5B_S_AT)               | 448 | 16001_at (AF035385.2_AT)      |
| 414 | 15617_s_at (ATHSAR1_S_AT)      |     |                               |

| #   | ProbeSet                          | #   | ProbeSet                          |
|-----|-----------------------------------|-----|-----------------------------------|
| 449 | 16003_s_at<br>(AL021749.64_S_AT)  | 481 | 16173_s_at (D78607_S_AT)          |
| 450 | 16021_s_at<br>(AL022224.182_S_AT) | 482 | 16203_at (AC007519.53_AT)         |
| 451 | 16031_at (X94248.1_AT)            | 483 | 16229_at (AL049638.148_AT)        |
| 452 | 16043_at (AC005489.17_AT)         | 484 | 16230_at (AL049655.78_AT)         |
| 453 | 16048_at (X78586.2_AT)            | 485 | 16232_s_at<br>(AL080252.77_S_AT)  |
| 454 | 16053_i_at (Y14251.4_I_AT)        | 486 | 16233_at (AL080254.83_AT)         |
| 455 | 16054_s_at (Y14251.4_S_AT)        | 487 | 16236_g_at (X92657.3_G_AT)        |
| 456 | 16058_s_at (ATU94495_S_AT)        | 488 | 16241_at (AL022604.84_AT)         |
| 457 | 16059_s_at (D88206_S_AT)          | 489 | 16272_at (AC006304.136_AT)        |
| 458 | 16062_s_at (AB007789_S_AT)        | 490 | 16288_at (AF024504.17_AT)         |
| 459 | 16063_s_at (AB008103_S_AT)        | 491 | 16298_at (AL021890.71_AT)         |
| 460 | 16073_f_at (AF062908_F_AT)        | 492 | 16299_at (AL024486.185_AT)        |
| 461 | 16077_s_at (AF085230_S_AT)        | 493 | 16301_s_at<br>(AL031018.105_S_AT) |
| 462 | 16080_f_at (AF118822_F_AT)        | 494 | 16306_at (AL049751.112_AT)        |
| 463 | 16083_s_at (AF153283_S_AT)        | 495 | 16327_at (AC002334.12_AT)         |
| 464 | 16087_s_at<br>(ATHATPK6A_S_AT)    | 496 | 16329_s_at<br>(AF013294.17_S_AT)  |
| 465 | 16090_s_at (ATHFAD8A_S_AT)        | 497 | 16335_at (AL079347.105_AT)        |
| 466 | 16091_s_at (ATHHSP83_S_AT)        | 498 | 16340_at (AC004255.15_AT)         |
| 467 | 16092_s_at (ATHKAT1_S_AT)         | 499 | 16351_at (AL021684.194_AT)        |
| 468 | 16103_s_at (ATU60445_S_AT)        | 500 | 16357_at (AF149413.38_AT)         |
| 469 | 16105_s_at (ATU68017_S_AT)        | 501 | 16363_at (AC004255.14_AT)         |
| 470 | 16108_s_at (D78604_S_AT)          | 502 | 16365_at (AC003974.136_AT)        |
| 471 | 16110_s_at (AB004822_S_AT)        | 503 | 16383_at (AC006300.64_AT)         |
| 472 | 16111_f_at (AB007788_F_AT)        | 504 | 16391_at (AL050351.194_AT)        |
| 473 | 16130_s_at (AF078683_S_AT)        | 505 | 16398_s_at (AL022603.3_S_AT)      |
| 474 | 16133_s_at (AF089810_S_AT)        | 506 | 16405_at (AC005850.9_AT)          |
| 475 | 16134_s_at (AF132016_S_AT)        | 507 | 16406_at (AC006921.33_AT)         |
| 476 | 16151_s_at<br>(ATHPRKINA_S_AT)    | 508 | 16409_at (AC004393.2_AT)          |
| 477 | 16153_s_at<br>(ATHRPRP1C_S_AT)    | 509 | 16434_at (AL021711.157_AT)        |
| 478 | 16159_s_at (ATU37697_S_AT)        | 510 | 16440_s_at<br>(AF002109.137_S_AT) |
| 479 | 16161_s_at (ATU39072_S_AT)        | 511 | 16442_s_at (AJ002551.2_S_AT)      |
| 480 | 16164_s_at (ATU47029_S_AT)        | 512 | 16457_s_at<br>(AC005397.17_S_AT)  |

| #   | ProbeSet                      | #   | ProbeSet                      |
|-----|-------------------------------|-----|-------------------------------|
| 513 | 16461_i_at (AC004683.79_I_AT) | 547 | 16649_s_at (ATHORF_S_AT)      |
|     | 16462_s_at                    | 548 | 16701_at (AC005312.61_AT)     |
| 514 | (AC004683.79_S_AT)            | 549 | 16712_at (AC006068.67_AT)     |
| 515 | 16465_at (Y08892.1_AT)        | 550 | 16721_at (AC006533.58_AT)     |
| 516 | 16468_at (AF049870.5_AT)      | 551 | 16747_at (AL021713.3_AT)      |
| 517 | 16470_s_at (AF068299.4_S_AT)  | 552 | 16753_at (AL031032.110_AT)    |
| 518 | 16483_at (X68053_AT)          | 553 | 16781_at (AC002392.100_AT)    |
| 519 | 16496_s_at (AF030386.1_S_AT)  | 554 | 16810_at (AC002339.46_AT)     |
| 520 | 16510_at (AL034567.198_AT)    | 555 | 16817_s_at (AL096882.91_S_AT) |
| 521 | 16522_at (X77500.2_AT)        | 556 | 16859_at (AL035523.135_AT)    |
| 522 | 16524_at (AC006577.38_AT)     | 557 | 16864_i_at (AF037367.4_I_AT)  |
| 523 | 16526_at (Z49227.1_AT)        | 558 | 16865_s_at (AF037367.4_S_AT)  |
| 524 | 16536_s_at (AB008107_S_AT)    | 559 | 16868_at (AL035679.123_AT)    |
| 525 | 16537_s_at (AB008111_S_AT)    |     | 16888_s_at                    |
| 526 | 16538_s_at (AB010259_S_AT)    | 560 | (AC004684.174_S_AT)           |
| 527 | 16539_s_at (AB013301_S_AT)    |     | 16891_s_at                    |
| 528 | 16541_s_at (AB023423_S_AT)    | 561 | (AF080120.33_S_AT)            |
| 529 | 16545_s_at (AF037229_S_AT)    | 562 | 16892_at (U37336.3_AT)        |
| 530 | 16548_s_at (AF054617_S_AT)    | 563 | 16902_at (AC007119.67_AT)     |
| 531 | 16553_f_at (AF078821_F_AT)    |     | 16903_g_at                    |
|     | 16568_s_at                    | 564 | (AC007119.67_G_AT)            |
| 532 | (ATHATCDPK_S_AT)              | 565 | 16908_at (AC002396.22_AT)     |
| 533 | 16570_s_at (ATHCDPKA_S_AT)    | 566 | 16914_s_at (AL049500.57_S_AT) |
| 534 | 16578_s_at (ATHRPRP1B_S_AT)   | 567 | 16916_s_at (X77199.8_S_AT)    |
| 535 | 16589_s_at (ATU26937_S_AT)    | 568 | 16927_s_at (AF035384.2_S_AT)  |
| 536 | 16594_s_at (ATU39783_S_AT)    | 569 | 16939_at (AC002334.110_AT)    |
| 537 | 16603_s_at (ATU81293_S_AT)    |     | 16940_g_at                    |
| 538 | 16609_s_at (AB008104_S_AT)    | 570 | (AC002334.110_G_AT)           |
| 539 | 16610_s_at (AB008490_S_AT)    | 571 | 16943_s_at (Z97339.466_S_AT)  |
| 540 | 16611_s_at (AB008782_S_AT)    | 572 | 16951_i_at (AC005662.30_I_AT) |
| 541 | 16613_s_at (AF012657_S_AT)    | 573 | 16952_s_at (AC005662.30_S_AT) |
| 542 | 16617_s_at (AF029980_S_AT)    | 574 | 16955_at (AL031326.215_AT)    |
| 543 | 16620_s_at (AF051338_S_AT)    | 575 | 16968_at (AL021961.93_AT)     |
| 544 | 16635_s_at (AF126057_S_AT)    | 576 | 16970_s_at (Y18291.5_S_AT)    |
| 545 | 16638_s_at (AF139098_S_AT)    | 577 | 16972_at (AC004261.89_AT)     |
| 546 | 16646_s_at (ATHDHS1_S_AT)     | 578 | 16978_g_at (AF076641.2_G_AT)  |
|     |                               | 579 | 16981_s_at (U35829.2_S_AT)    |

| #   | ProbeSet                     | #   | ProbeSet                     |
|-----|------------------------------|-----|------------------------------|
| 580 | 16989_at (AL030978.46_AT)    | 613 | 17362_s_at (Z97338.181_S_AT) |
| 581 | 16990_at (AC004684.91_AT)    | 614 | 17371_at (AF076243.44_AT)    |
| 582 | 16995_at (AC002391.188_AT)   | 615 | 17376_at (AL021890.218_AT)   |
| 583 | 17007_at (AC005896.26_AT)    | 616 | 17379_at (AF085279.9_AT)     |
| 584 | 17008_at (AC006585.212_AT)   | 617 | 17380_at (AL021961.39_AT)    |
| 585 | 17009_at (AL021633.163_AT)   | 618 | 17381_at (Z99708.402_AT)     |
| 586 | 17018_s_at (ATU18929_S_AT)   | 619 | 17398_at (AC002535.143_AT)   |
| 587 | 17027_s_at (ATU72958_S_AT)   | 620 | 17413_s_at (AJ006961.4_S_AT) |
| 588 | 17039_s_at (D78602_S_AT)     | 621 | 17451_at (AC002343.47_AT)    |
| 589 | 17041_s_at (D89631_S_AT)     |     | 17452_g_at                   |
| 590 | 17051_s_at (AF098947_S_AT)   | 622 | (AC002343.47_G_AT)           |
|     | 17066_s_at                   | 623 | 17458_at (AC006260.91_AT)    |
| 591 | (ATHLIPOXY_S_AT)             | 624 | 17464_at (AC000132.72_AT)    |
| 592 | 17073_s_at (ATTS4391_S_AT)   | 625 | 17477_s_at (X63443.2_S_AT)   |
| 593 | 17075_s_at (ATU09961_S_AT)   | 626 | 17482_s_at (Z97343.441_S_AT) |
| 594 | 17083_s_at (ATU18770_S_AT)   | 627 | 17484_at (X79052.2_AT)       |
| 595 | 17097_s_at (ATU66345_S_AT)   | 628 | 17485_s_at (Z97340.345_S_AT) |
| 596 | 17104_s_at (D88541_S_AT)     | 629 | 17487_s_at (U18993.2_S_AT)   |
| 597 | 17105_s_at (AF055357_S_AT)   | 630 | 17490_s_at (M90416.2_S_AT)   |
| 598 | 17111_s_at (ATHACSC_S_AT)    | 631 | 17494_s_at (ATU30478_S_AT)   |
| 599 | 17119_s_at (AF132212_S_AT)   |     | 17500_s_at                   |
|     | 17128_s_at                   | 632 | (ATHCALLGA_S_AT)             |
| 600 | (ATHRPRP1A_S_AT)             | 633 | 17511_s_at (AF067605_S_AT)   |
| 601 | 17134_at (AC000106.53_AT)    | 634 | 17514_s_at (AF076277_S_AT)   |
| 602 | 17180_at (AF007270.30_AT)    | 635 | 17516_s_at (AF072536_S_AT)   |
| 603 | 17187_at (AF128396.2_AT)     | 636 | 17522_s_at (D78606_S_AT)     |
|     | 17191_i_at                   | 637 | 17533_s_at (ATU43488_S_AT)   |
| 604 | (AL021961.178_I_AT)          | 638 | 17544_s_at (ATU40856_S_AT)   |
| 605 | 17300_at (X66017.2_AT)       | 639 | 17548_s_at (AF118823_S_AT)   |
|     | 17303_s_at                   | 640 | 17555_s_at (ATU89296_S_AT)   |
| 606 | (AC004683.25_S_AT)           | 641 | 17577_g_at (AF087820_G_AT)   |
| 607 | 17305_at (U25649.3_AT)       | 642 | 17578_at (AF093604_AT)       |
| 608 | 17323_at (U95973.69_AT)      | 643 | 17579_s_at (AF093753_S_AT)   |
| 609 | 17338_at (AC002535.97_AT)    | 644 | 17585_s_at (AF134487_S_AT)   |
| 610 | 17341_at (AL021713.89_AT)    | 645 | 17589_at (AF156783_AT)       |
| 611 | 17352_at (AC007127.33_AT)    | 646 | 17595_s_at (AF166352_S_AT)   |
| 612 | 17356_s_at (Z97338.190_S_AT) |     |                              |

| #   | ProbeSet                      | #   | ProbeSet                      |
|-----|-------------------------------|-----|-------------------------------|
| 647 | 17636_at (AF077409.7_AT)      | 679 | 17975_at (AF175998_AT)        |
| 648 | 17648_at (AL021684.43_AT)     | 680 | 17990_at (AF178075_AT)        |
| 649 | 17653_at (AL035679.144_AT)    | 681 | 18010_s_at (AJ001264_S_AT)    |
| 650 | 17702_at (AC005700.212_AT)    | 682 | 18012_s_at (AJ002295_S_AT)    |
| 651 | 17719_at (AC006592.17_AT)     | 683 | 18045_at (AJ011976_AT)        |
| 652 | 17743_at (AC002341.99_AT)     | 684 | 18054_at (AJ238846_AT)        |
|     | 17744_s_at                    |     | 18109_s_at                    |
| 653 | (AC004684.168_S_AT)           | 685 | (AC002391.206_S_AT)           |
| 654 | 17752_at (AC003974.37_AT)     | 686 | 18121_s_at (AC002337.21_S_AT) |
| 655 | 17758_at (AF076243.41_AT)     | 687 | 18122_at (AC002338.110_AT)    |
| 656 | 17775_at (AC004392.2_AT)      | 688 | 18140_at (Z97341.319_AT)      |
| 657 | 17781_at (AL049746.177_AT)    | 689 | 18148_at (AC004669.25_AT)     |
|     | 17823_s_at                    |     | 18167_s_at                    |
| 658 | (AC006555.10_S_AT)            | 690 | (AL021711.23_S_AT)            |
|     | 17840_s_at                    | 691 | 18176_at (AL035540.31_AT)     |
| 659 | (AC002333.223_S_AT)           |     | 18194_i_at                    |
| 660 | 17854_at (Z99707.366_AT)      | 692 | (AL096859.227_I_AT)           |
| 661 | 17860_at (AL078467.4_AT)      | 693 | 18213_at (AL022140.126_AT)    |
| 662 | 17876_at (AJ007587.2_AT)      | 694 | 18215_at (97335.114_AT)       |
| 663 | 17877_g_at (J007587.2_G_AT)   | 695 | 18216_at (X95573.2_AT)        |
| 664 | 17881_at (AC002391.54_AT)     | 696 | 18217_g_at (X95573.2_G_AT)    |
| 665 | 17882_at (AL035523.49_AT)     |     | 18224_s_at                    |
| 666 | 17886_at (AC004484.151_AT)    | 697 | (AL021890.57_S_AT)            |
| 667 | 17893_at (AC004401.135_AT)    |     | 18226_s_at                    |
| 668 | 17894_at (AC005724.44_AT)     | 698 | (AC002343.142_S_AT)           |
| 669 | 17899_at (Z97339.197_AT)      | 699 | 18228_at (X91259.1_AT)        |
|     | 17900_s_at                    | 700 | 18234_at (AC000348.3_AT)      |
| 670 | (AC000106.13_S_AT)            |     | 18236_s_at                    |
|     | 17907_s_at                    | 701 | (AC004683.69_S_AT)            |
| 671 | (AC004684.165_S_AT)           | 702 | 18241_at (AC006580.71_AT)     |
| 672 | 17916_at (U22428.2_AT)        |     | 18242_g_at                    |
| 673 | 17930_s_at (AJ006960.4_S_AT)  | 703 | (AC006580.71_G_AT)            |
| 674 | 17945_at (Z97341.411_AT)      | 704 | 18255_at (AC005770.25_AT)     |
| 675 | 17955_at (AL021768.242_AT)    |     | 18258_s_at                    |
| 676 | 17956_i_at (AC005967.32_I_AT) | 705 | (AC006439.222_S_AT)           |
| 677 | 17963_at (AL049730.88_AT)     | 706 | 18263_at (18263_AT)           |
| 678 | 17967_at (AL096859.32_AT)     | 707 | 18266_at (AC004684.33_AT)     |
|     |                               | 708 | 18267_at (AC006223.23_AT)     |

| #   | ProbeSet                      | #   | ProbeSet                     |
|-----|-------------------------------|-----|------------------------------|
| 709 | 18268_s_at (AC006418.38_S_AT) | 742 | 18681_at (L23573_AT)         |
| 710 | 18280_at (AC007369.2_AT)      | 743 | 18683_s_at (L27158_S_AT)     |
| 711 | 18284_at (AL021961.67_AT)     | 744 | 18686_s_at (U18126_S_AT)     |
| 712 | 18287_at (AC007661.142_AT)    | 745 | 18694_s_at (U89272_S_AT)     |
| 713 | 18299_s_at (M23872.2_S_AT)    | 746 | 18698_s_at (X17528_S_AT)     |
| 714 | 18301_s_at (AL022223.48_S_AT) | 747 | 18716_at (X91916_AT)         |
| 715 | 18314_i_at (AL078579.83_I_AT) | 748 | 18720_s_at (X92419_S_AT)     |
| 716 | 18348_at (AL022603.104_AT)    | 749 | 18735_s_at (Z29490_S_AT)     |
|     | 18456_s_at                    |     | 18753_s_at                   |
| 717 | (AC004697.159_S_AT)           | 750 | (AF118222.28_S_AT)           |
| 718 | 18471_at (AC006533.103_AT)    | 751 | 18782_at (AC003040.90_AT)    |
| 719 | 18508_s_at (AC006532.89_S_AT) | 752 | 18803_at (AC005315.94_AT)    |
| 720 | 18515_at (AC007063.215_AT)    | 753 | 18844_at (AC005315.131_AT)   |
| 721 | 18544_at (AC007060.14_AT)     | 754 | 18866_at (AC005917.178_AT)   |
| 722 | 18582_s_at (AC003671.36_S_AT) | 755 | 18885_at (AC006921.147_AT)   |
| 723 | 18587_s_at (AC007166.53_S_AT) | 756 | 18888_at (AC007591.68_AT)    |
| 724 | 18590_at (AJ222713.4_AT)      | 757 | 18896_at (AC002329.51_AT)    |
| 725 | 18591_at (X74756.2_AT)        | 758 | 18899_s_at (X13434.1_S_AT)   |
| 726 | 18596_at (AC005698.13_AT)     | 759 | 18908_i_at (AF055848.2_I_AT) |
| 727 | 18597_at (AL080282.74_AT)     | 760 | 18909_s_at (AF055848.2_S_AT) |
| 728 | 18600_at (L76926.1_AT)        | 761 | 18916_s_at (X92393.1_S_AT)   |
|     | 18601_s_at                    | 762 | 18928_at (AC002333.181_AT)   |
| 729 | (AC002387.279_S_AT)           | 763 | 18930_at (AC005990.57_AT)    |
| 730 | 18604_at (AF069298.31_AT)     | 764 | 18933_at (AC007020.48_AT)    |
| 731 | 18622_g_at (AJ005902.2_G_AT)  | 765 | 18936_at (AJ003119.4_AT)     |
| 732 | 18625_at (AC005278.22_AT)     | 766 | 18949_at (Z54136.1_AT)       |
| 733 | 18631_at (AC002510.112_AT)    | 767 | 18953_at (AF077955.1_AT)     |
| 734 | 18634_s_at (Z97343.468_S_AT)  | 768 | 18963_at (AC004561.99_AT)    |
| 735 | 18635_at (AC004005.44_AT)     | 769 | 18966_at (AC004561.106_AT)   |
| 736 | 18636_at (AC006577.22_AT)     | 770 | 18976_at (AC000106.31_AT)    |
|     | 18650_s_at                    | 771 | 18980_at (U78721.20_AT)      |
| 737 | (AF013294.25_S_AT)            | 772 | 18984_at (AC003096.100_AT)   |
|     | 18662_s_at                    | 773 | 19017_at (AL035709.69_AT)    |
| 738 | (AC002343.20_S_AT)            | 774 | 19019_i_at (X82623.2_I_AT)   |
| 739 | 18667_at (AJ249442_AT)        | 775 | 19044_at (AC004392.38_AT)    |
| 740 | 18668_at (AJ249794_AT)        | 776 | 19060_at (AC003671.34_AT)    |
| 741 | 18672_s_at (D13983_S_AT)      |     |                              |

| #   | ProbeSet                      | #   | ProbeSet                     |
|-----|-------------------------------|-----|------------------------------|
| 777 | 19092_at (AL078606.188_AT)    | 811 | 19451_at (AC004392.6_AT)     |
| 778 | 19110_s_at (X86947.2_S_AT)    |     | 19460_s_at                   |
|     | 19132_s_at                    | 812 | (AC000132.66_S_AT)           |
| 779 | (AL022603.298_S_AT)           | 813 | 19462_s_at (AF001168.2_S_AT) |
| 780 | 19137_at (X74755.2_AT)        | 814 | 19464_at (AC005560.51_AT)    |
| 781 | 19140_at (AC005170.24_AT)     | 815 | 19465_at (AL021768.96_AT)    |
| 782 | 19150_at (AC006577.20_AT)     | 816 | 19494_at (AC007296.26_AT)    |
| 783 | 19161_at (AL078579.9_AT)      | 817 | 19531_at (AL021960.91_AT)    |
| 784 | 19171_at (AC002335.160_AT)    | 818 | 19546_at (AC005398.172_AT)   |
| 785 | 19178_at (Y18227.2_AT)        | 819 | 19555_at (AF058919.48_AT)    |
| 786 | 19181_s_at (AF053065.2_S_AT)  | 820 | 19591_at (AJ010735.4_AT)     |
| 787 | 19182_at (AL031804.245_AT)    | 821 | 19614_at (AC003970.32_AT)    |
| 788 | 19207_at (AC006069.117_AT)    | 822 | 19623_at (AF000657.40_AT)    |
| 789 | 19219_at (AC007019.185_AT)    | 823 | 19624_at (AL049481.196_AT)   |
| 790 | 19230_at (AC003113.15_AT)     |     | 19625_s_at                   |
| 791 | 19247_at (AF071527.44_AT)     | 824 | (AC002311.26_S_AT)           |
| 792 | 19257_s_at (AC000104.57_S_AT) | 825 | 19635_at (AL049746.38_AT)    |
| 793 | 19284_at (AC003028.196_AT)    | 826 | 19639_at (AL080252.22_AT)    |
| 794 | 19288_at (AC005824.130_AT)    | 827 | 19640_at (AC004561.78_AT)    |
| 795 | 19325_at (AL022604.42_AT)     | 828 | 19641_at (AC004561.66_AT)    |
| 796 | 19364_at (AL022023.142_AT)    | 829 | 19645_at (AC004561.70_AT)    |
| 797 | 19368_at (AC000348.22_AT)     |     | 19646_s_at                   |
| 798 | 19376_at (AF024504.11_AT)     | 830 | (AC005819.55_S_AT)           |
| 799 | 19383_at (AC006200.203_AT)    | 831 | 19655_at (Y14199.1_AT)       |
| 800 | 19386_at (AC006592.51_AT)     | 832 | 19660_at (AC002336.29_AT)    |
| 801 | 19388_at (AC000104.61_AT)     | 833 | 19667_at (AL021710.5_AT)     |
| 802 | 19395_at (AF007270.32_AT)     | 834 | 19672_at (AC005687.19_AT)    |
| 803 | 19396_at (AJ001855.2_AT)      |     | 19673_g_at                   |
| 804 | 19405_at (AJ223803.1_AT)      | 835 | (AC005687.19_G_AT)           |
| 805 | 19407_at (AC004697.81_AT)     |     | 19700_s_at                   |
| 806 | 19409_at (AC007357.56_AT)     | 836 | (AL031326.154_S_AT)          |
| 807 | 19411_at (AC007661.104_AT)    |     | 19701_s_at                   |
| 808 | 19421_at (X70990.4_AT)        | 837 | (AC005724.67_S_AT)           |
| 809 | 19424_at (AC002396.44_AT)     | 838 | 19704_i_at (AJ005927.2_I_AT) |
|     | 19432_s_at                    | 839 | 19707_s_at (Z95768.3_S_AT)   |
| 810 | (AL035680.11_S_AT)            | 840 | 19741_at (AL049171.72_AT)    |
|     |                               | 841 | 19755_at (AC006593.64_AT)    |

| #   | ProbeSet                      |
|-----|-------------------------------|
| 842 | 19762_at (AL035527.204_AT)    |
| 843 | 19818_i_at (AL021749.33_I_AT) |
|     | 19819_s_at                    |
| 844 | (AL021749.33_S_AT)            |
| 845 | 19844_at (AJ007588.2_AT)      |
| 846 | 19845_g_at (AJ007588.2_G_AT)  |
|     | 19848_s_at                    |
| 847 | (AC004261.94_S_AT)            |
| 848 | 19851_at (U23794.3_AT)        |
|     | 19870_s_at                    |
| 849 | (AL021811.48_S_AT)            |
| 850 | 19878_at (AL080252.102_AT)    |
| 851 | 19879_s_at (Z97338.342_S_AT)  |
| 852 | 19881_at (AC004077.49_AT)     |
| 853 | 19892_at (AC005770.30_AT)     |
| 854 | 19894_at (AJ001809.1_AT)      |
| 855 | 19895_s_at (U77347.4_S_AT)    |
| 856 | 19903_at (AC007660.40_AT)     |
| 857 | 19916_at (AC006577.34_AT)     |
| 858 | 19944_at (AC002130.4_AT)      |
| 859 | 19946_at (AC004482.14_AT)     |
| 860 | 19956_at (AC006282.11_AT)     |
| 861 | 19960_at (AL035527.360_AT)    |
|     | 19970_s_at                    |
| 862 | (AC003674.10_S_AT)            |
| 863 | 19982_at (AC002986.28_AT)     |
| 864 | 19991_at (AC007017.124_AT)    |
| 865 | 20017_at (AC004521.66_AT)     |
| 866 | 20023_at (AC006577.46_AT)     |
| 867 | 20030_at (AL078637.51_AT)     |
| 868 | 20051_at (AC000106.38_AT)     |
| 869 | 20053_at (AC002292.27_AT)     |
| 870 | 20061_at (AC005508.23_AT)     |
| 871 | 20096_at (AC004238.31_AT)     |
| 872 | 20098_at (AC004697.123_AT)    |
| 873 | 20133_i_at (AC007178.71_I_AT) |
| 874 | 20134_s_at (AC007178.71_S_AT) |

| #   | ProbeSet                      |
|-----|-------------------------------|
| 875 | 20142_at (AL035521.155_AT)    |
| 876 | 20144_at (AL079350.68_AT)     |
| 877 | 20165_at (AC002311.16_AT)     |
| 878 | 20179_at (AL035538.229_AT)    |
| 879 | 20189_at (AC005489.2_AT)      |
| 880 | 20194_at (AC007584.48_AT)     |
| 881 | 20199_at (AL050300.89_AT)     |
| 882 | 20200_at (AL050400.67_AT)     |
| 883 | 20215_s_at (AF117125.2_S_AT)  |
| 884 | 20223_at (AL022347.145_AT)    |
| 885 | 20238_at (X74514.2_AT)        |
| 886 | 20239_g_at (X74514.2_G_AT)    |
|     | 20245_s_at                    |
| 887 | (AC005309.97_S_AT)            |
| 888 | 20246_s_at (AF084037.3_S_AT)  |
| 889 | 20247_at (AC004392.4_AT)      |
| 890 | 20258_at (AF130252.1_AT)      |
| 891 | 20262_at (AC002294.26_AT)     |
| 892 | 20263_at (AB004798.1_AT)      |
| 893 | 20269_at (AC002387.237_AT)    |
| 894 | 20271_at (Z99707.27_AT)       |
|     | 20285_s_at                    |
| 895 | (AC003674.18_S_AT)            |
| 896 | 20287_at (Y14590.5_AT)        |
| 897 | 20288_g_at (Y14590.5_G_AT)    |
| 898 | 20291_s_at (M92353.4_S_AT)    |
| 899 | 20297_at (AC007153.27_AT)     |
| 900 | 20323_at (AC004561.62_AT)     |
| 901 | 20335_s_at (Y14208.2_S_AT)    |
| 902 | 20345_at (AF104919.16_AT)     |
| 903 | 20346_at (L031135.156_AT)     |
| 904 | 20348_at (AC005967.35_AT)     |
| 905 | 20356_at (AC004561.74_AT)     |
| 906 | 20365_s_at (AC005850.19_S_AT) |
| 907 | 20370_at (AC004561.263_AT)    |
| 908 | 20372_at (AL021713.24_AT)     |



| #   | ProbeSet                          |
|-----|-----------------------------------|
| 909 | 20382_s_at<br>(AC002338.35_S_AT)  |
| 910 | 20409_g_at<br>(AC004077.132_G_AT) |
| 911 | 20420_at (AL024486.131_AT)        |
| 912 | 20421_at (U81294.2_AT)            |
| 913 | 20422_g_at (U81294.2_G_AT)        |
| 914 | 20432_at (U43486.2_AT)            |
| 915 | 20433_at (AC006232.147_AT)        |
| 916 | 20442_i_at (AC006341.42_I_AT)     |
| 917 | 20443_s_at (AC006341.42_S_AT)     |
| 918 | 20450_at (AJ005930.2_AT)          |
| 919 | 20461_at (20461_AT)               |
| 920 | 20462_at (U82399.2_AT)            |
| 921 | 20477_at (AC004238.154_AT)        |
| 922 | 20479_i_at (AF069495.2_I_AT)      |
| 923 | 20480_s_at (AF069495.2_S_AT)      |
| 924 | 20485_at (AC007660.131_AT)        |
| 925 | 20491_at (AC004561.146_AT)        |
| 926 | 20507_at (AL021635.67_AT)         |
| 927 | 20511_at (AC007290.24_AT)         |
| 928 | 20516_at (AL035523.64_AT)         |
| 929 | 20517_at (Y17722.7_AT)            |
| 930 | 20524_at (AC005698.12_AT)         |
| 931 | 20529_at (Z97341.125_AT)          |
| 932 | 20551_at (AC006081.211_AT)        |

| #   | ProbeSet                          |
|-----|-----------------------------------|
| 933 | 20572_s_at<br>(AC005560.229_S_AT) |
| 934 | 20577_at (AL078464.72_AT)         |
| 935 | 20584_at (AC004450.75_AT)         |
| 936 | 20586_i_at<br>(AC005824.195_I_AT) |
| 937 | 20587_s_at<br>(AC005824.195_S_AT) |
| 938 | 20589_at (AF081066.3_AT)          |
| 939 | 20590_at (AL035540.159_AT)        |
| 940 | 20591_at (AL080252.115_AT)        |
| 941 | 20619_at (AC005896.161_AT)        |
| 942 | 20620_g_at<br>(AC005896.161_G_AT) |
| 943 | 20646_at (AC002291.20_AT)         |
| 944 | 20656_at (AL035396.46_AT)         |
| 945 | 20658_s_at<br>(2AL050400.70_S_AT) |
| 946 | 20660_s_at (X97488.2_S_AT)        |
| 947 | 20669_s_at (AC002388.6_S_AT)      |
| 948 | 20675_at (AC006234.204_AT)        |
| 949 | 20678_at (AC007296.30_AT)         |
| 950 | 20685_at (AL049751.46_AT)         |
| 951 | 20686_at (Y14424.2_AT)            |
| 952 | 20689_s_at<br>(AC002335.19_S_AT)  |
| 953 | 20715_at (AF079183.1_AT)          |

**Table 20a:** cDNA-AFLP gene fragments with similarity to other known proteins

| cDNA***    | Known protein                               | Organism    | Related accession number** | Blast score* |
|------------|---|-------------|----------------------------|--------------|
| DESCA1     | no significant similarity                   | -           | -                          | -            |
| DESCA2     | ser/thr kinase                              | bean        | AF078082                   | 2e-12        |
| DESCA3     | endo-1,4-betaglucanase                      | rape        | AJ242807                   | 4e-19        |
| DESCA4     | pdr 5-abc transporter                       | duckweed    | Z70524                     | 1e-27        |
| DESCA5     | transcriptional regulator                   | yeast       | NP 014933                  | 9e-1         |
| DESCA6     | kinase                                      | Arabidopsis | T00502                     | 7e-1         |
| DESCA7     | salicylate-induced glucosyltransferase IS5a | tobacco     | T03747                     | 2e-21        |
| DESCA8     | nbs-LRR                                     | rice        | AAF82158                   | 5e-2         |
| DESCA9     | cytochrome p450 monooxygenase               | tobacco     | X96784                     | 7e-31        |
| DESCA10    | MRP-like ABC transporter                    | Arabidopsis | U96399                     | 8e-7         |
| DESCA11    | phosphoribosylanthranilate transferase      | Arabidopsis | AAF18518                   | 8e-17        |
| DESCA12    | hypersensitivity-related 201                | tobacco     | X95343                     | 3e-23        |
| DESCA13    | integral membrane glycoprotein              | puffer fish | AF013613                   | 9e-1         |
| 7a tgaa    | no significant similarity                   | -           | -                          | -            |
| 10d tcg    | hypothetical protein F3F9.18                | Arabidopsis | AC013430                   | 7e-15        |
| 11a tgca   | no significant similarity                   | -           | -                          | -            |
| c.r. actin | actin                                       | C. rubrum   | X92353                     | 7e-45        |

\* NCBI BLASTX translated search (Altschul et al., 1997)

\*\* Accession numbers are listed in SEQ ID Nos 1967-1980

\*\*\* DESCA sequences are listed in SEQ ID Nos 1954-1966

**Table 20b:** Relative gene expression fold changes of cDNA-AFLP gene fragments with similarity to other known proteins at different time points during viral infections as determined by quantitative RT-PCR

| cDNA       | Gene expression fold changes |      |       |                  |                |
|------------|------------------------------|------|-------|------------------|----------------|
|            | <i>C. amaranticolor</i>      |      |       | <i>C. quinoa</i> |                |
|            | TMV <sup>1</sup>             |      |       | TRV <sup>2</sup> |                |
|            | 4dai                         | 7dai | 11dai | 4dai             | 4dai           |
| DESCA1     | 200                          | 180  | 6.1   | 278              | not detectable |
| DESCA2     | 36                           | 25   | 27    | 10               | 700            |
| DESCA3     | 23                           | 15   | 2.7   | 19               | 46             |
| DESCA4     | 21                           | 9.2  | 4.5   | 6.8              | 52             |
| DESCA5     | 19                           | 8.0  | 9.1   | 15               | 1100           |
| DESCA6     | 8.4                          | 9.6  | 12    | 2.1              | not detectable |
| DESCA7     | 8.9                          | 2.8  | 5.6   | 53               | 150            |
| DESCA8     | 5.9                          | 3.2  | 2.0   | 5.8              | 120            |
| DESCA9     | 5.1                          | 5.9  | 4.2   | 3.1              | not detectable |
| DESCA10    | 5.6                          | 1.9  | 0.95  | 4.1              | not detectable |
| DESCA11    | 5.5                          | 3.6  | 3.0   | 3.9              | 230            |
| DESCA12    | 5.0                          | 2.4  | 1.7   | 5.6              | 7.9            |
| DESCA13    | 2.9                          | 1.8  | 1.9   | 3.3              | 34             |
| 7a tgaa    | 1.5                          | 1.4  | 1.7   | not tested       | not tested     |
| 10d tcg    | 2.2                          | 0.75 | 0.25  | not tested       | not tested     |
| 11a tgca   | 1.6                          | 0.54 | 0.18  | not tested       | not tested     |
| c.r. actin | 1.0                          | 1.0  | 1.0   | 1.0              | 1.0            |

<sup>1</sup> Values are the fold increases in gene expression of TMV-MGfus infected compared to mock-inoculated plants.

<sup>2</sup> Values are the fold increases in gene expression of TRV infected compared to mock-inoculated plants.

<sup>3</sup> Values are the fold increases in gene expression of TMV infected compared to mock-inoculated plants.

**Table 21** Known plant/pathogen interactions

| Plant        | Pathogen                      |
|--------------|-------------------------------|
| Tomato       | Cladosporium fulvum           |
| Maize        | Rust fungus                   |
| Antirrhinium | Rust fungus                   |
| Flax         | <i>Melampsora lini</i>        |
| Lettuce      | Downy mildew                  |
| Arabidopsis  | <i>Peronospora parasitica</i> |
| Tomato       | Nematode                      |
| Corn         | Cochliobolus carbonum         |
| Tomato       | Pseudomonas syringae          |
| Rice         | Xanthomonas oryzae pv. Oryzae |
| Rice         | Pyricularia oryzae            |
| Tobacco      | Tobacco Mosaic Virus          |

**Table 22** Rice Open Reading Frames (ORFs) orthologous to specific Arabidopsis ORFs and corresponding rice promoter regions

AORF ..... SEQ ID NO of a specific Arabidopsis ORF

RH ..... SEQ ID NO of homologous rice ORF(s)

RP ..... SEQ ID NO of the rice promoter linked to a homologous rice ORF

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
| 3    | 2808 | 4884 |      | 3475 | 5551 |      | 4307 | 6383 |
|      | 3066 | 5142 |      | 3567 | 5643 |      | 4458 | 6534 |
|      | 3820 | 5896 |      | 3632 | 5708 | 52   | 3035 | 5111 |
|      | 4655 | 6731 |      | 3769 | 5845 |      | 3157 | 5233 |
| 6    | 2757 | 4833 |      | 3916 | 5992 |      | 3379 | 5455 |
|      | 3013 | 5089 |      | 4298 | 6374 |      | 3626 | 5702 |
|      | 3393 | 5469 | 19   | 3184 | 5260 |      | 3990 | 6066 |
|      | 3899 | 5975 | 20   | 2678 | 4754 | 55   | 2687 | 4763 |
| 7    | 3227 | 5303 |      | 2926 | 5002 |      | 2869 | 4945 |
|      | 3579 | 5655 |      | 3736 | 5812 |      | 2984 | 5060 |
|      | 3851 | 5927 |      | 3864 | 5940 |      | 3054 | 5130 |
|      | 3890 | 5966 |      | 4124 | 6200 |      | 3180 | 5256 |
|      | 4516 | 6592 |      | 4414 | 6490 |      | 3561 | 5637 |
| 8    | 3086 | 5162 | 22   | 2805 | 4881 |      | 4306 | 6382 |
|      | 3341 | 5417 |      | 3997 | 6073 |      | 4326 | 6402 |
|      | 3762 | 5838 |      | 4126 | 6202 |      | 4460 | 6536 |
|      | 3767 | 5843 |      | 4153 | 6229 |      | 4476 | 6552 |
|      | 4285 | 6361 |      | 4311 | 6387 |      | 4521 | 6597 |
|      | 4487 | 6563 |      | 4639 | 6715 |      | 4603 | 6679 |
|      | 4488 | 6564 | 32   | 3718 | 5794 | 57   | 4005 | 6081 |
|      | 4489 | 6565 |      | 4345 | 6421 | 59   | 2922 | 4998 |
| 10   | 3629 | 5705 | 37   | 4493 | 6569 |      | 3833 | 5909 |
|      | 3914 | 5990 | 42   | 3479 | 5555 |      | 3967 | 6043 |
|      | 4291 | 6367 |      | 4208 | 6284 |      | 4202 | 6278 |
|      | 4424 | 6500 |      | 4236 | 6312 | 60   | 2706 | 4782 |
| 11   | 3996 | 6072 | 46   | 3891 | 5967 |      | 4125 | 6201 |
|      | 4555 | 6631 | 49   | 4645 | 6721 |      | 4720 | 6796 |
| 14   | 2809 | 4885 | 50   | 4210 | 6286 | 65   | 2815 | 4891 |
|      | 3160 | 5236 | 51   | 2864 | 4940 |      | 4112 | 6188 |
|      | 3201 | 5277 |      | 2964 | 5040 |      | 4288 | 6364 |
|      | 3455 | 5531 |      | 4218 | 6294 | 67   | 2716 | 4792 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 2928 | 5004 | 75   | 2920 | 4996 |      | 3198 | 5274 |
|      | 2940 | 5016 |      | 3417 | 5493 |      | 3319 | 5395 |
|      | 3182 | 5258 |      | 3742 | 5818 |      | 3797 | 5873 |
|      | 3333 | 5409 |      | 3927 | 6003 |      | 4048 | 6124 |
|      | 3351 | 5427 |      | 4061 | 6137 |      | 4184 | 6260 |
|      | 3386 | 5462 | 78   | 2705 | 4781 |      | 4259 | 6335 |
|      | 3387 | 5463 |      | 2737 | 4813 | 102  | 3563 | 5639 |
|      | 3445 | 5521 |      | 2738 | 4814 |      | 3683 | 5759 |
|      | 3530 | 5606 |      | 4654 | 6730 |      | 4108 | 6184 |
|      | 3581 | 5657 |      | 4681 | 6757 |      | 4168 | 6244 |
|      | 3727 | 5803 | 85   | 3163 | 5239 |      | 4232 | 6308 |
|      | 3825 | 5901 |      | 3525 | 5601 | 110  | 2876 | 4952 |
|      | 3915 | 5991 |      | 3965 | 6041 |      | 3099 | 5175 |
|      | 4120 | 6196 |      | 4114 | 6190 |      | 3367 | 5443 |
|      | 4170 | 6246 | 91   | 3138 | 5214 |      | 3610 | 5686 |
|      | 4267 | 6343 |      | 3745 | 5821 |      | 4363 | 6439 |
|      | 4280 | 6356 |      | 3803 | 5879 |      | 4618 | 6694 |
|      | 4350 | 6426 |      | 4233 | 6309 |      | 4656 | 6732 |
|      | 4479 | 6555 |      | 4275 | 6351 | 111  | 2747 | 4823 |
|      | 4713 | 6789 | 93   | 4325 | 6401 |      | 2766 | 4842 |
| 68   | 2764 | 4840 | 95   | 2903 | 4979 |      | 2955 | 5031 |
|      | 3072 | 5148 |      | 3230 | 5306 |      | 3450 | 5526 |
|      | 3192 | 5268 |      | 3676 | 5752 |      | 3988 | 6064 |
|      | 3290 | 5366 |      | 4480 | 6556 | 112  | 2779 | 4855 |
|      | 3546 | 5622 |      | 4631 | 6707 |      | 3345 | 5421 |
|      | 3553 | 5629 | 97   | 2959 | 5035 |      | 3799 | 5875 |
|      | 3658 | 5734 |      | 3105 | 5181 |      | 3801 | 5877 |
|      | 3747 | 5823 |      | 3222 | 5298 |      | 4216 | 6292 |
|      | 3972 | 6048 |      | 3838 | 5914 | 113  | 3371 | 5447 |
|      | 4543 | 6619 |      | 4026 | 6102 |      | 3372 | 5448 |
| 70   | 3173 | 5249 |      | 4224 | 6300 |      | 3373 | 5449 |
| 72   | 3245 | 5321 |      | 4523 | 6599 |      | 4022 | 6098 |
|      | 4329 | 6405 |      | 4646 | 6722 |      | 4319 | 6395 |
|      | 4412 | 6488 | 100  | 2667 | 4743 | 115  | 2707 | 4783 |
| 73   | 2862 | 4938 |      | 2751 | 4827 |      | 2871 | 4947 |
|      | 3012 | 5088 |      | 2948 | 5024 |      | 3261 | 5337 |
|      | 3506 | 5582 |      | 3167 | 5243 |      | 3452 | 5528 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 4513 | 6589 |      | 4496 | 6572 |      | 3761 | 5837 |
| 116  | 2833 | 4909 |      | 4593 | 6669 |      | 3869 | 5945 |
|      | 2939 | 5015 |      | 4629 | 6705 |      | 4609 | 6685 |
|      | 3200 | 5276 | 127  | 2981 | 5057 | 137  | 2723 | 4799 |
|      | 3571 | 5647 |      | 3199 | 5275 |      | 3220 | 5296 |
|      | 4031 | 6107 |      | 3467 | 5543 |      | 4173 | 6249 |
| 118  | 2944 | 5020 |      | 3963 | 6039 | 138  | 3024 | 5100 |
|      | 3186 | 5262 |      | 4053 | 6129 |      | 3487 | 5563 |
|      | 3631 | 5707 | 128  | 3267 | 5343 |      | 3583 | 5659 |
|      | 3633 | 5709 |      | 4182 | 6258 |      | 4503 | 6579 |
|      | 3697 | 5773 |      | 4530 | 6606 |      | 4591 | 6667 |
|      | 3726 | 5802 | 130  | 2890 | 4966 | 139  | 2699 | 4775 |
|      | 4009 | 6085 |      | 3293 | 5369 |      | 3463 | 5539 |
|      | 4597 | 6673 |      | 3312 | 5388 |      | 3584 | 5660 |
| 119  | 3169 | 5245 |      | 3326 | 5402 |      | 4451 | 6527 |
|      | 3637 | 5713 |      | 3812 | 5888 |      | 4595 | 6671 |
|      | 3638 | 5714 |      | 3889 | 5965 | 140  | 3042 | 5118 |
|      | 3639 | 5715 |      | 4134 | 6210 |      | 3175 | 5251 |
|      | 3656 | 5732 |      | 4254 | 6330 |      | 4337 | 6413 |
| 121  | 4692 | 6768 | 133  | 2799 | 4875 | 148  | 3558 | 5634 |
| 122  | 3041 | 5117 |      | 3008 | 5084 |      | 3655 | 5731 |
|      | 3179 | 5255 |      | 3208 | 5284 |      | 4084 | 6160 |
|      | 3291 | 5367 |      | 3706 | 5782 | 150  | 3107 | 5183 |
|      | 4269 | 6345 |      | 3765 | 5841 |      | 3560 | 5636 |
|      | 4633 | 6709 | 134  | 3005 | 5081 | 151  | 4146 | 6222 |
| 124  | 3388 | 5464 |      | 3145 | 5221 |      | 4398 | 6474 |
|      | 3410 | 5486 |      | 3146 | 5222 |      | 4399 | 6475 |
|      | 3469 | 5545 |      | 3328 | 5404 | 152  | 4368 | 6444 |
|      | 4316 | 6392 |      | 3511 | 5587 |      | 4483 | 6559 |
|      | 4449 | 6525 |      | 4572 | 6648 |      | 4550 | 6627 |
| 126  | 3215 | 5291 |      | 4584 | 6660 |      | 4551 | 6628 |
|      | 3557 | 5633 | 135  | 2832 | 4908 |      | 4616 | 6692 |
|      | 3575 | 5651 |      | 3073 | 5149 | 158  | 3023 | 5099 |
|      | 3847 | 5923 |      | 3257 | 5333 |      | 3880 | 5956 |
|      | 4044 | 6120 |      | 3352 | 5428 |      | 4237 | 6313 |
|      | 4400 | 6476 |      | 3378 | 5454 |      | 4492 | 6568 |
|      | 4444 | 6520 |      | 3555 | 5631 | 159  | 2720 | 4796 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 3115 | 5191 | 172  | 3389 | 5465 | 189  | 2778 | 4854 |
|      | 3383 | 5459 |      | 3509 | 5585 |      | 3514 | 5590 |
|      | 3844 | 5920 |      | 3603 | 5679 |      | 4662 | 6738 |
|      | 3989 | 6065 |      | 3664 | 5740 | 190  | 3020 | 5096 |
|      | 4157 | 6233 |      | 3665 | 5741 |      | 3088 | 5164 |
|      | 4627 | 6703 | 173  | 4608 | 6684 |      | 3605 | 5681 |
| 162  | 2813 | 4889 | 175  | 3754 | 5830 |      | 3606 | 5682 |
|      | 2927 | 5003 |      | 3835 | 5911 |      | 3992 | 6068 |
|      | 3346 | 5422 |      | 4015 | 6091 | 199  | 2873 | 4949 |
|      | 3423 | 5499 |      | 4198 | 6274 |      | 2942 | 5018 |
|      | 3682 | 5758 |      | 4420 | 6496 |      | 3492 | 5568 |
|      | 3781 | 5857 | 179  | 2684 | 4760 |      | 3671 | 5747 |
|      | 3830 | 5906 |      | 4394 | 6470 |      | 3856 | 5932 |
|      | 4225 | 6301 | 180  | 3158 | 5234 |      | 4030 | 6106 |
|      | 4475 | 6551 |      | 3635 | 5711 |      | 4100 | 6176 |
|      | 4622 | 6698 |      | 4557 | 6633 |      | 4253 | 6329 |
| 164  | 3232 | 5308 |      | 4558 | 6634 |      | 4680 | 6756 |
|      | 3491 | 5567 | 181  | 2696 | 4772 | 202  | 2711 | 4787 |
|      | 4590 | 6666 |      | 3850 | 5926 |      | 3015 | 5091 |
|      | 4625 | 6701 |      | 4549 | 6625 |      | 3774 | 5850 |
| 165  | 3271 | 5347 | 185  | 3110 | 5186 |      | 3906 | 5982 |
|      | 3322 | 5398 |      | 3304 | 5380 |      | 3943 | 6019 |
|      | 3406 | 5482 |      | 3518 | 5594 | 203  | 3135 | 5211 |
|      | 3681 | 5757 |      | 3594 | 5670 |      | 3318 | 5394 |
|      | 3973 | 6049 |      | 3853 | 5929 |      | 3443 | 5519 |
|      | 4231 | 6307 | 187  | 2730 | 4806 |      | 3523 | 5599 |
|      | 4324 | 6400 |      | 2893 | 4969 |      | 3911 | 5987 |
| 166  | 2767 | 4843 |      | 3087 | 5163 |      | 3924 | 6000 |
|      | 2781 | 4857 |      | 3176 | 5252 |      | 4596 | 6672 |
|      | 3347 | 5423 |      | 4234 | 6310 | 204  | 2851 | 4927 |
|      | 4093 | 6169 | 188  | 3068 | 5144 |      | 2852 | 4928 |
|      | 4130 | 6206 |      | 3082 | 5158 |      | 3151 | 5227 |
| 171  | 2733 | 4809 |      | 3644 | 5720 |      | 3152 | 5228 |
|      | 2761 | 4837 |      | 4344 | 6420 |      | 4186 | 6262 |
|      | 3049 | 5125 |      | 4427 | 6503 | 205  | 3078 | 5154 |
|      | 3402 | 5478 |      | 4468 | 6544 |      | 3984 | 6060 |
|      | 3945 | 6021 |      | 4636 | 6712 |      | 4085 | 6161 |



| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
| 206  | 3051 | 5127 | 219  | 2789 | 4865 | 247  | 3971 | 6047 |
|      | 3075 | 5151 |      | 2801 | 4877 | 250  | 2971 | 5047 |
|      | 3117 | 5193 |      | 2802 | 4878 |      | 3696 | 5772 |
|      | 3527 | 5603 |      | 2803 | 4879 |      | 3719 | 5795 |
|      | 3756 | 5832 |      | 3703 | 5779 |      | 4035 | 6111 |
| 209  | 3091 | 5167 |      | 4222 | 6298 |      | 4273 | 6349 |
|      | 3505 | 5581 | 220  | 2885 | 4961 |      | 4675 | 6751 |
|      | 3602 | 5678 |      | 3394 | 5470 | 252  | 2909 | 4985 |
|      | 3921 | 5997 |      | 3395 | 5471 |      | 3738 | 5814 |
|      | 4679 | 6755 |      | 3396 | 5472 |      | 3831 | 5907 |
| 210  | 3628 | 5704 |      | 3499 | 5575 |      | 3904 | 5980 |
|      | 4129 | 6205 | 221  | 2918 | 4994 |      | 4039 | 6115 |
| 212  | 2736 | 4812 |      | 3225 | 5301 |      | 4219 | 6295 |
|      | 2966 | 5042 |      | 4453 | 6529 |      | 4246 | 6322 |
|      | 3888 | 5964 |      | 4534 | 6610 |      | 4467 | 6543 |
|      | 4240 | 6316 |      | 4553 | 6630 |      | 4495 | 6571 |
|      | 4484 | 6560 | 225  | 3634 | 5710 | 254  | 2771 | 4847 |
| 213  | 2822 | 4898 |      | 3673 | 5749 |      | 2874 | 4950 |
|      | 2916 | 4992 |      | 3920 | 5996 |      | 3413 | 5489 |
|      | 2961 | 5037 |      | 4440 | 6516 |      | 3510 | 5586 |
|      | 3102 | 5178 |      | 4668 | 6744 |      | 4143 | 6219 |
|      | 3255 | 5331 | 229  | 2731 | 4807 | 255  | 2664 | 4740 |
| 215  | 2709 | 4785 |      | 3466 | 5542 |      | 2740 | 4816 |
|      | 2804 | 4880 |      | 3897 | 5973 |      | 2792 | 4868 |
|      | 2993 | 5069 |      | 4395 | 6471 |      | 3171 | 5247 |
|      | 3174 | 5250 |      | 4664 | 6740 |      | 3368 | 5444 |
|      | 3852 | 5928 | 237  | 2847 | 4923 |      | 3425 | 5501 |
| 216  | 2662 | 4738 |      | 3126 | 5202 |      | 3788 | 5864 |
|      | 2695 | 4771 |      | 3237 | 5313 |      | 4183 | 6259 |
|      | 3457 | 5533 |      | 3508 | 5584 |      | 4206 | 6282 |
|      | 3504 | 5580 |      | 3704 | 5780 |      | 4439 | 6515 |
|      | 3842 | 5918 |      | 3930 | 6006 | 256  | 2798 | 4874 |
|      | 4580 | 6656 |      | 4006 | 6082 |      | 3109 | 5185 |
|      | 4583 | 6659 | 239  | 2924 | 5000 |      | 3566 | 5642 |
| 218  | 3338 | 5414 |      | 4707 | 6783 |      | 3901 | 5977 |
|      | 3951 | 6027 | 240  | 3211 | 5287 |      | 4190 | 6266 |
|      | 4051 | 6127 |      | 3751 | 5827 |      | 4501 | 6577 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 4579 | 6655 |      | 4333 | 6409 |      | 3953 | 6029 |
|      | 4601 | 6677 |      | 4377 | 6453 |      | 3966 | 6042 |
| 258  | 2952 | 5028 | 271  | 3979 | 6055 |      | 4025 | 6101 |
|      | 3283 | 5359 |      | 4570 | 6646 |      | 4161 | 6237 |
|      | 4001 | 6077 |      | 4571 | 6647 |      | 4544 | 6620 |
|      | 4040 | 6116 | 274  | 3426 | 5502 | 289  | 2668 | 4744 |
|      | 4397 | 6473 |      | 3534 | 5610 |      | 2746 | 4822 |
| 260  | 3438 | 5514 | 275  | 2951 | 5027 |      | 3147 | 5223 |
|      | 3886 | 5962 |      | 3268 | 5344 |      | 3284 | 5360 |
|      | 4070 | 6146 |      | 3909 | 5985 |      | 3285 | 5361 |
|      | 4215 | 6291 |      | 4277 | 6353 |      | 3878 | 5954 |
|      | 4703 | 6779 |      | 4490 | 6566 |      | 4373 | 6449 |
| 263  | 2752 | 4828 | 285  | 3336 | 5412 |      | 4457 | 6533 |
|      | 3098 | 5174 |      | 3987 | 6063 | 290  | 2780 | 4856 |
|      | 3276 | 5352 |      | 4431 | 6507 |      | 2896 | 4972 |
|      | 3625 | 5701 | 286  | 3430 | 5506 |      | 3569 | 5645 |
|      | 4309 | 6385 | 287  | 2703 | 4779 |      | 4486 | 6562 |
| 265  | 2701 | 4777 |      | 3028 | 5104 |      | 4731 | 6807 |
|      | 2861 | 4937 |      | 3329 | 5405 | 295  | 2785 | 4861 |
|      | 3164 | 5240 |      | 3595 | 5671 |      | 2807 | 4883 |
|      | 4187 | 6263 |      | 3667 | 5743 |      | 2921 | 4997 |
|      | 4270 | 6346 |      | 3895 | 5971 |      | 2957 | 5033 |
| 267  | 2866 | 4942 |      | 4598 | 6674 |      | 3281 | 5357 |
|      | 2941 | 5017 |      | 4602 | 6678 |      | 3303 | 5379 |
|      | 3002 | 5078 |      | 4688 | 6764 |      | 3477 | 5553 |
|      | 4330 | 6406 | 288  | 2820 | 4896 |      | 3792 | 5868 |
| 270  | 2875 | 4951 |      | 2977 | 5053 |      | 3925 | 6001 |
|      | 3001 | 5077 |      | 2991 | 5067 |      | 3946 | 6022 |
|      | 3061 | 5137 |      | 3007 | 5083 |      | 4175 | 6251 |
|      | 3282 | 5358 |      | 3057 | 5133 | 296  | 2689 | 4765 |
|      | 3590 | 5666 |      | 3097 | 5173 |      | 2995 | 5071 |
|      | 3608 | 5684 |      | 3155 | 5231 |      | 2996 | 5072 |
|      | 3618 | 5694 |      | 3170 | 5246 |      | 3133 | 5209 |
|      | 3817 | 5893 |      | 3464 | 5540 |      | 4058 | 6134 |
|      | 3969 | 6045 |      | 3528 | 5604 |      | 4185 | 6261 |
|      | 4211 | 6287 |      | 3532 | 5608 |      | 4695 | 6771 |
|      | 4265 | 6341 |      | 3872 | 5948 | 298  | 2685 | 4761 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 2769 | 4845 |      | 3256 | 5332 |      | 4381 | 6457 |
|      | 2777 | 4853 |      | 3385 | 5461 |      | 4430 | 6506 |
|      | 3122 | 5198 |      | 3539 | 5615 |      | 4533 | 6609 |
|      | 3153 | 5229 |      | 3541 | 5617 | 331  | 2760 | 4836 |
|      | 3187 | 5263 |      | 4145 | 6221 |      | 2848 | 4924 |
|      | 3323 | 5399 | 314  | 2839 | 4915 |      | 3103 | 5179 |
|      | 4542 | 6618 |      | 3240 | 5316 |      | 3419 | 5495 |
| 299  | 2915 | 4991 |      | 3485 | 5561 |      | 3740 | 5816 |
|      | 3705 | 5781 |      | 3643 | 5719 |      | 3999 | 6075 |
|      | 4313 | 6389 |      | 3848 | 5924 |      | 4023 | 6099 |
|      | 4353 | 6429 |      | 3957 | 6033 |      | 4352 | 6428 |
|      | 4415 | 6491 |      | 4406 | 6482 |      | 4735 | 6811 |
|      | 4600 | 6676 |      | 4514 | 6590 | 333  | 2796 | 4872 |
|      | 4721 | 6797 | 316  | 3119 | 5195 |      | 2958 | 5034 |
| 300  | 3380 | 5456 |      | 3142 | 5218 |      | 3191 | 5267 |
|      | 3381 | 5457 |      | 3251 | 5327 |      | 3365 | 5441 |
|      | 3456 | 5532 |      | 3274 | 5350 |      | 3687 | 5763 |
|      | 3755 | 5831 |      | 3619 | 5695 |      | 3932 | 6008 |
|      | 4027 | 6103 | 318  | 2972 | 5048 |      | 4342 | 6418 |
| 301  | 2679 | 4755 |      | 3875 | 5951 |      | 4360 | 6436 |
|      | 2865 | 4941 |      | 4152 | 6228 | 339  | 3011 | 5087 |
|      | 3565 | 5641 |      | 4375 | 6451 |      | 3296 | 5372 |
|      | 4029 | 6105 |      | 4461 | 6537 |      | 3624 | 5700 |
|      | 4272 | 6348 | 319  | 4197 | 6273 |      | 3948 | 6024 |
| 302  | 2836 | 4912 |      | 4334 | 6410 |      | 4620 | 6696 |
|      | 3236 | 5312 | 324  | 3033 | 5109 |      | 4733 | 6809 |
| 304  | 2960 | 5036 |      | 3771 | 5847 | 341  | 3248 | 5324 |
|      | 3111 | 5187 |      | 3910 | 5986 |      | 3942 | 6018 |
|      | 3252 | 5328 | 326  | 2934 | 5010 | 346  | 4299 | 6375 |
|      | 3884 | 5960 |      | 3114 | 5190 | 348  | 3000 | 5076 |
|      | 3885 | 5961 |      | 3806 | 5882 |      | 3137 | 5213 |
| 305  | 2899 | 4975 |      | 4359 | 6435 |      | 3866 | 5942 |
|      | 3074 | 5150 |      | 4520 | 6596 |      | 4038 | 6114 |
|      | 3431 | 5507 | 330  | 3401 | 5477 |      | 4327 | 6403 |
|      | 3713 | 5789 |      | 3804 | 5880 |      | 4425 | 6501 |
|      | 4008 | 6084 |      | 4098 | 6174 | 360  | 2989 | 5065 |
| 312  | 2721 | 4797 |      | 4331 | 6407 |      | 2990 | 5066 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 3090 | 5166 |      | 4546 | 6622 | 391  | 2671 | 4747 |
|      | 3100 | 5176 |      | 4587 | 6663 |      | 3862 | 5938 |
|      | 3436 | 5512 | 370  | 4056 | 6132 |      | 3863 | 5939 |
|      | 3515 | 5591 | 372  | 3414 | 5490 |      | 3955 | 6031 |
|      | 3522 | 5598 |      | 3471 | 5547 |      | 3956 | 6032 |
|      | 3636 | 5712 |      | 4247 | 6323 |      | 4133 | 6209 |
|      | 3883 | 5959 |      | 4482 | 6558 | 393  | 3010 | 5086 |
|      | 3977 | 6053 |      | 4650 | 6726 |      | 3802 | 5878 |
|      | 4028 | 6104 | 374  | 2776 | 4852 |      | 3839 | 5915 |
|      | 4199 | 6275 |      | 3497 | 5573 |      | 3923 | 5999 |
|      | 4201 | 6277 |      | 3645 | 5721 |      | 4421 | 6497 |
|      | 4223 | 6299 |      | 3728 | 5804 |      | 4556 | 6632 |
|      | 4317 | 6393 |      | 3837 | 5913 |      | 4614 | 6690 |
|      | 4354 | 6430 | 378  | 3808 | 5884 | 394  | 3069 | 5145 |
|      | 4408 | 6484 |      | 4045 | 6121 |      | 3159 | 5235 |
|      | 4450 | 6526 |      | 4308 | 6384 |      | 3870 | 5946 |
|      | 4509 | 6585 |      | 4525 | 6601 |      | 4147 | 6223 |
| 365  | 3678 | 5754 | 380  | 3207 | 5283 |      | 4512 | 6588 |
|      | 3941 | 6017 |      | 3773 | 5849 | 398  | 2976 | 5052 |
|      | 4547 | 6623 |      | 4701 | 6777 |      | 3680 | 5756 |
|      | 4577 | 6653 |      | 4715 | 6791 | 404  | 4217 | 6293 |
|      | 4684 | 6760 |      | 4716 | 6792 | 405  | 4116 | 6192 |
|      | 4714 | 6790 | 381  | 4034 | 6110 | 410  | 2754 | 4830 |
| 366  | 2691 | 4767 | 382  | 4405 | 6481 |      | 3260 | 5336 |
|      | 3614 | 5690 |      | 4537 | 6613 | 411  | 3824 | 5900 |
|      | 3711 | 5787 | 385  | 3490 | 5566 |      | 4004 | 6080 |
|      | 3763 | 5839 |      | 4065 | 6141 | 412  | 2967 | 5043 |
|      | 3975 | 6051 |      | 4104 | 6180 |      | 2988 | 5064 |
| 369  | 2708 | 4784 |      | 4454 | 6530 |      | 3887 | 5963 |
|      | 3125 | 5201 |      | 4456 | 6532 |      | 3935 | 6011 |
|      | 3144 | 5220 |      | 4730 | 6806 |      | 4462 | 6538 |
|      | 3190 | 5266 | 389  | 2750 | 4826 | 418  | 3223 | 5299 |
|      | 3370 | 5446 |      | 3529 | 5605 |      | 4376 | 6452 |
|      | 3432 | 5508 |      | 3620 | 5696 |      | 4504 | 6580 |
|      | 4067 | 6143 |      | 4575 | 6651 |      | 4505 | 6581 |
|      | 4355 | 6431 |      | 4694 | 6770 |      | 4663 | 6739 |
|      | 4410 | 6486 |      | 4723 | 6799 | 419  | 2854 | 4930 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 3085 | 5161 |      | 4640 | 6716 |      | 4049 | 6125 |
|      | 3442 | 5518 | 438  | 2818 | 4894 |      | 4158 | 6234 |
|      | 3537 | 5613 |      | 2863 | 4939 |      | 4164 | 6240 |
|      | 3552 | 5628 |      | 3344 | 5420 |      | 4466 | 6542 |
|      | 4200 | 6276 |      | 3454 | 5530 | 457  | 4382 | 6458 |
|      | 4378 | 6454 |      | 3613 | 5689 | 460  | 3166 | 5242 |
| 420  | 3321 | 5397 |      | 4683 | 6759 |      | 3592 | 5668 |
|      | 3517 | 5593 | 439  | 2819 | 4895 |      | 4107 | 6183 |
|      | 3544 | 5620 |      | 2905 | 4981 |      | 4128 | 6204 |
|      | 3689 | 5765 |      | 2935 | 5011 | 463  | 2850 | 4926 |
|      | 4243 | 6319 |      | 3949 | 6025 |      | 3488 | 5564 |
| 422  | 3228 | 5304 |      | 3958 | 6034 |      | 3489 | 5565 |
|      | 3249 | 5325 |      | 4413 | 6489 |      | 3578 | 5654 |
|      | 3250 | 5326 |      | 4417 | 6493 |      | 3849 | 5925 |
|      | 3709 | 5785 |      | 4465 | 6541 |      | 4517 | 6593 |
|      | 4589 | 6665 | 441  | 3330 | 5406 |      | 4624 | 6700 |
| 426  | 2919 | 4995 |      | 3708 | 5784 | 464  | 3657 | 5733 |
|      | 2943 | 5019 |      | 3836 | 5912 |      | 3752 | 5828 |
|      | 3022 | 5098 |      | 3898 | 5974 |      | 4251 | 6327 |
|      | 3273 | 5349 |      | 4419 | 6495 |      | 4372 | 6448 |
|      | 3739 | 5815 | 448  | 2715 | 4791 |      | 4726 | 6802 |
|      | 4409 | 6485 |      | 2773 | 4849 | 465  | 3071 | 5147 |
| 428  | 2753 | 4829 |      | 3053 | 5129 |      | 3270 | 5346 |
|      | 2897 | 4973 |      | 3408 | 5484 |      | 3533 | 5609 |
|      | 3096 | 5172 |      | 3415 | 5491 |      | 3796 | 5872 |
|      | 3500 | 5576 |      | 3538 | 5614 |      | 3905 | 5981 |
|      | 4438 | 6514 |      | 3900 | 5976 |      | 4418 | 6494 |
| 431  | 3193 | 5269 |      | 4358 | 6434 |      | 4605 | 6681 |
| 432  | 3029 | 5105 | 450  | 3398 | 5474 |      | 4674 | 6750 |
|      | 3219 | 5295 |      | 3795 | 5871 | 467  | 3640 | 5716 |
|      | 3375 | 5451 | 451  | 2992 | 5068 |      | 3734 | 5810 |
|      | 3376 | 5452 |      | 3183 | 5259 |      | 4447 | 6523 |
|      | 3623 | 5699 |      | 3269 | 5345 |      | 4464 | 6540 |
|      | 3968 | 6044 |      | 3300 | 5376 |      | 4515 | 6591 |
| 437  | 2855 | 4931 |      | 3478 | 5554 | 472  | 3462 | 5538 |
|      | 2937 | 5013 |      | 4000 | 6076 |      | 3591 | 5667 |
|      | 4341 | 6417 |      | 4018 | 6094 |      | 3823 | 5899 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 4212 | 6288 |      | 3503 | 5579 |      | 3929 | 6005 |
|      | 4349 | 6425 |      | 3776 | 5852 |      | 4052 | 6128 |
| 475  | 2739 | 4815 |      | 4685 | 6761 |      | 4081 | 6157 |
|      | 2772 | 4848 | 497  | 3669 | 5745 |      | 4089 | 6165 |
|      | 3168 | 5244 | 500  | 3444 | 5520 |      | 4314 | 6390 |
|      | 3382 | 5458 |      | 4226 | 6302 |      | 4628 | 6704 |
|      | 3451 | 5527 |      | 4370 | 6446 | 525  | 4162 | 6238 |
|      | 3865 | 5941 |      | 4621 | 6697 | 526  | 2840 | 4916 |
|      | 3974 | 6050 |      | 4651 | 6727 |      | 2994 | 5070 |
|      | 3982 | 6058 | 501  | 3481 | 5557 |      | 3003 | 5079 |
|      | 4364 | 6440 |      | 4239 | 6315 |      | 4020 | 6096 |
| 476  | 2829 | 4905 | 502  | 4610 | 6686 |      | 4653 | 6729 |
|      | 3343 | 5419 | 503  | 3857 | 5933 | 528  | 3004 | 5080 |
|      | 3593 | 5669 |      | 4611 | 6687 |      | 3433 | 5509 |
|      | 3741 | 5817 | 505  | 3165 | 5241 |      | 3674 | 5750 |
|      | 3829 | 5905 |      | 3242 | 5318 |      | 3789 | 5865 |
| 482  | 2853 | 4929 |      | 3749 | 5825 |      | 4638 | 6714 |
|      | 4062 | 6138 |      | 4383 | 6459 | 533  | 3044 | 5120 |
|      | 4106 | 6182 |      | 4718 | 6794 |      | 3241 | 5317 |
|      | 4127 | 6203 | 507  | 2763 | 4839 |      | 3646 | 5722 |
| 484  | 2702 | 4778 |      | 3026 | 5102 |      | 3961 | 6037 |
|      | 3302 | 5378 |      | 3409 | 5485 |      | 4518 | 6594 |
|      | 3327 | 5403 |      | 3468 | 5544 | 535  | 3859 | 5935 |
|      | 3701 | 5777 |      | 3983 | 6059 |      | 4328 | 6404 |
|      | 4502 | 6578 | 509  | 2759 | 4835 | 536  | 3286 | 5362 |
| 487  | 3143 | 5219 |      | 3418 | 5494 |      | 3545 | 5621 |
|      | 3677 | 5753 |      | 3912 | 5988 |      | 3659 | 5735 |
|      | 4075 | 6151 |      | 4079 | 6155 |      | 4117 | 6193 |
|      | 4443 | 6519 |      | 4470 | 6546 |      | 4149 | 6225 |
|      | 4710 | 6786 | 517  | 2672 | 4748 | 539  | 3641 | 5717 |
| 489  | 2841 | 4917 |      | 2690 | 4766 |      | 4248 | 6324 |
|      | 3188 | 5264 |      | 2727 | 4803 |      | 4286 | 6362 |
|      | 3686 | 5762 |      | 2846 | 4922 | 540  | 3093 | 5169 |
|      | 3867 | 5943 |      | 3278 | 5354 |      | 3127 | 5203 |
|      | 4673 | 6749 |      | 3337 | 5413 |      | 3325 | 5401 |
| 493  | 3032 | 5108 |      | 3526 | 5602 | 541  | 2888 | 4964 |
|      | 3480 | 5556 |      | 3730 | 5806 |      | 3037 | 5113 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
| 543  | 3116 | 5192 |      | 4136 | 6212 |      | 3822 | 5898 |
|      | 3434 | 5510 | 553  | 2666 | 4742 |      | 4010 | 6086 |
|      | 3435 | 5511 |      | 2945 | 5021 | 581  | 2765 | 4841 |
|      | 4087 | 6163 |      | 3101 | 5177 |      | 3065 | 5141 |
|      | 4088 | 6164 |      | 3224 | 5300 |      | 3513 | 5589 |
| 546  | 2791 | 4867 |      | 4459 | 6535 |      | 4043 | 6119 |
|      | 2933 | 5009 | 556  | 2929 | 5005 |      | 4416 | 6492 |
|      | 3735 | 5811 |      | 3810 | 5886 | 582  | 3212 | 5288 |
|      | 3970 | 6046 |      | 3811 | 5887 |      | 3411 | 5487 |
|      | 4154 | 6230 |      | 4046 | 6122 |      | 3841 | 5917 |
|      | 4213 | 6289 |      | 4548 | 6624 |      | 4169 | 6245 |
|      | 4292 | 6368 | 559  | 2734 | 4810 |      | 4293 | 6369 |
|      | 4401 | 6477 |      | 3611 | 5687 | 583  | 2712 | 4788 |
| 549  | 3369 | 5445 |      | 3737 | 5813 |      | 2713 | 4789 |
|      | 3420 | 5496 |      | 4179 | 6255 |      | 2725 | 4801 |
|      | 3421 | 5497 |      | 4647 | 6723 |      | 2812 | 4888 |
|      | 3441 | 5517 | 560  | 3598 | 5674 |      | 3031 | 5107 |
|      | 3559 | 5635 |      | 3599 | 5675 |      | 3794 | 5870 |
|      | 4069 | 6145 |      | 3600 | 5676 |      | 4274 | 6350 |
|      | 4300 | 6376 |      | 3712 | 5788 |      | 4699 | 6775 |
|      | 4301 | 6377 |      | 4388 | 6464 | 586  | 3746 | 5822 |
|      | 4561 | 6637 | 566  | 2953 | 5029 |      | 4672 | 6748 |
| 550  | 3141 | 5217 |      | 2954 | 5030 | 595  | 2821 | 4897 |
|      | 3361 | 5437 |      | 3615 | 5691 |      | 3275 | 5351 |
|      | 3377 | 5453 |      | 4386 | 6462 |      | 3473 | 5549 |
|      | 3939 | 6015 |      | 4387 | 6463 |      | 3877 | 5953 |
|      | 3960 | 6036 | 569  | 2786 | 4862 |      | 4097 | 6173 |
|      | 4091 | 6167 |      | 3019 | 5095 |      | 4312 | 6388 |
| 551  | 3340 | 5416 |      | 3233 | 5309 |      | 4362 | 6438 |
|      | 3363 | 5439 |      | 3307 | 5383 |      | 4554 | 6626 |
|      | 4142 | 6218 |      | 4469 | 6545 | 599  | 3055 | 5131 |
|      | 4159 | 6235 | 572  | 2674 | 4750 |      | 3213 | 5289 |
|      | 4436 | 6512 |      | 4099 | 6175 |      | 4166 | 6242 |
| 552  | 3805 | 5881 |      | 4366 | 6442 |      | 4167 | 6243 |
|      | 3993 | 6069 | 577  | 2825 | 4901 |      | 4562 | 6638 |
|      | 4016 | 6092 |      | 3006 | 5082 | 601  | 3079 | 5155 |
|      | 4077 | 6153 |      | 3136 | 5212 |      | 3424 | 5500 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 4241 | 6317 |      | 2999 | 5075 | 642  | 3039 | 5115 |
|      | 4321 | 6397 |      | 3353 | 5429 |      | 4082 | 6158 |
| 602  | 4083 | 6159 |      | 3362 | 5438 |      | 4238 | 6314 |
| 603  | 2775 | 4851 |      | 3472 | 5548 |      | 4437 | 6513 |
|      | 3685 | 5761 |      | 3723 | 5799 |      | 4563 | 6639 |
|      | 3934 | 6010 | 628  | 2898 | 4974 | 647  | 2758 | 4834 |
|      | 4235 | 6311 |      | 3106 | 5182 |      | 2879 | 4955 |
|      | 4351 | 6427 |      | 3108 | 5184 |      | 3214 | 5290 |
|      | 4657 | 6733 |      | 4539 | 6615 |      | 3476 | 5552 |
| 605  | 2724 | 4800 |      | 4540 | 6616 |      | 3564 | 5640 |
|      | 2784 | 4860 | 631  | 3247 | 5323 |      | 3661 | 5737 |
|      | 3052 | 5128 | 633  | 3294 | 5370 |      | 3772 | 5848 |
|      | 3288 | 5364 |      | 3295 | 5371 | 651  | 2867 | 4943 |
|      | 3422 | 5498 |      | 3391 | 5467 |      | 3244 | 5320 |
|      | 3731 | 5807 |      | 3453 | 5529 |      | 4017 | 6093 |
|      | 3944 | 6020 |      | 3621 | 5697 |      | 4191 | 6267 |
|      | 4007 | 6083 | 634  | 2710 | 4786 |      | 4242 | 6318 |
|      | 4346 | 6422 |      | 2756 | 4832 | 653  | 3077 | 5153 |
|      | 4389 | 6465 |      | 3148 | 5224 | 654  | 2663 | 4739 |
| 609  | 3311 | 5387 |      | 3470 | 5546 |      | 2835 | 4911 |
|      | 3791 | 5867 |      | 3572 | 5648 |      | 3459 | 5535 |
| 610  | 4011 | 6087 |      | 4078 | 6154 |      | 3908 | 5984 |
|      | 4522 | 6598 |      | 4221 | 6297 |      | 4582 | 6658 |
| 612  | 2795 | 4871 |      | 4365 | 6441 | 656  | 3660 | 5736 |
|      | 3130 | 5206 |      | 4404 | 6480 |      | 3893 | 5969 |
|      | 3574 | 5650 | 635  | 3040 | 5116 |      | 4012 | 6088 |
|      | 4315 | 6391 |      | 3258 | 5334 |      | 4021 | 6097 |
|      | 4529 | 6605 |      | 3339 | 5415 |      | 4508 | 6584 |
| 613  | 3162 | 5238 | 640  | 3568 | 5644 | 657  | 4139 | 6215 |
| 619  | 2974 | 5050 |      | 3790 | 5866 | 658  | 2878 | 4954 |
|      | 3717 | 5793 |      | 3821 | 5897 |      | 3437 | 5513 |
| 620  | 3650 | 5726 |      | 4511 | 6587 |      | 3486 | 5562 |
|      | 3653 | 5729 | 641  | 2719 | 4795 |      | 4002 | 6078 |
|      | 3675 | 5751 |      | 2793 | 4869 |      | 4160 | 6236 |
|      | 4526 | 6602 |      | 3556 | 5632 | 660  | 3132 | 5208 |
| 623  | 2849 | 4925 |      | 3714 | 5790 |      | 3292 | 5368 |
|      | 2936 | 5012 |      | 3991 | 6067 |      | 4426 | 6502 |



| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
| 662  | 4632 | 6708 | 676  | 3622 | 5698 | 686  | 2907 | 4983 |
|      | 4687 | 6763 |      | 3940 | 6016 |      | 2917 | 4993 |
|      | 2969 | 5045 |      | 4367 | 6443 |      | 3540 | 5616 |
|      | 3535 | 5611 |      | 2963 | 5039 |      | 3892 | 5968 |
|      | 4260 | 6336 |      | 3084 | 5160 |      | 3937 | 6013 |
| 664  | 4261 | 6337 | 677  | 3112 | 5188 | 687  | 3779 | 5855 |
|      | 4262 | 6338 |      | 3221 | 5297 |      | 3902 | 5978 |
|      | 2704 | 4780 |      | 3332 | 5408 |      | 4569 | 6645 |
|      | 3766 | 5842 |      | 3507 | 5583 |      | 4615 | 6691 |
|      | 4094 | 6170 |      | 3570 | 5646 |      | 4722 | 6798 |
| 665  | 4689 | 6765 | 678  | 3699 | 5775 | 688  | 3297 | 5373 |
|      | 4732 | 6808 |      | 4138 | 6214 |      | 4095 | 6171 |
|      | 2741 | 4817 |      | 4390 | 6466 |      | 4282 | 6358 |
|      | 2827 | 4903 |      | 3439 | 5515 |      | 4433 | 6509 |
|      | 3056 | 5132 |      | 3440 | 5516 |      | 4446 | 6522 |
| 672  | 3181 | 5257 | 680  | 3952 | 6028 | 691  | 2837 | 4913 |
|      | 3448 | 5524 |      | 4072 | 6148 |      | 3238 | 5314 |
|      | 3520 | 5596 |      | 4073 | 6149 |      | 3364 | 5440 |
|      | 3882 | 5958 |      | 2810 | 4886 |      | 3542 | 5618 |
|      | 4080 | 6156 |      | 2883 | 4959 |      | 3768 | 5844 |
| 675  | 4101 | 6177 | 683  | 3195 | 5271 | 692  | 3585 | 5661 |
|      | 4171 | 6247 |      | 3266 | 5342 |      | 3586 | 5662 |
|      | 4176 | 6252 |      | 3427 | 5503 |      | 4658 | 6734 |
|      | 4205 | 6281 |      | 2797 | 4873 |      | 4659 | 6735 |
|      | 4264 | 6340 |      | 3750 | 5826 | 693  | 2806 | 4882 |
| 677  | 4278 | 6354 | 684  | 4588 | 6664 |      | 2923 | 4999 |
|      | 4507 | 6583 |      | 2735 | 4811 |      | 3043 | 5119 |
|      | 4676 | 6752 |      | 2970 | 5046 |      | 3229 | 5305 |
|      | 4691 | 6767 |      | 3760 | 5836 |      | 3320 | 5396 |
|      | 4734 | 6810 |      | 3778 | 5854 |      | 4258 | 6334 |
| 678  | 3104 | 5180 | 685  | 4560 | 6636 | 695  | 2842 | 4918 |
|      | 3907 | 5983 |      | 4320 | 6396 |      | 2843 | 4919 |
|      | 4141 | 6217 |      | 3140 | 5216 |      | 2844 | 4920 |
|      | 4612 | 6688 |      | 3587 | 5663 |      | 2908 | 4984 |
|      | 4669 | 6745 |      | 3876 | 5952 |      | 2949 | 5025 |
| 679  | 2717 | 4793 | 686  | 4119 | 6195 | 696  | 3554 | 5630 |
|      | 3149 | 5225 |      | 4727 | 6803 |      | 3670 | 5746 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 3926 | 6002 |      | 4538 | 6614 |      | 3725 | 5801 |
|      | 4338 | 6414 |      | 4607 | 6683 |      | 4189 | 6265 |
| 700  | 2947 | 5023 |      | 4704 | 6780 |      | 4402 | 6478 |
|      | 3246 | 5322 |      | 4705 | 6781 | 731  | 2688 | 4764 |
|      | 4180 | 6256 |      | 4706 | 6782 |      | 2900 | 4976 |
|      | 4441 | 6517 | 715  | 3757 | 5833 |      | 2962 | 5038 |
|      | 4666 | 6742 |      | 3936 | 6012 |      | 2979 | 5055 |
|      | 4667 | 6743 |      | 4229 | 6305 |      | 3016 | 5092 |
| 702  | 3216 | 5292 |      | 4230 | 6306 |      | 3048 | 5124 |
|      | 3549 | 5625 |      | 4528 | 6604 |      | 3070 | 5146 |
|      | 3654 | 5730 | 718  | 3047 | 5123 |      | 3092 | 5168 |
|      | 3733 | 5809 | 719  | 2817 | 4893 |      | 3589 | 5665 |
|      | 4132 | 6208 |      | 2894 | 4970 |      | 3617 | 5693 |
| 706  | 2697 | 4773 |      | 3076 | 5152 |      | 3720 | 5796 |
|      | 2770 | 4846 |      | 3314 | 5390 |      | 3782 | 5858 |
|      | 2788 | 4864 |      | 3759 | 5835 |      | 3962 | 6038 |
|      | 2845 | 4921 |      | 4092 | 6168 |      | 4155 | 6231 |
|      | 3156 | 5232 |      | 4271 | 6347 |      | 4174 | 6250 |
|      | 3234 | 5310 |      | 4302 | 6378 |      | 4474 | 6550 |
|      | 3744 | 5820 |      | 4303 | 6379 |      | 4510 | 6586 |
|      | 4013 | 6089 |      | 4304 | 6380 |      | 4545 | 6621 |
| 707  | 2983 | 5059 |      | 4305 | 6381 |      | 4729 | 6805 |
|      | 3582 | 5658 |      | 4455 | 6531 | 732  | 3027 | 5103 |
|      | 3642 | 5718 |      | 4473 | 6549 |      | 3064 | 5140 |
|      | 4059 | 6135 | 720  | 2930 | 5006 |      | 3710 | 5786 |
|      | 4076 | 6152 |      | 3062 | 5138 |      | 4524 | 6600 |
|      | 4194 | 6270 |      | 3172 | 5248 |      | 4566 | 6642 |
|      | 4686 | 6762 |      | 3289 | 5365 | 740  | 3861 | 5937 |
| 712  | 3038 | 5114 |      | 4384 | 6460 | 741  | 3196 | 5272 |
|      | 3392 | 5468 | 725  | 4228 | 6304 |      | 3871 | 5947 |
|      | 3400 | 5476 | 729  | 2787 | 4863 | 744  | 2732 | 4808 |
|      | 3512 | 5588 |      | 2975 | 5051 |      | 3210 | 5286 |
|      | 3758 | 5834 |      | 3095 | 5171 |      | 4255 | 6331 |
|      | 4110 | 6186 |      | 3494 | 5570 |      | 4256 | 6332 |
|      | 4118 | 6194 |      | 3495 | 5571 |      | 4347 | 6423 |
|      | 4177 | 6253 |      | 3496 | 5572 |      | 4374 | 6450 |
|      | 4422 | 6498 |      | 3702 | 5778 |      | 4500 | 6576 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 4626 | 6702 |      | 4144 | 6220 | 782  | 3783 | 5859 |
| 745  | 2892 | 4968 |      | 4196 | 6272 |      | 3834 | 5910 |
|      | 4428 | 6504 |      | 4295 | 6371 |      | 3854 | 5930 |
| 752  | 2686 | 4762 |      | 4396 | 6472 |      | 4165 | 6241 |
|      | 2830 | 4906 |      | 4724 | 6800 |      | 4586 | 6662 |
|      | 2831 | 4907 | 765  | 3231 | 5307 | 788  | 2677 | 4753 |
|      | 2932 | 5008 |      | 3627 | 5703 |      | 3573 | 5649 |
|      | 4623 | 6699 |      | 4268 | 6344 |      | 4192 | 6268 |
| 756  | 3094 | 5170 |      | 4361 | 6437 |      | 4434 | 6510 |
|      | 3484 | 5560 |      | 4661 | 6737 | 790  | 3349 | 5425 |
|      | 3597 | 5673 | 767  | 3777 | 5853 |      | 3793 | 5869 |
|      | 3652 | 5728 | 768  | 3813 | 5889 |      | 4263 | 6339 |
|      | 3860 | 5936 |      | 3814 | 5890 | 792  | 2877 | 4953 |
|      | 3879 | 5955 | 770  | 2913 | 4989 |      | 2901 | 4977 |
|      | 3964 | 6040 |      | 3354 | 5430 |      | 3516 | 5592 |
| 757  | 2826 | 4902 |      | 3355 | 5431 |      | 3918 | 5994 |
|      | 2910 | 4986 |      | 3903 | 5979 |      | 3947 | 6023 |
|      | 3017 | 5093 |      | 4527 | 6603 |      | 4019 | 6095 |
|      | 3030 | 5106 | 771  | 2906 | 4982 |      | 4279 | 6355 |
|      | 3067 | 5143 |      | 2911 | 4987 |      | 4296 | 6372 |
|      | 3080 | 5156 |      | 3131 | 5207 |      | 4356 | 6432 |
|      | 3113 | 5189 |      | 3458 | 5534 |      | 4643 | 6719 |
|      | 3178 | 5254 |      | 3874 | 5950 | 793  | 2698 | 4774 |
|      | 3272 | 5348 | 772  | 2856 | 4932 |      | 2794 | 4870 |
|      | 3482 | 5558 |      | 2973 | 5049 |      | 2811 | 4887 |
|      | 4585 | 6661 |      | 3083 | 5159 |      | 3350 | 5426 |
|      | 4671 | 6747 |      | 3197 | 5273 | 794  | 3919 | 5995 |
|      | 4711 | 6787 |      | 3217 | 5293 |      | 3954 | 6030 |
| 763  | 2868 | 4944 |      | 3502 | 5578 |      | 3986 | 6062 |
|      | 3124 | 5200 |      | 4003 | 6079 |      | 4448 | 6524 |
|      | 3679 | 5755 |      | 4565 | 6641 |      | 4665 | 6741 |
|      | 4203 | 6279 |      | 4604 | 6680 | 795  | 2676 | 4752 |
|      | 4432 | 6508 | 776  | 2670 | 4746 |      | 3058 | 5134 |
|      | 4498 | 6574 |      | 3209 | 5285 |      | 3139 | 5215 |
|      | 4712 | 6788 |      | 3780 | 5856 |      | 3690 | 5766 |
| 764  | 3301 | 5377 |      | 4209 | 6285 |      | 3691 | 5767 |
|      | 3715 | 5791 |      | 4485 | 6561 |      | 3692 | 5768 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 3693 | 5769 |      | 4559 | 6635 |      | 4369 | 6445 |
|      | 3928 | 6004 | 816  | 2768 | 4844 | 832  | 2790 | 4866 |
|      | 4214 | 6290 |      | 3150 | 5226 | 833  | 2824 | 4900 |
| 797  | 2700 | 4776 |      | 3846 | 5922 |      | 2889 | 4965 |
|      | 2828 | 4904 |      | 3894 | 5970 |      | 2956 | 5032 |
|      | 2882 | 4958 |      | 4057 | 6133 |      | 3360 | 5436 |
|      | 3306 | 5382 | 820  | 3648 | 5724 |      | 3397 | 5473 |
|      | 3695 | 5771 |      | 3855 | 5931 |      | 3521 | 5597 |
|      | 4135 | 6211 |      | 4257 | 6333 |      | 3647 | 5723 |
| 798  | 3358 | 5434 | 823  | 3800 | 5876 |      | 3698 | 5774 |
|      | 3405 | 5481 | 825  | 2870 | 4946 |      | 4357 | 6433 |
|      | 3407 | 5483 |      | 4096 | 6172 |      | 4423 | 6499 |
|      | 3460 | 5536 |      | 4283 | 6359 |      | 4637 | 6713 |
|      | 3596 | 5672 |      | 4284 | 6360 | 836  | 3243 | 5319 |
| 803  | 3194 | 5270 |      | 4564 | 6640 |      | 3262 | 5338 |
|      | 3239 | 5315 | 826  | 2931 | 5007 |      | 3263 | 5339 |
|      | 3299 | 5375 |      | 3331 | 5407 |      | 3404 | 5480 |
|      | 3588 | 5664 |      | 3429 | 5505 |      | 4131 | 6207 |
|      | 3612 | 5688 |      | 3601 | 5677 |      | 4322 | 6398 |
|      | 4736 | 6812 |      | 4103 | 6179 |      | 4348 | 6424 |
| 804  | 2692 | 4768 |      | 4494 | 6570 |      | 4592 | 6668 |
|      | 3059 | 5135 |      | 4541 | 6617 | 838  | 3045 | 5121 |
|      | 3399 | 5475 | 827  | 2814 | 4890 |      | 3154 | 5230 |
|      | 3721 | 5797 |      | 3707 | 5783 |      | 3357 | 5433 |
|      | 3729 | 5805 |      | 3784 | 5860 | 840  | 2800 | 4876 |
| 808  | 2904 | 4980 |      | 3785 | 5861 |      | 3843 | 5919 |
|      | 3280 | 5356 |      | 4188 | 6264 |      | 3933 | 6009 |
|      | 3827 | 5903 | 829  | 3786 | 5862 |      | 3978 | 6054 |
|      | 4429 | 6505 | 830  | 2673 | 4749 |      | 3998 | 6074 |
|      | 4660 | 6736 |      | 2982 | 5058 | 841  | 3089 | 5165 |
| 809  | 2872 | 4948 |      | 3390 | 5466 |      | 3279 | 5355 |
|      | 4717 | 6793 |      | 4435 | 6511 |      | 3335 | 5411 |
| 811  | 4033 | 6109 |      | 4649 | 6725 |      | 4641 | 6717 |
| 814  | 2880 | 4956 |      | 4690 | 6766 | 848  | 2749 | 4825 |
|      | 2987 | 5063 | 831  | 3334 | 5410 |      | 2912 | 4988 |
|      | 3416 | 5492 |      | 3543 | 5619 |      | 3254 | 5330 |
|      | 4055 | 6131 |      | 3743 | 5819 |      | 3976 | 6052 |

| AORF | RH   | RP   | AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|------|------|------|
|      | 4708 | 6784 |      | 4452 | 6528 |      | 3985 | 6061 |
| 852  | 2680 | 4756 |      | 4599 | 6675 |      | 4391 | 6467 |
|      | 2859 | 4935 | 868  | 2681 | 4757 |      | 4617 | 6693 |
|      | 2860 | 4936 |      | 2895 | 4971 | 901  | 3265 | 5341 |
|      | 3118 | 5194 |      | 2980 | 5056 |      | 3672 | 5748 |
|      | 3313 | 5389 |      | 4193 | 6269 |      | 3775 | 5851 |
|      | 4047 | 6123 |      | 4477 | 6553 |      | 3873 | 5949 |
|      | 4379 | 6455 |      | 4478 | 6554 |      | 4635 | 6711 |
|      | 4532 | 6608 |      | 4696 | 6772 | 904  | 4266 | 6342 |
| 856  | 3366 | 5442 | 871  | 3308 | 5384 | 908  | 2891 | 4967 |
|      | 4481 | 6557 |      | 3309 | 5385 |      | 3009 | 5085 |
| 857  | 3501 | 5577 |      | 3310 | 5386 |      | 3536 | 5612 |
|      | 3609 | 5685 |      | 4567 | 6643 |      | 3950 | 6026 |
|      | 3931 | 6007 |      | 4576 | 6652 |      | 4071 | 6147 |
|      | 4336 | 6412 | 876  | 3021 | 5097 | 909  | 3348 | 5424 |
|      | 4702 | 6778 |      | 3161 | 5237 |      | 3651 | 5727 |
| 859  | 3412 | 5488 |      | 3206 | 5282 |      | 4102 | 6178 |
|      | 3694 | 5770 |      | 3607 | 5683 |      | 4578 | 6654 |
|      | 4156 | 6232 |      | 4024 | 6100 |      | 4613 | 6689 |
|      | 4499 | 6575 | 883  | 3384 | 5460 | 911  | 2823 | 4899 |
|      | 4568 | 6644 |      | 3819 | 5895 |      | 2857 | 4933 |
| 861  | 2665 | 4741 |      | 3832 | 5908 |      | 3226 | 5302 |
|      | 3120 | 5196 |      | 3868 | 5944 |      | 3938 | 6014 |
|      | 3121 | 5197 |      | 4318 | 6394 |      | 4294 | 6370 |
|      | 4109 | 6185 | 887  | 3018 | 5094 | 912  | 2718 | 4794 |
|      | 4140 | 6216 |      | 3913 | 5989 |      | 2816 | 4892 |
|      | 4163 | 6239 |      | 4066 | 6142 |      | 3185 | 5261 |
|      | 4574 | 6650 |      | 4407 | 6483 |      | 3662 | 5738 |
| 863  | 2729 | 4805 |      | 4725 | 6801 |      | 4393 | 6469 |
|      | 2925 | 5001 | 889  | 4204 | 6280 | 914  | 2728 | 4804 |
|      | 3576 | 5652 | 891  | 2834 | 4910 |      | 3177 | 5253 |
|      | 4148 | 6224 |      | 3298 | 5374 |      | 3483 | 5559 |
|      | 4195 | 6271 |      | 3815 | 5891 |      | 3809 | 5885 |
| 867  | 2726 | 4802 |      | 4151 | 6227 |      | 4339 | 6415 |
|      | 4310 | 6386 |      | 4252 | 6328 | 918  | 2968 | 5044 |
|      | 4332 | 6408 | 892  | 3129 | 5205 |      | 3577 | 5653 |
|      | 4442 | 6518 |      | 3342 | 5418 |      | 3787 | 5863 |

| AORF | RH   | RP   | AORF | RH   | RP   |
|------|------|------|------|------|------|
|      | 4297 | 6373 |      | 3981 | 6057 |
|      | 4445 | 6521 | 944  | 2997 | 5073 |
| 919  | 2985 | 5061 |      | 3046 | 5122 |
|      | 3356 | 5432 | 950  | 3519 | 5595 |
|      | 3881 | 5957 |      | 3980 | 6056 |
|      | 4090 | 6166 |      | 4250 | 6326 |
|      | 4652 | 6728 |      | 4471 | 6547 |
| 921  | 2887 | 4963 |      | 4535 | 6611 |
| 926  | 2946 | 5022 | 952  | 2714 | 4790 |
|      | 3128 | 5204 |      | 2774 | 4850 |
|      | 3649 | 5725 |      | 3287 | 5363 |
|      | 3858 | 5934 |      | 3716 | 5792 |
| 928  | 4121 | 6197 |      | 4122 | 6198 |
|      | 4392 | 6468 |      | 4123 | 6199 |
|      | 4552 | 6629 |      | 4531 | 6607 |
| 929  | 2743 | 4819 |      | 4536 | 6612 |
|      | 2745 | 4821 |      | 4573 | 6649 |
|      | 3666 | 5742 |      | 4634 | 6710 |
|      | 4276 | 6352 | 953  | 2683 | 4759 |
|      | 4343 | 6419 |      | 3531 | 5607 |
|      | 4630 | 6706 |      | 3668 | 5744 |
| 930  | 4249 | 6325 |      | 3826 | 5902 |
| 932  | 2965 | 5041 |      | 4150 | 6226 |
|      | 3014 | 5090 |      | SEQ  |      |
|      | 3840 | 5916 |      | ID   |      |
|      | 4041 | 6117 |      | NO:  |      |
|      | 4137 | 6213 |      |      |      |
|      | 4506 | 6582 |      |      |      |
| 934  | 3616 | 5692 |      |      |      |
|      | 3816 | 5892 |      |      |      |
|      | 4105 | 6181 |      |      |      |
|      | 4111 | 6187 |      |      |      |
|      | 4172 | 6248 |      |      |      |
| 943  | 2881 | 4957 |      |      |      |
|      | 2998 | 5074 |      |      |      |
|      | 3203 | 5279 |      |      |      |
|      | 3818 | 5894 |      |      |      |

**Table 23** Plant Open Reading Frames (ORFs) orthologous to specific Arabidopsis ORFs

| SEQ ID NO: 2        |   |
|---------------------|---|
| AAD17487.1 AF049347 | Berberis stolonifera<br>Catalyzes the oxidative cyclization of the N-methyl group of (S)-reticuline into the berberine bridge carbon (C-8) of (S)-scoulerine. berberine bridge enzyme. bbe1. Covalently flavinylated oxidase of isoquinoline alkaloid biosynthesis in plants.         |
| AAB20352.1 S65550   | Eschscholzia californica<br>(S)-reticuline:oxygen oxidoreductase (methylene-bridge-forming). /gene="(S)-reticuline:oxygen oxidoreductase. This sequence comes from Fig 2; berberine bridge enzyme.  |
| AAC39358.1 AF005655 | Eschscholzia californica<br>oxidizes the N-methyl group of (S)-reticuline to the berberine bridge carbon C-8 of (S)-scoulerine in berberine and benzophenanthridine alkaloid biosynthesis. berberine bridge enzyme. bbe1. covalently-bound FAD-dependent oxidase; elicitor-inducible. |
| AAC61839.1 AF025430 | Papaver somniferum<br>berberine bridge enzyme. bbe1. (S)-reticuline:oxygen oxidoreductase (methylene bridge forming).   |
| SEQ ID NO: 19       |   |
| AAC83458.1 AF023140 | Thlaspi goesingense<br>imidazoleglycerol phosphate dehydratase. THB1.   |
| AAB67738.1 U49978   | Pisum sativum<br>imidazoleglycerol-phosphate dehydratase. PSHIS3.   |
| AAA93197.1 U02690   | Triticum aestivum<br>imidazoleglycerolphosphate dehydratase. partial interval represents the confirmed coding region based on homology to sequence U02689, maybe a chimeric molecule.   |
| CAB55393.1 AL117264 | Oryza sativa<br>zwh0009.1. similar to wheat imidazoleglycerol-phosphate dehydratase (P34048); Method: conceptual translation with partial peptide sequencing.   |
| SEQ ID NO: 32       |   |
| CAA45066.2 X63464   | Pisum sativum<br>dihydrolipoamide dehydrogenase. lpd.   |
| CAA44729.1 X62995   | Pisum sativum<br>lipoamide dehydrogenase.   |
| AAG17888.1 AF295339 | Solanum tuberosum<br>dihydrolipoamide dehydrogenase precursor. lpd2. NADH; E3 isoform.  |
| BAB39219.1 AP002869 | Oryza sativa<br>putative dihydrolipoamide dehydrogenase precursor. P0554D10.3. contains ESTs C98433(E0148),AU101058(E0148).   |
| SEQ ID NO: 33       |   |
| CAC03581.1 AJ297566 | Zea mays<br>putative 1-deoxy-D-xylulose 5-phosphate reductoisomerase. dxr.  |
| AAD24768.1 AF116825 | Mentha x piperita<br>1-deoxy-D-xylulose-5-phosphate reductoisomerase. DXR.  |

|               |          |                             |  |
|---------------|----------|-----------------------------|--|
| AAF65154.1    | AF250235 | Catharanthus roseus         | catalyzes the formation of 2-C-methyl-D-erythritol 4-phosphate from 1-deoxy-D-xylulose-5-phosphate. 1-deoxy-D-xylulose-5-phosphate reductoisomerase. dxr.  |
| AAD56391.2    | AF182287 | Artemisia annua             | catalyzes the formation of 2-C-methyl-D-erythritol 4-phosphate from 1-deoxy-D-xylulose-5-phosphate. 1-deoxy-D-xylulose-5-phosphate reductoisomerase. DXR1. |
| BAB16915.1    | AP002863 | Oryza sativa                | 1-deoxy-d-xylulose-5-phosphate reductoisomerase. P0005A05.19. contains ESTs AU108198(S11168),D46469(S11168).   |
| SEQ ID NO: 37 |          |                             |  |
| AAD21872.1    | AF078082 | Phaseolus vulgaris          | receptor-like protein kinase homolog RK20-1.   |
| CAA73134.1    | Y12531   | Brassica oleracea           | serine/threonine kinase. BRLK.   |
| AAB93834.1    | U82481   | Zea mays                    | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.   |
| CAA67145.1    | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.  |
| AAC23542.1    | U20948   | Ipomoea trifida             | receptor protein kinase. IRK1.   |
| CAA73133.1    | Y12530   | Brassica oleracea           | serine /threonine kinase. ARLK.  |
| CAB41878.1    | Y18259   | Brassica oleracea           | SRK5 protein. SRK5. receptor-like kinase.  |
| CAB41879.1    | Y18260   | Brassica oleracea           | SRK15 protein. SRK15. receptor-like kinase.  |
| CAA74661.1    | Y14285   | Brassica oleracea           | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.   |
| CAA74662.1    | Y14286   | Brassica oleracea           | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.   |
| AAA33000.1    | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| BAA23676.1    | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.  |
| AAA33008.1    | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| AAA62232.1    | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| CAB89179.1    | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |



|            |          |                   |  |
|------------|----------|-------------------|--|
| BAA92836.1 | AB032473 | Brassica oleracea | S18 S-locus receptor kinase. SRK18.  |
| CAA79355.1 | Z18921   | Brassica oleracea | S-receptor kinase-like protein.  |
| BAA92837.1 | AB032474 | Brassica oleracea | S60 S-locus receptor kinase. SRK60.  |
| BAA07577.2 | D38564   | Brassica rapa     | receptor protein kinase SRK12.   |
| BAA07576.1 | D38563   | Brassica rapa     | receptor protein kinase SRK8.  |
| BAB21001.1 | AB054061 | Brassica rapa     | S locus receptor kinase. SRK22.  |
| BAA06285.1 | D30049   | Brassica rapa     | S-receptor kinase SRK9.  |
| BAA21132.1 | D88193   | Brassica rapa     | S-receptor kinase. SRK9 (B.c).   |
| AAD52097.1 | AF088885 | Nicotiana tabacum | receptor-like kinase CHRK1. Chrk1.   |
| BAB18292.1 | AP002860 | Oryza sativa      | putative receptor-like protein kinase. P0409B08.19.  |
| BAB16871.1 | AP002537 | Oryza sativa      | putative protein kinase APK1A Arabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).                          |
| BAB39873.1 | AP002882 | Oryza sativa      | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| BAA87853.1 | AP000816 | Oryza sativa      | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308). |
| CAA79324.1 | Z18884   | Brassica oleracea | S-receptor kinase related protein.   |
| AAK21965.1 | AY028699 | Brassica napus    | receptor protein kinase PERK1.   |
| AAK00425.1 | AC069324 | Oryza sativa      | Putative protein kinase. OSJNBa0071K19.11.   |
| AAF66615.1 | AF142596 | Nicotiana tabacum | LRR receptor-like protein kinase.  |
| BAB21240.1 | AP002953 | Oryza sativa      | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).   |
| BAB39409.1 | AP002901 | Oryza sativa      | putative protein kinase. P0456F08.9. contains EST C23560(R0290).   |
| AAG59657.1 | AC084319 | Oryza sativa      | putative protein kinase. OSJNBa0004B24.20.   |

AAG03090.1 AC073405 *Oryza sativa*  
Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).

AAG16628.1 AY007545 *Brassica napus*  
protein serine/threonine kinase BNK1.

BAA94509.1 AB041503 *Populus nigra*  
protein kinase 1. PnPK1.

---

SEQ ID NO: 38

---

CAA58750.1 X83869 *Daucus carota*  
CDPK-related protein kinase. CRK (or PK421).

BAA12692.1 D84508 *Zea mays*  
CDPK-related protein kinase. Does not require calcium for its activity.

AAG01179.1 AF289237 *Zea mays*  
calcium/calmodulin dependent protein kinase MCK2. MCK2.

AAB47181.1 S82324 *Zea mays*  
/gene="calcium/calmodulin-dependent protein kinase. This sequence comes from Fig. 1.

BAA12691.1 D84507 *Zea mays*  
CDPK-related protein kinase. Does not require calcium for its activity (by similarity).

BAA22410.1 D38452 *Zea mays*  
calcium-dependent protein kinase-related kinase.

AAC24961.1 AF009337 *Tradescantia virginiana*  
CDPK-related protein kinase. CRK1.

AAF23901.2 AF194414 *Oryza sativa*  
calcium-dependent protein kinase. CDPK5. OsCDPK5.

AAF23900.1 AF194413 *Oryza sativa*  
calcium-dependent protein kinase. CDPK1. OsCDPK1.

AAC78558.1 AF030879 *Solanum tuberosum*  
protein kinase CPK1.

CAA57157.1 X81394 *Oryza sativa*  
calcium-dependent protein kinase. OSCPK2.

BAB21081.1 AP002819 *Oryza sativa*  
putative calcium-dependent protein kinase. P0501G01.10.

AAC25423.1 AF072908 *Nicotiana tabacum*  
calcium-dependent protein kinase. CDPK1.

BAA12715.1 D85039 *Zea mays*  
calcium-dependent protein kinase.

AAB49984.1 U90262 *Cucurbita pepo*  
calcium-dependent calmodulin-independent protein kinase CDPK. cpCPK1. serine/threonine protein kinase that is activated by direct binding of calcium.

AAA69507.1 U28376 *Zea mays*  
calcium-dependent protein kinase. MZECDPK2.

AAD17800.1 AF090835 *Mesembryanthemum crystallinum*  
Ca<sup>2+</sup>-dependent protein kinase. CPK1. serine/threonine protein kinase.

---

|            |          |                       |   |
|------------|----------|-----------------------|---|
| AAB80693.1 | U69174   | Glycine max           | calmodulin-like domain protein kinase isoenzyme gamma. CDPK gamma.  |
| BAA81751.1 | AB017517 | Marchantia polymorpha | calcium-dependent protein kinase. CDPK1. CDPK-B; alternative splicing.  |
| BAA81749.1 | AB017515 | Marchantia polymorpha | calcium-dependent protein kinase. CDPK1. CDPK-B; alternative splicing.  |
| CAA39936.1 | X56599   | Daucus carota         | calcium- dependent protein kinase. DcPK431.   |
| BAA81748.1 | AB017515 | Marchantia polymorpha | calcium-dependent protein kinase. CDPK1. CDPK-A; alternative splicing.  |
| BAA81750.1 | AB017516 | Marchantia polymorpha | calcium-dependent protein kinase. CDPK1. CDPK-A; alternative splicing.  |
| AAA61682.1 | L27484   | Zea mays              | calcium-dependent protein kinase. CDPK.   |
| BAB16888.1 | AB042550 | Oryza sativa          | OsCDPK7. oscdpk7.   |
| AAB88537.1 | AF035944 | Fragaria x ananassa   | calcium-dependent protein kinase. MAX17.  |
| AAD28192.2 | AF115406 | Solanum tuberosum     | calcium-dependent protein kinase. CDPK; catalytic domain.   |
| CAA07481.1 | AJ007366 | Zea mays              | calcium-dependent protein kinase.   |
| BAA12338.1 | D84408   | Zea mays              | calcium dependent protein kinase. ZmCDPK1.  |
| BAA13440.1 | D87707   | Ipomoea batatas       | calcium dependent protein kinase. CDPK.   |
| BAA13232.1 | D87042   | Zea mays              | Calcium-dependent protein kinase.   |
| CAA65500.1 | X96723   | Medicago sativa       | protein kinase. CDPK.   |
| AAB70706.1 | U82087   | Tortula ruralis       | calmodulin-like domain protein kinase. TrCPK1.  |
| AAC49405.1 | U08140   | Vigna radiata         | calcium dependent protein kinase. CDPK.   |
| BAA85396.1 | AP000615 | Oryza sativa          | ESTs C22369(C12239),C22370(C12239), AU057852(S21844),AU057853(S21844) correspond to a region of the predicted gene.; similar to calcium dependent protein kinase. (AF048691). |
| CAA57156.1 | X81393   | Oryza sativa          | calcium-dependent protein kinase. OSCPKII.  |
| AAC05270.1 | AF048691 | Oryza sativa          | calcium dependent protein kinase. CDPK12.   |

|               |          |                         |   |
|---------------|----------|-------------------------|---|
| AAF21062.1    | AF216527 | Dunaliella tertiolecta  | calcium-dependent protein kinase. CPK1; CDPK.   |
| AAB80692.1    | U69173   | Glycine max             | calmodulin-like domain protein kinase isoenzyme beta. CDPK beta.  |
| AAK26164.1    | AY027885 | Cucumis sativus         | calcium-dependent calmodulin-independent protein kinase 5. CDPK5. CsCDPK5.  |
| AAA33443.1    | L15390   | Zea mays                | calcium-dependent protein kinase. CDPK.   |
| CAA89202.1    | Z49233   | Chlamydomonas eugametos | calcium-stimulated protein kinase.  |
| AAG46110.1    | AC073166 | Oryza sativa            | calcium-dependent protein kinase. OSJNBb0064P21.2.  |
| BAA02698.1    | D13436   | Oryza sativa            | calcium-dependent protein kinase. spk.  |
| BAA90814.1    | AP001168 | Oryza sativa            | ESTs AU030197(E50746),AU030196(E50746) correspond to a region of the predicted gene.; Similar to calcium-dependent calmodulin-independent protein kinase CDPK (U90262). |
| CAB46228.1    | Y18055   | Arachis hypogaea        | calcium dependent protein kinase. CDPK.   |
| AAC49008.1    | U24188   | Lilium longiflorum      | calcium/calmodulin-dependent phosphorylation activity. calcium/calmodulin-dependent protein kinase. CcCaMK. serine/threonine kinase; binds to calcium and calmodulin.   |
| AAF19401.1    | AF203479 | Glycine max             | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                    |
| AAF06970.1    | AF162662 | Kalanchoe fedtschenkoi  | phosphoenolpyruvate carboxylase kinase. calcium-independent protein kinase.   |
| SEQ ID NO: 39 |          |                         |   |
| AAG00510.1    | AF285172 | Phaseolus vulgaris      | leaf senescence-associated receptor-like protein kinase. SARK.  |
| BAB21175.1    | AP002909 | Oryza sativa            | putative serine/threonine-specific receptor protein kinase. P0044F08.3.   |
| CAB51480.1    | Y14600   | Sorghum bicolor         | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| AAF59905.1    | AF197946 | Glycine max             | receptor protein kinase-like protein. CLV1A.  |
| AAF59906.1    | AF197947 | Glycine max             | receptor protein kinase-like protein. CLV1B.  |
| AAB61708.1    | U93048   | Daucus carota           | somatic embryogenesis receptor-like kinase. SERK.   |
| BAB39873.1    | AP002882 | Oryza sativa            | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAC36318.1 | AF053127 | Malus x domestica             | leucine-rich receptor-like protein kinase. LRPKm1.  |
| AAK21965.1 | AY028699 | Brassica napus                | receptor protein kinase PERK1.  |
| AAF91324.1 | AF244890 | Glycine max                   | receptor-like protein kinase 3. RLK3. GmRLK3.   |
| AAK00425.1 | AC069324 | Oryza sativa                  | Putative protein kinase. OSJNBa0071K19.11.  |
| AAF91323.1 | AF244889 | Glycine max                   | receptor-like protein kinase 2. RLK2. GmRLK2.   |
| AAG59657.1 | AC084319 | Oryza sativa                  | putative protein kinase. OSJNBa0004B24.20.  |
| CAA97692.1 | Z73295   | Catharanthus roseus           | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.  |
| BAB40094.1 | AP003210 | Oryza sativa                  | putative receptor protein kinase. OSJNBa0010K01.7.  |
| CAA61510.1 | X89226   | Oryza sativa                  | leucine-rich repeat/receptor protein kinase. lrk2.  |
| BAA87852.1 | AP000816 | Oryza sativa                  | Similar to putative Ser/Thr protein kinase. (AC004218).   |
| BAA92221.1 | AP001278 | Oryza sativa                  | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).  |
| AAF91322.1 | AF244888 | Glycine max                   | receptor-like protein kinase 1. RLK1. GmRLK1.   |
| AAK11569.1 | AF318493 | Lycopersicon hirsutum         | Pto-like protein kinase D. LhirPtoD.  |
| CAB51834.1 | 00069    | Oryza sativa                  | l1332.5. contains eukaryotic protein kinase domain PF.  |
| BAA90808.1 | AP001168 | Oryza sativa                  | Similar to putative receptor-like protein kinase (AL035679).  |
| AAC48932.1 | U13923   | Lycopersicon pimpinellifolium | Fen. putative serine/threonine protein kinase; similar to product encoded by Lycopersicon pimpinellifolium Pto gene, GenBank Accession Number U02271; Fen is a member of the Pto gene family. |
| AAF76307.1 | AF220602 | Lycopersicon pimpinellifolium | Fen kinase.   |
| AAB47424.1 | U59317   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Fen. Fen. fenthion sensitivity gene from tomato.  |
| AAK11566.1 | AF318490 | Lycopersicon hirsutum         | Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.   |

|               |          |                               |  |
|---------------|----------|-------------------------------|--|
| AAB47423.1    | U59315   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Pto. Pto. bacterial speck disease resistance in tomato; disease resistance gene.                     |
| AAC48914.1    | U02271   | Lycopersicon pimpinellifolium | protein kinase.  |
| AAF76306.1    | AF220602 | Lycopersicon pimpinellifolium | Pto kinase.  |
| BAA87853.1    | AP000816 | Oryza sativa                  | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308). |
| AAF76313.1    | AF220603 | Lycopersicon esculentum       | Pto kinase. LescPth5.  |
| AAB47421.1    | U59316   | Lycopersicon esculentum       | serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.                           |
| AAG03090.1    | AC073405 | Oryza sativa                  | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).   |
| AAA33915.1    | L27821   | Oryza sativa                  | receptor type serine/threonine kinase. protein kinase.   |
| AAF76314.1    | AF220603 | Lycopersicon esculentum       | Fen kinase. Lescfen.   |
| AAK11568.1    | AF318492 | Lycopersicon hirsutum         | Pto-like protein kinase B. LhirPtoB.   |
| AAB47422.1    | U59318   | Lycopersicon esculentum       | serine/threonine protein kinase Fen. fen. allele of Fen, fenthion sensitivity gene from tomato.                                      |
| BAB16871.1    | AP002537 | Oryza sativa                  | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).                           |
| BAA06538.1    | D31737   | Nicotiana tabacum             | protein-serine/threonine kinase.   |
| AAK11567.1    | AF318491 | Lycopersicon hirsutum         | Pto-like protein kinase F. LhirPtoF.   |
| AAF34426.1    | AF172282 | Oryza sativa                  | leucine rich repeat containing protein kinase. DUPR11.16.  |
| AAG25966.1    | AF302082 | Nicotiana tabacum             | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.                |
| BAB21240.1    | AP002953 | Oryza sativa                  | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).   |
| SEQ ID NO: 41 |          |                               |  |
| AAG12987.1    | AF166498 | Lycopersicon esculentum       | sucrose transporter-like protein. SUT2. sucrose sensor.  |

|            |          |                            |  |
|------------|----------|----------------------------|--|
| AAF08330.1 | AF021809 | Vitis vinifera             | putative sucrose transporter. VvSUC12.                                   |
| BAA24071.1 | D87819   | Oryza sativa               | sucrose transporter. OsSUT1.   |
| AAF90181.1 | AF280050 | Oryza sativa subsp. indica | sucrose proton symporter. sucrose transporter. SUT1.                     |
| BAA83501.1 | AB008464 | Zea mays                   | Sucrose Transporter. ZmSUT1.   |
| CAB75882.1 | AJ272309 | Hordeum vulgare            | Transport sucrose through membrane. sucrose transporter 1. sut1.         |
| AAD55269.1 | AF182445 | Vitis vinifera             | sucrose transporter.   |
| CAA83436.1 | Z31561   | Ricinus communis           | sucrose transport. sucrose carrier. Scr1.                                |
| CAA76368.1 | Y16767   | Daucus carota              | sucrose/H <sup>+</sup> symporter. SUT1b.                                 |
| CAA76367.1 | Y16766   | Daucus carota              | sucrose/H <sup>+</sup> symporter. SUT1a.                                 |
| CAC19688.1 | AJ303198 | Daucus carota              | essential for sucrose transport. sucrose/proton symporter. sut1a.        |
| AAG25923.1 | AF237780 | Solanum tuberosum          | sucrose transporter SUT4.  |
| AAF08329.1 | AF021808 | Vitis vinifera             | putative sucrose transporter. VvSUC11.                                   |
| AAG09270.1 | AF176950 | Lycopersicon esculentum    | sucrose transporter. SUT4.   |
| AAF65765.1 | AF242307 | Euphorbia esula            | sucrose transport protein.   |
| AAF04295.1 | AF191025 | Alonsoa meridionalis       | sucrose transporter 1. SUT1.   |
| CAA57727.1 | X82276   | Nicotiana tabacum          | sucrose transporter. NtSUT1a. Var. SNN bp 1-1346, Var. SR1 bp 1172-1614. |
| BAA89458.1 | AB036758 | Daucus carota              | sucrose transporter protein. cSUT.                                       |
| CAB75881.1 | AJ272308 | Hordeum vulgare            | transport sucrose through membrane. sucrose transporter 2. sut2.         |
| CAA76369.1 | Y16768   | Daucus carota              | sucrose/H <sup>+</sup> symporter. SUT2.                                  |
| CAC19689.1 | AJ303199 | Daucus carota              | essential for sucrose transport. sucrose/proton symporter. sut2.         |
| CAB07811.1 | Z93774   | Vicia faba                 | sugar transport. sucrose transport protein. sut.                         |

|               |          |                         |   |
|---------------|----------|-------------------------|---|
| CAA47604.1    | X67125   | Spinacia oleracea       | sucrose permease. S21.                    |
| CAA48915.1    | X69165   | Solanum tuberosum       | sucrose transport protein. SUT1.          |
| AAD41024.1    | AF109922 | Pisum sativum           | sucrose transport protein SUT1.           |
| AAD53000.1    | U64967   | Beta vulgaris           | sucrose-proton symporter. Bv8-6.          |
| AAF08331.1    | AF021810 | Vitis vinifera          | putative sucrose transporter. VvSUC27.    |
| CAA58730.1    | X83850   | Beta vulgaris           | sucrose/proton-symporter. sut1.           |
| AAC99332.1    | AF063400 | Apium graveolens        | sucrose transporter. SUT1.                |
| AAD45391.1    | AF167416 | Apium graveolens        | sucrose transporter SUT2B. SUT2B.         |
| AAD45390.1    | AF167415 | Apium graveolens        | sucrose transporter SUT2A. SUT2A.         |
| AAF04294.1    | AF191024 | Asarina barclaiiana     | sucrose transporter 1. SUT1.              |
| CAA59113.1    | X84379   | Plantago major          | SUC1-sucrose proton symporter. SUC1.      |
| AAD34610.1    | AF149981 | Nicotiana tabacum       | sucrose transporter-like protein. SUT3.   |
| CAA57726.1    | X82275   | Lycopersicon esculentum | sucrose transporter. LeSUT1.              |
| CAA12256.1    | AJ224961 | Ricinus communis        | Sucrose transport. Sucrose carrier. SUT1. |
| CAC33492.1    | AJ310643 | Ricinus communis        | sucrose carrier. sucrose carrier. scr1.   |
| AAD45932.1    | AF168771 | Betula pendula          | sucrose transport protein. SUC1.          |
| CAA53390.1    | X75764   | Plantago major          | sucrose transporter. ptp1.                |
| AAF22281.1    | AF167417 | Apium graveolens        | putative sucrose transporter SUT1. SUT1.  |
| BAA76434.1    | AB025006 | Cicer arietinum         | sucrose transport protein.                |
| SEQ ID NO: 42 |          |                         |   |
| AAC64184.1    | AF095577 | Prunus persica          | endopolygalacturonase.                    |



|            |          |  |
|------------|----------|--|
| CAA54150.1 | X76735   | <i>Prunus persica</i><br>endopolygalacturonase.  |
| AAC28906.1 | AF001003 | <i>Lycopersicon esculentum</i><br>hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 5.<br>TAPG5. expressed in abscission.    |
| AAC28947.1 | AF029230 | <i>Lycopersicon esculentum</i><br>polygalacturonase. TPG6.   |
| AAC70951.1 | AF072732 | <i>Lycopersicon esculentum</i><br>hydrolyzes pectin in the cell wall and middle lamella. polygalacturonase 7. specifically and<br>abundantly expressed in pistils. |
| AAC28905.1 | AF001002 | <i>Lycopersicon esculentum</i><br>hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 4.<br>TAPG4. expressed in abscission.    |
| AAA80489.1 | U23053   | <i>Lycopersicon esculentum</i><br>polygalacturonase precursor.   |
| AAC28903.1 | AF001000 | <i>Lycopersicon esculentum</i><br>hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 1.<br>TAPG1. expressed in abscission.    |
| AAB09575.1 | U70480   | <i>Lycopersicon esculentum</i><br>abscission polygalacturonase. TAPG2.   |
| AAC28904.1 | AF001001 | <i>Lycopersicon esculentum</i><br>hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 2.<br>TAPG2. expressed in abscission.    |
| AAB09576.1 | U70481   | <i>Lycopersicon esculentum</i><br>abscission polygalacturonase. TAPG4.   |
| AAC28902.2 | AF000999 | <i>Lycopersicon esculentum</i><br>hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 3.<br>TAPG3. expressed in abscission.    |
| AAC26510.1 | AF062465 | <i>Cucumis melo</i><br>polygalacturonase precursor. MPG1.  |
| AAC26511.1 | AF062466 | <i>Cucumis melo</i><br>polygalacturonase precursor. MPG2.  |
| BAB21092.1 | AP002819 | <i>Oryza sativa</i><br>putative endopolygalacturonase. P0501G01.21.  |
| AAA62286.1 | U20431   | <i>Medicago sativa</i><br>polygalacturonase.   |
| AAA82167.1 | U09717   | <i>Gossypium hirsutum</i><br>polygalacturonase.  |
| AAA58322.1 | U09805   | <i>Gossypium barbadense</i><br>polygalacturonase.  |
| CAA50336.1 | X71018   | <i>Nicotiana tabacum</i><br>polygalacturonase. NPG. Protein sequence is in conflict with the conceptual translation;<br>G27Y.                                      |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAA50334.1 | X71016   | Nicotiana tabacum       | polygalacturonase. NPG. Protein sequence is in conflict with the conceptual translation; G27.X. |
| CAA50337.1 | X71019   | Nicotiana tabacum       | polygalacturonase. NPG. Protein sequence is in conflict with the conceptual translation; G27W.  |
| CAA65072.1 | X95800   | Brassica napus          | polygalacturonase.  |
| BAA89478.1 | AB029459 | Salix gilgiana          | polygalacturonase. SgPG3.   |
| CAA90272.1 | Z49971   | Brassica napus          | Hydrolytic enzyme. Polygalacturonase. pga.  |
| CAA50335.1 | X71017   | Nicotiana tabacum       | polygalacturonase. Npg1. Protein sequence is in conflict with the conceptual translation.       |
| CAA50338.1 | X71020   | Nicotiana tabacum       | polygalacturonase. Npg1.  |
| BAA89479.1 | AB029460 | Salix gilgiana          | polygalacturonase. SgPG4.   |
| AAC14453.1 | L12019   | Actinidia deliciosa     | polygalacturonase.  |
| BAA89476.1 | AB029457 | Salix gilgiana          | polygalacturonase. SgPG1.   |
| AAC26512.1 | AF062467 | Cucumis melo            | polygalacturonase precursor. MPG3.  |
| CAA72003.1 | Y11118   | Medicago sativa         | polygalacturonase. PG3.   |
| CAB42886.1 | AJ238848 | Phleum pratense         | polygalacturonase. pg.  |
| BAA89477.1 | AB029458 | Salix gilgiana          | polygalacturonase. SgPG2.   |
| CAC05658.1 | AJ250919 | Brassica napus          | endopolygalacturonase. pegaz.   |
| CAC05657.1 | AJ250918 | Brassica napus          | endopolygalacturonase. pgaz.  |
| AAA32914.1 | L06094   | Persea americana        | cell wall degradation. polygalacturonase.   |
| AAF71160.1 | AF152758 | Actinidia chinensis     | polygalacturonase A. PGA.   |
| AAF61444.1 | AF138858 | Lycopersicon esculentum | hydrolyses polygalacturonic acid. polygalacturonase. XOPG1.                                     |
| AAA34178.1 | M37304   | Lycopersicon esculentum | polygalacturonase.  |

|               |          |   |
|---------------|----------|---|
| CAA32235.1    | X14074   | Lycopersicon esculentum<br>polygalacturonase.   |
| CAA29148.1    | X05656   | Lycopersicon esculentum<br>polygalacturonase (AA 1-457).  |
| AAD46483.1    | AF128266 | Glycine max<br>polygalacturonase PG1.   |
| BAA88472.1    | AB035890 | Cucumis sativus<br>polygalacturonase. CUPG1.  |
| AAD46484.1    | AF128267 | Glycine max<br>polygalacturonase PG2.   |
| CAA54448.1    | X77231   | Prunus persica<br>polygalacturonase. PG.  |
| CAA47055.1    | X66426   | Persea americana<br>polygalacturonase.  |
| CAA47052.1    | X66422   | Zea mays<br>polygalacturonase. PG.  |
| SEQ ID NO: 44 |          |   |
| AAC61805.1    | U28007   | Lycopersicon esculentum<br>serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.   |
| BAB21241.1    | AP002953 | Oryza sativa<br>Putative Pto kinase interactor 1. P0426D06.21. contains ESTs<br>AU108280(E0721),D48017(S13927).   |
| AAF91337.1    | AF249318 | Glycine max<br>Pti1 kinase-like protein. Pti1b. protein kinase.   |
| AAF91336.1    | AF249317 | Glycine max<br>Pti1 kinase-like protein. Pti1a. protein kinase.   |
| BAA87853.1    | AP000816 | Oryza sativa<br>EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative<br>NAK-like Ser/Thr protein kinase. (AF001308). |
| BAB16871.1    | AP002537 | Oryza sativa<br>putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs<br>C22608(R3192),D25110(R3192).                           |
| AAG16628.1    | AY007545 | Brassica napus<br>protein serine/threonine kinase BNK1.   |
| BAB39409.1    | AP002901 | Oryza sativa<br>putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| BAB39873.1    | AP002882 | Oryza sativa<br>putative protein kinase. P0439B06.8. contains ESTs<br>AU056701(S20808),AU056702(S20808).  |
| AAC27894.1    | AF023164 | Zea mays<br>leucine-rich repeat transmembrane protein kinase 1. ltk1.   |
| BAA94509.1    | AB041503 | Populus nigra<br>protein kinase 1. PnPK1.   |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| BAB07999.1 | AP002525 | Oryza sativa            | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB03429.1 | AP002817 | Oryza sativa            | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).             |
| AAK21965.1 | AY028699 | Brassica napus          | receptor protein kinase PERK1.  |
| AAC27895.1 | AF023165 | Zea mays                | leucine-rich repeat transmembrane protein kinase 2. ltk2.   |
| AAK00425.1 | AC069324 | Oryza sativa            | Putative protein kinase. OSJNBa0071K19.11.  |
| BAB21240.1 | AP002953 | Oryza sativa            | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| BAA94510.1 | AB041504 | Populus nigra           | protein kinase 2. PnPK2.  |
| AAG03090.1 | AC073405 | Oryza sativa            | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| AAG59657.1 | AC084319 | Oryza sativa            | putative protein kinase. OSJNBa0004B24.20.  |
| BAA78764.1 | AB023482 | Oryza sativa            | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAG25966.1 | AF302082 | Nicotiana tabacum       | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.   |
| AAF43496.1 | AF131222 | Lophopyrum elongatum    | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| AAK11674.1 | AF339747 | Lophopyrum elongatum    | protein kinase. ESI47.  |
| AAB09771.1 | U67422   | Zea mays                | CRINKLY4 precursor. cr4. receptor kinase homolog.   |
| CAA97692.1 | Z73295   | Catharanthus roseus     | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.  |
| BAA90808.1 | AP001168 | Oryza sativa            | Similar to putative receptor-like protein kinase (AL035679).  |
| AAB47421.1 | U59316   | Lycopersicon esculentum | serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.  |
| AAF76313.1 | AF220603 | Lycopersicon esculentum | Pto kinase. LescPth5.   |

|               |          |                               |  |
|---------------|----------|-------------------------------|--|
| CAB51834.1    | 00069    | Oryza sativa                  | 11332.5. contains eukaryotic protein kinase domain PF.   |
| AAB47423.1    | U59315   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Pto. Pto. bacterial speck disease resistance in tomato; disease resistance gene. |
| AAF76306.1    | AF220602 | Lycopersicon pimpinellifolium | Pto kinase.  |
| AAC48914.1    | U02271   | Lycopersicon pimpinellifolium | protein kinase.  |
| BAA92221.1    | AP001278 | Oryza sativa                  | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).   |
| BAA87852.1    | AP000816 | Oryza sativa                  | Similar to putative Ser/Thr protein kinase. (AC004218).  |
| AAF66615.1    | AF142596 | Nicotiana tabacum             | LRR receptor-like protein kinase.  |
| AAK11566.1    | AF318490 | Lycopersicon hirsutum         | Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.                              |
| AAK11567.1    | AF318491 | Lycopersicon hirsutum         | Pto-like protein kinase F. LhirPtoF.   |
| CAB51836.1    | AJ243961 | Oryza sativa                  | Putative Ser/Thr protein kinase. 11332.7.  |
| AAD38286.1    | AC007789 | Oryza sativa                  | putative protein kinase. OSJNBa0049B20.13.   |
| BAB40081.1    | AP003074 | Oryza sativa                  | putative receptor protein kinase. OSJNBa0004G10.30.  |
| BAB18321.1    | AP002865 | Oryza sativa                  | putative receptor protein kinase. P0034C11.11.   |
| BAA92836.1    | AB032473 | Brassica oleracea             | S18 S-locus receptor kinase. SRK18.  |
| SEQ ID NO: 45 |          |                               |  |
| AAB72110.1    | U79958   | Pisum sativum                 | BP-80 vacuolar sorting receptor.   |
| AAF80450.1    | AF161719 | Triticum aestivum             | vacuolar targeting receptor bp-80.   |
| AAG60258.1    | AY017377 | Physcomitrella patens         | EGF receptor-like protein. ELP. PPELP; similar to putative vacuolar sorting receptor.                            |
| AAF22842.1    | AF209910 | Prunus dulcis                 | vacuolar sorting receptor protein. BP-80 type protein.   |
| AAK31596.1    | AY029172 | Helianthus annuus             | EGF receptor-like protein. ELP.  |

|               |          |   |   |
|---------------|----------|---|---|
| AAB72113.1    | U79961   | <i>Zea mays</i>                                     | vacuolar sorting receptor homolog. similar to <i>Pisum sativum</i> BP-80 vacuolar sorting receptor, GenBank Accession Number U79958.  |
| BAA92985.1    | AP001550 | <i>Oryza sativa</i>                                 | EST D15617(C0960A) corresponds to a region of the predicted gene.; Similar to <i>Arabidopsis thaliana</i> chromosome 4, BAC clone F18F4; vacuolar sorting receptor-like protein (AL021637). |
| SEQ ID NO: 47 |          |   |   |
| BAA00885.1    | D10001   | <i>Pisum sativum</i>                                | phenylalanine ammonia-lyase.  |
| AAK15640.1    | AF326116 | <i>Agastache rugosa</i>                             | phenylalanine ammonia-lyase. PAL.   |
| CAB42794.1    | AJ238754 | <i>Citrus clementina</i> x <i>Citrus reticulata</i> | phenylalanine-ammonia lyase. pal2.  |
| BAA00886.1    | D10002   | <i>Pisum sativum</i>                                | phenylalanine ammonia-lyase. PAL1.  |
| BAA00887.1    | D10003   | <i>Pisum sativum</i>                                | phenylalanine ammonia-lyase. PAL2.  |
| AAB67733.1    | U43338   | <i>Citrus limon</i>                                 | phenylalanine ammonia-lyase. pal6.  |
| AAF40224.1    | AF237955 | <i>Rubus idaeus</i>                                 | phenylalanine ammonia-lyase 2. PAL2. PAL; phenylpropanoid; multigene; flavonoid.  |
| AAA17993.1    | M91192   | <i>Trifolium subterraneum</i>                       | phenylalanine ammonia-lyase. PAL1.  |
| CAA57057.1    | X81159   | <i>Petroselinum crispum</i>                         | phenylalanine ammonia-lyase 3. PAL3. tetramere subunit.   |
| AAF40223.1    | AF237954 | <i>Rubus idaeus</i>                                 | phenylalanine ammonia-lyase 1. PAL1. PAL; phenylpropanoid; multigene; flavonoid.  |
| CAA68938.1    | Y07654   | <i>Petroselinum crispum</i>                         | PAL1 protein. pal1 gene.  |
| CAA57056.1    | X81158   | <i>Petroselinum crispum</i>                         | phenylalanine ammonia-lyase 2. PAL2. deaminase subunit.   |
| CAB60719.1    | AJ250836 | <i>Cicer arietinum</i>                              | phenylpropanoid pathway. phenylalanine ammonia-lyase. pal.  |
| CAA05251.1    | AJ002221 | <i>Digitalis lanata</i>                             | phenylalanine ammonia lyase.  |
| AAC78457.1    | AF036948 | <i>Prunus avium</i>                                 | phenylalanine ammonia-lyase. PAL1.  |
| BAA23367.1    | D85850   | <i>Daucus carota</i>                                | phenylalanine ammonia-lyase. gDcPAL1.   |
| CAA68256.1    | X99997   | <i>Bromheadia finlaysoniana</i>                     | phenylalanine ammonia-lyase. pal.   |

|            |          |                                       |                                       |
|------------|----------|---------------------------------------|---------------------------------------|
| AAA33805.1 | L11747   | Populus x generosa                    | phenylalanine ammonia lyase. PAL.     |
| BAA24928.1 | D83075   | Lithospermum erythrorhizon            | phenylalanine ammonia-lyase.          |
| CAA34226.1 | X16099   | Oryza sativa subsp. japonica          | phenylalanine ammonia-lyase.          |
| CAA41169.1 | X58180   | Medicago sativa                       | phenylalanine ammonia-lyase. PAL.     |
| AAA34176.1 | M90692   | Lycopersicon esculentum               | phenylalanine ammonia-lyase. PAL5.    |
| BAA07860.1 | D43802   | Populus kitakamiensis                 | phenylalanine ammonia-lyase.          |
| AAA84889.1 | U39792   | Pinus taeda                           | phenylalanine ammonia-lyase. lpPAL.   |
| CAA73065.1 | Y12461   | Helianthus annuus                     | phenylalanine ammonia lyase. PAL.     |
| BAA95629.1 | AB042520 | Catharanthus roseus                   | phenylalanine ammonia lyase.          |
| BAA05643.1 | D26596   | Camellia sinensis                     | phenylalanine ammonia-lyase.          |
| BAA24929.1 | D83076   | Lithospermum erythrorhizon            | phenylalanine ammonia-lyase.          |
| BAA21643.1 | D30656   | Populus kitakamiensis                 | phenylalanine ammonia-lyase.          |
| CAA37129.1 | X52953   | Glycine max                           | phenylalanine ammonia-lyase. PAL1.    |
| AAA34122.1 | M84466   | Nicotiana tabacum                     | phenylalanine ammonia lyase. tpal.    |
| BAA22948.1 | AB008200 | Nicotiana tabacum                     | phenylalanine ammonia-lyase. palB.    |
| AAA99500.1 | L36822   | Stylosanthes humilis                  | phenylalanine ammonia lyase. PAL17.1. |
| CAB42793.1 | AJ238753 | Citrus clementina x Citrus reticulata | phenylalanine-ammonia lyase. pal1.    |
| AAG49585.1 | AF325496 | Ipomoea nil                           | phenylalanine ammonia-lyase.          |
| CAA55075.1 | X78269   | Nicotiana tabacum                     | phenylalanine ammonia-lyase.          |
| BAA22963.1 | D17467   | Nicotiana tabacum                     | phenylalanine ammonia-lyase. TOBPAL1. |
| BAA22947.1 | AB008199 | Nicotiana tabacum                     | phenylalanine ammonia-lyase. palA.    |

|               |          |  |  |
|---------------|----------|--|--|
| AAA34179.2    | M83314   | Lycopersicon esculentum                | deamination of phenylalanine to coumarate. phenylalanine ammonia lyase. pal.   |
| BAA11459.1    | D78640   | Ipomoea batatas                        | Phenylalanine Ammonia-Lyase.   |
| AAA33389.1    | M29232   | Ipomoea batatas                        | phenylalanine ammonia-lyase.   |
| CAA68036.1    | X99705   | Triticum aestivum                      | phenylalanine ammonia-lyase. PAL.  |
| CAA61198.1    | X87946   | Oryza sativa                           | phenylalanine ammonia-lyase. ZB8.  |
| BAA06337.1    | D30657   | Populus kitakamiensis                  | phenylalanine ammonia-lyase.   |
| AAD45384.1    | AF165998 | Vigna unguiculata                      | phenylalanine ammonia-lyase.   |
| CAA53733.1    | X76130   | Cucumis melo                           | phenylalanine ammonia-lyase. pal.  |
| AAA51873.1    | U16130   | Persea americana                       | phenylalanine ammonia lyase. PAL.  |
| BAB19128.1    | AB041361 | Dianthus caryophyllus                  | phenylalanine ammonia-lyase. Dcpal1.   |
| CAA34715.1    | X16772   | Petroselinum crispum                   | phenylalanine ammonia-lyase (AA 137 - 716) (1 is 3rd base in codon).   |
| BAA07861.1    | D43803   | Populus kitakamiensis                  | phenylalanine ammonia-lyase.   |
| SEQ ID NO: 48 |          |  |  |
| CAB94692.1    | AJ242742 | Ipomoea batatas                        | Removal of H <sub>2</sub> O <sub>2</sub> , oxidation of toxic reductants, defence response toward wounding. peroxidase. pod. |
| AAD37430.1    | AF149280 | Phaseolus vulgaris                     | peroxidase 5 precursor. FBP5. secretory peroxidase.  |
| CAA66037.1    | X97351   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.   |
| BAA06335.1    | D30653   | Populus kitakamiensis                  | peroxidase.  |
| AAA34108.1    | J02979   | Nicotiana tabacum                      | lignin-forming peroxidase precursor (EC 1.11.1.7).   |
| BAA01992.1    | D11396   | Nicotiana tabacum                      | 'peroxidase'.  |
| CAA50597.1    | X71593   | Lycopersicon esculentum                | peroxidase. CEVI-1.  |
| CAB67121.1    | Y19023   | Lycopersicon esculentum                | peroxidase. cevi-1.  |



|            |          |  |
|------------|----------|--|
| BAA11853.1 | D83225   | Populus nigra<br>peroxidase.   |
| CAA66034.1 | X97348   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA01877.1 | D11102   | Populus kitakamiensis<br>peroxidase. prxA1.  |
| BAA11852.1 | D83224   | Populus nigra<br>peroxidase.   |
| CAA66035.1 | X97349   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| AAB47602.1 | L07554   | Linum usitatissimum<br>peroxidase. FLXPER1.  |
| AAD37427.1 | AF149277 | Phaseolus vulgaris<br>peroxidase 1 precursor. FBP1. secretory peroxidase.  |
| CAA66036.1 | X97350   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| AAB97734.1 | AF014502 | Glycine max<br>seed coat peroxidase precursor. Ep. H2O2 oxidoreductase; class III plant peroxidase.  |
| BAA06334.1 | D30652   | Populus kitakamiensis<br>peroxidase.   |
| BAA07241.1 | D38051   | Populus kitakamiensis<br>peroxidase. prxA4a.   |
| CAA62226.1 | X90693   | Medicago sativa<br>peroxidase1B. prx1B.  |
| CAA62225.1 | X90692   | Medicago sativa<br>peroxidase1A. prx1A.  |
| CAA62227.1 | X90694   | Medicago sativa<br>peroxidase1C. prx1C.  |
| AAB41811.1 | L36157   | Medicago sativa<br>peroxidase. pxdC. amino acid feature: conserved domains, aa 123 .. 129, 191 .. 198; amino acid feature: heme-binding domain, aa 68 .. 73. |
| AAB41810.1 | L36156   | Medicago sativa<br>peroxidase. pxdA. amino acid feature: conserved motifs, aa 181 .. 188; amino acid feature: heme-binding domain, aa 60 .. 65.              |
| AAC98519.1 | AF007211 | Glycine max<br>peroxidase precursor. GMIPER1. pathogen-induced.  |
| BAA02840.1 | D13683   | Populus kitakamiensis<br>peroxidase. HPOX14.   |
| BAA14144.1 | D90116   | Armoracia rusticana<br>peroxidase isozyme.   |
| AAA33129.1 | M91372   | Cucumis sativus<br>peroxidase. pre-peroxidase.   |

|               |          |                         |  |
|---------------|----------|-------------------------|--|
| BAA14143.1    | D90115   | Armoracia rusticana     | peroxidase isozyme.  |
| BAA08499.1    | D49551   | Oryza sativa            | peroxidase. poxN.  |
| BAA03373.1    | D14482   | Oryza sativa            | putative peroxidase.   |
| AAA34101.1    | L02124   | Nicotiana tabacum       | peroxidase.  |
| BAA82306.1    | AB027752 | Nicotiana tabacum       | peroxidase.  |
| CAA40796.1    | X57564   | Armoracia rusticana     | peroxidase. peroxidase precursor.  |
| AAA33121.1    | M32742   | Cucumis sativus         | peroxidase (CuPer2).   |
| CAA76680.1    | Y17192   | Cucurbita pepo          | peroxidase. aprx. type III peroxidase.   |
| BAA92500.1    | AP001383 | Oryza sativa            | ESTs D39300(R3292),AU030751(E60187) correspond to a region of the predicted gene. Similar to peroxidase ATP6a. (X98774). |
| BAA77388.1    | AB024438 | Scutellaria baicalensis | peroxidase 2.  |
| AAA33127.1    | M91373   | Cucumis sativus         | peroxidase. pre-peroxidase. putative.  |
| AAD43561.1    | AF155124 | Gossypium hirsutum      | bacterial-induced peroxidase precursor. Perx_Goshiko.  |
| AAB06183.1    | M37636   | Arachis hypogaea        | cationic peroxidase. PNC1.   |
| AAF63027.1    | AF244924 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.   |
| CAA71492.1    | Y10466   | Spinacia oleracea       | peroxidase. prxr5.   |
| AAB02554.1    | L37790   | Stylosanthes humilis    | cationic peroxidase.   |
| BAA94962.1    | AB042103 | Asparagus officinalis   | peroxidase. AspPOX1.   |
| BAA77389.1    | AB024439 | Scutellaria baicalensis | peroxidase 3.  |
| SEQ ID NO: 50 |          |                         |  |
| CAA40796.1    | X57564   | Armoracia rusticana     | peroxidase. peroxidase precursor.  |
| BAA03373.1    | D14482   | Oryza sativa            | putative peroxidase.   |

|            |          |  |
|------------|----------|--|
| BAA08499.1 | D49551   | Oryza sativa<br>peroxidase. poxN.  |
| CAA66037.1 | X97351   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA84764.1 | D84400   | Oryza sativa<br>peroxidase. poxA.  |
| BAA03372.1 | D14481   | Oryza sativa<br>putative peroxidase.   |
| CAB94692.1 | AJ242742 | Ipomoea batatas<br>Removal of H <sub>2</sub> O <sub>2</sub> , oxidation of toxic reductants, defence response toward wounding.<br>peroxidase. pod. |
| BAA06335.1 | D30653   | Populus kitakamiensis<br>peroxidase.   |
| BAA11853.1 | D83225   | Populus nigra<br>peroxidase.   |
| AAD37427.1 | AF149277 | Phaseolus vulgaris<br>peroxidase 1 precursor. FBP1. secretory peroxidase.  |
| AAD37430.1 | AF149280 | Phaseolus vulgaris<br>peroxidase 5 precursor. FBP5. secretory peroxidase.  |
| BAA11852.1 | D83224   | Populus nigra<br>peroxidase.   |
| CAA66035.1 | X97349   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| CAA66034.1 | X97348   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA07241.1 | D38051   | Populus kitakamiensis<br>peroxidase. prxA4a.   |
| BAA92500.1 | AP001383 | Oryza sativa<br>ESTs D39300(R3292),AU030751(E60187) correspond to a region of the predicted gene.<br>Similar to peroxidase ATP6a. (X98774).        |
| CAA50597.1 | X71593   | Lycopersicon esculentum<br>peroxidase. CEVI-1.   |
| CAB67121.1 | Y19023   | Lycopersicon esculentum<br>peroxidase. cevi-1.   |
| CAA66036.1 | X97350   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| AAB47602.1 | L07554   | Linum usitatissimum<br>peroxidase. FLXPER1.  |
| CAA62226.1 | X90693   | Medicago sativa<br>peroxidase1B. prx1B.  |
| AAA34108.1 | J02979   | Nicotiana tabacum<br>lignin-forming peroxidase precursor (EC 1.11.1.7).  |

|            |          |  |
|------------|----------|--|
| CAA62227.1 | X90694   | Medicago sativa<br>peroxidase1C. prx1C.  |
| BAA06334.1 | D30652   | Populus kitakamiensis<br>peroxidase.   |
| BAA01992.1 | D11396   | Nicotiana tabacum<br>'peroxidase'.   |
| AAD43561.1 | AF155124 | Gossypium hirsutum<br>bacterial-induced peroxidase precursor. Perx_Goshiko.  |
| BAA82306.1 | AB027752 | Nicotiana tabacum<br>peroxidase.   |
| AAC98519.1 | AF007211 | Glycine max<br>peroxidase precursor. GMIPER1. pathogen-induced.  |
| BAA14143.1 | D90115   | Armoracia rusticana<br>peroxidase isozyme.   |
| AAB97734.1 | AF014502 | Glycine max<br>seed coat peroxidase precursor. Ep. H2O2 oxidoreductase; class III plant peroxidase.  |
| AAF63027.1 | AF244924 | Spinacia oleracea<br>hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.  |
| CAA62225.1 | X90692   | Medicago sativa<br>peroxidase1A. prx1A.  |
| BAA01877.1 | D11102   | Populus kitakamiensis<br>peroxidase. prxA1.  |
| AAF63026.1 | AF244923 | Spinacia oleracea<br>hydrogen peroxide catabolism. peroxidase prx14 precursor. type III peroxidase.  |
| BAA14144.1 | D90116   | Armoracia rusticana<br>peroxidase isozyme.   |
| AAB41810.1 | L36156   | Medicago sativa<br>peroxidase. pxdA. amino acid feature: conserved motifs, aa 181 .. 188; amino acid feature:<br>heme-binding domain, aa 60 .. 65. |
| AAA34050.1 | M74103   | Nicotiana sylvestris<br>anionic peroxidase.  |
| CAA62597.1 | X91172   | Raphanus sativus<br>korean-radish isoperoxidase. prxk1.  |
| AAB02554.1 | L37790   | Stylosanthes humilis<br>cationic peroxidase.   |
| CAA76680.1 | Y17192   | Cucurbita pepo<br>peroxidase. aprx. type III peroxidase.   |
| CAA71492.1 | Y10466   | Spinacia oleracea<br>peroxidase. prxr5.  |
| BAA94962.1 | AB042103 | Asparagus officinalis<br>peroxidase. AspPOX1.  |
| BAA77389.1 | AB024439 | Scutellaria baicalensis<br>peroxidase 3.   |

|   |          |                         |
|---|----------|-------------------------|
| AAD37428.1  | AF149278 | Phaseolus vulgaris      |
| peroxidase 3 precursor. FBP3. secretory peroxidase.   |          |                         |
| AAB41811.1  | L36157   | Medicago sativa         |
| peroxidase. pxdC. amino acid feature: conserved domains, aa 123 .. 129, 191 .. 198; amino acid feature: heme-binding domain, aa 68 .. 73. |          |                         |
| AAB06183.1  | M37636   | Arachis hypogaea        |
| cationic peroxidase. PNC1.  |          |                         |
| SEQ ID NO: 52   |          |                         |
| BAA02112.1  | D12544   | Pisum sativum           |
| GTP-binding protein.  |          |                         |
| CAA98184.1  | Z73956   | Lotus japonicus         |
| GTP-binding protein. RAB11H. rab11H.  |          |                         |
| AAK15703.1  | AF327517 | Oryza sativa            |
| GTP-binding protein.  |          |                         |
| BAA02904.1  | D13758   | Oryza sativa            |
| ras-related GTP binding protein. ss230.   |          |                         |
| BAA02111.1  | D12543   | Pisum sativum           |
| GTP-binding protein.  |          |                         |
| BAA02113.1  | D12545   | Pisum sativum           |
| GTP-binding protein.  |          |                         |
| CAA98180.1  | Z73952   | Lotus japonicus         |
| GTP-binding protein. RAB11D. rab11D.  |          |                         |
| CAB65172.1  | AJ245570 | Lycopersicon esculentum |
| putative role in secretion of cell wall modifying enzymes. Rab11 GTPase. Rab11a.  |          |                         |
| BAA02114.1  | D12546   | Pisum sativum           |
| GTP-binding protein.  |          |                         |
| CAA98181.1  | Z73953   | Lotus japonicus         |
| GTP-binding protein. RAB11E. rab11E.  |          |                         |
| CAA95859.1  | Z71276   | Mangifera indica        |
| small GTPase. rabX. homologous to Rab11.  |          |                         |
| CAA55865.1  | X79278   | Medicago sativa         |
| GTP binding protein. Rab.   |          |                         |
| CAA89049.1  | Z49190   | Beta vulgaris           |
| GTP-binding. small G protein.   |          |                         |
| CAA98179.1  | Z73951   | Lotus japonicus         |
| GTP-binding protein. RAB11C. rab11C.  |          |                         |
| BAA02437.1  | D13152   | Oryza sativa            |
| GTP binding protein. rgp2.  |          |                         |
| BAA06701.1  | D31905   | Zea mays                |
| mgp1 GTP-binding protein. mgp1.   |          |                         |
| BAA06702.1  | D31906   | Zea mays                |
| mgp2 GTP-binding protein. mgp2.   |          |                         |

|            |          |                    |  |
|------------|----------|--------------------|--|
| BAA02110.1 | D12542   | Pisum sativum      | GTP-binding protein.   |
| CAA98177.1 | Z73949   | Lotus japonicus    | GTP-binding protein. RAB11A. rab11A.   |
| CAA41966.1 | X59276   | Oryza sativa       | GTP-binding protein. rgp1.   |
| AAB97114.1 | U58853   | Glycine max        | small GTP-binding protein. sra1.   |
| CAA98185.1 | Z73957   | Lotus japonicus    | GTP-binding protein. RAB11I. rab11I.   |
| CAA67153.1 | X98540   | Fagus sylvatica    | FSGTP1.  |
| CAA98183.1 | Z73955   | Lotus japonicus    | GTP-binding protein. RAB11G. rab11G.   |
| CAA98182.1 | Z73954   | Lotus japonicus    | GTP-binding protein. RAB11F. rab11F.   |
| CAA54506.1 | X77301   | Glycine max        | GTPase. gmr2.  |
| BAA02108.1 | D12540   | Pisum sativum      | GTP-binding protein.   |
| AAA68983.1 | L12395   | Brassica napus     | signal transduction, membrane vehicle traffic. small GTP-binding protein. bra. putative.       |
| AAD48018.1 | AF165095 | Gossypium hirsutum | Rab GTP-binding protein Rab11a.  |
| BAA02109.1 | D12541   | Pisum sativum      | GTP-binding protein.   |
| CAA98186.1 | Z73958   | Lotus japonicus    | GTP-binding protein. RAB11J. rab11J.   |
| BAA84640.1 | AB007911 | Pisum sativum      | PRA2. pra2. light-repressible GTP binding protein.   |
| AAD48019.1 | AF165096 | Gossypium hirsutum | Rab GTP-binding protein Rab11b.  |
| CAA98178.1 | Z73950   | Lotus japonicus    | GTP-binding protein. RAB11B. rab11B.   |
| AAA63901.1 | U22432   | Zea mays           | GTP binding protein. rab2.   |
| CAA98165.1 | Z73937   | Lotus japonicus    | GTP-binding protein. RAB2A. rab2A.   |
| AAA34253.1 | L08130   | Volvox carteri     | GTP-binding protein. yptV4.  |
| AAA90955.1 | U32185   | Glycine max        | vesicular transport. guanine nucleotide regulatory protein. rab2. GTP-binding protein; soyrab. |

|  |          |                            |
|--|----------|----------------------------|
| AAA63902.1   | U22433   | Zea mays                   |
| GTP binding protein. rab2.   |          |                            |
| SEQ ID NO: 53  |          |                            |
| CAA64327.1   | X94624   | Brassica napus             |
| acyl-CoA synthetase.   |          |                            |
| CAA96523.1   | Z72153   | Brassica napus             |
| acyl CoA synthetase.   |          |                            |
| CAC19877.1   | AJ401089 | Brassica napus             |
| activation of free fatty acids. long chain acyl-CoA synthetase. acs6. activity confirmed by expression in E. coli. |          |                            |
| CAA06820.1   | AJ006025 | Cicer arietinum            |
| acyl-coA synthetase.   |          |                            |
| BAA08365.1   | D49366   | Lithospermum erythrorhizon |
| 4-coumarate:CoA ligase.  |          |                            |
| CAA36850.1   | X52623   | Oryza sativa               |
| 4-coumarate-CoA ligase.  |          |                            |
| AAF37734.1   | AF052223 | Lolium perenne             |
| 4-coumarate--CoA ligase 4CL3.  |          |                            |
| AAF91309.1   | AF239686 | Rubus idaeus               |
| 4-coumarate:coA ligase 2. adenylate-forming enzyme; 4CL2.  |          |                            |
| AAC24503.1   | AF041049 | Populus tremuloides        |
| 4-coumarate:CoA ligase.  |          |                            |
| BAA08366.2   | D49367   | Lithospermum erythrorhizon |
| 4-coumarate:CoA ligase.  |          |                            |
| AAF91308.1   | AF239685 | Rubus idaeus               |
| 4-coumarate:coA ligase 3. adenylate-forming enzyme; 4CL3.  |          |                            |
| AAC39366.1   | AF008184 | Populus x generosa         |
| 4-coumarate:CoA ligase 1. 4CL1.  |          |                            |
| CAA31696.1   | X13324   | Petroselinum crispum       |
| 4-coumarate:CoA ligase Pc4Cl-1 (AA 1-544).   |          |                            |
| AAC39365.1   | AF008183 | Populus x generosa         |
| 4-coumarate:CoA ligase 2. 4CL2.  |          |                            |
| AAA33842.1   | M62755   | Solanum tuberosum          |
| 4-coumarate--CoA ligase. St4Cl-1.  |          |                            |
| CAA31697.1   | X13325   | Petroselinum crispum       |
| 4-coumarate:CoA ligase Pc4Cl-2 (AA 1-544).   |          |                            |
| AAB42383.1   | U39405   | Pinus taeda                |
| 4-coumarate:CoA ligase. lp4CL-1.   |          |                            |
| AAB42382.1   | U39404   | Pinus taeda                |
| 4-coumarate:CoA ligase. lp4CL-2.   |          |                            |
| AAA92669.1   | U12013   | Pinus taeda                |
| 4-coumarate-CoA ligase enzyme.   |          |                            |

|               |          |                         |   |
|---------------|----------|-------------------------|---|
| AAF91310.1    | AF239687 | Rubus idaeus            | 4-coumarate:coA ligase 1. adenylate-forming enzyme; 4CL1. |
| AAA92668.1    | U12012   | Pinus taeda             | 4-coumarate-CoA ligase enzyme.                            |
| AAF37732.1    | AF052221 | Lolium perenne          | 4-coumarate--CoA ligase 4CL1.                             |
| CAA49575.1    | X69954   | Glycine max             | 4-coumarate--CoA ligase.                                  |
| AAC24504.1    | AF041050 | Populus tremuloides     | 4-coumarate:CoA ligase.                                   |
| AAF37733.1    | AF052222 | Lolium perenne          | 4-coumarate--CoA ligase 4CL2.                             |
| CAB97359.1    | AJ278455 | Juglans nigra           | 4-coumarate-CoA ligase. 4CL.                              |
| AAA69580.1    | L43362   | Oryza sativa            | 4-coumarate:CoA ligase isoform 2. 4cl.2. putative.        |
| AAG46175.1    | AC018727 | Oryza sativa            | putative 4-coumarate CoA ligase. OSJNBa0056G17.30.        |
| AAF73997.2    | AF144504 | Picea smithiana         | 4-coumarate:CoA ligase. 4CL.                              |
| AAF73995.2    | AF144502 | Pinus armandii          | 4-coumarate:CoA ligase. 4CL.                              |
| AAF73998.2    | AF144505 | Cathaya argyrophylla    | 4-coumarate:CoA ligase. 4CL.                              |
| AAF73994.2    | AF144501 | Pinus armandii          | 4-coumarate:CoA ligase. 4CL.                              |
| AAF73996.2    | AF144503 | Pinus armandii          | 4-coumarate:CoA ligase. 4CL.                              |
| SEQ ID NO: 55 |          |                         |   |
| CAA87068.1    | Z46944   | Citrus sinensis         | non-photosynthetic ferredoxin.                            |
| AAK15005.1    | AF233452 | Impatiens balsamina     | ferredoxin. plastidal protein.                            |
| AAA33461.1    | M73831   | Zea mays                | ferredoxin.   |
| CAB65696.1    | AJ270962 | Lycopersicon esculentum | electron transfer. putative ferredoxin. ferredoxin.       |
| BAA90760.1    | AB038037 | Ipomoea nil             | non-photosynthetic ferredoxin.                            |
| BAA06456.1    | D30794   | Oryza sativa            | ferredoxin.   |



|               |          |                               |  |
|---------------|----------|-------------------------------|--|
| AAB61593.1    | AF003125 | Mesembryanthemum crystallinum | ferredoxin I precursor.  |
| AAA33665.1    | M31713   | Pisum sativum                 | ferredoxin I precursor.  |
| AAD02175.1    | AF039662 | Capsicum annuum               | delays the harpin-mediated hypersensitive response. ferredoxin-like protein. ap1.  |
| CAA73265.1    | Y12734   | Physcomitrella patens         | ferredoxin.  |
| CAA52980.1    | X75089   | Triticum aestivum             | ferredoxin. petF.  |
| BAA06436.1    | D30763   | Oryza sativa                  | ferredoxin.  |
| CAA99756.1    | Z75520   | Lycopersicon esculentum       | transfer of electrons in a wide variety of metabolic reactions. ferredoxin-I. precursor.                                     |
| AAA33462.1    | M73828   | Zea mays                      | ferredoxin.  |
| AAA34028.1    | M35660   | Spinacia oleracea             | ferredoxin I precursor.  |
| CAA26281.1    | X02432   | Silene latifolia subsp. alba  | ferredoxin precursor.  |
| AAA33459.1    | M73829   | Zea mays                      | ferredoxin.  |
| AAA33460.1    | M73830   | Zea mays                      | ferredoxin.  |
| BAA32348.1    | AB016810 | Zea mays                      | ferredoxin. pFD2.  |
| BAA19865.1    | D83660   | Oryza sativa                  | root ferredoxin.   |
| AAA33085.1    | L10349   | Chlamydomonas reinhardtii     | ferredoxin. precursor.   |
| AAC49171.1    | U29516   | Chlamydomonas reinhardtii     | ferredoxin precursor. Method: conceptual translation supplied by author.   |
| AAB65699.1    | AF010320 | Oryza sativa                  | ferredoxin.  |
| AAK14422.1    | AC087851 | Oryza sativa                  | putative ferredoxin. OSJNBb0072E24.16.   |
| SEQ ID NO: 57 |          |                               |  |
| CAB94692.1    | AJ242742 | Ipomoea batatas               | Removal of H <sub>2</sub> O <sub>2</sub> , oxidation of toxic reductants, defence response toward wounding. peroxidase. pod. |
| AAD37430.1    | AF149280 | Phaseolus vulgaris            | peroxidase 5 precursor. FBP5. secretory peroxidase.  |

|            |          |  |
|------------|----------|--|
| CAA66037.1 | X97351   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA06335.1 | D30653   | Populus kitakamiensis<br>peroxidase.   |
| AAA34108.1 | J02979   | Nicotiana tabacum<br>lignin-forming peroxidase precursor (EC 1.11.1.7).  |
| BAA01992.1 | D11396   | Nicotiana tabacum<br>'peroxidase'.   |
| CAA66034.1 | X97348   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| AAB47602.1 | L07554   | Linum usitatissimum<br>peroxidase. FLXPER1.  |
| BAA11853.1 | D83225   | Populus nigra<br>peroxidase.   |
| CAA50597.1 | X71593   | Lycopersicon esculentum<br>peroxidase. CEVI-1.   |
| CAB67121.1 | Y19023   | Lycopersicon esculentum<br>peroxidase. cevi-1.   |
| BAA11852.1 | D83224   | Populus nigra<br>peroxidase.   |
| CAA66035.1 | X97349   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| AAD37427.1 | AF149277 | Phaseolus vulgaris<br>peroxidase 1 precursor. FBP1. secretory peroxidase.  |
| BAA01877.1 | D11102   | Populus kitakamiensis<br>peroxidase. prxA1.  |
| CAA66036.1 | X97350   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| CAA62225.1 | X90692   | Medicago sativa<br>peroxidase1A. prx1A.  |
| AAB97734.1 | AF014502 | Glycine max<br>seed coat peroxidase precursor. Ep. H2O2 oxidoreductase; class III plant peroxidase.  |
| CAA62226.1 | X90693   | Medicago sativa<br>peroxidase1B. prx1B.  |
| CAA62227.1 | X90694   | Medicago sativa<br>peroxidase1C. prx1C.  |
| BAA06334.1 | D30652   | Populus kitakamiensis<br>peroxidase.   |
| BAA07241.1 | D38051   | Populus kitakamiensis<br>peroxidase. prxA4a.   |
| AAB41810.1 | L36156   | Medicago sativa<br>peroxidase. pxdA. amino acid feature: conserved motifs, aa 181 .. 188; amino acid feature:<br>heme-binding domain, aa 60 .. 65. |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAB41811.1 | L36157   | Medicago sativa         | peroxidase. pxdC. amino acid feature: conserved domains, aa 123 .. 129, 191 .. 198; amino acid feature: heme-binding domain, aa 68 .. 73. |
| BAA14144.1 | D90116   | Armoracia rusticana     | peroxidase isozyme.   |
| AAC98519.1 | AF007211 | Glycine max             | peroxidase precursor. GMIPER1. pathogen-induced.  |
| BAA02840.1 | D13683   | Populus kitakamiensis   | peroxidase. HPOX14.   |
| BAA14143.1 | D90115   | Armoracia rusticana     | peroxidase isozyme.   |
| AAA33129.1 | M91372   | Cucumis sativus         | peroxidase. pre-peroxidase.   |
| BAA08499.1 | D49551   | Oryza sativa            | peroxidase. poxN.   |
| BAA03373.1 | D14482   | Oryza sativa            | putative peroxidase.  |
| AAA34101.1 | L02124   | Nicotiana tabacum       | peroxidase.   |
| CAA76680.1 | Y17192   | Cucurbita pepo          | peroxidase. aprx. type III peroxidase.  |
| CAA40796.1 | X57564   | Armoracia rusticana     | peroxidase. peroxidase precursor.   |
| AAA33121.1 | M32742   | Cucumis sativus         | peroxidase (CuPer2).  |
| BAA82306.1 | AB027752 | Nicotiana tabacum       | peroxidase.   |
| AAD43561.1 | AF155124 | Gossypium hirsutum      | bacterial-induced peroxidase precursor. Perx_Goshiko.   |
| AAB06183.1 | M37636   | Arachis hypogaea        | cationic peroxidase. PNC1.  |
| AAA33127.1 | M91373   | Cucumis sativus         | peroxidase. pre-peroxidase. putative.   |
| CAA71492.1 | Y10466   | Spinacia oleracea       | peroxidase. prxr5.  |
| AAF63027.1 | AF244924 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.  |
| AAA33128.1 | M91374   | Cucumis sativus         | peroxidase. putative.   |
| BAA77389.1 | AB024439 | Scutellaria baicalensis | peroxidase 3.   |
| BAA77388.1 | AB024438 | Scutellaria baicalensis | peroxidase 2.   |

|               |          |                      |  |
|---------------|----------|----------------------|--|
| AAB02554.1    | L37790   | Stylosanthes humilis | cationic peroxidase.   |
| AAF63026.1    | AF244923 | Spinacia oleracea    | hydrogen peroxide catabolism. peroxidase prx14 precursor. type III peroxidase.   |
| SEQ ID NO: 58 |          |                      |  |
| AAG49002.1    | AY013246 | Hordeum vulgare      | putative ABC transporter. 635P2.4a; GC splice donor confirmed by cDNA and comparative sequencing.                            |
| AAG45492.1    | AY013245 | Oryza sativa         | 36I5.4. putative ABC transporter; GC splice donor confirmed by cDNA alignment and comparative sequence.                      |
| AAG49003.1    | AY013246 | Hordeum vulgare      | putative ABC transporter. 635P2.4b; GC splice donor confirmed by cDNA alignment and comparative sequence.                    |
| BAA83352.1    | AP000391 | Oryza sativa         | ESTs AU067992(C11433),AU077424(C11433) correspond to a region of the predicted gene.; Similar to ABC transporter-7 (U43892). |
| BAB17113.1    | AP002866 | Oryza sativa         | putative white protein; ATP-binding cassette transporter. P0410E01.34.   |
| BAA90508.1    | AP001111 | Oryza sativa         | similar to ABC transporter of Arabidopsis thaliana (AC004697).   |
| BAA90507.1    | AP001111 | Oryza sativa         | similar to ABC transporter of Arabidopsis thaliana (AC004697).   |
| BAB40032.1    | AP003046 | Oryza sativa         | putative ABC transporter. P0445D12.3.  |
| BAB16495.1    | AP002861 | Oryza sativa         | putative ABC transporter ATP-binding protein. P0665D10.21.   |
| BAA94511.1    | AB041505 | Populus nigra        | ABC transporter homolog. PnATH.  |
| BAB21275.1    | AP002844 | Oryza sativa         | putative ABC transporter protein. P0410E03.6.  |
| BAB21276.1    | AP002844 | Oryza sativa         | putative ABC transporter protein. P0410E03.7. contains EST D22472(C1173).  |
| BAB21279.1    | AP002844 | Oryza sativa         | putative ABC transporter protein. P0410E03.10. contains ESTs AU065360(R3463),AU101680(R3463).                                |
| CAA94437.1    | Z70524   | Spirodela polyrrhiza | multidrug resistance protein. PDR5-like ABC transporter.   |
| BAB21273.1    | AP002844 | Oryza sativa         | putative ABC transporter protein. P0410E03.4.  |
| SEQ ID NO: 59 |          |                      |  |
| CAA71369.1    | Y10338   | Solanum tuberosum    | chloride channel Stclcl. putative.   |

|               |          |                   |  |
|---------------|----------|-------------------|--|
| AAD29679.1    | AF133209 | Nicotiana tabacum | CLC-Nt2 protein. putative chloride channel.  |
| SEQ ID NO: 60 |          |                   |  |
| BAB40094.1    | AP003210 | Oryza sativa      | putative receptor protein kinase. OSJNBa0010K01.7.   |
| AAG52992.1    | U77888   | Ipomoea nil       | receptor-like protein kinase INRPK1a. inrpk1.  |
| AAF59906.1    | AF197947 | Glycine max       | receptor protein kinase-like protein. CLV1B.   |
| AAC36318.1    | AF053127 | Malus x domestica | leucine-rich receptor-like protein kinase. LRPKml.   |
| AAF91323.1    | AF244889 | Glycine max       | receptor-like protein kinase 2. RLK2. GmRLK2.  |
| AAF91324.1    | AF244890 | Glycine max       | receptor-like protein kinase 3. RLK3. GmRLK3.  |
| AAB36558.1    | U77888   | Ipomoea nil       | receptor-like protein kinase INRPK1. inrpk1. leucine-rich repeat.  |
| BAA83373.1    | AP000391 | Oryza sativa      | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| BAA84787.1    | AP000559 | Oryza sativa      | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| AAF91322.1    | AF244888 | Glycine max       | receptor-like protein kinase 1. RLK1. GmRLK1.  |
| AAF59905.1    | AF197946 | Glycine max       | receptor protein kinase-like protein. CLV1A.   |
| AAB61708.1    | U93048   | Daucus carota     | somatic embryogenesis receptor-like kinase. SERK.  |
| CAB51480.1    | Y14600   | Sorghum bicolor   | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.   |
| CAC20842.1    | AJ250467 | Pinus sylvestris  | receptor protein kinase. upk.  |
| CAA61510.1    | X89226   | Oryza sativa      | leucine-rich repeat/receptor protein kinase. lrk2.   |
| BAB19337.1    | AP003044 | Oryza sativa      | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).   |
| AAG52994.1    | U77888   | Ipomoea nil       | receptor-like protein kinase INRPK1c. inrpk1.  |
| BAB40081.1    | AP003074 | Oryza sativa      | putative receptor protein kinase. OSJNBa0004G10.30.  |

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAB18321.1 | AP002865 | Oryza sativa         | putative receptor protein kinase. P0034C11.11.  |
| AAD38286.1 | AC007789 | Oryza sativa         | putative protein kinase. OSJNBa0049B20.13.  |
| BAB07903.1 | AP002835 | Oryza sativa         | putative receptor kinase. P0417G05.10. contains ESTs AU032341(R3918),AU071016(R10613).  |
| AAF34426.1 | AF172282 | Oryza sativa         | leucine rich repeat containing protein kinase. DUPR11.16.   |
| BAA94519.1 | AP001800 | Oryza sativa         | ESTs AU032341(R3918),AU071016(R10613) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 4, BAC F9D16; putative receptor kinase (AL035394). |
| AAF66615.1 | AF142596 | Nicotiana tabacum    | LRR receptor-like protein kinase.   |
| BAB21240.1 | AP002953 | Oryza sativa         | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| BAA82393.1 | AP000367 | Oryza sativa         | EST C96716(C10608) corresponds to a region of the predicted gene.; Similar to putative receptor protein kinase. (AC002334).   |
| BAB03631.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.32.  |
| AAB82755.1 | U72725   | Oryza longistaminata | receptor kinase-like protein. Xa21 gene family member A1; downstream of microsatellite region; disease resistance gene family member.   |
| AAK00425.1 | AC069324 | Oryza sativa         | Putative protein kinase. OSJNBa0071K19.11.  |
| AAK27817.1 | AC022457 | Oryza sativa         | putative protein kinase. OSJNBa0006L06.16.  |
| BAA87853.1 | AP000816 | Oryza sativa         | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB03621.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.21.  |
| AAG59657.1 | AC084319 | Oryza sativa         | putative protein kinase. OSJNBa0004B24.20.  |
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| BAB03429.1 | AP002817 | Oryza sativa         | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).             |
| BAB07999.1 | AP002525 | Oryza sativa         | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |

|               |          |                   |  |
|---------------|----------|-------------------|--|
| AAK21965.1    | AY028699 | Brassica napus    | receptor protein kinase PERK1.   |
| AAG16628.1    | AY007545 | Brassica napus    | protein serine/threonine kinase BNK1.  |
| BAA94509.1    | AB041503 | Populus nigra     | protein kinase 1. PnPK1.   |
| BAB39421.1    | AP002901 | Oryza sativa      | putative receptor kinase. P0456F08.21.   |
| BAA94510.1    | AB041504 | Populus nigra     | protein kinase 2. PnPK2.   |
| CAB51836.1    | AJ243961 | Oryza sativa      | Putitive Ser/Thr protein kinase. I1332.7.  |
| BAB17139.1    | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.31.   |
| BAB07904.1    | AP002835 | Oryza sativa      | putative S-receptor kinase. P0417G05.12.   |
| BAB39409.1    | AP002901 | Oryza sativa      | putative protein kinase. P0456F08.9. contains EST C23560(R0290).                                     |
| SEQ ID NO: 62 |          |                   |  |
| BAA25753.1    | AB012932 | Vigna radiata     | Ca <sup>2+</sup> /H <sup>+</sup> exchanger. VCAX1.   |
| AAF91350.1    | AF256229 | Zea mays          | calcium/proton exchanger CAX1-like protein.  |
| BAA75232.1    | AB018526 | Ipomoea nil       | H <sup>+</sup> /Ca <sup>2+</sup> exchanger 2. CAX2.  |
| SEQ ID NO: 63 |          |                   |  |
| BAA85655.1    | AB026297 | Pisum sativum     | elicitor-responsive Dof protein ERDP.  |
| AAB70119.1    | U82230   | Zea mays          | transcription factor. prolamin box binding factor. PBF. endosperm-specific DOF protein; DNA binding. |
| CAB89831.1    | AJ242853 | Solanum tuberosum | DNA binding protein. Dof zinc finger protein. dof1.  |
| CAA08755.1    | AJ009594 | Nicotiana tabacum | Dof zinc finger protein. BBF1.   |
| CAA66601.1    | X97942   | Nicotiana tabacum | Zn finger protein. NtBBF1.1.   |
| CAA66604.1    | X97945   | Nicotiana tabacum | Zn finger protein. NtBBF2a.  |
| BAA78575.1    | AB028132 | Oryza sativa      | Dof zinc finger protein.   |
| BAA78572.1    | AB028129 | Oryza sativa      | Dof zinc finger protein.   |

|               |          |                    |  |
|---------------|----------|--------------------|--|
| BAA78573.1    | AB028130 | Oryza sativa       | Dof zinc finger protein.   |
| CAA66606.1    | X97947   | Nicotiana tabacum  | Zn finger protein. NtBBF3.   |
| CAA66605.1    | X97946   | Nicotiana tabacum  | Zn finger protein. NtBBF2b.  |
| BAA78574.1    | AB028131 | Oryza sativa       | Dof zinc finger protein. Rice cDNA for protein that binds to AT-rich sequence of rice carboxypeptidase-3 promoter. |
| CAA56287.1    | X79934   | Zea mays           | Dof2. No start codon.  |
| CAA56288.1    | X79935   | Zea mays           | Dof3 gene. no start codon.   |
| BAA78576.1    | AB028133 | Oryza sativa       | Dof zinc finger protein.   |
| SEQ ID NO: 64 |          |                    |  |
| CAA11219.1    | AJ223281 | Manihot esculenta  | alpha-hydroxynitrile lyase. HNL4.  |
| BAB19413.1    | AP002870 | Oryza sativa       | putative acetone-cyanohydrin lyase. P0458A05.22.   |
| AAC49184.1    | U40402   | Hevea brasiliensis | hydroxynitrile lyase. hnl.   |
| CAA82334.1    | Z29091   | Manihot esculenta  | alpha-hydroxynitrile lyase.  |
| CAA11428.1    | AJ223506 | Manihot esculenta  | alpha-hydroxynitrile lyase. HNL24.   |
| SEQ ID NO: 67 |          |                    |  |
| CAB89831.1    | AJ242853 | Solanum tuberosum  | DNA binding protein. Dof zinc finger protein. dof1.  |
| CAA56288.1    | X79935   | Zea mays           | Dof3 gene. no start codon.   |
| BAA85655.1    | AB026297 | Pisum sativum      | elicitor-responsive Dof protein ERDP.  |
| BAA78575.1    | AB028132 | Oryza sativa       | Dof zinc finger protein.   |
| AAB70119.1    | U82230   | Zea mays           | transcription factor. prolamin box binding factor. PBF. endosperm-specific DOF protein; DNA binding.               |
| CAA08755.1    | AJ009594 | Nicotiana tabacum  | Dof zinc finger protein. BBF1.   |
| CAA66601.1    | X97942   | Nicotiana tabacum  | Zn finger protein. NtBBF1.1.   |



---

CAA66606.1 X97947 *Nicotiana tabacum*

Zn finger protein. NtBBF3.

---

BAA78573.1 AB028130 *Oryza sativa*

Dof zinc finger protein.

---

BAA78572.1 AB028129 *Oryza sativa*

Dof zinc finger protein.

---

CAA56287.1 X79934 *Zea mays*

Dof2. No start codon.

---

CAA66604.1 X97945 *Nicotiana tabacum*

Zn finger protein. NtBBF2a.

---

CAA66605.1 X97946 *Nicotiana tabacum*

Zn finger protein. NtBBF2b.

---

BAA78574.1 AB028131 *Oryza sativa*

Dof zinc finger protein. Rice cDNA for protein that binds to AT-rich sequence of rice carboxypeptidase-3 promoter.

---

BAA78576.1 AB028133 *Oryza sativa*

Dof zinc finger protein.

---

SEQ ID NO: 68

---

CAA83453.1 Z31581 *Pisum sativum*

chloroplast outer envelope protein 86.

---

AAA53276.1 L36857 *Pisum sativum*

component of chloroplast outer membrane protein import apparatus. GTP-binding protein. IAP86.

---

AAF75761.1 AF262939 *Pisum sativum*

chloroplast protein import component Toc159. major receptor of the chloroplast outer envelope membrane protein import apparatus.

---

AAG48839.1 AC084218 *Oryza sativa*

similar to *Arabidopsis thaliana* putative chloroplast outer envelope 86-like protein (AC002330).

---

AAC25785.1 L36856 *Pisum sativum*

component of chloroplast outer membrane protein import apparatus. GTP-binding protein. IAP34.

---

CAB77551.1 AJ271049 *Zea mays*

a component of the protein translocon at the outer envelope of chloroplast. Toc34-2 protein. toc34B.

---

CAB65537.1 AJ245968 *Zea mays*

a component of the protein translocon at the outer envelope of chloroplast. Toc34-1 protein. toc34A.

---

SEQ ID NO: 70

---

AAB71887.1 AF020791 *Hordeum vulgare*

inserts Fe<sup>2+</sup> into protoporphyrin IX. ferrochelatase. hemH. protoheme IX ferro-lyase.

---

BAA05101.1 D26105 *Hordeum vulgare*

ferrochelatase. hemH.

---

|               |          |                                     |   |
|---------------|----------|-------------------------------------|---|
| BAA05102.1    | D26106   | Cucumis sativus                     | ferrochelata.se. hemH.  |
| BAB20760.1    | AB037113 | Cucumis sativus                     | ferrochelata.se. hemH.  |
| BAA22284.1    | AB007120 | Oryza sativa                        | ferrochelata.se. hemH.  |
| CAA06705.1    | AJ005802 | Solanum tuberosum                   | tetrapyrrole biosynthesis. ferrochelata.se.   |
| AAK16728.1    | AF332962 | Chlamydomonas reinhardtii           | ferrochelata.se.  |
| AAK16729.1    | AF332963 | Polytomella sp. 'Pringsheim 198.80' | ferrochelata.se.  |
| AAC84139.1    | AF101426 | Cichorium intybus                   | ferrochelata.se.  |
| SEQ ID NO: 71 |          |                                     |   |
| CAA64442.1    | X94986   | Manihot esculenta                   | beta glucosidase. bglA.   |
| AAB22162.1    | S35175   | Manihot esculenta                   | linamarase. linamarase. beta-glucosidase; Method: conceptual translation with partial peptide sequencing; This sequence comes from Fig. 4.  |
| AAA93032.1    | U50201   | Prunus serotina                     | hydrolysis of the cyanogenic glucoside (R)-prunasin. prunasin hydrolase precursor. located in protein bodies of Prunus seeds; encodes 8 putative N-glycosylation sites (N-X-S/T); encodes NEP and ITENG motifs characteristic of the BGA family of beta-glucosidases. |
| AAA91166.1    | U39228   | Prunus avium                        | beta-glucosidase.   |
| AAF34650.1    | AF221526 | Prunus serotina                     | hydrolysis of the cyanogenic glucoside (R)-prunasin. prunasin hydrolase isoform PHA precursor. beta-glucosidase; contains 6 potential N-glycosylation sites (N-X-S/T); glycosyl hydrolase family 1 member.  |
| AAF03675.1    | AF149311 | Rauvolfia serpentina                | hydrolyses the glucoalkaloid raucaffricine. raucaffricine-O-beta-D-glucosidase. beta glucosidase; RG; part of the Ajmaline biosynthesis pathway; belongs to family 1 of the glucosyl hydrolases.  |
| CAA57913.1    | X82577   | Brassica napus                      | beta-glucosidase. bgl.  |
| BAA78708.1    | AB003089 | Polygonum tinctorium                | beta-glucosidase.   |
| BAA11831.1    | D83177   | Costus speciosus                    | saponin metabolite. furostanol glycoside 26-O-beta-glucosidase (F26G). functional expression in E. coli; one of the F26G isozymes.  |
| AAG25897.1    | AF170087 | Cucurbita pepo                      | silverleaf whitefly-induced protein 3. SLW3. similar to beta-glucosidase.   |

|            |          |   |
|------------|----------|---|
| AAF04007.1 | AF163097 | <i>Dalbergia cochinchinensis</i><br>beta-fucosidase beta-glucosidase. dalcocinin 8'-O-beta-glucoside beta-glucosidase precursor.<br>BGLU1. rotenoid beta-glucosidase.   |
| AAC69619.1 | AF072736 | <i>Pinus contorta</i><br>beta-glucosidase.  |
| AAB38784.1 | U72154   | <i>Brassica nigra</i><br>beta-glucosidase. psr3.1. PSR3.1; phosphate-starvation responsive enzyme.  |
| AAD02839.1 | AF082991 | <i>Avena sativa</i><br>beta-D-glucosidase beta subunit precursor. P60b. avenacosidase.  |
| AAA87339.1 | L41869   | <i>Hordeum vulgare</i><br>beta-glucosidase. BGQ60. expression specific to starchy endosperm of seed.  |
| AAB71381.1 | U95298   | <i>Manihot esculenta</i><br>linamarase. pLIN-GEN. beta-glucosidase.   |
| CAA55196.1 | X78433   | <i>Avena sativa</i><br>beta-D-glucosidase.  |
| AAC49177.1 | U33817   | <i>Sorghum bicolor</i><br>beta-glucosidase, catalyzes the hydrolysis of the cyanogenic beta-glucoside dhurrin.<br>dhurrinase.   |
| AAD09850.1 | U44087   | <i>Zea mays</i><br>beta-D-glucosidase precursor. glu2. product subunit structure: autodimer of 58.4 kDa<br>monomers Allele: glu2-B73.   |
| AAG00614.1 | AF293849 | <i>Secale cereale</i><br>beta-glucosidase.  |
| AAF28800.1 | AF112888 | <i>Catharanthus roseus</i><br>plays a role in secondary metabolism by hydrolyzing strictosidine to cathenamine during<br>indole alkaloid biosynthesis. strictosidine beta-glucosidase. localized in the endoplasmic<br>reticulum. |
| CAA40058.1 | X56734   | <i>Trifolium repens</i><br>beta-glucosidase. non-cyanogenic.  |
| CAA40057.1 | X56733   | <i>Trifolium repens</i><br>beta-glucosidase. Li.  |
| AAD10503.1 | U33816   | <i>Zea mays</i><br>functions in defense of young plant parts against pests via the production of hydroxamic acids<br>from hydroxamic acid glucosides. beta-D-glucosidase.   |
| AAA65946.1 | U25157   | <i>Zea mays</i><br>functions in defense of young plant parts against pests via the production of hydroxamic acids<br>from hydroxamic acid glucosides. beta-D-glucosidase.   |
| CAA52293.1 | X74217   | <i>Zea mays</i><br>beta-glucosidase. p60.1.   |
| AAB03266.1 | U44773   | <i>Zea mays</i><br>beta-D-glucosidase. glu1. autodimer of 58.4 kD monomers.   |
| AAK07429.1 | AF321287 | <i>Musa acuminata</i><br>beta-glucosidase.  |

---

CAA79989.2 Z21977 *Brassica napus*  
beta thioglucosidase. myrosinase, thioglucoside glucohydrolase. Myr1.Bn1.

---

AAF34651.1 AF221527 *Prunus serotina*  
putative prunasin hydrolase precursor. beta-glucosidase; glycosyl hydrolase family 1 member.

---

AAA84906.1 U28047 *Oryza sativa*  
catalyzes the release of either giberellin or cyanogenic substances from their glucoconjugates.  
beta glucosidase. beta-D-glucoside glucohydrolase; dimer of 60 kDa monomers; localized in  
the plastid.

---

CAC08209.1 AJ005950 *Cicer arietinum*  
beta-glucosidase.

---

SEQ ID NO: 72

---

CAA56570.1 X80301 *Nicotiana tabacum*  
axi 1.

---

AAB72114.1 U81288 *Pisum sativum*  
PsRT17-1. similar to the *Nicotiana tabacum* axi 1 gene product encoded by the sequence  
presented in GenBank Accession Number X80301.

---

SEQ ID NO: 74

---

AAG23130.1 AF198260 *Lycopersicon esculentum*  
diacylglycerol kinase variant A. Dgk1. DGK-1; alternatively spliced.

---

AAG23129.1 AF198259 *Lycopersicon esculentum*  
phosphorylates diacylglycerol to yield phosphatidic acid. diacylglycerol kinase. DGK1.  
LeDGK1; lipid kinase; non-calmodulin-binding isoform; alternatively spliced product.

---

AAG23131.1 AF198260 *Lycopersicon esculentum*  
diacylglycerol kinase variant B. Dgk1. CBDGK; calmodulin-binding; alternatively spliced.

---

AAG23128.1 AF198258 *Lycopersicon esculentum*  
phosphorylates diacylglycerol to yield phosphatidic acid. calmodulin-binding diacylglycerol  
kinase. DGK1. LeCBDGK; lipid kinase; alternatively spliced product.

---

SEQ ID NO: 75

---

CAC09580.1 AJ298992 *Fagus sylvatica*  
Absciscic acid (ABA) and calcium induced protein kinase. protein kinase (PK). pk1.

---

AAA34002.1 M67449 *Glycine max*  
protein kinase. PK6.

---

AAG31141.1 AF305911 *Oryza sativa*  
EDR1. EDR1. MAP kinase kinase kinase; similar to *Arabidopsis thaliana* EDR1.

---

AAG31142.1 AF305912 *Hordeum vulgare*  
EDR1. EDR1. MAP kinase kinase kinase; similar to *Arabidopsis thaliana* EDR1.

---

CAA73722.1 Y13273 *Lycopersicon esculentum*  
putative protein kinase.

---

AAD46406.1 AF096250 *Lycopersicon esculentum*  
ethylene-responsive protein kinase TCTR1. ER50. serine/threonine kinase; similar to  
*Arabidopsis thaliana* negative regulator of the ethylene response pathway encoded by  
GenBank Accession Number L08789.

---

|            |          |  |
|------------|----------|--|
| AAD10056.1 | AF110518 | <i>Lycopersicon esculentum</i><br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1.  |
| AAD10057.1 | AF110519 | <i>Lycopersicon esculentum</i><br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1v. |
| CAA06334.1 | AJ005077 | <i>Lycopersicon esculentum</i><br>protein kinase. TCTR2 protein. TCTR2.  |
| AAK30005.1 | AY029067 | <i>Rosa hybrid cultivar</i><br>CTR2 protein kinase.  |
| AAK11734.1 | AY027437 | <i>Arachis hypogaea</i><br>serine/threonine/tyrosine kinase.   |
| CAA73067.1 | Y12464   | <i>Sorghum bicolor</i><br>serine/threonine kinase. SNFL1.  |
| CAA73068.1 | Y12465   | <i>Sorghum bicolor</i><br>serine/threonine kinase. SNFL2.  |
| AAK31267.1 | AC079890 | <i>Oryza sativa</i><br>putative protein kinase. OSJNBb0089A17.2.   |
| AAF66615.1 | AF142596 | <i>Nicotiana tabacum</i><br>LRR receptor-like protein kinase.  |
| BAB39451.1 | AP003338 | <i>Oryza sativa</i><br>putative receptor kinase. OJ1212_B09.24.  |
| BAB17126.1 | AP002867 | <i>Oryza sativa</i><br>putative receptor kinase. P0463F06.16.  |
| BAB17321.1 | AP002747 | <i>Oryza sativa</i><br>putative receptor kinase. P0698G03.1.   |
| AAF68398.1 | AF237568 | <i>Oryza sativa</i><br>receptor-like protein kinase. RLG2.   |
| BAB17129.1 | AP002867 | <i>Oryza sativa</i><br>putative receptor kinase. P0463F06.20.  |
| BAB17348.1 | AP002747 | <i>Oryza sativa</i><br>putative receptor kinase. P0698G03.32.  |
| CAA57898.1 | X82548   | <i>Hordeum vulgare</i><br>SNF1-related protein kinase. BKin2.  |
| BAB17345.1 | AP002747 | <i>Oryza sativa</i><br>putative receptor kinase. P0698G03.29.  |
| BAB17116.1 | AP002867 | <i>Oryza sativa</i><br>putative receptor kinase. P0463F06.3.   |
| BAB39441.1 | AP003338 | <i>Oryza sativa</i><br>putative receptor kinase. OJ1212_B09.11.  |
| CAC20842.1 | AJ250467 | <i>Pinus sylvestris</i><br>receptor protein kinase. upk.   |

---

AAD52097.1 AF088885 *Nicotiana tabacum*  
receptor-like kinase CHRK1. Chrkl.

---

BAB16918.1 AP002863 *Oryza sativa*  
putative protein kinase. P0005A05.22.

---

AAF91322.1 AF244888 *Glycine max*  
receptor-like protein kinase 1. RLK1. GmRLK1.

---

BAB17342.1 AP002747 *Oryza sativa*  
putative receptor kinase. P0698G03.26.

---

AAF78044.1 AF248493 *Oryza sativa*  
receptor-like kinase. RLG18. protein kinase.

---

AAF91324.1 AF244890 *Glycine max*  
receptor-like protein kinase 3. RLK3. GmRLK3.

---

AAF59905.1 AF197946 *Glycine max*  
receptor protein kinase-like protein. CLV1A.

---

AAF91323.1 AF244889 *Glycine max*  
receptor-like protein kinase 2. RLK2. GmRLK2.

---

AAD46917.1 AF164021 *Oryza sativa*  
receptor kinase.

---

AAF59906.1 AF197947 *Glycine max*  
receptor protein kinase-like protein. CLV1B.

---

AAF78018.1 AF238474 *Oryza sativa*  
receptor-like kinase. RLG16. protein kinase.

---

BAB39438.1 AP003338 *Oryza sativa*  
putative receptor kinase. OJ1212\_B09.7.

---

BAA05649.1 D26602 *Nicotiana tabacum*  
protein kinase.

---

AAC01746.1 AF044489 *Oryza sativa*  
receptor-like protein kinase. drpk1.

---

CAA71142.1 Y10036 *Cucumis sativus*  
SNF1-related protein kinase.

---

AAA62232.1 U00443 *Brassica napus*  
S-receptor kinase. protein contains an immunoglobulin-like domain.

---

CAA61510.1 X89226 *Oryza sativa*  
leucine-rich repeat/receptor protein kinase. lrk2.

---

SEQ ID NO: 77

---

AAD03693.1 AF084554 *Brassica napus*  
fibrillin.

---

CAA10372.1 AJ131455 *Plastid Solanum demissum*  
carotenoid-associated. fibrillin. c40.4.

---

CAA50750.1 X71952 *Capsicum annuum*  
fibrillin.

---

SEQ ID NO: 79

---

|            |          |                               |  |
|------------|----------|-------------------------------|--|
| AAB53155.1 | U43629   | Beta vulgaris                 | putative sugar transporter. integral membrane protein. member of major facilitator superfamily.              |
| AAF74566.1 | AF215852 | Nicotiana tabacum             | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                  |
| AAG00995.1 | AF286906 | Mesembryanthemum crystallinum | putative glucose translocator. metabolite transporter; targeted to plastid inner envelope membrane.          |
| AAF74565.1 | AF215851 | Spinacia oleracea             | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                  |
| AAF74567.1 | AF215853 | Solanum tuberosum             | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                  |
| AAF74568.1 | AF215854 | Zea mays                      | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                  |
| AAG46179.1 | AC018727 | Oryza sativa                  | putative sugar transporter protein. OSJNBa0056G17.3.   |
| AAB88879.1 | AF000952 | Prunus armeniaca              | putative sugar transporter.  |
| AAG43998.1 | AF215837 | Apium graveolens var. dulce   | mannitol transporter. Mat1.  |
| CAA68813.1 | Y07520   | Chlorella kessleri            | H(+)/hexose cotransporter (AA 1-533).  |
| CAA53192.1 | X75440   | Chlorella kessleri            | hexose transporter like protein. HUP3.   |
| CAB52689.1 | AJ132224 | Lycopersicon esculentum       | hexose transporter. ht2.   |
| CAA39036.1 | X55349   | Chlorella kessleri            | H(+)/hexose-cotransporter. HUP1.   |
| CAA47324.1 | X66856   | Nicotiana tabacum             | monosaccharid transporter. MST1.   |
| AAB68028.1 | U64902   | Beta vulgaris                 | BvcDNA-205. putative sugar transporter; member of major facilitative superfamily; integral membrane protein. |
| AAB68029.1 | U64903   | Beta vulgaris                 | BvcDNA-397. putative sugar transporter; member of major facilitative superfamily; integral membrane protein. |
| CAA09419.1 | AJ010942 | Lycopersicon esculentum       | hexose transporter protein.  |
| CAB07812.1 | Z93775   | Vicia faba                    | sugar transport. monosaccharid transport protein. hexT.  |
| BAB19864.1 | AB052885 | Oryza sativa                  | monosaccharide transporter 3. OsMST3.  |

|            |          |                                    |   |
|------------|----------|------------------------------------|---|
| AAA79761.1 | L08196   | Ricinus communis                   | hexose transport. sugar carrier protein. RCSTC.   |
| AAK13147.1 | AC083945 | Oryza sativa                       | Putative sugar transporter. OSJNBa0058E19.22.   |
| AAB06594.1 | U38651   | Medicago truncatula                | sugar transporter.  |
| CAA04511.1 | AJ001061 | Vitis vinifera                     | hexose uptake. hexose transporter.  |
| AAC61852.1 | AF061106 | Petunia x hybrida                  | putative monosaccharide transporter 1. pmt1. similar to hexose transporter protein; PMT1. |
| CAA70777.1 | Y09590   | Vitis vinifera                     | hexose transporter.   |
| AAA79769.1 | L08197   | Ricinus communis                   | sugar transport. sugar carrier protein. RCSTA. putative.                                  |
| CAB06079.1 | Z83829   | Picea abies                        | monosaccharide transporter. PaMst-1. PaMst-1.   |
| AAA18534.1 | L21753   | Saccharum hybrid cultivar H65-7052 | glucose transporter. putative.  |
| CAC00697.1 | AJ278765 | Lycopersicon esculentum            | putative sugar transporter. st3.  |
| AAA79857.1 | L08188   | Ricinus communis                   | hexose transport. hexose carrier protein. HEX6.   |
| BAB19863.1 | AB052884 | Oryza sativa                       | monosaccharide transporter 2. OsMST2.   |
| BAB19862.1 | AB052883 | Oryza sativa                       | monosaccharide transporter 1. OsMST1.   |
| AAF91432.1 | AF280432 | Mesembryanthemum crystallinum      | putative Na <sup>+</sup> /myo-inositol symporter. Itr2. membrane transport protein.       |
| CAB52688.1 | AJ132223 | Lycopersicon esculentum            | hexose transporter. ht1.  |
| BAA85398.1 | AP000615 | Oryza sativa                       | similar to sugar transporter protein. (AL022604).   |
| AAG46115.1 | AC073166 | Oryza sativa                       | putative sugar transporter. OSJNBb0064P21.3.  |
| CAB52690.1 | AJ132225 | Lycopersicon esculentum            | hexose transporter. ht3.  |
| AAA18533.1 | L21752   | Saccharum hybrid cultivar H65-7052 | glucose transporter. putative.  |
| AAD55054.1 | AF173655 | Beta vulgaris                      | glucose transporter. Gt.  |
| BAB39246.1 | AP002869 | Oryza sativa                       | putative transport protein homolog. P0554D10.30.  |



|               |          |                                |  |
|---------------|----------|--------------------------------|--|
| AAA79764.1    | L08191   | <i>Ricinus communis</i>        | sugar transport. sugar carrier protein. RCSTC. Nucleotides 1 to 23 and 226 to 246 are originated from degenerated primers; putative.   |
| AAA79762.1    | L08189   | <i>Ricinus communis</i>        | sugar transport. sugar carrier protein. RCSTA. Nucleotides 1 to 23 and 226 to 246 are originated from degenerated primers; putative.   |
| AAA79767.1    | L08194   | <i>Ricinus communis</i>        | sugar transport. sugar carrier protein. RCSTG. Nucleotides 1 to 23 and 226 to 246 are originated from degenerated primers; putative.   |
| AAA79766.1    | L08193   | <i>Ricinus communis</i>        | sugar transport. sugar carrier protein. RCSTF. Nucleotides 1 to 23 and 226 to 246 are originated from degenerated primers; putative.   |
| SEQ ID NO: 82 |          |                                |  |
| CAA78386.1    | Z13996   | <i>Petunia x hybrida</i>       | DNA binding protein; transcriptional activator. protein 1. myb.Ph3. Product related to animal myb proto-oncoproteins. Sequence from nucleotide 992 is not included in clone cPF1 and has been obtained by PCR amplification of cDNA. |
| CAB43399.1    | AJ006292 | <i>Antirrhinum majus</i>       | Myb-related transcription factor mixta-like 1. mybml1.   |
| CAA67600.1    | X99210   | <i>Lycopersicon esculentum</i> | myb-related transcription factor. THM16.   |
| BAA93038.1    | AP001552 | <i>Oryza sativa</i>            | EST AU082058(C12976) corresponds to a region of the predicted gene. Similar to <i>Arabidopsis thaliana</i> putative transcription factor (AF062916).   |
| BAA23337.1    | D88617   | <i>Oryza sativa</i>            | transfactor. OSMYB1. Osmyb1.   |
| CAA64614.1    | X95296   | <i>Lycopersicon esculentum</i> | transcription factor. THM27. myb-related.  |
| CAA72186.1    | Y11351   | <i>Oryza sativa</i>            | myb factor. myb.   |
| AAF22256.1    | AF161711 | <i>Pimpinella brachycarpa</i>  | myb-related transcription factor.  |
| AAC04720.1    | AF034134 | <i>Gossypium hirsutum</i>      | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-O. similar to MYB A encoded by GenBank Accession Number L04497.   |
| AAA82943.1    | U39448   | <i>Picea mariana</i>           | MYB-like transcriptional factor MBF1. putative DNA binding region highly similar to the maize C1.  |
| BAB39987.1    | AP003020 | <i>Oryza sativa</i>            | putative transcription factor (myb). P0498A12.16. contains ESTs AU097474(S5087),D40175(S1959).   |
| BAB39972.1    | AP003018 | <i>Oryza sativa</i>            | putative transcription factor (myb). OSJNBa0004B13.27. contains ESTs AU097474(S5087),D40175(S1959).  |

|               |          |                         |   |
|---------------|----------|-------------------------|---|
| BAA23338.1    | D88618   | Oryza sativa            | transfactor. OSMYB2. Osmbyb2.   |
| CAA67575.1    | X99134   | Lycopersicon esculentum | transcription factor. THM6. myb-related.  |
| CAA78387.1    | Z13997   | Petunia x hybrida       | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.                   |
| AAA19821.1    | L19495   | Zea mays                | transcriptional activator for anthocyanin synthesis. transcriptional activator.   |
| AAC49394.1    | U57002   | Zea mays                | P protein. P. allele P-wr; Myb-like transcriptional regulator with a putative zinc-finger at the C-terminal.                    |
| AAA33500.1    | M73028   | Zea mays                | myb-like transcription factor. P.   |
| AAG36774.1    | AF210616 | Zea mays                | P2 protein. P2. myb-like transcriptional factor; similar to Zea mays P gene.  |
| BAA88222.1    | AB028650 | Nicotiana tabacum       | myb-related transcription factor LBM2. lbm2.  |
| CAA72185.1    | Y11350   | Oryza sativa            | myb factor. myb.  |
| AAG13574.1    | AC037425 | Oryza sativa            | myb factor. OSJNBa0055P24.4.  |
| AAB41101.1    | U72762   | Nicotiana tabacum       | transcription factor Myb1. myb1. TMV-inducible Myb homolog; contains helix-turn-helix motif; contains redox-sensitive cysteine. |
| BAA88223.1    | AB028651 | Nicotiana tabacum       | myb-related transcription factor LBM3. lbm3.  |
| AAA19819.1    | L19496   | Zea mays                | transcriptional activator for anthocyanin synthesis. transcriptional activator.   |
| BAA88224.1    | AB028652 | Nicotiana tabacum       | myb-related transcription factor LBM4. lbm4.  |
| AAB67720.1    | AF015268 | Zea mays                | activator of anthocyanin structural genes. PL transcription factor. Pl.   |
| AAA33492.1    | L13454   | Zea mays                | transcriptional activator for anthocyanin biosynthesis. Pl-Bh (Blotched1).  |
| SEQ ID NO: 83 |          |                         |   |
| CAA78387.1    | Z13997   | Petunia x hybrida       | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.                   |
| BAA81736.1    | AB029165 | Glycine max             | GmMYB29B2.  |
| BAA81732.1    | AB029161 | Glycine max             | GmMYB29A2.  |

|            |          |   |
|------------|----------|---|
| BAA81731.1 | AB029160 | Glycine max<br>GmMYB29A1.   |
| BAA81730.1 | AB029159 | Glycine max<br>GmMYB29A1.   |
| BAA88221.1 | AB028649 | Nicotiana tabacum<br>myb-related transcription factor LBM1. lbm1.   |
| BAA88224.1 | AB028652 | Nicotiana tabacum<br>myb-related transcription factor LBM4. lbm4.   |
| CAA66952.1 | X98308   | Lycopersicon esculentum<br>THM18. myb-related transcription factor.   |
| BAA81733.2 | AB029162 | Glycine max<br>GmMYB29A2.   |
| CAA72217.1 | Y11414   | Oryza sativa<br>myb.  |
| BAA88222.1 | AB028650 | Nicotiana tabacum<br>myb-related transcription factor LBM2. lbm2.   |
| AAB41101.1 | U72762   | Nicotiana tabacum<br>transcription factor Myb1. myb1. TMV-inducible Myb homolog; contains helix-turn-helix motif; contains redox-sensitive cysteine.  |
| BAA88223.1 | AB028651 | Nicotiana tabacum<br>myb-related transcription factor LBM3. lbm3.   |
| CAA72185.1 | Y11350   | Oryza sativa<br>myb factor. myb.  |
| AAG13574.1 | AC037425 | Oryza sativa<br>myb factor. OSJNBa0055P24.4.  |
| AAK19616.1 | AF336283 | Gossypium hirsutum<br>GHMYB25. ghmyb25. similar to myb; contains an unspliced intron.   |
| CAA78386.1 | Z13996   | Petunia x hybrida<br>DNA binding protein; transcriptional activator. protein 1. myb.Ph3. Product related to animal myb proto-oncoproteins. Sequence from nucleotide 992 is not included in clone cPF1 and has been obtained by PCR amplification of cDNA. |
| CAA72218.1 | Y11415   | Oryza sativa<br>myb.  |
| AAA33067.1 | L04497   | Gossypium hirsutum<br>MYB A; putative.  |
| CAA68235.1 | X99973   | Hordeum vulgare<br>myb4 transcription factor. myb4.   |
| AAC49394.1 | U57002   | Zea mays<br>P protein. P. allele P-wr; Myb-like transcriptional regulator with a putative zinc-finger at the C-terminal.  |
| AAA33500.1 | M73028   | Zea mays<br>myb-like transcription factor. P.   |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| CAA67600.1 | X99210   | Lycopersicon esculentum | myb-related transcription factor. THM16.   |
| AAG36774.1 | AF210616 | Zea mays                | P2 protein. P2. myb-like transcriptional factor; similar to Zea mays P gene.   |
| AAF22256.1 | AF161711 | Pimpinella brachycarpa  | myb-related transcription factor.  |
| CAA72187.1 | Y11352   | Oryza sativa            | myb factor. myb.   |
| AAC04718.1 | AF034132 | Gossypium hirsutum      | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-J. similar to MYB A encoded by GenBank Accession Number L04497. |
| CAB43399.1 | AJ006292 | Antirrhinum majus       | Myb-related transcription factor mixta-like 1. mybml1.   |
| AAK19615.1 | AF336282 | Gossypium hirsutum      | GHMYB10. ghmyb10. similar to myb.  |
| AAK19618.1 | AF336285 | Gossypium hirsutum      | GHMYB38. ghmyb38. similar to myb.  |
| CAA72186.1 | Y11351   | Oryza sativa            | myb factor. myb.   |
| CAA64614.1 | X95296   | Lycopersicon esculentum | transcription factor. THM27. myb-related.  |
| CAA50221.1 | X70876   | Hordeum vulgare         | MybHv5. myb2.  |
| BAA23338.1 | D88618   | Oryza sativa            | transfactor. OSMYB2. Osmyb2.   |
| AAK19619.1 | AF336286 | Gossypium hirsutum      | GHMYB9. ghmyb9. similar to myb.  |
| AAA82943.1 | U39448   | Picea mariana           | MYB-like transcriptional factor MBF1. putative DNA binding region highly similar to the maize C1.  |
| AAC04720.1 | AF034134 | Gossypium hirsutum      | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-O. similar to MYB A encoded by GenBank Accession Number L04497. |
| AAK19611.1 | AF336278 | Gossypium hirsutum      | BNLGH1233. bnlghi6233. similar to myb.   |
| BAA23337.1 | D88617   | Oryza sativa            | transfactor. OSMYB1. Osmyb1.   |
| BAB39987.1 | AP003020 | Oryza sativa            | putative transcription factor (myb). P0498A12.16. contains ESTs AU097474(S5087),D40175(S1959).   |
| BAB39972.1 | AP003018 | Oryza sativa            | putative transcription factor (myb). OSJNBa0004B13.27. contains ESTs AU097474(S5087),D40175(S1959).  |

|               |          |                         |   |
|---------------|----------|-------------------------|---|
| CAA65525.1    | X96749   | Oryza sativa            | myb7.   |
| AAK19617.1    | AF336284 | Gossypium hirsutum      | GHMYB36. ghmyb36. similar to myb.   |
| CAA67575.1    | X99134   | Lycopersicon esculentum | transcription factor. THM6. myb-related.  |
| CAA50224.1    | X70879   | Hordeum vulgare         | MybHv1. myb1.   |
| SEQ ID NO: 84 |          |                         |   |
| BAA88222.1    | AB028650 | Nicotiana tabacum       | myb-related transcription factor LBM2. lbm2.  |
| BAA88221.1    | AB028649 | Nicotiana tabacum       | myb-related transcription factor LBM1. lbm1.  |
| BAA88224.1    | AB028652 | Nicotiana tabacum       | myb-related transcription factor LBM4. lbm4.  |
| CAA78387.1    | Z13997   | Petunia x hybrida       | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.                   |
| CAA66952.1    | X98308   | Lycopersicon esculentum | THM18. myb-related transcription factor.  |
| AAB41101.1    | U72762   | Nicotiana tabacum       | transcription factor Myb1. myb1. TMV-inducible Myb homolog; contains helix-turn-helix motif; contains redox-sensitive cysteine. |
| BAA88223.1    | AB028651 | Nicotiana tabacum       | myb-related transcription factor LBM3. lbm3.  |
| BAA81733.2    | AB029162 | Glycine max             | GmMYB29A2.  |
| BAA81731.1    | AB029160 | Glycine max             | GmMYB29A1.  |
| BAA81730.1    | AB029159 | Glycine max             | GmMYB29A1.  |
| BAA81736.1    | AB029165 | Glycine max             | GmMYB29B2.  |
| CAA72217.1    | Y11414   | Oryza sativa            | myb.  |
| BAA81732.1    | AB029161 | Glycine max             | GmMYB29A2.  |
| CAA72185.1    | Y11350   | Oryza sativa            | myb factor. myb.  |
| AAG13574.1    | AC037425 | Oryza sativa            | myb factor. OSJNBa0055P24.4.  |
| CAA72218.1    | Y11415   | Oryza sativa            | myb.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| CAA78386.1 | Z13996   | Petunia x hybrida       | DNA binding protein; transcriptional activator. protein 1. myb.Ph3. Product related to animal myb proto-oncoproteins. Sequence from nucleotide 992 is not included in clone cPF1 and has been obtained by PCR amplification of cDNA. |
| AAC49394.1 | U57002   | Zea mays                | P protein. P. allele P-wr; Myb-like transcriptional regulator with a putative zinc-finger at the C-terminal.   |
| CAB43399.1 | AJ006292 | Antirrhinum majus       | Myb-related transcription factor mixta-like 1. mybml1.   |
| AAK19616.1 | AF336283 | Gossypium hirsutum      | GHMYB25. ghmyb25. similar to myb; contains an unspliced intron.  |
| AAG36774.1 | AF210616 | Zea mays                | P2 protein. P2. myb-like transcriptional factor; similar to Zea mays P gene.   |
| AAA33500.1 | M73028   | Zea mays                | myb-like transcription factor. P.  |
| CAA68235.1 | X99973   | Hordeum vulgare         | myb4 transcription factor. myb4.   |
| CAA72187.1 | Y11352   | Oryza sativa            | myb factor. myb.   |
| AAA33067.1 | L04497   | Gossypium hirsutum      | MYB A; putative.   |
| CAA72186.1 | Y11351   | Oryza sativa            | myb factor. myb.   |
| CAA67600.1 | X99210   | Lycopersicon esculentum | myb-related transcription factor. THM16.   |
| AAK19618.1 | AF336285 | Gossypium hirsutum      | GHMYB38. ghmyb38. similar to myb.  |
| AAK19611.1 | AF336278 | Gossypium hirsutum      | BNLGH1233. bnlghi6233. similar to myb.   |
| AAC04718.1 | AF034132 | Gossypium hirsutum      | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. Cmy-J. similar to MYB A encoded by GenBank Accession Number L04497.   |
| AAK19619.1 | AF336286 | Gossypium hirsutum      | GHMYB9. ghmyb9. similar to myb.  |
| AAK19615.1 | AF336282 | Gossypium hirsutum      | GHMYB10. ghmyb10. similar to myb.  |
| CAA64614.1 | X95296   | Lycopersicon esculentum | transcription factor. THM27. myb-related.  |
| BAA23338.1 | D88618   | Oryza sativa            | transfactor. OSMYB2. Osmyb2.   |
| BAA23337.1 | D88617   | Oryza sativa            | transfactor. OSMYB1. Osmyb1.   |

|               |          |                         |   |
|---------------|----------|-------------------------|---|
| CAA67575.1    | X99134   | Lycopersicon esculentum | transcription factor. THM6. myb-related.  |
| CAA65525.1    | X96749   | Oryza sativa            | myb7.   |
| BAB39987.1    | AP003020 | Oryza sativa            | putative transcription factor (myb). P0498A12.16. contains ESTs AU097474(S5087),D40175(S1959).      |
| BAB39972.1    | AP003018 | Oryza sativa            | putative transcription factor (myb). OSJNBa0004B13.27. contains ESTs AU097474(S5087),D40175(S1959). |
| CAA50221.1    | X70876   | Hordeum vulgare         | MybHv5. myb2.   |
| AAA82943.1    | U39448   | Picea mariana           | MYB-like transcriptional factor MBF1. putative DNA binding region highly similar to the maize C1.   |
| AAF22256.1    | AF161711 | Pimpinella brachycarpa  | myb-related transcription factor.   |
| AAK19617.1    | AF336284 | Gossypium hirsutum      | GHMYB36. ghmyb36. similar to myb.   |
| CAA50222.1    | X70877   | Hordeum vulgare         | MybHv1. myb1.   |
| CAA50224.1    | X70879   | Hordeum vulgare         | MybHv1. myb1.   |
| SEQ ID NO: 85 |          |                         |   |
| AAD11575.1    | AF064029 | Helianthus tuberosus    | lectin 1. LECHeltubal; agglutinin.  |
| AAD11578.1    | AF064030 | Helianthus tuberosus    | lectin 2. LECHeltuba2; agglutinin.  |
| AAD11577.1    | AF064032 | Helianthus tuberosus    | lectin HE17.  |
| AAG10403.1    | AF233284 | Convolvulus arvensis    | mannose-binding lectin. cr8. Conarva.   |
| AAD11576.1    | AF064031 | Helianthus tuberosus    | lectin 3.   |
| AAB82776.2    | AF001527 | Musa acuminata          | ripening-associated protein. similar to lectin.   |
| AAC49564.1    | U56820   | Calystegia sepium       | lectin.   |
| CAB40792.1    | AJ237754 | Hordeum vulgare         | putative lectin. hl#2.  |
| SEQ ID NO: 88 |          |                         |   |
| AAB65163.1    | AF002692 | Solanum commersonii     | glutathione S-transferase, class-phi. GST1. low temperature induced.                                |

|            |          |   |
|------------|----------|---|
| CAA55039.1 | X78203   | Hyoscyamus muticus<br>glutathione transferase.                              |
| AAA33930.1 | M84968   | Silene vulgaris<br>glutathione-S-transferase.                               |
| AAA33931.1 | M84969   | Silene vulgaris<br>glutathione-S-transferase.                               |
| AAF65767.1 | AF242309 | Euphorbia esula<br>glutathione S-transferase. putative auxin-binding GST.   |
| BAA01394.1 | D10524   | Nicotiana tabacum<br>glutathione S-transferase. parB.                       |
| CAA96431.1 | Z71749   | Nicotiana plumbaginifolia<br>glutathione S-transferase.                     |
| AAF61392.1 | AF133894 | Persea americana<br>glutathione S-transferase. GTH.                         |
| CAB38119.1 | AJ010296 | Zea mays<br>Glutathione transferase III(b). gst3b.                          |
| CAB38118.1 | AJ010295 | Zea mays<br>Glutathione transferase III(a). gst3a.                          |
| BAB39935.1 | AP002914 | Oryza sativa<br>putative glutathione S-transferase. P0493G01.17.            |
| CAA09190.1 | AJ010451 | Alopecurus myosuroides<br>glutathione transferase. GST2a.                   |
| CAA09192.1 | AJ010453 | Alopecurus myosuroides<br>glutathione transferase. GST2c.                   |
| CAA09193.1 | AJ010454 | Alopecurus myosuroides<br>glutathione transferase. GST2d.                   |
| CAA09191.1 | AJ010452 | Alopecurus myosuroides<br>glutathione transferase. GST2b.                   |
| BAB39941.1 | AP002914 | Oryza sativa<br>putative glutathione S-transferase. P0493G01.23.            |
| AAG34811.1 | AF243376 | Glycine max<br>glutathione S-transferase GST 21.                            |
| BAB39939.1 | AP002914 | Oryza sativa<br>putative glutathione S-transferase. P0493G01.21.            |
| AAG32476.1 | AF309383 | Oryza sativa subsp. japonica<br>putative glutathione S-transferase OsGSTF4. |
| AAG34814.1 | AF243379 | Glycine max<br>glutathione S-transferase GST 24.                            |
| AAG34812.1 | AF243377 | Glycine max<br>glutathione S-transferase GST 22.                            |
| BAB39929.1 | AP002914 | Oryza sativa<br>putative glutathione transferase. P0493G01.7.               |



|            |          |                              |   |
|------------|----------|------------------------------|---|
| CAA39487.1 | X56012   | Triticum aestivum            | glutathione transferase. gstA1.   |
| AAD56395.1 | AF184059 | Triticum aestivum            | glutathione S-transferase. GST1.  |
| BAB39927.1 | AP002914 | Oryza sativa                 | putative glutathione S-transferase. P0493G01.1. contains ESTs AU031696(R0596), C97559(C60386), C28218(C60386), D28287(R0596). |
| AAA20585.1 | U12679   | Zea mays                     | glutathione S-transferase IV. GSTIV.  |
| CAA56047.1 | X79515   | Zea mays                     | glutathione transferase. GST27.   |
| CAA39480.1 | X56004   | Triticum aestivum            | glutathione transferase. gstA2.   |
| AAA33469.1 | M16902   | Zea mays                     | glutathione S-transferase I.  |
| AAG32475.1 | AF309382 | Oryza sativa subsp. japonica | putative glutathione S-transferase OsGSTF5.   |
| AAA33470.1 | M16901   | Zea mays                     | glutathione S-transferase I.  |
| AAG32477.1 | AF309384 | Oryza sativa subsp. japonica | putative glutathione S-transferase OsGSTF3.   |
| CAA68993.1 | Y07721   | Petunia x hybrida            | conjugates glutathione to anthocyanin to facilitate transport to the vacuole. glutathione S-transferase. an9 locus.           |
| BAB39940.1 | AP002914 | Oryza sativa                 | putative glutathione S-transferase. P0493G01.22.  |
| AAC64007.1 | AF062403 | Oryza sativa                 | glutathione S-transferase II.   |
| AAG34823.1 | AF244680 | Zea mays                     | glutathione S-transferase GST 15.   |
| AAG34817.1 | AF244674 | Zea mays                     | glutathione S-transferase GST 9.  |
| CAB66333.1 | AJ279691 | Betula pendula               | glutathione-S-transferase. gst.   |
| AAG34820.1 | AF244677 | Zea mays                     | glutathione S-transferase GST 11.   |
| AAG34821.1 | AF244678 | Zea mays                     | glutathione S-transferase GST 13.   |
| CAA05354.1 | AJ002380 | Oryza sativa                 | glutathione S-transferase. Rgst I.  |
| AAG34816.1 | AF244673 | Zea mays                     | glutathione S-transferase GST 8.  |

|               |          |                      |  |
|---------------|----------|----------------------|--|
| AAG34818.1    | AF244675 | Zea mays             | glutathione S-transferase GST 10.  |
| CAA05355.1    | AJ002381 | Oryza sativa         | glutathione S-transferase. Rgst II.  |
| AAG34824.1    | AF244681 | Zea mays             | glutathione S-transferase GST 16.  |
| SEQ ID NO: 91 |          |                      |  |
| AAG34695.1    | AF313492 | Matthiola incana     | putative cytochrome P450.  |
| CAA71516.1    | Y10492   | Glycine max          | putative cytochrome P450.  |
| BAA12159.1    | D83968   | Glycine max          | Cytochrome P-450 (CYP93A1).  |
| AAC32274.1    | AF081575 | Petunia x hybrida    | flavonoid 3',5'-hydroxylase. Hfl. P450 enzyme.   |
| AAA32913.1    | M32885   | Persea americana     | cytochrome P-450LXXIA1 (cyp71A1).  |
| BAA13076.1    | D86351   | Glycine max          | cytochrome P-450 (CYP93A2).  |
| CAA64635.1    | X95342   | Nicotiana tabacum    | cytochrome P450. hsr515. hypersensitivity-related gene.  |
| CAA65580.1    | X96784   | Nicotiana tabacum    | cytochrome P450. hsr515.   |
| AAD56282.1    | AF155332 | Petunia x hybrida    | flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.   |
| BAB12433.1    | AB025030 | Coptis japonica      | p450.  |
| AAG44132.1    | AF218296 | Pisum sativum        | cytochrome P450. P450 isolog.  |
| CAA50155.1    | X70824   | Solanum melongena    | flavonoid hydroxylase (P450). CYP75.   |
| AAB17562.1    | U72654   | Eustoma grandiflorum | flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.  |
| AAF05621.1    | AF191772 | Papaver somniferum   | hydroxylase involved in benzyloquinoline alkaloid biosynthesis. (S)-N-methylcoclaurine 3'-hydroxylase. CYP80B1. cytochrome P450-dependent monooxygenase. |
| CAB56503.1    | AJ238612 | Catharanthus roseus  | cytochrome P450.   |
| CAA70575.1    | Y09423   | Nepeta racemosa      | cytochrome P450. CYP71A5.  |
| BAA74466.1    | AB022733 | Glycyrrhiza echinata | cytochrome P450. CYP Ge-51.  |

|               |          |   |
|---------------|----------|---|
| CAA50648.1    | X71657   | <i>Solanum melongena</i><br>P450 hydroxylase.   |
| BAA22423.1    | AB001380 | <i>Glycyrrhiza echinata</i><br>cytochrome P450. CYP93B1.  |
| BAB40324.1    | AB037245 | <i>Asparagus officinalis</i><br>cytochrome P450. ASPI-2.  |
| AAB94588.1    | AF022459 | <i>Glycine max</i><br>CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.   |
| AAG14961.1    | AF214007 | <i>Brassica napus</i><br>cytochrome p450-dependent monooxygenase. BNF5H1.   |
| BAB40323.1    | AB037244 | <i>Asparagus officinalis</i><br>cytochrome P450. ASPI-1.  |
| AAB61965.1    | U48435   | <i>Solanum chacoense</i><br>putative cytochrome P450.   |
| AAC39452.1    | AF014800 | <i>Eschscholzia californica</i><br>hydroxylase involved in the biosynthesis of tetrahydrobenzylisoquinoline alkaloids in plants.<br>(S)-N-methylcoclaurine 3'-hydroxylase. CYP80B1. cytochrome P-450-dependent<br>monooxygenase; CYP80B1v1. |
| AAG14962.1    | AF214008 | <i>Brassica napus</i><br>cytochrome p450-dependent monooxygenase. BNF5H2.   |
| BAA84072.1    | AB028152 | <i>Torenia hybrida</i><br>flavone synthase II. cytochrome P450. TFNS5.  |
| AAC39453.1    | AF014801 | <i>Eschscholzia californica</i><br>hydroxylase involved in the biosynthesis of tetrahydrobenzylisoquinoline alkaloids in plants.<br>(S)-N-methylcoclaurine 3'-hydroxylase. CYP80B1. cytochrome P-450-dependent<br>monooxygenase; CYP80B1v2. |
| AAD47832.1    | AF166332 | <i>Nicotiana tabacum</i><br>cytochrome P450.  |
| SEQ ID NO: 92 |          |   |
| AAG42490.1    | AF321001 | <i>Suaeda maritima</i> subsp. <i>salsa</i><br>S-adenosylmethionine synthetase 2.  |
| AAG17666.1    | AF271220 | <i>Brassica juncea</i><br>S-adenosylmethionine synthetase. MSAMS2.  |
| BAA96637.1    | AP002482 | <i>Oryza sativa</i><br>Similar to <i>Oryza sativa</i> S-adenosylmethionine synthetase 1 (P46611).   |
| AAG17036.1    | AF187821 | <i>Pinus contorta</i><br>catalyzes the reaction between methionine and ATP to S-adenosylmethionine. S-<br>adenosylmethionine synthetase. sams2.   |
| BAA94605.1    | AB041534 | <i>Camellia sinensis</i><br>s-adenosylmethionine synthetase. SAM.   |
| AAA81377.1    | U17239   | <i>Actinidia chinensis</i><br>S-adenosylmethionine synthetase.  |

|               |          |                               |   |
|---------------|----------|-------------------------------|---|
| AAB38500.1    | U79767   | Mesembryanthemum crystallinum | S-adenosylmethionine synthetase. methionine adenosyltransferase.  |
| AAA81378.1    | U17240   | Actinidia chinensis           | S-adenosylmethionine synthetase.  |
| BAA09895.1    | D63835   | Hordeum vulgare               | S-adenosylmethionine synthetase.  |
| AAA33274.1    | M61882   | Dianthus caryophyllus         | S-adenosylmethionine synthetase. CARSAM2.   |
| AAA58773.1    | L36681   | Pisum sativum                 | S-adenosylmethionine synthase. precursor for ethylene and polyamine biosynthesis.   |
| AAA58772.1    | L36680   | Pisum sativum                 | precursor for ethylene and polyamine biosynthesis. S-adenosylmethionine synthase.   |
| AAA81379.1    | U17241   | Actinidia chinensis           | S-adenosylmethionine synthetase.  |
| AAA33857.1    | M62758   | Petroselinum crispum          | S-adenosylmethionine synthetase. SMS-1.   |
| AAG17035.1    | AF187820 | Pinus contorta                | catalyzes the reaction between methionine and ATP to S-adenosylmethionine. S-adenosylmethionine synthetase. sams1.  |
| AAB71833.1    | AF008568 | Chlamydomonas reinhardtii     | S-adenosylmethionine synthetase. CHRSAMS.   |
| AAA33858.1    | M62757   | Petroselinum crispum          | S-adenosylmethionine synthetase. SMS-2.   |
| BAA21726.1    | AB006187 | Nicotiana tabacum             | S-adenosylmethionine synthase. BYJ90.   |
| AAF78525.1    | AF195233 | Pyrus pyrifolia               | S-adenosylmethionine synthase. SAMS.  |
| SEQ ID NO: 94 |          |                               |   |
| AAB36543.1    | U77935   | Phaseolus vulgaris            | DnaJ-like protein. synthesis and expression are regulated by heavy metal stress, virus infection and wounding treatment, suggesting that DnaJ-like protein plays a role in plant defense. |
| SEQ ID NO: 95 |          |                               |   |
| AAB36546.1    | U77940   | Phaseolus vulgaris            | polyubiquitin. expression is regulated by heavy metal stress, UV, virus infection, heat shock and wounding treatment, this suggests that ubiquitin plays a role in plant defense.         |
| SEQ ID NO: 98 |          |                               |   |
| AAF22109.1    | AF119411 | Lupinus albus                 | ethylene biosynthesis enzyme. 1-aminocyclopropane-1-carboxylate synthase 1. ACS1. ACC synthase; S-adenosyl-L-methionine methylthioadenosine-lyase.  |
| AAC83146.1    | AF057562 | Nicotiana glutinosa           | 1-aminocyclopropane-1-carboxylate synthase. ACS1. ACC synthase.   |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAB06724.1 | U64804   | Petunia x hybrida       | ethylene biosynthesis enzyme. 1-aminocyclopropane-1-carboxylate synthase. PH-ACS1.  |
| CAA79478.1 | Z18953   | Petunia x hybrida       | 1-aminocyclopropane-1-carboxylate synthase.   |
| AAB17279.1 | U72390   | Lycopersicon esculentum | converts S-adenosylmethionine to 1-aminocyclopropane-1-carboxylate hormone biosynthesis. 1-aminocyclopropane-1-carboxylate synthase. LE-ACS1B.  |
| BAA94600.1 | AB033503 | Populus euramericana    | 1-aminocyclopropane-1-carboxylate synthase. peacs-2.  |
| BAA96743.1 | AB044662 | Prunus persica          | 1-aminocyclopropane-1-carboxylate synthase. PP-ACS1.  |
| CAB60831.1 | AJ012696 | Citrus sinensis         | ACC synthase. acs2.   |
| CAA09477.1 | AJ011095 | Citrus sinensis         | ACC synthase. acs-1.  |
| CAA41855.1 | X59139   | Lycopersicon esculentum | 1-aminocyclopropane 1-carboxylate synthase. LE-Acc2.  |
| BAA90549.1 | AB031026 | Prunus mume             | ACC synthase. PM-ACS1. 1-aminocyclopropane-1-carboxylic acid synthase.  |
| AAA81580.1 | M34289   | Lycopersicon esculentum | 1-aminocyclopropane-1-carboxylate synthase.   |
| CAA44397.1 | X62536   | Lycopersicon esculentum | 1-aminocyclopropane 1-carboxylate synthase.   |
| CAA46797.1 | X65982   | Nicotiana tabacum       | 1-aminocyclopropane 1-carboxylate synthase.   |
| BAA78333.1 | AB015625 | Pyrus pyrifolia         | 1-aminocyclopropane-1-carboxylic acid synthase. pPPACS3.  |
| CAA41856.1 | X59145   | Lycopersicon esculentum | 1-aminocyclopropane 1-carboxylate synthase. LE-ACC2.  |
| AAB17278.1 | U72389   | Lycopersicon esculentum | converts S-adenosylmethionine to 1-aminocyclopropane-1-carboxylate hormone biosynthesis. 1-aminocyclopropane-1-carboxylate synthase. LE-ACS1A.  |
| AAF97614.1 | U18056   | Lycopersicon esculentum | converts SAM to ACC; plant hormone biosynthesis. 1-amino-cyclopropane-1-carboxylate synthase. LE-ACS1A. ACC synthase; direct precursor of ethylene; one member of a multigene family. |
| BAB16433.1 | AB041521 | Solanum tuberosum       | ACC synthase. ppACS1.   |
| BAA34923.1 | AB013100 | Lycopersicon esculentum | 1-aminocyclopropane-1-carboxylate synthase. LE-ACS6. ACC synthase.  |
| BAA25916.1 | AB013346 | Lycopersicon esculentum | 1-aminocyclopropane-1-carboxylate synthase.   |

|            |          |                                |   |
|------------|----------|--------------------------------|---|
| AAF97615.1 | U18057   | <i>Lycopersicon esculentum</i> | converts SAM to ACC; plant hormone biosynthesis. 1-amino-cyclopropane-1-carboxylate synthase. LE-ACS1B. ACC synthase; direct precursor of ethylene; one member of a multigene family. |
| BAA92350.1 | AB034992 | <i>Malus x domestica</i>       | 1-aminocyclopropane-1-carboxylate synthase. MdACS-5A.   |
| BAA92351.1 | AB034993 | <i>Malus x domestica</i>       | 1-aminocyclopropane-1-carboxylate synthase. MdACS-5B.   |
| BAA93712.1 | AB032935 | <i>Cucumis melo</i>            | 1-aminocyclopropane-1-carboxylate synthase. CMe-ACS1.   |
| BAA94599.1 | AB033502 | <i>Populus euphratica</i>      | 1-aminocyclopropane-1-carboxylate synthase. peacs-1.  |
| BAA33374.1 | AB006803 | <i>Cucumis sativus</i>         | ACC synthase. CS-ACS1.  |
| BAA93714.1 | AB032937 | <i>Cucumis sativus</i>         | 1-aminocyclopropane-1-carboxylate synthase. CS-ACS1.  |
| AAC49153.1 | U17229   | <i>Pelargonium x hortorum</i>  | 1-aminocyclopropane 1-carboxylate synthase. ACC synthase.   |
| CAA06288.1 | AJ005002 | <i>Nicotiana tabacum</i>       | 1-cyclopropane-1-carboxylate synthase. acs2.  |
| CAB65314.1 | AJ131836 | <i>Nicotiana tabacum</i>       | 1-aminocyclopropane-1-carboxylate synthase. acs2.   |
| AAC15777.1 | AF061605 | <i>Nicotiana glutinosa</i>     | ACC synthase. ACS3.   |
| AAC83147.1 | AF057563 | <i>Nicotiana glutinosa</i>     | 1-aminocyclopropane-1-carboxylate synthase. ACS2. ACC synthase.   |
| BAA06464.1 | D30805   | <i>Cucumis melo</i>            | 1-aminocyclopropane-1-carboxylate synthase.   |
| BAA83618.1 | AB025906 | <i>Cucumis melo</i>            | 1-aminocyclopropane-1-carboxylate synthase. CMe-ACS1 (ME-ACS1).   |
| AAB70885.1 | U88971   | <i>Pelargonium x hortorum</i>  | 1-aminocyclopropane-1-carboxylate synthase. GACS2. similar to <i>Pelargonium hortorum</i> clone pGAC-2, Genbank Accession Number U17231.  |
| AAA33113.1 | M58323   | <i>Cucurbita pepo</i>          | 1-aminocyclopropane-1-carboxylate synthase.   |
| AAA33111.1 | M61195   | <i>Cucurbita pepo</i>          | 1-aminocyclopropane-1-carboxylate synthase. CP-ACC1A.   |
| CAA57724.1 | X82273   | <i>Brassica oleracea</i>       | 1-aminocyclopropane-1-carboxylate synthase.   |
| CAA47474.1 | X67100   | <i>Glycine max</i>             | 1-aminocyclopropane 1-carboxylate synthase.   |
| AAA33112.1 | M61195   | <i>Cucurbita pepo</i>          | 1-aminocyclopropane-1-carboxylate synthase. CP-ACC1B.   |

|               |          |                                      |  |
|---------------|----------|--------------------------------------|--|
| AAC98809.1    | U68216   | Carica papaya                        | ACC synthase. fruit specific; ripening related.      |
| CAA51227.1    | X72676   | Brassica juncea                      | 1-aminocyclopropane-1-carboxylate synthase. MACC.    |
| BAA00838.1    | D01032   | Cucurbita maxima                     | 1-aminocyclopropane-1-carboxylate synthase. accW.    |
| AAA34131.1    | M63490   | Lycopersicon esculentum              | enzyme. 1-aminocyclopropane-1-carboxylate synthase.  |
| AAA03164.1    | M88487   | Lycopersicon esculentum              | 1-aminocyclopropane-1-carboxylate synthase. ACC4.    |
| CAA41857.1    | X59146   | Lycopersicon esculentum              | 1-aminocyclopropane 1-carboxylate synthase. LE-ACC4. |
| CAA77688.1    | Z11613   | Vigna radiata                        | 1-aminocyclopropane 1-carboxylate synthase.          |
| CAA67118.1    | X98492   | Nicotiana tabacum                    | ACC synthase. ACCS2.                                 |
| SEQ ID NO: 99 |          |                                      |  |
| BAA22976.1    | D63457   | Arabis gemmifera                     | alcohol dehydrogenase. Adh. ADH.                     |
| BAA22973.1    | D63454   | Arabis gemmifera                     | alcohol dehydrogenase. Adh. ADH.                     |
| AAF23537.1    | AF110439 | Arabis glabra                        | alcohol dehydrogenase.                               |
| BAA22978.1    | D63459   | Arabis gemmifera                     | alcohol dehydrogenase. Adh. ADH.                     |
| BAA22974.1    | D63455   | Arabis gemmifera                     | alcohol dehydrogenase. Adh. ADH.                     |
| AAF23540.1    | AF110442 | Arabidopsis halleri                  | alcohol dehydrogenase.                               |
| AAF23551.1    | AF110453 | Arabidopsis lyrata subsp. petraea    | alcohol dehydrogenase.                               |
| AAF23539.1    | AF110441 | Halimolobos perplexa var. lemhiensis | alcohol dehydrogenase.                               |
| BAA22975.1    | D63456   | Arabis gemmifera                     | alcohol dehydrogenase. Adh. ADH.                     |
| AAF23546.1    | AF110448 | Arabis lyallii                       | alcohol dehydrogenase.                               |
| AAF23548.1    | AF110450 | Arabis parishii                      | alcohol dehydrogenase.                               |
| AAF23550.1    | AF110452 | Arabidopsis lyrata subsp. petraea    | alcohol dehydrogenase.                               |

|            |          |  |
|------------|----------|--|
| BAA22972.1 | D63453   | <i>Arabis gemmifera</i><br>alcohol dehydrogenase. Adh. ADH.              |
| BAA22971.1 | D63452   | <i>Arabis gemmifera</i><br>alcohol dehydrogenase. Adh. ADH.              |
| BAA22977.1 | D63458   | <i>Arabis gemmifera</i><br>alcohol dehydrogenase. Adh. ADH.              |
| AAF23549.1 | AF110451 | <i>Arabis pauciflora</i><br>alcohol dehydrogenase.                       |
| AAF23545.1 | AF110447 | <i>Arabis lignifera</i><br>alcohol dehydrogenase.                        |
| AAF23536.1 | AF110438 | <i>Arabis fendleri</i><br>alcohol dehydrogenase.                         |
| AAF23541.1 | AF110443 | <i>Arabis hirsuta</i><br>alcohol dehydrogenase.                          |
| AAF23531.1 | AF110433 | <i>Arabis blepharophylla</i><br>alcohol dehydrogenase.                   |
| AAF23530.1 | AF110432 | <i>Arabis blepharophylla</i><br>alcohol dehydrogenase.                   |
| AAF23523.1 | AF110425 | <i>Aubrieta deltoidea</i><br>alcohol dehydrogenase.                      |
| AAF23529.1 | AF110431 | <i>Arabis blepharophylla</i><br>alcohol dehydrogenase.                   |
| AAF23547.1 | AF110449 | <i>Arabidopsis lyrata</i> subsp. <i>lyrata</i><br>alcohol dehydrogenase. |
| AAF23533.1 | AF110435 | <i>Capsella rubella</i><br>alcohol dehydrogenase.                        |
| AAF23534.1 | AF110436 | <i>Arabis drummondii</i><br>alcohol dehydrogenase.                       |
| AAF23532.1 | AF110434 | <i>Brassica oleracea</i><br>alcohol dehydrogenase.                       |
| AAF23556.1 | AF110458 | <i>Barbarea vulgaris</i><br>alcohol dehydrogenase.                       |
| AAF23555.1 | AF110457 | <i>Arabis turrita</i><br>alcohol dehydrogenase.                          |
| AAF23538.1 | AF110440 | <i>Arabidopsis griffithiana</i><br>alcohol dehydrogenase.                |
| AAF23524.1 | AF110426 | <i>Arabis alpina</i><br>alcohol dehydrogenase.                           |
| AAF23543.1 | AF110445 | <i>Arabis hirsuta</i><br>alcohol dehydrogenase.                          |
| AAF23525.1 | AF110427 | <i>Arabis alpina</i><br>alcohol dehydrogenase.                           |



|                |          |   |
|----------------|----------|---|
| AAF23527.1     | AF110429 | <i>Arabis alpina</i><br>alcohol dehydrogenase.                                  |
| AAF23535.1     | AF110437 | <i>Arabis drummondii</i><br>alcohol dehydrogenase.                              |
| AAF23553.1     | AF110455 | <i>Arabis procurrens</i><br>alcohol dehydrogenase.                              |
| AAF23544.1     | AF110446 | <i>Arabis jacquini</i><br>alcohol dehydrogenase.                                |
| AAF23526.1     | AF110428 | <i>Arabis alpina</i><br>alcohol dehydrogenase.                                  |
| AAF23528.1     | AF110430 | <i>Cardamine amara</i><br>alcohol dehydrogenase.                                |
| AAF23552.1     | AF110454 | <i>Arabis procurrens</i><br>alcohol dehydrogenase.                              |
| AAF23542.1     | AF110444 | <i>Arabis hirsuta</i><br>alcohol dehydrogenase.                                 |
| AAC79418.1     | AF037560 | <i>Leavenworthia stylosa</i><br>alcohol dehydrogenase 3. Adh3.                  |
| BAA34682.1     | AB015504 | <i>Arabidopsis griffithiana</i><br>alcohol dehydrogenase. Adh.                  |
| BAA34685.1     | AB015507 | <i>Arabidopsis suecica</i><br>alcohol dehydrogenase. Adh.                       |
| BAA34683.1     | AB015505 | <i>Arabidopsis korshinskyi</i><br>alcohol dehydrogenase. Adh.                   |
| CAB72921.1     | AJ251281 | <i>Arabidopsis lyrata</i> subsp. <i>petraea</i><br>alcohol dehydrogenase. adh.  |
| CAB72920.1     | AJ251280 | <i>Arabidopsis lyrata</i> subsp. <i>petraea</i><br>alcohol dehydrogenase. adh.  |
| CAB72919.1     | AJ251279 | <i>Arabidopsis lyrata</i> subsp. <i>petraea</i><br>alcohol dehydrogenase. adh.  |
| CAB72918.1     | AJ251278 | <i>Arabidopsis lyrata</i> subsp. <i>petraea</i><br>alcohol dehydrogenase. adh.  |
| CAB72917.1     | AJ251277 | <i>Arabidopsis lyrata</i> subsp. <i>petraea</i><br>alcohol dehydrogenase. adh.  |
| SEQ ID NO: 100 |          |   |
| AAD09518.1     | U64925   | <i>Nicotiana tabacum</i><br>NTGP4, similar to AIG1; geranylgeranylated protein. |
| SEQ ID NO: 102 |          |   |
| BAA12039.1     | D83669   | <i>Spinacia oleracea</i><br>stromal ascorbate peroxidase.                       |
| BAA19611.1     | D77997   | <i>Spinacia oleracea</i><br>thylakoid-bound ascorbate peroxidase.               |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| BAA24610.1 | AB002467 | Spinacia oleracea             | stromal ascorbate peroxidase. APX2.   |
| BAA24609.1 | AB002467 | Spinacia oleracea             | thylakoid-bound ascorbate peroxidase. APX2.   |
| AAC19394.1 | AF069316 | Mesembryanthemum crystallinum | stromal L-ascorbate peroxidase precursor.   |
| AAC19393.1 | AF069315 | Mesembryanthemum crystallinum | thylakoid-bound L-ascorbate peroxidase precursor.   |
| BAA78553.1 | AB022274 | Chloroplast Nicotiana tabacum | stromal ascorbate peroxidase.   |
| BAA78552.1 | AB022273 | Chloroplast Nicotiana tabacum | thylakoid-bound ascorbate peroxidase.   |
| BAA22196.1 | D88420   | Cucurbita sp.                 | a hydrogen peroxide-scavenging enzyme. stromal ascorbate peroxidase.  |
| BAA12029.1 | D83656   | Cucurbita sp.                 | 2AsA+H2O2 → 2MDA+2H2O. thylakoid-bound ascorbate peroxidase.  |
| CAA11265.1 | AJ223325 | Chlamydomonas reinhardtii     | ascorbate peroxidase. apx1.   |
| BAA83595.1 | AB009084 | Chlamydomonas sp. W80         | chloroplast ascorbate peroxidase.   |
| AAD30294.1 | AF139190 | Mesembryanthemum crystallinum | cytosolic ascorbate peroxidase. similar to L0-328, R6-3, L72-2.   |
| AAB52954.1 | U37060   | Gossypium hirsutum            | ascorbate peroxidase. glyoxysomal membrane-bound protein.   |
| AAD43334.1 | AF159254 | Zantedeschia aethiopica       | ascorbate peroxidase. apx2. putative glyoxysomal membrane-bound protein.  |
| AAC08576.1 | AF053474 | Zantedeschia aethiopica       | cytosolic ascorbate peroxidase. apx.  |
| AAD43338.1 | AF158654 | Fragaria x ananassa           | cytosolic ascorbate peroxidase APX26.   |
| AAD43336.1 | AF158652 | Fragaria x ananassa           | cytosolic ascorbate peroxidase. ApxSC. corresponds to mRNA sequence deposited in GenBank Accession Number AF039953. |
| AAB95222.1 | AF039953 | Fragaria x ananassa           | cytosolic ascorbate peroxidase. ApxSC.  |
| AAD41408.1 | AF159633 | Fragaria x ananassa           | cytosolic ascorbate peroxidase. APX.  |
| AAD41407.1 | AF159632 | Fragaria x ananassa           | cytosolic ascorbate peroxidase. APX.  |
| AAD41403.1 | AF159628 | Fragaria x ananassa           | cytosolic ascorbate peroxidase. APX.  |
| AAD41402.1 | AF159627 | Fragaria x ananassa           | cytosolic ascorbate peroxidase. APX.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAD43337.1 | AF158653 | Fragaria x ananassa     | cytosolic ascorbate peroxidase APX19.                                    |
| AAD41406.1 | AF159631 | Fragaria x ananassa     | cytosolic ascorbate peroxidase. APX.                                     |
| AAD41404.1 | AF159629 | Fragaria x ananassa     | cytosolic ascorbate peroxidase. APX.                                     |
| AAB94574.1 | AF022213 | Fragaria x ananassa     | cytosolic ascorbate peroxidase. APX-c.                                   |
| AAD41405.1 | AF159630 | Fragaria x ananassa     | cytosolic ascorbate peroxidase. APX.                                     |
| BAA08264.1 | D45423   | Oryza sativa            | ascorbate peroxidase.  |
| BAA13671.1 | D88649   | Cucumis sativus         | cytosolic ascorbate peroxidase.  |
| CAA55209.1 | X78452   | Raphanus sativus        | L-ascorbate peroxidase. APX.   |
| BAA12890.1 | D85864   | Spinacia oleracea       | cytosolic ascorbate peroxidase.  |
| AAA99518.1 | L20864   | Spinacia oleracea       | ascorbate peroxidase.  |
| BAA12918.1 | D85912   | Nicotiana tabacum       | cytosolic ascorbate peroxidase.  |
| AAB03844.1 | U61379   | Vigna unguiculata       | ascorbate peroxidase.  |
| CAB58361.1 | Y16773   | Lycopersicon esculentum | ascorbate peroxidase. Apx. H2O2 scavenger, cytosolic protein.            |
| CAA57140.1 | X81376   | Capsicum annuum         | L-ascorbate peroxidase.  |
| CAA06996.1 | AJ006358 | Hordeum vulgare         | ascorbate peroxidase. apx.   |
| AAF22246.1 | AF159380 | Pimpinella brachycarpa  | ascorbate peroxidase. APX.   |
| AAA86689.1 | U15933   | Nicotiana tabacum       | reduction of hydrogen peroxide. ascorbate peroxidase. APX.               |
| CAA84406.1 | Z34934   | Zea mays                | cytosolic ascorbate peroxidase. apx.                                     |
| CAA43992.1 | X62077   | Pisum sativum           | L-ascorbate peroxidase. AppxI.   |
| AAA33645.1 | M93051   | Pisum sativum           | hydrogen peroxide removal; oxidative stress. ascorbate peroxidase. ApxI. |
| AAB01221.1 | U56634   | Glycine max             | ascorbate peroxidase 2. APx2.  |

---

AAA61779.1 L10292 Glycine max  
ascorbate peroxidase. Automated Edman sequencing of the N-terminal amino acids confirmed that the protein was authentic ascorbate peroxidase; putative.

---

BAB20889.1 AB053297 Oryza sativa  
L-ascorbate peroxidase. APXb.

---

CAA72247.1 Y11461 Brassica napus  
L-ascorbate peroxidase. APX.

---

BAB17666.1 AB050724 Oryza sativa subsp. japonica  
ascorbate peroxidase. APXb.

---

AAD20022.1 AF127804 Glycine max  
ascorbate peroxidase. apx1. AsPOX; cytosolic.

---

AAB94927.1 AF038839 Brassica juncea  
ascorbate peroxidase.

---

SEQ ID NO: 103

---

AAA74900.1 L34343 Ruta graveolens  
anthranilate synthase alpha subunit.

---

AAA74901.1 L34344 Ruta graveolens  
anthranilate synthase alpha subunit.

---

AAC27795.1 AF079168 Nicotiana tabacum  
feedback-insensitive anthranilate synthase alpha-2 chain. ASA2.

---

BAA82095.1 AB022603 Oryza sativa  
anthranilate synthase alpha 2 subunit. OsASA2.

---

BAA82094.1 AB022602 Oryza sativa  
anthranilate synthase alpha 1 subunit. OsASA1.

---

CAC29060.1 AJ250008 Catharanthus roseus  
enzyme in tryptophan biosynthesis. anthranilate synthase alpha subunit. asa.

---

CAA06837.1 AJ006065 Catharanthus roseus  
isochorismate synthase.

---

SEQ ID NO: 104

---

CAA57724.1 X82273 Brassica oleracea  
1-aminocyclopropane-1-carboxylate synthase.

---

CAA51227.1 X72676 Brassica juncea  
1-aminocyclopropane-1-carboxylate synthase. MACC.

---

AAC83147.1 AF057563 Nicotiana glutinosa  
1-aminocyclopropane-1-carboxylate synthase. ACS2. ACC synthase.

---

CAB60722.1 AJ012551 Citrus sinensis  
ACC synthase. acs1.

---

CAB60721.1 AJ012550 Citrus sinensis  
ACC synthase. acs1.

---

CAA06288.1 AJ005002 Nicotiana tabacum  
1-cyclopropane-1-carboxylate synthase. acs2.

---

CAB65314.1 AJ131836 Nicotiana tabacum  
1-aminocyclopropane-1-carboxylate synthase. acs2.

---

|            |          |                                |   |
|------------|----------|--------------------------------|---|
| BAA78333.1 | AB015625 | <i>Pyrus pyrifolia</i>         | 1-aminocyclopropane-1-carboxylic acid synthase. pPPACS3.  |
| AAB17279.1 | U72390   | <i>Lycopersicon esculentum</i> | converts S-adenosylmethionine to 1-aminocyclopropane-1-carboxylate hormone biosynthesis. 1-aminocyclopropane-1-carboxylate synthase. LE-ACS1B.  |
| AAF97614.1 | U18056   | <i>Lycopersicon esculentum</i> | converts SAM to ACC; plant hormone biosynthesis. 1-amino-cyclopropane-1-carboxylate synthase. LE-ACS1A. ACC synthase; direct precursor of ethylene; one member of a multigene family. |
| AAF97615.1 | U18057   | <i>Lycopersicon esculentum</i> | converts SAM to ACC; plant hormone biosynthesis. 1-amino-cyclopropane-1-carboxylate synthase. LE-ACS1B. ACC synthase; direct precursor of ethylene; one member of a multigene family. |
| AAB17278.1 | U72389   | <i>Lycopersicon esculentum</i> | converts S-adenosylmethionine to 1-aminocyclopropane-1-carboxylate hormone biosynthesis. 1-aminocyclopropane-1-carboxylate synthase. LE-ACS1A.  |
| CAA46797.1 | X65982   | <i>Nicotiana tabacum</i>       | 1-aminocyclopropane 1-carboxylate synthase.   |
| BAA06464.1 | D30805   | <i>Cucumis melo</i>            | 1-aminocyclopropane-1-carboxylate synthase.   |
| AAB06724.1 | U64804   | <i>Petunia x hybrida</i>       | ethylene biosynthesis enzyme. 1-aminocyclopropane-1-carboxylate synthase. PH-ACS1.  |
| CAA79478.1 | Z18953   | <i>Petunia x hybrida</i>       | 1-aminocyclopropane-1-carboxylate synthase.   |
| BAA94599.1 | AB033502 | <i>Populus euphratica</i>      | 1-aminocyclopropane-1-carboxylate synthase. peacs-1.  |
| CAA67118.1 | X98492   | <i>Nicotiana tabacum</i>       | ACC synthase. ACCS2.  |
| CAA79477.1 | Z18952   | <i>Dianthus caryophyllus</i>   | 1-aminocyclopropane 1-carboxylate synthase.   |
| AAA33275.1 | M66619   | <i>Dianthus caryophyllus</i>   | 1-aminocyclopropane-1-carboxylate synthase. CARACC.   |
| BAA83618.1 | AB025906 | <i>Cucumis melo</i>            | 1-aminocyclopropane-1-carboxylate synthase. CMe-ACS1 (ME-ACS1).   |
| BAA34923.1 | AB013100 | <i>Lycopersicon esculentum</i> | 1-aminocyclopropane-1-carboxylate synthase. LE-ACS6. ACC synthase.  |
| BAA25916.1 | AB013346 | <i>Lycopersicon esculentum</i> | 1-aminocyclopropane-1-carboxylate synthase.   |
| CAB60831.1 | AJ012696 | <i>Citrus sinensis</i>         | ACC synthase. acs2.   |
| CAA41855.1 | X59139   | <i>Lycopersicon esculentum</i> | 1-aminocyclopropane 1-carboxylate synthase. LE-Acc2.  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAA44397.1 | X62536   | Lycopersicon esculentum | 1-aminocyclopropane 1-carboxylate synthase.   |
| CAA41856.1 | X59145   | Lycopersicon esculentum | 1-aminocyclopropane 1-carboxylate synthase. LE-ACC2.  |
| AAA81580.1 | M34289   | Lycopersicon esculentum | 1-aminocyclopropane-1-carboxylate synthase.   |
| AAC98809.1 | U68216   | Carica papaya           | ACC synthase. fruit specific; ripening related.   |
| BAA92350.1 | AB034992 | Malus x domestica       | 1-aminocyclopropane-1-carboxylate synthase. MdACS-5A.   |
| BAA92351.1 | AB034993 | Malus x domestica       | 1-aminocyclopropane-1-carboxylate synthase. MdACS-5B.   |
| BAA94600.1 | AB033503 | Populus euramericana    | 1-aminocyclopropane-1-carboxylate synthase. peacs-2.  |
| CAA77688.1 | Z11613   | Vigna radiata           | 1-aminocyclopropane 1-carboxylate synthase.   |
| CAA47474.1 | X67100   | Glycine max             | 1-aminocyclopropane 1-carboxylate synthase.   |
| AAB70885.1 | U88971   | Pelargonium x hortorum  | 1-aminocyclopropane-1-carboxylate synthase. GACS2. similar to Pelargonium hortorum clone pGAC-2, Genbank Accession Number U17231. |
| AAD04199.1 | AF016459 | Pisum sativum           | 1-aminocyclopropane-1-carboxylate synthase. ACS2. ACC synthase.   |
| BAB18464.1 | D86242   | Cucumis melo            | 1-aminocyclopropane-1-carboxylate synthase. CMe-ACS2 (ME-ACS2).   |
| BAA93713.1 | AB032936 | Cucumis melo            | 1-aminocyclopropane-1-carboxylate synthase. CMe-ACS2.   |
| BAA93715.1 | AB032938 | Cucumis sativus         | 1-aminocyclopropane-1-carboxylate synthase. CS-ACS2.  |
| BAA33375.1 | AB006804 | Cucumis sativus         | ACC synthase. CS-ACS2.  |
| BAB16433.1 | AB041521 | Solanum tuberosum       | ACC synthase. ppACS1.   |
| AAC15777.1 | AF061605 | Nicotiana glutinosa     | ACC synthase. ACS3.   |
| CAA72191.1 | Y11357   | Carica papaya           | 1-aminocyclopropane-1-carboxylate synthase.   |
| BAA96743.1 | AB044662 | Prunus persica          | 1-aminocyclopropane-1-carboxylate synthase. PP-ACS1.  |
| BAA90549.1 | AB031026 | Prunus mume             | ACC synthase. PM-ACS1. 1-aminocyclopropane-1-carboxylic acid synthase.  |
| BAA93712.1 | AB032935 | Cucumis melo            | 1-aminocyclopropane-1-carboxylate synthase. CMe-ACS1.   |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAF22109.1     | AF119411 | Lupinus albus                 | ethylene biosynthesis enzyme. 1-aminocyclopropane-1-carboxylate synthase 1. ACS1. ACC synthase; S-adenosyl-L-methionine methylthioadenosine-lyase.                    |
| AAC83146.1     | AF057562 | Nicotiana glutinosa           | 1-aminocyclopropane-1-carboxylate synthase. ACS1. ACC synthase.   |
| CAA09477.1     | AJ011095 | Citrus sinensis               | ACC synthase. acs-1.  |
| BAA93714.1     | AB032937 | Cucumis sativus               | 1-aminocyclopropane-1-carboxylate synthase. CS-ACS1.  |
| SEQ ID NO: 106 |          |                               |   |
| AAF63205.1     | AF245119 | Mesembryanthemum crystallinum | AP2-related transcription factor. CDBP. stress induced transcription factor.  |
| BAA97122.1     | AB016264 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2. |
| AAC50047.1     | U89255   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti4.  |
| BAA07321.1     | D38123   | Nicotiana tabacum             | ERF1. ethylene-responsive transcription factor.   |
| AAC62619.1     | AF057373 | Nicotiana tabacum             | transcription factor. ethylene response element binding protein 1. EREBP1.  |
| BAA87068.1     | AB035270 | Matricaria chamomilla         | ethylene-responsive element binding protein1 homolog. McEREBP1.   |
| AAC49740.1     | U89256   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti5.  |
| AAG43545.1     | AF211527 | Nicotiana tabacum             | Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.  |
| BAA97124.1     | AB016266 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4. |
| CAB96900.1     | AJ251250 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| CAB96899.1     | AJ251249 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| AAB38748.1     | U81157   | Nicotiana tabacum             | S25-XP1 DNA binding protein.  |
| AAG60182.1     | AC084763 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBa0027P10.12.   |
| AAK31279.1     | AC079890 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBb0089A17.16.   |
| CAB93940.1     | AJ238740 | Catharanthus roseus           | putative transcription factor. AP2-domain DNA-binding protein. orca2.   |

|                |          |                                |  |
|----------------|----------|--------------------------------|--|
| AAC49741.1     | U89257   | <i>Lycopersicon esculentum</i> | DNA-binding protein binds the GCC box pathogenesis-related promoter element. Pti6.   |
| CAC12822.1     | AJ299252 | <i>Nicotiana tabacum</i>       | AP2 domain-containing transcription factor. ap2.   |
| BAB03248.1     | AB037183 | <i>Oryza sativa</i>            | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.  |
| AAC14323.1     | AF058827 | <i>Nicotiana tabacum</i>       | TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.   |
| BAA97123.1     | AB016265 | <i>Nicotiana sylvestris</i>    | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.                |
| AAD00708.1     | U91857   | <i>Stylosanthes hamata</i>     | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively. |
| AAF05606.1     | AF190770 | <i>Oryza sativa</i>            | EREBP-like protein. tsh1. TSH1; induced by ethylene.   |
| BAA76734.1     | AB024575 | <i>Nicotiana tabacum</i>       | ethylene responsive element binding factor.  |
| AAC29516.1     | U77655   | <i>Solanum tuberosum</i>       | DNA binding protein homolog. STWAAEIRD.  |
| AAF23899.1     | AF193803 | <i>Oryza sativa</i>            | transcription factor EREBP1. EREBP/AP2-like transcription factor.  |
| CAB93939.1     | AJ238739 | <i>Catharanthus roseus</i>     | putative transcription factor. AP2-domain DNA-binding protein. orca1.  |
| AAD45623.1     | AF084185 | <i>Brassica napus</i>          | dehydration responsive element binding protein. DNA binding protein; DRE binding protein.  |
| AAK31271.1     | AC079890 | <i>Oryza sativa</i>            | putative transcriptional factor. OSJNBb0089A17.22.   |
| AAG43548.1     | AF211530 | <i>Nicotiana tabacum</i>       | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.   |
| BAB21218.1     | AP002913 | <i>Oryza sativa</i>            | putative DNA binding protein RAV2. P0480E02.24.  |
| AAG43549.1     | AF211531 | <i>Nicotiana tabacum</i>       | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.   |
| BAB21211.1     | AP002913 | <i>Oryza sativa</i>            | putative DNA binding protein RAV2. P0480E02.17.  |
| AAK01088.1     | AF298230 | <i>Hordeum vulgare</i>         | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.   |
| AAK01089.1     | AF298231 | <i>Hordeum vulgare</i>         | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.  |
| SEQ ID NO: 107 |          |                                |  |



|            |          |                               |  |
|------------|----------|-------------------------------|--|
| AAF63205.1 | AF245119 | Mesembryanthemum crystallinum | AP2-related transcription factor. CDBP. stress induced transcription factor.   |
| AAC50047.1 | U89255   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti4.   |
| BAA97122.1 | AB016264 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2.                |
| BAA07321.1 | D38123   | Nicotiana tabacum             | ERF1. ethylene-responsive transcription factor.  |
| AAC62619.1 | AF057373 | Nicotiana tabacum             | transcription factor. ethylene response element binding protein 1. EREBP1.   |
| BAA87068.1 | AB035270 | Matricaria chamomilla         | ethylene-responsive element binding protein1 homolog. McEREBP1.  |
| CAB96899.1 | AJ251249 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| CAB96900.1 | AJ251250 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| AAB38748.1 | U81157   | Nicotiana tabacum             | S25-XP1 DNA binding protein.   |
| AAC49740.1 | U89256   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti5.   |
| CAB93940.1 | AJ238740 | Catharanthus roseus           | putative transcription factor. AP2-domain DNA-binding protein. orca2.  |
| AAG43545.1 | AF211527 | Nicotiana tabacum             | Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.   |
| BAA97124.1 | AB016266 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4.                |
| AAG60182.1 | AC084763 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBa0027P10.12.  |
| AAK31279.1 | AC079890 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBb0089A17.16.  |
| BAA97123.1 | AB016265 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.                |
| AAD00708.1 | U91857   | Stylosanthes hamata           | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively. |
| AAC14323.1 | AF058827 | Nicotiana tabacum             | TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.   |
| CAC12822.1 | AJ299252 | Nicotiana tabacum             | AP2 domain-containing transcription factor. ap2.   |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| BAB03248.1     | AB037183 | Oryza sativa            | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.   |
| AAF05606.1     | AF190770 | Oryza sativa            | EREBP-like protein. tsh1. TSH1; induced by ethylene.  |
| BAA76734.1     | AB024575 | Nicotiana tabacum       | ethylene responsive element binding factor.   |
| AAC49741.1     | U89257   | Lycopersicon esculentum | DNA-binding protein binds the GCC box pathogenesis-related promoter element. Pti6.  |
| AAF23899.1     | AF193803 | Oryza sativa            | transcription factor EREBP1. EREBP/AP2-like transcription factor.   |
| AAC29516.1     | U77655   | Solanum tuberosum       | DNA binding protein homolog. STWAAEIRD.   |
| CAB93939.1     | AJ238739 | Catharanthus roseus     | putative transcription factor. AP2-domain DNA-binding protein. orca1.   |
| BAA78738.1     | AB023482 | Oryza sativa            | EST AU055776(S20048) corresponds to a region of the predicted gene.; Similar to Arabidopsis thaliana AP2 domain containing protein RAP2.10 mRNA, partial cds.(AF003103).          |
| AAD45623.1     | AF084185 | Brassica napus          | dehydration responsive element binding protein. DNA binding protein; DRE binding protein.   |
| AAG43548.1     | AF211530 | Nicotiana tabacum       | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.  |
| AAG43549.1     | AF211531 | Nicotiana tabacum       | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.  |
| BAA99376.1     | AP002526 | Oryza sativa            | ESTs AU093391(E60370),AU091593(C60458), AU093392(E60370) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana BAC F21J9; AP2 domain protein. (AC000103). |
| BAB21218.1     | AP002913 | Oryza sativa            | putative DNA binding protein RAV2. P0480E02.24.   |
| AAK31271.1     | AC079890 | Oryza sativa            | putative transcriptional factor. OSJNBb0089A17.22.  |
| BAB21211.1     | AP002913 | Oryza sativa            | putative DNA binding protein RAV2. P0480E02.17.   |
| AAK01088.1     | AF298230 | Hordeum vulgare         | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.  |
| AAG59618.1     | AF239616 | Hordeum vulgare         | CRT/DRE-binding factor. CBF.  |
| AAK01089.1     | AF298231 | Hordeum vulgare         | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.   |
| SEQ ID NO: 109 |          |                         |   |

|            |          |                               |  |
|------------|----------|-------------------------------|--|
| BAA97124.1 | AB016266 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4.                |
| AAG43545.1 | AF211527 | Nicotiana tabacum             | Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.   |
| AAK31279.1 | AC079890 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBb0089A17.16.  |
| AAG60182.1 | AC084763 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBa0027P10.12.  |
| BAA87068.1 | AB035270 | Matricaria chamomilla         | ethylene-responsive element binding protein1 homolog. McEREBP1.  |
| AAF63205.1 | AF245119 | Mesembryanthemum crystallinum | AP2-related transcription factor. CDBP. stress induced transcription factor.   |
| AAC50047.1 | U89255   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti4.   |
| BAA97122.1 | AB016264 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2.                |
| BAA07321.1 | D38123   | Nicotiana tabacum             | ERF1. ethylene-responsive transcription factor.  |
| CAB93940.1 | AJ238740 | Catharanthus roseus           | putative transcription factor. AP2-domain DNA-binding protein. orca2.  |
| AAC49740.1 | U89256   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti5.   |
| AAC62619.1 | AF057373 | Nicotiana tabacum             | transcription factor. ethylene response element binding protein 1. EREBP1.   |
| AAB38748.1 | U81157   | Nicotiana tabacum             | S25-XP1 DNA binding protein.   |
| CAC12822.1 | AJ299252 | Nicotiana tabacum             | AP2 domain-containing transcription factor. ap2.   |
| CAB96900.1 | AJ251250 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| CAB96899.1 | AJ251249 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| BAA97123.1 | AB016265 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.                |
| BAB03248.1 | AB037183 | Oryza sativa                  | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.  |
| AAD00708.1 | U91857   | Stylosanthes hamata           | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively. |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAC14323.1 | AF058827 | Nicotiana tabacum       | TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.  |
| AAC49741.1 | U89257   | Lycopersicon esculentum | DNA-binding protein binds the GCC box pathogenesis-related promoter element. Pti6.  |
| AAC29516.1 | U77655   | Solanum tuberosum       | DNA binding protein homolog. STWAAEIRD.   |
| AAF05606.1 | AF190770 | Oryza sativa            | EREBP-like protein. tsh1. TSH1; induced by ethylene.  |
| BAA76734.1 | AB024575 | Nicotiana tabacum       | ethylene responsive element binding factor.   |
| AAF23899.1 | AF193803 | Oryza sativa            | transcription factor EREBP1. EREBP/AP2-like transcription factor.   |
| BAA78738.1 | AB023482 | Oryza sativa            | EST AU055776(S20048) corresponds to a region of the predicted gene.; Similar to Arabidopsis thaliana AP2 domain containing protein RAP2.10 mRNA, partial cds.(AF003103).  |
| CAB93939.1 | AJ238739 | Catharanthus roseus     | putative transcription factor. AP2-domain DNA-binding protein. orca1.   |
| AAD45623.1 | AF084185 | Brassica napus          | dehydration responsive element binding protein. DNA binding protein; DRE binding protein.   |
| AAG43549.1 | AF211531 | Nicotiana tabacum       | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.  |
| AAG43548.1 | AF211530 | Nicotiana tabacum       | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.  |
| AAK31271.1 | AC079890 | Oryza sativa            | putative transcriptional factor. OSJNBb0089A17.22.  |
| AAK01088.1 | AF298230 | Hordeum vulgare         | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.  |
| BAA99376.1 | AP002526 | Oryza sativa            | ESTs AU093391(E60370), AU091593(C60458), AU093392(E60370) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana BAC F21J9; AP2 domain protein. (AC000103).  |
| AAG59619.1 | AF243384 | Oryza sativa            | CRT/DRE binding factor. CBF. DREB.  |
| BAB21218.1 | AP002913 | Oryza sativa            | putative DNA binding protein RAV2. P0480E02.24.   |
| AAG32659.1 | AF253971 | Picea abies             | APETALA2-related transcription factor 2. AP2L2. PaAP2L2.  |
| AAC49567.1 | U41466   | Zea mays                | Glossy15. Glossy15. AP2 DNA-binding domain protein; similar to the Arabidopsis floral homeotic gene APETALA2; maize AP2-domain regulator of leaf epidermal traits; homeotic regulator of leaf epidermal cell identity; allele: W64A; putative transcription factor. |

SEQ ID NO: 110

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAG43545.1 | AF211527 | Nicotiana tabacum             | Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.  |
| BAA97124.1 | AB016266 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4. |
| AAK31279.1 | AC079890 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBb0089A17.16.   |
| AAG60182.1 | AC084763 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBa0027P10.12.   |
| AAF63205.1 | AF245119 | Mesembryanthemum crystallinum | AP2-related transcription factor. CDBP. stress induced transcription factor.  |
| BAA97122.1 | AB016264 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2. |
| BAA87068.1 | AB035270 | Matricaria chamomilla         | ethylene-responsive element binding protein1 homolog. McEREBP1.   |
| AAC50047.1 | U89255   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti4.  |
| BAA07321.1 | D38123   | Nicotiana tabacum             | ERF1. ethylene-responsive transcription factor.   |
| AAC49740.1 | U89256   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti5.  |
| CAB93940.1 | AJ238740 | Catharanthus roseus           | putative transcription factor. AP2-domain DNA-binding protein. orca2.   |
| AAC62619.1 | AF057373 | Nicotiana tabacum             | transcription factor. ethylene response element binding protein 1. EREBP1.  |
| CAB96899.1 | AJ251249 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| CAB96900.1 | AJ251250 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| AAB38748.1 | U81157   | Nicotiana tabacum             | S25-XP1 DNA binding protein.  |
| CAC12822.1 | AJ299252 | Nicotiana tabacum             | AP2 domain-containing transcription factor. ap2.  |
| AAF05606.1 | AF190770 | Oryza sativa                  | EREBP-like protein. tsh1. TSH1; induced by ethylene.  |
| AAC29516.1 | U77655   | Solanum tuberosum             | DNA binding protein homolog. STWAAEIRD.   |
| AAF23899.1 | AF193803 | Oryza sativa                  | transcription factor EREBP1. EREBP/AP2-like transcription factor.   |
| BAB03248.1 | AB037183 | Oryza sativa                  | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAA97123.1     | AB016265 | Nicotiana sylvestris    | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.                |
| BAA76734.1     | AB024575 | Nicotiana tabacum       | ethylene responsive element binding factor.  |
| AAD00708.1     | U91857   | Stylosanthes hamata     | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively. |
| AAC49741.1     | U89257   | Lycopersicon esculentum | DNA-binding protein binds the GCC box pathogenesis-related promoter element. Pti6.   |
| AAC14323.1     | AF058827 | Nicotiana tabacum       | TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.   |
| CAB93939.1     | AJ238739 | Catharanthus roseus     | putative transcription factor. AP2-domain DNA-binding protein. orca1.  |
| AAD45623.1     | AF084185 | Brassica napus          | dehydration responsive element binding protein. DNA binding protein; DRE binding protein.  |
| AAG43548.1     | AF211530 | Nicotiana tabacum       | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.   |
| AAG43549.1     | AF211531 | Nicotiana tabacum       | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.   |
| AAK31271.1     | AC079890 | Oryza sativa            | putative transcriptional factor. OSJNBb0089A17.22.   |
| BAB21218.1     | AP002913 | Oryza sativa            | putative DNA binding protein RAV2. P0480E02.24.  |
| AAK01088.1     | AF298230 | Hordeum vulgare         | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.   |
| AAG59619.1     | AF243384 | Oryza sativa            | CRT/DRE binding factor. CBF. DREB.   |
| BAB21211.1     | AP002913 | Oryza sativa            | putative DNA binding protein RAV2. P0480E02.17.  |
| AAK01089.1     | AF298231 | Hordeum vulgare         | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.  |
| SEQ ID NO: 111 |          |                         |  |
| AAF87216.1     | AF231351 | Nicotiana tabacum       | plastidic glucose 6-phosphate dehydrogenase. G6PDHP2.  |
| CAA67782.1     | X99405   | Nicotiana tabacum       | glucose-6-phosphate dehydrogenase. G6PD.   |
| AAB69317.1     | AF012861 | Petroselinum crispum    | plastidic glucose-6-phosphate dehydrogenase. pG6PDH.   |
| CAB52708.1     | AJ010712 | Solanum tuberosum       | glucose-6-phosphate 1-dehydrogenase. g6pd.   |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| CAB52685.1     | AJ132346 | Dunaliella bioculata          | plastidic glucose-6-phosphate dehydrogenase. g6PD.  |
| CAA04994.1     | AJ001772 | Nicotiana tabacum             | glucose-6-phosphate dehydrogenase. TCG18.   |
| CAA58775.1     | X83923   | Solanum tuberosum             | glucose-6-phosphate dehydrogenase.  |
| CAA03941.1     | AJ000184 | Spinacia oleracea             | Glucose-6-phosphate dehydrogenase. G6PD.  |
| CAA03939.1     | AJ000182 | Spinacia oleracea             | Glucose-6-phosphate dehydrogenase. G6PD.  |
| CAA03940.1     | AJ000183 | Spinacia oleracea             | Glucose-6-phosphate dehydrogenase. G6PD.  |
| AAB41552.1     | U18238   | Medicago sativa subsp. sativa | glucose-6-phosphate dehydrogenase.  |
| AAD11426.1     | AF097663 | Mesembryanthemum crystallinum | cytoplasmic glucose-6-phosphate 1-dehydrogenase. G6PD.  |
| AAB69319.1     | AF012863 | Petroselinum crispum          | cytosolic glucose-6-phosphate dehydrogenase 2. cG6PDH2.                                       |
| CAA52442.1     | X74421   | Solanum tuberosum             | glucose-6-phosphate 1-dehydrogenase. g6pdh. homotetramer.                                     |
| AAB69318.1     | AF012862 | Petroselinum crispum          | cytosolic glucose-6-phosphate dehydrogenase 1. cG6PDH1.                                       |
| BAA97662.1     | AB029454 | Triticum aestivum             | glucose-6-phosphate dehydrogenase. g6pdh.   |
| BAA97663.1     | AB029455 | Triticum aestivum             | glucose-6-phosphate dehydrogenase. g6pdh.   |
| CAA04993.1     | AJ001770 | Nicotiana tabacum             | glucose-6-phosphate dehydrogenase. TCG9.  |
| CAA04992.1     | AJ001769 | Nicotiana tabacum             | glucose-6-phosphate dehydrogenase. TCG6.  |
| BAA97664.1     | AB029456 | Triticum aestivum             | glucose-6-phosphate dehydrogenase. g6pdh.   |
| AAG23802.1     | AF260736 | Cucurbita pepo                | plastidic glucose-6-phosphate dehydrogenase.  |
| CAB66330.1     | AJ279688 | Betula pendula                | glucose-6-phosphate dehydrogenase. g6pd.  |
| BAA82155.1     | AB011441 | Triticum aestivum             | glucose-6-phosphate dehydrogenase. WESR5. salt-stress responding gene.                        |
| CAA06200.1     | AJ004900 | Glycine max                   | pentose phosphate pathway oxidoreductase generating NADPH. glucose-6-phosphate-dehydrogenase. |
| SEQ ID NO: 112 |          |                               |   |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| CAA48611.1 | X68652   | Raphanus sativus        | hydroxymethylglutaryl-CoA reductase (NADPH). HMG2.   |
| CAA48610.1 | X68651   | Raphanus sativus        | hydroxymethylglutaryl-CoA reductase (NADPH). HMG1.   |
| AAC05089.1 | AF038046 | Gossypium hirsutum      | catalyzes the synthesis of mevalonate from hmg-coA. 3-hydroxy-3-methylglutaryl-coenzyme A reductase 2. hmg2. hmg-coA reductase 2; HMGR2.   |
| AAA33108.1 | M96068   | Catharanthus roseus     | hydroxymethylglutaryl-CoA reductase. HMGR.   |
| BAA36291.1 | AB021862 | Cucumis melo            | HMG-CoA reductase. Cm-HMGR. putative.  |
| AAB52551.1 | U51985   | Solanum tuberosum       | HMG-CoA reductase.   |
| AAA93498.1 | L01400   | Solanum tuberosum       | convert HMG-CoA into mevalonate. hydroxymethylglutaryl coenzyme A reductase. hmgr. putative.   |
| AAB87727.1 | U60452   | Nicotiana tabacum       | hydroxy-methylglutaryl-coenzyme A reductase. HMGR1.  |
| CAA70440.1 | Y09238   | Zea mays                | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| CAA45181.1 | X63649   | Nicotiana glauca        | catalyzes synthesis of mevalonate. 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR. endoplasmic reticulum location.  |
| AAD28179.1 | AF110383 | Capsicum annuum         | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR2M. HMGR.   |
| AAB52552.1 | U51986   | Solanum tuberosum       | HMG-CoA reductase.   |
| BAA93631.1 | AB022690 | Solanum tuberosum       | 3-hydroxy-3-methylglutaryl coenzyme A reductase. HMG3.   |
| AAB62581.1 | U68072   | Lycopersicon esculentum | 3-hydroxy-3-methylglutaryl CoA reductase 2. HMG2.  |
| AAB69726.1 | U72145   | Camptotheca acuminata   | converts HMGCofA to mevalonate. 3-hydroxy-3-methylglutaryl coenzyme a reductase. hmg3. HMGR.   |
| AAB53748.1 | U95816   | Oryza sativa            | 3-hydroxy-3-methylglutaryl-CoA reductase. HMGR.  |
| BAB20771.1 | AB041031 | Solanum tuberosum       | 3-hydroxy-3-methylglutaryl coenzyme A reductase. HMG2.   |
| AAB69727.1 | U72146   | Camptotheca acuminata   | 3-hydroxy-3-methylglutaryl coenzyme A reductase. converts HMGCofA in mevalonate precursor for isoprenoid compounds; HMGR; hmg2; similar to protein encoded by GenBank Accession Number L10390. |



|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAD47596.1 | AF142473 | Artemisia annua         | HMG-CoA reductase. HMGR1.  |
| AAA34169.1 | M63642   | Lycopersicon esculentum | 3-hydroxy-3-methylglutaryl coenzyme A reductase. hmg2.   |
| AAA68966.1 | U14625   | Artemisia annua         | 3-hydroxy-3-methylglutaryl coenzyme A reductase. AHM4.   |
| AAA33358.1 | M74798   | Hevea brasiliensis      | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. hmgr3.  |
| AAA68965.1 | U14624   | Artemisia annua         | 3-hydroxy-3-methylglutaryl coenzyme A reductase. AHM1.   |
| AAD08820.1 | U43961   | Oryza sativa            | 3-hydroxy-3-methylglutaryl-CoA reductase. HMGR.  |
| AAA33360.1 | M74800   | Hevea brasiliensis      | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. hmgr3.  |
| CAA92821.1 | Z68504   | Oryza sativa            | 3-hydroxy-3-methylglutaryl-CoA reductase.  |
| AAC05088.1 | AF038045 | Gossypium hirsutum      | catalyzes the synthesis of mevalonate from hmg-coA. 3-hydroxy-3-methylglutaryl-coenzyme A reductase 1. hmg1. hmg-coA reductase 1; HMGR1. |
| AAC15475.1 | AF034760 | Tagetes erecta          | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| AAC15476.1 | AF034761 | Tagetes erecta          | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| CAA38469.1 | X54659   | Hevea brasiliensis      | hydroxymethylglutaryl-CoA reductase. HMGR1.  |
| CAA38467.1 | X54657   | Hevea brasiliensis      | hydroxymethylglutaryl-CoA reductase. HMGR1.  |
| AAD38873.1 | AF110382 | Oryza sativa            | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR3G. HMG-CoA reductase.  |
| AAA33040.1 | L10390   | Camptotheca acuminata   | 3-hydroxy-3-methylglutaryl coA reductase.  |
| AAD03789.1 | U43711   | Morus alba              | catalyzes the final step in mevalonate pathway. 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR.                                   |
| AAA21720.1 | L28995   | Oryza sativa            | conversion of hydroxymethylglutaryl coenzyme A to mevalonate. 3-hydroxy-3-methylglutaryl coenzyme A reductase. putative.                 |
| AAC72378.1 | AF096838 | Solanum tuberosum       | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| AAB04043.1 | L40938   | Lycopersicon esculentum | HMGR CoA reductase. HMGR1.   |
| CAA38468.1 | X54658   | Hevea brasiliensis      | hydroxymethylglutaryl-CoA reductase. HMGR2.  |

|                |          |                            |  |
|----------------|----------|----------------------------|--|
| CAA52787.1     | X74783   | Lithospermum erythrorhizon | 3-hydroxy-3-methyl glutaryl coenzyme A reductase. Lehmgr1.   |
| AAD09278.1     | U97683   | Glycine max                | catalyzes the synthesis of mevalonate, the specific precursor of all isoprenoid compounds present in plants. 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMG-CoA reductase.           |
| AAG43469.1     | AF196964 | Bixa orellana              | catalyzes mevalonate synthesis from hmg-CoA. 3-hydroxy-3-methylglutaryl-coenzyme A reductase.  |
| BAA09705.1     | D63389   | Cucumis sativus            | 3-hydroxy-3-methylglutaryl CoA reductase.  |
| AAB47161.1     | S82272   | Gossypium barbadense       | 3-hydroxy-3-methylglutaryl coenzyme A reductase. /gene="3-hydroxy-3-methylglutaryl coenzyme A reductase,. This sequence comes from Fig. 2; 3-hydroxy-3-methylglutaryl CoA reductase; HMGR. |
| AAA33359.1     | M74799   | Hevea brasiliensis         | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. hmgr3.  |
| AAC37432.1     | L34825   | Solanum tuberosum          | HMG-CoA reductase. hmgl gene family.   |
| AAC37434.1     | L34827   | Solanum tuberosum          | HMG-CoA reductase. hmgl gene family.   |
| AAC37431.1     | L34823   | Solanum tuberosum          | HMG-CoA reductase. hmgl gene family.   |
| AAC37433.1     | L34826   | Solanum tuberosum          | HMG-CoA reductase. hmgl gene family.   |
| AAC37435.1     | L34828   | Solanum tuberosum          | HMG-CoA reductase. hmgl gene family.   |
| AAC37436.1     | L34829   | Solanum tuberosum          | HMG-CoA reductase. hmgl gene family.   |
| SEQ ID NO: 113 |          |                            |  |
| AAC49676.1     | U77345   | Zea mays                   | lethal leaf-spot 1. lls1. Allele: wild-type; LLS1; similar to bacterial ring-hydroxylating dioxygenase.  |
| AAG03051.1     | AF284781 | Oryza sativa               | LLS1 protein. Lls1.  |
| SEQ ID NO: 115 |          |                            |  |
| BAA82379.1     | AP000367 | Oryza sativa               | Similar to putative receptor kinase. (AC002332).   |
| BAA87853.1     | AP000816 | Oryza sativa               | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).   |
| AAG25966.1     | AF302082 | Nicotiana tabacum          | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.  |

|            |          |                     |   |
|------------|----------|---------------------|---|
| BAB39873.1 | AP002882 | Oryza sativa        | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| AAC27894.1 | AF023164 | Zea mays            | leucine-rich repeat transmembrane protein kinase 1. ltk1.   |
| AAC27895.1 | AF023165 | Zea mays            | leucine-rich repeat transmembrane protein kinase 2. ltk2.   |
| BAB21240.1 | AP002953 | Oryza sativa        | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| BAB16871.1 | AP002537 | Oryza sativa        | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).  |
| BAB39409.1 | AP002901 | Oryza sativa        | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| AAF66615.1 | AF142596 | Nicotiana tabacum   | LRR receptor-like protein kinase.   |
| CAB41878.1 | Y18259   | Brassica oleracea   | SRK5 protein. SRK5. receptor-like kinase.   |
| AAG59657.1 | AC084319 | Oryza sativa        | putative protein kinase. OSJNBa0004B24.20.  |
| BAB03429.1 | AP002817 | Oryza sativa        | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587). |
| BAB07999.1 | AP002525 | Oryza sativa        | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| AAK00425.1 | AC069324 | Oryza sativa        | Putative protein kinase. OSJNBa0071K19.11.  |
| CAB41879.1 | Y18260   | Brassica oleracea   | SRK15 protein. SRK15. receptor-like kinase.   |
| AAK21965.1 | AY028699 | Brassica napus      | receptor protein kinase PERK1.  |
| AAD21872.1 | AF078082 | Phaseolus vulgaris  | receptor-like protein kinase homolog RK20-1.  |
| CAA97692.1 | Z73295   | Catharanthus roseus | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.                                    |
| CAA74661.1 | Y14285   | Brassica oleracea   | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.  |
| AAC23542.1 | U20948   | Ipomoea trifida     | receptor protein kinase. IRK1.  |
| AAG16628.1 | AY007545 | Brassica napus      | protein serine/threonine kinase BNK1.   |

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAA94509.1 | AB041503 | Populus nigra        | protein kinase 1. PnPK1.  |
| BAA23676.1 | AB000970 | Brassica rapa        | receptor kinase 1. BcRK1.   |
| CAA47962.1 | X67733   | Zea mays             | receptor-like protein kinase. PK1.  |
| CAA67145.1 | X98520   | Brassica oleracea    | receptor-like kinase. SFR2.   |
| AAB09771.1 | U67422   | Zea mays             | CRINKLY4 precursor. cr4. receptor kinase homolog.   |
| CAA74662.1 | Y14286   | Brassica oleracea    | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.  |
| BAA94510.1 | AB041504 | Populus nigra        | protein kinase 2. PnPK2.  |
| AAB93834.1 | U82481   | Zea mays             | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.  |
| CAA73133.1 | Y12530   | Brassica oleracea    | serine /threonine kinase. ARLK.   |
| BAA06538.1 | D31737   | Nicotiana tabacum    | protein-serine/threonine kinase.  |
| AAF43496.1 | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| AAK11674.1 | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| BAB21001.1 | AB054061 | Brassica rapa        | S locus receptor kinase. SRK22.   |
| CAB51480.1 | Y14600   | Sorghum bicolor      | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| BAA87852.1 | AP000816 | Oryza sativa         | Similar to putative Ser/Thr protein kinase. (AC004218).   |
| BAA92221.1 | AP001278 | Oryza sativa         | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).  |
| BAA78764.1 | AB023482 | Oryza sativa         | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| BAA90808.1 | AP001168 | Oryza sativa         | Similar to putative receptor-like protein kinase (AL035679).  |
| BAA07576.1 | D38563   | Brassica rapa        | receptor protein kinase SRK8.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAB47422.1     | U59318   | Lycopersicon esculentum | serine/threonine protein kinase Fen. fen. allele of Fen, fenthion sensitivity gene from tomato.  |
| BAA21132.1     | D88193   | Brassica rapa           | S-receptor kinase. SRK9 (B.c).   |
| SEQ ID NO: 117 |          |                         |  |
| AAA34002.1     | M67449   | Glycine max             | protein kinase. PK6.   |
| AAG31141.1     | AF305911 | Oryza sativa            | EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.  |
| AAG31142.1     | AF305912 | Hordeum vulgare         | EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.  |
| CAC09580.1     | AJ298992 | Fagus sylvatica         | Abscisic acid (ABA) and calcium induced protein kinase. protein kinase (PK). pk1.  |
| CAA06334.1     | AJ005077 | Lycopersicon esculentum | protein kinase. TCTR2 protein. TCTR2.  |
| AAD10057.1     | AF110519 | Lycopersicon esculentum | ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1v.   |
| AAD10056.1     | AF110518 | Lycopersicon esculentum | ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1.  |
| AAD46406.1     | AF096250 | Lycopersicon esculentum | ethylene-responsive protein kinase TCTR1. ER50. serine/threonine kinase; similar to Arabidopsis thaliana negative regulator of the ethylene response pathway encoded by GenBank Accession Number L08789. |
| CAA73722.1     | Y13273   | Lycopersicon esculentum | putative protein kinase.   |
| AAK11734.1     | AY027437 | Arachis hypogaea        | serine/threonine/tyrosine kinase.  |
| AAK30005.1     | AY029067 | Rosa hybrid cultivar    | CTR2 protein kinase.   |
| AAF66615.1     | AF142596 | Nicotiana tabacum       | LRR receptor-like protein kinase.  |
| CAA73068.1     | Y12465   | Sorghum bicolor         | serine/threonine kinase. SNFL2.  |
| CAA61510.1     | X89226   | Oryza sativa            | leucine-rich repeat/receptor protein kinase. lrk2.   |
| BAA83688.1     | AB011967 | Oryza sativa            | OsPK4. OsPK4. protein kinase.  |
| AAF34436.1     | AF172282 | Oryza sativa            | similar to mitogen-activated protein kinases. DUPR11.32.   |
| AAF22219.1     | AF141378 | Zea mays                | protein kinase PK4. ZmPK4.   |

|            |          |                   |   |
|------------|----------|-------------------|---|
| BAA87852.1 | AP000816 | Oryza sativa      | Similar to putative Ser/Thr protein kinase. (AC004218).   |
| BAA92221.1 | AP001278 | Oryza sativa      | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).  |
| AAF91323.1 | AF244889 | Glycine max       | receptor-like protein kinase 2. RLK2. GmRLK2.   |
| BAB40094.1 | AP003210 | Oryza sativa      | putative receptor protein kinase. OSJNBa0010K01.7.  |
| AAF91324.1 | AF244890 | Glycine max       | receptor-like protein kinase 3. RLK3. GmRLK3.   |
| BAA34675.1 | AB011670 | Triticum aestivum | wpk4 protein kinase. wpk4.  |
| BAA92970.1 | AP001551 | Oryza sativa      | Similar to Arabidopsis thaliana chromosome 4 BAC clone F28A21 ; putative protein kinase. (AL035526).  |
| AAF91322.1 | AF244888 | Glycine max       | receptor-like protein kinase 1. RLK1. GmRLK1.   |
| BAB07999.1 | AP002525 | Oryza sativa      | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB03429.1 | AP002817 | Oryza sativa      | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587). |
| CAA73067.1 | Y12464   | Sorghum bicolor   | serine/threonine kinase. SNFL1.   |
| AAF59906.1 | AF197947 | Glycine max       | receptor protein kinase-like protein. CLV1B.  |
| AAF59905.1 | AF197946 | Glycine max       | receptor protein kinase-like protein. CLV1A.  |
| BAB16918.1 | AP002863 | Oryza sativa      | putative protein kinase. P0005A05.22.   |
| AAC36318.1 | AF053127 | Malus x domestica | leucine-rich receptor-like protein kinase. LRPKml.  |
| BAA83689.1 | AB011968 | Oryza sativa      | OsPK7. OsPK7. protein kinase.   |
| CAA74646.1 | Y14274   | Sorghum bicolor   | putative serine/threonine protein kinase. SNFL3.  |
| BAA95893.1 | AP002071 | Oryza sativa      | Similar to Arabidopsis thaliana wak4 gene; wall-associated kinase 4. (AJ009695).  |
| BAA92954.1 | AP001551 | Oryza sativa      | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).   |
| AAC83393.1 | U83625   | Zea mays          | protein kinase ZmMEK1. mitogen-activated; ERK-activating protein kinase (MEK) homolog.  |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAA92972.1     | AP001551 | Oryza sativa            | ESTs AU056183(S20356),AU056881(S20950) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 4 BAC clone F6I18 ; putative protein kinase. (AL022198). |
| BAB18292.1     | AP002860 | Oryza sativa            | putative receptor-like protein kinase. P0409B08.19.  |
| BAB40015.1     | AP003021 | Oryza sativa            | putative wall-associated kinase 1. P0503E05.18.  |
| BAA05648.1     | D26601   | Nicotiana tabacum       | protein kinase.  |
| AAC27489.1     | AF077130 | Oryza sativa            | receptor-like protein kinase.  |
| AAC02535.1     | AF044260 | Oryza sativa            | receptor serine/threonine kinase. protein kinase.  |
| AAG40578.1     | AF216314 | Oryza sativa            | MAP kinase kinase 1. protein kinase; MEK1.   |
| CAA08997.1     | AJ010093 | Brassica napus          | MAP3K beta 1 protein kinase. MAP3K beta 1.   |
| SEQ ID NO: 118 |          |                         |  |
| AAD21199.1     | AF127797 | Capsicum chinense       | putative bZIP DNA-binding protein.   |
| CAC00658.1     | AJ292745 | Petroselinum crispum    | bZIP type transcription factor. common plant regulatory factor 7. cprf7.   |
| CAC00657.1     | AJ292744 | Petroselinum crispum    | bZIP type transcription factor. common plant regulatory factor 6. cprf6.   |
| CAA74023.1     | Y13676   | Antirrhinum majus       | bZIP DNA-binding protein.  |
| CAA74022.1     | Y13675   | Antirrhinum majus       | bZIP DNA-binding protein.  |
| BAA22204.1     | D63951   | Nicotiana tabacum       | TBZ17. tbz17. bZIP protein.  |
| AAD55394.1     | AF176641 | Lycopersicon esculentum | bZIP DNA-binding protein.  |
| CAA44607.1     | X62745   | Zea mays                | ocs-binding factor 1. OBF1.  |
| AAK25822.1     | AF350505 | Phaseolus vulgaris      | bZip transcription factor.   |
| AAK01953.1     | AY026054 | Phaseolus acutifolius   | bZIP. transcription factor.  |
| CAA71687.1     | Y10685   | Glycine max             | bZIP DNA-binding protein. G/HBF-1. G/HBF-1.  |
| AAC37418.1     | L34551   | Oryza sativa            | transcriptional activator protein. RITA-1.   |

|                |          |                      |   |
|----------------|----------|----------------------|---|
| BAA36492.1     | AB021736 | Oryza sativa         | bZIP protein.   |
| BAA11431.1     | D78609   | Oryza sativa         | bZIP protein.   |
| CAA41453.1     | X58577   | Petroselinum crispum | DNA-binding protein; bZIP type. CPRF2.  |
| CAA71768.1     | Y10809   | Petroselinum crispum | bZIP DNA-binding protein. CPRF4a.   |
| AAD42938.1     | AF084972 | Catharanthus roseus  | G-Box binding protein 2. GBF2. basic leucine zipper; trans-regulatory factor.   |
| AAC49556.1     | U04295   | Oryza sativa         | DNA-binding factor of bZIP class. osZIP-1a.   |
| BAA07289.1     | D38111   | Triticum aestivum    | transcription factor HBP-1a(17).  |
| CAA71795.1     | Y10834   | Hordeum vulgare      | bZIP transcription factor 2. Blz2.  |
| CAA70216.1     | Y09013   | Triticum aestivum    | transcriptional activator. SPA.   |
| CAA40101.1     | X56781   | Triticum aestivum    | transcription factor. HBP-1a. la-17.  |
| AAA80169.1     | U10270   | Zea mays             | G-box binding factor 1. GBF1.   |
| CAA11499.1     | AJ223624 | Spinacia oleracea    | basic leucine zipper protein. bZIP.   |
| AAC49474.1     | U41817   | Phaseolus vulgaris   | regulator of MAT2. ROM2. Repressor of seed-specific lectin (phytohemagglutinin) gene; bZIP transcriptional repressor.   |
| AAB40291.1     | U42208   | Oryza sativa         | OSBZ8. G-box binding protein; GBF type bZIP protein.  |
| BAA02304.1     | D12920   | Triticum aestivum    | transcription factor HBP-1a(c14).   |
| SEQ ID NO: 119 |          |                      |   |
| BAA96200.1     | AP002093 | Oryza sativa         | EST D40368(S2303) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome II putative cytochrome P450 (AC004077).                             |
| BAA96158.1     | AP002092 | Oryza sativa         | EST D40368(S2303) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome II BAC T31E10; putative cytochrome P450 (AC004077).                 |
| BAA96196.1     | AP002093 | Oryza sativa         | ESTs AU086027(S2303),D40339(S2251) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome II BAC T31E10; putative cytochrome P450 (AC004077). |



|            |          |                         |   |
|------------|----------|-------------------------|---|
| BAA96154.1 | AP002092 | Oryza sativa            | ESTs AU086027(S2303),D40339(S2251) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome II BAC T31E10; putative cytochrome P450 (AC004077). |
| BAA96193.1 | AP002093 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC T31E10; putative cytochrome P450 (AC004077).  |
| BAA96151.1 | AP002092 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC T31E10; putative cytochrome P450 (AC004077).  |
| BAA96194.1 | AP002093 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC T31E10; putative cytochrome P450 (AC004077).  |
| BAA96152.1 | AP002092 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC T31E10; putative cytochrome P450 (AC004077).  |
| AAG41777.1 | AF212991 | Cucurbita maxima        | ent-kaurene oxidase. CYP88A2. cytochrome P450; similar to maize Dwarf3 protein.   |
| AAK11616.1 | AF326277 | Hordeum vulgare         | ent-kaurenoic acid oxidase. KAO1. cytochrome P450; CYP88A; HvKAO1.  |
| AAK00946.1 | AF318211 | Taxus cuspidata         | 5-alpha-taxadienol-10-beta-hydroxylase. cytochrome P450-like protein.   |
| AAC49659.1 | U74319   | Sorghum bicolor         | obtusifolios 14-alpha demethylase CYP51. CYP51. cytochrome P450 catalyzing the 14-alpha demethylation of obtusifolios in plants.  |
| AAA17746.1 | L19075   | Catharanthus roseus     | cytochrome P450. CYP72C. putative.  |
| AAA33106.1 | L10081   | Catharanthus roseus     | cytochrome P-450 protein. CYP72. putative; CYP72 protein.   |
| AAA17732.1 | L19074   | Catharanthus roseus     | cytochrome P450. CYP72B.  |
| CAB56503.1 | AJ238612 | Catharanthus roseus     | cytochrome P450.  |
| AAB17070.1 | U54770   | Lycopersicon esculentum | cytochrome P450 homolog. dwarf.   |
| CAB41490.1 | AJ238439 | Cicer arietinum         | cytochrome P450 monooxygenase. cyp81E3v2.   |
| BAB19089.1 | AP002744 | Oryza sativa            | putative cytochrome P450. P0006C01.31. contains ESTs D24685(R2374),AU031882(R2374).   |
| AAD44150.1 | AF124815 | Mentha spicata          | cytochrome p450.  |
| BAB19110.1 | AP002839 | Oryza sativa            | putative cytochrome P450. P0688A04.16. contains ESTs D24685(R2374),AU031882(R2374).   |

|                |          |   |
|----------------|----------|---|
| BAA74465.1     | AB022732 | Glycyrrhiza echinata<br>cytochrome P450. CYP Ge-31.   |
| BAA22422.1     | AB001379 | Glycyrrhiza echinata<br>cytochrome P450. CYP81E1.   |
| CAA10067.1     | AJ012581 | Cicer arietinum<br>cytochrome P450. cyp81E3.  |
| BAB40322.1     | AB036772 | Triticum aestivum<br>cytochrome P450. N-1.  |
| CAB43505.1     | AJ239051 | Cicer arietinum<br>cytochrome P450. cyp81E2.  |
| CAB56742.1     | AJ249800 | Cicer arietinum<br>cytochrome P450 monooxygenase. cyp81E5.  |
| AAF89209.1     | AF279252 | Vigna radiata<br>cytochrome P450. CipCYP.   |
| BAA93634.1     | AB025016 | Lotus japonicus<br>cytochrome P450.   |
| BAB19107.1     | AP002839 | Oryza sativa<br>putative cytochrome P450. P0688A04.13. contains ESTs<br>AU100635(C10787),D22354(C10787).                  |
| BAB19086.1     | AP002744 | Oryza sativa<br>putative cytochrome P450. P0006C01.28. contains ESTs<br>AU100635(C10787),D22354(C10787).                  |
| CAA04116.1     | AJ000477 | Helianthus tuberosus<br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.                                       |
| CAA04117.1     | AJ000478 | Helianthus tuberosus<br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-<br>race). |
| CAA71517.1     | Y10493   | Glycine max<br>putative cytochrome P450.  |
| BAB21156.1     | AP002899 | Oryza sativa<br>putative cytochrome P450. P0456A01.12.  |
| AAG13498.1     | AC068924 | Oryza sativa<br>putative cytochrome P450. OSJNBa0026L12.20.   |
| AAF34534.1     | AF195813 | Lupinus albus<br>isoflavone synthase 1. ifs1. cytochrome P450.  |
| CAA83941.1     | Z33875   | Mentha x piperita<br>cytochrome P-450 oxidase.  |
| AAF34533.1     | AF195812 | Pisum sativum<br>isoflavone synthase 1. ifs1. cytochrome P450.  |
| AAF45142.1     | AF195818 | Glycine max<br>isoflavone synthase 1. ifs1. cytochrome P450.  |
| SEQ ID NO: 120 |          |   |

---

CAB42052.1 AJ242045 *Lycopersicon esculentum*  
nicotianamine synthase. chln.

---

BAA74581.1 AB011264 *Hordeum vulgare*  
nicochianamine synthase 3. hvnas3.

---

AAD32651.1 AF136942 *Hordeum vulgare*  
nicotianamine synthase 2. nashor2.

---

BAA74582.1 AB011265 *Hordeum vulgare*  
nicochianamine synthase 2. hvnas2.

---

BAB17824.1 AB023819 *Oryza sativa*  
nicotianamine synthase 3. osnas3.

---

AAD32650.1 AF136941 *Hordeum vulgare*  
nicotianamine synthase 1. nashor1.

---

BAA74583.1 AB011266 *Hordeum vulgare*  
nicotianamine Synthase 4. hvnas4.

---

BAA74586.1 AB011269 *Hordeum vulgare*  
nicotianamine Synthase 6. hvnas6.

---

BAA74587.1 AB019525 *Hordeum vulgare*  
nicotianamine synthase 7. hvnas7.

---

BAB17826.1 AB046401 *Oryza sativa*  
nicotianamine synthase 2. OsNAS2.

---

BAB17823.1 AB023818 *Oryza sativa*  
nicotianamine synthase 2. osnas2.

---

BAB17825.1 AB046401 *Oryza sativa*  
nicotianamine synthase 1. OsNAS1.

---

BAA74588.2 AB021746 *Oryza sativa*  
nicotianamine synthase 1. osnas1.

---

BAA74580.1 AB010086 *Hordeum vulgare*  
nicotianamine synthase 1. hvnas1.

---

BAA74585.1 AB011268 *Hordeum vulgare*  
nicotianamine Synthase 5-2. hvnas5-2.

---

BAA74584.1 AB011267 *Hordeum vulgare*  
nicotianamine synthase 5-1. hvnas5-1.

---

SEQ ID NO: 122

---

AAD01804.1 AF026480 *Dianthus caryophyllus*  
lipase. lipid-protein-particle associated.

---

BAB39417.1 AP002901 *Oryza sativa*  
putative lipase. P0456F08.17. contains ESTs  
C99390(E11001),AU101109(E0858),AU101332(E11001).

---

AAB07724.1 U55867 *Ipomoea nil*  
Pn47p. lipase-like protein.

---

AAK31273.1 AC079890 *Oryza sativa*  
putative lipase. OSJNBb0089A17.13.

---

SEQ ID NO: 123

---

|               |  |                   |
|---------------|--|-------------------|
| BAA21923.1    | AB006601   | Petunia x hybrida |
| ZPT2-14.      | C2H2 zinc finger protein, 2 finger.                                    |                   |
| BAA21922.1    | AB006600   | Petunia x hybrida |
| ZPT2-13.      | C2H2 zinc finger protein, 2finger.                                     |                   |
| BAA21921.1    | AB006599   | Petunia x hybrida |
| ZPT2-12.      | C2H2 zinc finger protein, 2 finger.                                    |                   |
| BAA19110.1    | AB000451   | Petunia x hybrida |
| PETHy;ZPT2-5. | Cys(2) His(2) zinc finger protein, 2 fingers.                          |                   |
| BAA21926.1    | AB006604   | Petunia x hybrida |
| ZPT2-9.       | C2H2 zinc finger protein, 2 finger.                                    |                   |
| BAA21925.1    | AB006603   | Petunia x hybrida |
| ZPT2-8.       | C2H2 zinc finger protein, 2 finger.                                    |                   |
| BAA21924.1    | AB006602   | Petunia x hybrida |
| ZPT2-7.       | C2H2 zinc finger protein, 2finger.                                     |                   |
| BAA21920.1    | AB006598   | Petunia x hybrida |
| ZPT2-11.      | C2H2 zinc finger protein, 2finger.                                     |                   |
| CAA60828.1    | X87374   | Pisum sativum     |
|               | putative zinc finger protein.  |                   |
| BAA19111.1    | AB000452   | Petunia x hybrida |
| PETHy;ZPT2-6. | Cys(2) His(2) zinc finger protein, 2 fingers.                          |                   |
| CAB77055.1    | Y18788   | Medicago sativa   |
|               | putative TFIIIA (or kruppel)-like zinc finger protein.                 |                   |
| BAA96071.1    | AB035133   | Petunia x hybrida |
|               | C2H2 zinc-finger protein ZPT3-3. ZPT3-3.                               |                   |
| BAA21927.1    | AB006605   | Petunia x hybrida |
| ZPT3-3.       | C2H2 zinc finger protein, 3 finger.                                    |                   |
| BAA96070.1    | AB035132   | Petunia x hybrida |
|               | C2H2 zinc-finger protein ZPT2-10. PETHy;ZPT2-10.                       |                   |
| BAA21919.1    | AB006597   | Petunia x hybrida |
| ZPT2-10.      | C2H2 zinc finger protein, 2 finger.                                    |                   |
| AAK01713.1    | AF332876   | Oryza sativa      |
|               | zinc finger transcription factor ZF1.                                  |                   |
| AAC06243.1    | AF053077   | Nicotiana tabacum |
|               | transcription factor. osmotic stress-induced zinc-finger protein. zfp. |                   |
| BAA05079.1    | D26086   | Petunia x hybrida |
|               | zinc-finger protein.   |                   |
| BAA05078.1    | D26085   | Petunia x hybrida |
|               | zinc-finger DNA binding protein.                                       |                   |
| AAB39638.1    | U68763   | Glycine max       |
|               | putative transcription factor. SCOF-1. scof-1. zinc-finger protein.    |                   |
| BAA05076.1    | D26083   | Petunia x hybrida |
|               | zinc-finger DNA binding protein.                                       |                   |

|                |          |                                |  |
|----------------|----------|--------------------------------|--|
| BAA05077.1     | D26084   | Petunia x hybrida              | zinc-finger DNA binding protein.   |
| BAA21928.1     | AB006606 | Petunia x hybrida              | ZPT4-4. C2H2 zinc finger protein, 4 finger.  |
| BAA19114.1     | AB000455 | Petunia x hybrida              | PETHy;ZPT4-1. Cys(2) His(2) zinc finger protein, 4 fingers.  |
| AAD26942.1     | AF119050 | Datisca glomerata              | zinc-finger protein 1. zfp1. DgZFP1.   |
| BAA19112.1     | AB000453 | Petunia x hybrida              | PETHy;ZPT3-1. Cys(2) His(2) zinc finger protein, 3 fingers.  |
| AAB53260.1     | U76554   | Brassica rapa                  | transcription factor. zinc-finger protein-1. BR42.   |
| AAB53261.1     | U76555   | Brassica rapa                  | zinc-finger protein BcZFP1. BcZFP1(3-2z).  |
| BAA19926.1     | AB000456 | Petunia x hybrida              | PETHy; ZPT4-2. C2H2 zinc finger protein, 4 finger.   |
| SEQ ID NO: 126 |          |                                |  |
| CAA10134.1     | AJ012693 | Cicer arietinum                | basic blue copper protein.   |
| CAB65280.1     | AJ248323 | Medicago sativa subsp. x varia | basic blue protein. babl.  |
| AAC32448.1     | U76296   | Spinacia oleracea              | plantacyanin. member of a strictly plant-specific family of mononuclear blue copper proteins known as phytocyanins, which are further classified into three distinct subfamilies: Uclacyanins, Stellacyanins, and Plantacyanins; member of the subfamily of Plantacyanins; non-glycosylated. |
| AAF66243.1     | AF243181 | Lycopersicon esculentum        | plantacyanin. naturally occurring cupredoxin with a Val residue in the position of the axial ligand Met; member of the plantacyanin subfamily of a strictly plant-specific family of mononuclear blue copper proteins known as phytocyanins.   |
| AAD10251.1     | AF031195 | Triticum aestivum              | blue copper-binding protein homolog. S85.  |
| AAF66242.1     | AF243180 | Lycopersicon esculentum        | dicyanin. binuclear blue copper protein; contains two stellacyanins linked together.   |
| CAA80963.1     | Z25471   | Pisum sativum                  | blue copper protein.   |
| AAC64163.1     | AF093537 | Zea mays                       | blue copper protein. similar to pea blue copper protein in GenBank Accession Number Z25471.  |

AAC32421.1 U65511 *Cucumis sativus*  
 putative oxygen activation and/or lignin formation. stellacyanin. member of a strictly plant-specific family of mononuclear blue copper proteins known as phytoeyanins, which are further classified into three distinct subfamilies: Uclacyanins, Stellacyanins, and Plantacyanins; member of the subfamily of Stellacyanins; similar to *Rhus vernicifera* stellacyanin: SwissProt Accession Number P00302; similar to umecyanin: SwissProt Accession Number P42849; similar to mavicyanin; SwissProt Accession Number P80728; similar to BCB encoded by GenBank Accession Number Z15058; similar to F18A8.9, encoded by GenBank Accession Number AC003105; similar to F7F1.27 encoded by GenBank Accession Number AC004669; similar to F9D12.16 encoded by GenBank Accession Number AF077407; glycoprotein.

## SEQ ID NO: 127

AAA17000.1 L08632 *Glycine max*  
 pyruvate kinase.

CAA37727.1 X53688 *Solanum tuberosum*  
 pyruvate kinase.

CAA82628.1 Z29492 *Nicotiana tabacum*  
 glycolytic enzyme. pyruvate kinase.

AAF44707.1 AF242871 *Lilium longiflorum*  
 cytosolic pyruvate kinase.

BAA88185.1 AP000836 *Oryza sativa*  
 ESTs AU081247(C50004), AU068940(C51113) correspond to a region of the predicted gene. Similar to pyruvate kinase (Q42954).

BAA76433.1 AB025005 *Cicer arietinum*  
 pyruvate kinase.

CAA49996.1 X70653 *Nicotiana tabacum*  
 pyruvate kinase. PKTL7. monomer.

CAA82223.1 Z28374 *Nicotiana tabacum*  
 glycolytic enzyme. Pyruvate kinase; plastid isozyme.

CAA82222.1 Z28373 *Nicotiana tabacum*  
 Glycolytic enzyme. pyruvate kinase; plastid isozyme.

AAA33871.1 M64737 *Ricinus communis*  
 ATP:pyruvate phosphotransferase.

AAA33870.1 M64736 *Ricinus communis*  
 ATP:pyruvate phosphotransferase.

## SEQ ID NO: 129

BAB12433.1 AB025030 *Coptis japonica*  
 p450.

AAF05621.1 AF191772 *Papaver somniferum*  
 hydroxylase involved in benzylisoquinoline alkaloid biosynthesis. (S)-N-methylcoclaurine 3'-hydroxylase. CYP80B1. cytochrome P450-dependent monooxygenase.

|            |          |   |
|------------|----------|---|
| AAC39452.1 | AF014800 | <i>Eschscholzia californica</i><br>hydroxylase involved in the biosynthesis of tetrahydrobenzylisoquinoline alkaloids in plants.<br>(S)-N-methylcoclaaurine 3'-hydroxylase. CYP80B1. cytochrome P-450-dependent<br>monooxygenase; CYP80B1v1.  |
| AAC39453.1 | AF014801 | <i>Eschscholzia californica</i><br>hydroxylase involved in the biosynthesis of tetrahydrobenzylisoquinoline alkaloids in plants.<br>(S)-N-methylcoclaaurine 3'-hydroxylase. CYP80B1. cytochrome P-450-dependent<br>monooxygenase; CYP80B1v2.  |
| AAA32913.1 | M32885   | <i>Persea americana</i><br>cytochrome P-450LXXIA1 (cyp71A1).  |
| AAA19701.1 | L24438   | <i>Thlaspi arvense</i><br>cytochrome P450.  |
| CAA50648.1 | X71657   | <i>Solanum melongena</i><br>P450 hydroxylase.   |
| AAG34695.1 | AF313492 | <i>Matthiola incana</i><br>putative cytochrome P450.  |
| AAG44132.1 | AF218296 | <i>Pisum sativum</i><br>cytochrome P450. P450 isolog.   |
| BAA12159.1 | D83968   | <i>Glycine max</i><br>Cytochrome P-450 (CYP93A1).   |
| AAC48987.1 | U09610   | <i>Berberis stolonifera</i><br>cytochrome P-450 CYP80.  |
| AAD56282.1 | AF155332 | <i>Petunia x hybrida</i><br>flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.  |
| AAC32274.1 | AF081575 | <i>Petunia x hybrida</i><br>flavonoid 3',5'-hydroxylase. Hf1. P450 enzyme.  |
| AAB17562.1 | U72654   | <i>Eustoma grandiflorum</i><br>flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.  |
| BAA84916.1 | AB032833 | <i>Cicer arietinum</i><br>cytochrome P450. CYP76D1.   |
| AAB94588.1 | AF022459 | <i>Glycine max</i><br>CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.   |
| BAB40324.1 | AB037245 | <i>Asparagus officinalis</i><br>cytochrome P450. ASPI-2.  |
| BAB40323.1 | AB037244 | <i>Asparagus officinalis</i><br>cytochrome P450. ASPI-1.  |
| CAA50155.1 | X70824   | <i>Solanum melongena</i><br>flavonoid hydroxylase (P450). CYP75.  |
| AAC39318.1 | AF029858 | <i>Sorghum bicolor</i><br>second multifunctional cytochrome P450 in the biosynthetic pathway of the cyanogenic<br>glucoside dhurrin. Catalyzes the conversion of p-hydroxyphenylacetaldoxime to p-<br>hydroxymandelonitrile. cytochrome P450 CYP71E1. CYP71E1. No EST#s identified. |

CAB56503.1 AJ238612 *Catharanthus roseus*  
cytochrome P450.

CAA71514.1 Y10490 *Glycine max*  
putative cytochrome P450.

BAA84071.1 AB028151 *Antirrhinum majus*  
flavone synthase II. cytochrome P450. AFNS2.

CAA70575.1 Y09423 *Nepeta racemosa*  
cytochrome P450. CYP71A5.

CAA70576.1 Y09424 *Nepeta racemosa*  
cytochrome P450. CYP71A6.

CAA71513.1 Y10489 *Glycine max*  
putative cytochrome P450.

BAA84072.1 AB028152 *Torenia hybrida*  
flavone synthase II. cytochrome P450. TFNS5.

CAA71516.1 Y10492 *Glycine max*  
putative cytochrome P450.

BAA13076.1 D86351 *Glycine max*  
cytochrome P-450 (CYP93A2).

CAA71517.1 Y10493 *Glycine max*  
putative cytochrome P450.

SEQ ID NO: 132

CAB55396.1 AL117264 *Oryza sativa*  
zwh12.1. similar to Arabidopsis putative UDP-galactase-4-epimerase (AC007060); Method:  
conceptual translation with partial peptide sequencing.

SEQ ID NO: 133

AAG43835.1 AF213455 *Zea mays*  
protein phosphatase type-2C. pp2c-1. PP2C-1.

AAG13599.1 AC051633 *Oryza sativa*  
putative protein phosphatase-2C. OSJNBb0015I11.26.

AAG46118.1 AC073166 *Oryza sativa*  
putative protein phosphatase-2C. OSJNBb0064P21.12. tRNA-Lys.

BAB12036.1 AP002820 *Oryza sativa*  
putative protein phosphatase. P0702D12.18.

AAC36698.1 AF075580 *Mesembryanthemum crystallinum*  
protein phosphatase-2C. PP2C.

CAC10359.1 AJ277087 *Nicotiana tabacum*  
protein phosphatase 2C. PP2C2.

CAB90633.1 AJ277743 *Fagus sylvatica*  
protein phosphatase 2C (PP2C). pp2C1. ABA-induced protein.

CAC10358.1 AJ277086 *Nicotiana tabacum*  
protein phosphatase 2C. PP2C1.

CAA72341.1 Y11607 *Medicago sativa*  
protein phosphatase 2C. MP2C.



|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAC36697.1     | AF075579 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| CAB61839.1     | AJ242803 | Sporobolus stapfianus         | putative serine/threonine phosphatase type 2c.  |
| AAC36700.1     | AF075582 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| AAD17804.1     | AF092431 | Lotus japonicus               | nodule-enhanced protein phosphatase type 2C. NPP2C1.  |
| AAD17805.1     | AF092432 | Lotus japonicus               | protein phosphatase type 2C. PP2C2.   |
| CAC09575.1     | AJ298987 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2Cf1.  |
| AAC36699.1     | AF075581 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| AAD11430.1     | AF097667 | Mesembryanthemum crystallinum | protein phosphatase 2C homolog. PP2C.   |
| CAB90634.1     | AJ277744 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2C2. ABA and calcium induced protein.  |
| AAC26828.1     | AF075603 | Oryza sativa                  | kinase interaction domain mediates phosphorylation-dependent binding to a subset of active receptor-like protein kinases. kinase associated protein phosphatase. kapp. type 2C protein phosphatase. |
| AAC35951.1     | AF079355 | Mesembryanthemum crystallinum | protein phosphatase-2c. PP2C.   |
| AAK20060.1     | AC025783 | Oryza sativa                  | putative protein phosphatase 2C. OSJNBa0001O14.1.   |
| AAB93832.1     | U81960   | Zea mays                      | kinase interaction domain mediates phosphorylation-dependent binding to a subset of active receptor-like protein kinases; type 2C protein phosphatase. kinase associated protein phosphatase. KAPP. |
| CAC09576.1     | AJ298988 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2Cf2.  |
| SEQ ID NO: 134 |          |                               |   |
| AAD26116.1     | AF106954 | Brassica napus                | galactinol synthase.-GS. UDP-D-galactose:myo-inositol-D- galactosyltransferase.   |
| CAB51130.1     | AJ243815 | Pisum sativum                 | role in alpha galactoside synthesis. putative galactinol synthase.  |
| CAB51533.1     | AJ237693 | Ajuga reptans                 | galactosyl transfer from UDP-galactose to myo-inositol to form galactinol. galactinol synthase, isoform GolS-1. GolS.   |
| CAB51534.1     | AJ237694 | Ajuga reptans                 | galactosyl transfer from UDP-galactose to myo-inositol. galactinol synthase, isoform GolS-2. GolS.  |

|   |          |                                |
|---|----------|--------------------------------|
| AAD55726.1  | AF178569 | <i>Vitis riparia</i>           |
| galactinol synthase. WSI76. water stress induced protein.   |          |                                |
| SEQ ID NO: 135  |          |                                |
| AAB57734.1  | U64818   | <i>Lycopersicon esculentum</i> |
| fructokinase. Frk2.   |          |                                |
| AAB51108.1  | U62329   | <i>Lycopersicon esculentum</i> |
| fructokinase. FK.   |          |                                |
| AAA80675.1  | U37838   | <i>Beta vulgaris</i>           |
| fructokinase.   |          |                                |
| CAA78283.1  | Z12823   | <i>Solanum tuberosum</i>       |
| fructokinase.   |          |                                |
| AAB57733.1  | U64817   | <i>Lycopersicon esculentum</i> |
| fructokinase. Frk1.   |          |                                |
| SEQ ID NO: 138  |          |                                |
| BAA94601.1  | AB033504 | <i>Populus euramericana</i>    |
| 1-aminocyclopropane-1-carboxylate oxidase. peaco-1.   |          |                                |
| AAA33697.1  | L21978   | <i>Petunia x hybrida</i>       |
| 1-aminocyclopropane-1-carboxylate oxidase. ACO3.  |          |                                |
| AAC48977.1  | U07953   | <i>Pelargonium x hortorum</i>  |
| 1-aminocyclopropane-1-carboxylate oxidase.  |          |                                |
| CAA54449.1  | X77232   | <i>Prunus persica</i>          |
| 1-aminocyclopropane-1-carboxylate oxidase. PAO1.  |          |                                |
| AAF36483.1  | AF129073 | <i>Prunus persica</i>          |
| 1-aminocyclopropane-1-carboxylate oxidase. ACO1.  |          |                                |
| AAC33524.1  | AF026793 | <i>Prunus armeniaca</i>        |
| 1-aminocyclopropane-1-carboxylate oxidase. ACC oxidase.   |          |                                |
| AAB70884.1  | U67861   | <i>Pelargonium x hortorum</i>  |
| 1-aminocyclopropane-1-carboxylate oxidase. GACO3.   |          |                                |
| AAG49361.1  | AF321533 | <i>Citrus sinensis</i>         |
| ACC oxidase.  |          |                                |
| BAA90550.1  | AB031027 | <i>Prunus mume</i>             |
| ACC oxidase. PM-ACO1. 1-aminocyclopropane-1-carboxylic acid oxidase.  |          |                                |
| AAA99792.1  | U54565   | <i>Nicotiana glutinosa</i>     |
| oxidation of 1-aminocyclopropane-1-carboxylic acid. 1-aminocyclopropane-1-carboxylic acid oxidase. NGACO1. ACC oxidase. |          |                                |
| AAB05171.1  | U62764   | <i>Nicotiana glutinosa</i>     |
| oxidation of 1-aminocyclopropane-1-carboxylic acid. ACC oxidase. NGACO3.  |          |                                |
| AAC37381.1  | L21976   | <i>Petunia x hybrida</i>       |
| 1-aminocyclopropane-1-carboxylate oxidase. ACO1.  |          |                                |
| CAA71738.1  | Y10749   | <i>Betula pendula</i>          |
| 1-aminocyclopropane-1-carboxylate oxidase. ACO.   |          |                                |
| CAA86468.1  | Z46349   | <i>Nicotiana tabacum</i>       |
| 1-aminocyclopropane-1-carboxylate deaminase.  |          |                                |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| BAA83466.1 | AB012857 | Nicotiana tabacum       | ACC oxidase.  |
| AAC98808.1 | U68215   | Carica papaya           | ACC oxidase. fruit specific; ripening related.  |
| BAA06526.1 | D31727   | Cucumis melo            | 1-aminocyclopropane-1-carboxylate oxidase.  |
| CAA64797.1 | X95551   | Cucumis melo            | ACC oxidase.  |
| CAA58232.1 | X83229   | Nicotiana tabacum       | ethylene forming enzyme. 1-aminocyclopropane-1-carboxylate oxidase.   |
| BAA34924.1 | AB013101 | Lycopersicon esculentum | 1-aminocyclopropane-1-carboxylate oxidase. LE-ACO4. ACC oxidase.  |
| AAF64528.1 | AF254125 | Carica papaya           | 1-aminocyclopropane-1-carboxylate oxidase. ACC oxidase.   |
| AAA33698.1 | L21979   | Petunia x hybrida       | 1-aminocyclopropane-1-carboxylate oxidase. ACO4.  |
| CAA41212.1 | X58273   | Lycopersicon esculentum | conversion of ACC to ethylene. 1-Aminocyclopropane-1-carboxylic acid oxidase. LEACO1.                                   |
| BAA21541.1 | AB003514 | Actinidia deliciosa     | 1-aminocyclopropane-1-carboxylic acid oxidase.  |
| AAB71421.1 | L29405   | Helianthus annuus       | 1-aminocyclopropapne-1-carboxylic acid oxidase. ACC oxidase.  |
| AAA99793.1 | U54566   | Nicotiana glutinosa     | oxidation of 1-aminocyclopropane-1-carboxylic acid. 1-aminocyclopropane-1-carboxylic acid oxidase. NGACO2. ACC oxidase. |
| AAF36484.1 | AF129074 | Prunus persica          | 1-aminocyclopropane-1-carboxylate oxidase. ACO2.  |
| CAA64799.1 | X95553   | Cucumis melo            | ACC oxidase.  |
| CAA90904.1 | Z54199   | Lycopersicon esculentum | catalyses the final step in ethylene biosynthesis. 1-aminocyclopropane-1-carboxylic acid oxidase. ACO3.                 |
| CAA68538.1 | Y00478   | Lycopersicon esculentum | conversion of ACC to ethylene. 1-aminocyclopropane-1-carboxylate oxidase. LEACO2.                                       |
| CAB97173.1 | AJ297435 | Mangifera indica        | ethene biosynthesis. putative 1-aminocyclopropane-1-carboxylic acid oxidase. aco1.                                      |
| CAA82646.1 | Z29529   | Nicotiana tabacum       | oxidation of 1-aminocyclopropane-1-carboxylic acid. ethylene forming enzyme (EFE).                                      |
| AAC12934.1 | AF053354 | Phaseolus vulgaris      | 1-aminocyclopropane-1-carboxylic acid oxidase. ACO1. ACC oxidase.   |
| AAB70883.1 | U19856   | Pelargonium x hortorum  | 1-aminocyclopropane-1-carboxylate oxidase.  |

|                |          |                       |   |
|----------------|----------|-----------------------|---|
| AAC67233.1     | AF033582 | Cucumis sativus       | ACC oxidase 2. Cs-ACO2.   |
| AAB02051.1     | L76283   | Carica papaya         | formation of ethylene. 1-aminocyclopropane-1-carboxylate oxidase. putative. |
| BAA33377.1     | AB006806 | Cucumis sativus       | ACC oxidase. CS-ACO1.   |
| BAA33378.1     | AB006807 | Cucumis sativus       | ACC oxidase. CS-ACO2.   |
| CAA71140.1     | Y10034   | Rumex palustris       | 1-aminocyclopropane-1-carboxylic acid oxidase.                              |
| AAA33644.1     | M98357   | Pisum sativum         | convert ACC to ethylene. 1-aminocyclopropane-1-carboxylate oxidase.         |
| AAC48921.1     | U06046   | Vigna radiata         | 1-aminocyclopropane-1-carboxylate oxidase homolog.                          |
| AAK07883.1     | AF315316 | Vigna radiata         | ACC oxidase. ACO.   |
| AAA33273.1     | L35152   | Dianthus caryophyllus | amino-cyclopropane carboxylic acid oxidase.                                 |
| CAA74328.1     | Y14005   | Malus x domestica     | Converts ACC into ethylene in apple fruit. ACC oxidase.                     |
| SEQ ID NO: 139 |          |                       |   |
| AAB65777.1     | U97522   | Vitis vinifera        | class IV endochitinase. VvChi4B.  |
| AAB65776.1     | U97521   | Vitis vinifera        | class IV endochitinase. VvChi4A.  |
| CAC17793.1     | AJ301671 | Nicotiana sylvestris  | hydrolysis of chitin. endochitinase. chnb. class I chitinase.               |
| AAA34070.1     | M15173   | Nicotiana tabacum     | endochitinase precursor (EC 3.2.1.14).                                      |
| CAA30142.1     | X07130   | Solanum tuberosum     | endochitinase.  |
| CAA53626.1     | X76041   | Triticum aestivum     | endochitinase. CHI.   |
| SEQ ID NO: 140 |          |                       |   |
| AAB94587.1     | AF022458 | Glycine max           | CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.                           |
| BAA92894.1     | AB006790 | Petunia x hybrida     | cytochrome P450. IMT-2.   |
| AAD56282.1     | AF155332 | Petunia x hybrida     | flavonoid 3'-hydroxylase. Htl. cytochrome P450; CYP75B2.                    |
| AAG44132.1     | AF218296 | Pisum sativum         | cytochrome P450. P450 isolog.   |

|            |          |  |
|------------|----------|--|
| CAA65580.1 | X96784   | Nicotiana tabacum<br>cytochrome P450. hsr515.  |
| AAA32913.1 | M32885   | Persea americana<br>cytochrome P-450LXXIA1 (cyp71A1).  |
| CAA64635.1 | X95342   | Nicotiana tabacum<br>cytochrome P450. hsr515. hypersensitivity-related gene.   |
| CAA70575.1 | Y09423   | Nepeta racemosa<br>cytochrome P450. CYP71A5.   |
| BAB40323.1 | AB037244 | Asparagus officinalis<br>cytochrome P450. ASPI-1.  |
| BAB40324.1 | AB037245 | Asparagus officinalis<br>cytochrome P450. ASPI-2.  |
| CAA50312.1 | X70981   | Solanum melongena<br>P450 hydroxylase. CYPEG2.   |
| CAA50155.1 | X70824   | Solanum melongena<br>flavonoid hydroxylase (P450). CYP75.  |
| AAB17562.1 | U72654   | Eustoma grandiflorum<br>flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.  |
| BAA84071.1 | AB028151 | Antirrhinum majus<br>flavone synthase II. cytochrome P450. AFNS2.  |
| AAC32274.1 | AF081575 | Petunia x hybrida<br>flavonoid 3',5'-hydroxylase. Hfl. P450 enzyme.  |
| AAG34695.1 | AF313492 | Matthiola incana<br>putative cytochrome P450.  |
| AAB94588.1 | AF022459 | Glycine max<br>CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.   |
| AAD37433.1 | AF150881 | Lycopersicon esculentum x Lycopersicon peruvianum<br>catalyzes the hydroxylation of ferulic acid to 5-hydroxyferulic acid. ferulate-5-hydroxylase.<br>CYP84. cytochrome P450-dependent monooxygenase; F5H; FAH1. |
| CAA71517.1 | Y10493   | Glycine max<br>putative cytochrome P450.   |
| BAA13414.1 | D87520   | Glycyrrhiza echinata<br>putative trans-cinnamic acid 4-hydroxylase. cytochrome P450 (CYP73A14). CYP Ge-1.  |
| AAA19701.1 | L24438   | Thlaspi arvense<br>cytochrome P450.  |
| CAA50645.1 | X71654   | Solanum melongena<br>P450 hydroxylase.   |
| BAA03635.1 | D14990   | Solanum melongena<br>Cytochrome P-450EG4.  |
| BAA93634.1 | AB025016 | Lotus japonicus<br>cytochrome P450.  |
| AAC05148.1 | AF049067 | Pinus radiata<br>cytochrome P450. PRE74.   |

|                |          |                                   |   |
|----------------|----------|-----------------------------------|---|
| CAA70576.1     | Y09424   | Nepeta racemosa                   | cytochrome P450. CYP71A6.   |
| AAG10196.1     | AF286647 | Gossypium arboreum                | cinnamate-4-hydroxylase. LP89. P450.  |
| AAB94584.1     | AF022157 | Glycine max                       | capable of catalyzing the metabolism of phenylurea herbicides. CYP71A10. CYP71A10. cytochrome P450 monooxygenase. |
| CAA50648.1     | X71657   | Solanum melongena                 | P450 hydroxylase.   |
| CAB43505.1     | AJ239051 | Cicer arietinum                   | cytochrome P450. cyp81E2.   |
| SEQ ID NO: 141 |          |                                   |   |
| AAB97167.1     | AF030882 | Zea mays                          | SU1 isoamylase. sugary1. starch debranching enzyme.   |
| AAA91298.1     | U18908   | Zea mays                          | Sulp. Sugary1. similar to Pseudomonas sp. isoamylase, Swiss-Prot Accession Number P26501.                         |
| AAD33889.1     | AF142589 | Hordeum vulgare                   | isoamylase 1.   |
| BAA29041.1     | AB015615 | Oryza sativa                      | isoamylase.   |
| AAD33891.1     | AF142591 | Solanum tuberosum                 | isoamylase 1.   |
| AAD33890.1     | AF142590 | Triticum aestivum                 | isoamylase 1.   |
| AAD53260.1     | AF142588 | Hordeum vulgare                   | isoamylase 1.   |
| SEQ ID NO: 145 |          |                                   |   |
| AAG35777.1     | AF273844 | Brassica oleracea var. alboglabra | thioredoxin-h-like protein 1. THL1.   |
| AAB53694.1     | U59379   | Brassica napus                    | thioredoxin-h-like-1. THL-1. thioredoxin-h homolog.   |
| CAA61908.1     | X89759   | Brassica oleracea                 | pollen coat protein. bopcl7.  |
| BAA25681.1     | AB010434 | Brassica rapa                     | Thioredoxin. PEC-2.   |
| BAB20886.1     | AB053294 | Oryza sativa                      | thioredoxin h. RTRXH2.  |
| AAB53695.1     | U59380   | Brassica napus                    | thioredoxin-h-like-2. THL-2. thioredoxin-h homolog.   |
| AAF88067.1     | AF286593 | Triticum aestivum                 | thioredoxin H. similar to wheat thioredoxin H.  |

|            |          |   |
|------------|----------|---|
| CAA94534.1 | Z70677   | Ricinus communis<br>thioredoxin.  |
| CAA05081.1 | AJ001903 | Triticum turgidum subsp. durum<br>thioredoxin H.  |
| CAA49540.1 | X69915   | Triticum aestivum<br>unnamed protein product.   |
| BAA13524.1 | D87984   | Fagopyrum esculentum<br>thioredoxin.  |
| CAA41415.1 | X58527   | Nicotiana tabacum<br>thioredoxin.   |
| CAA77847.1 | Z11803   | Nicotiana tabacum<br>THIOREDOXIN.   |
| AAC32111.1 | AF051206 | Picea mariana<br>probable thioredoxin H. Sb09. similar to Nicotiana tabacum thioredoxin H1 encoded by<br>GenBank Accession Number X58527. |
| BAA05546.1 | D26547   | Oryza sativa<br>rice thioredoxin h.   |
| BAA04864.1 | D21836   | Oryza sativa<br>thioredoxin h. encoding rice phloem sap 13kD protein-1.   |
| AAB51522.1 | U92541   | Oryza sativa<br>thioredoxin h.  |
| AAD49232.1 | AF159387 | Lolium perenne<br>thioredoxin-like protein. Trx.  |
| AAD56954.1 | AF186240 | Secale cereale<br>thioredoxin-like protein. Trx.  |
| AAD49231.1 | AF159386 | Secale cereale<br>thioredoxin-like protein. Trx.  |
| AAD49230.1 | AF159385 | Hordeum bulbosum<br>thioredoxin-like protein. Trx.  |
| AAD49234.1 | AF159389 | Phalaris coerulescens<br>thioredoxin-like protein. Trx.   |
| AAD49233.1 | AF159388 | Phalaris coerulescens<br>thioredoxin-like protein. Trx.   |
| BAB39913.1 | AP002912 | Oryza sativa<br>thioredoxin-like protein. P0028E10.17. contains EST C72705(E2091).  |
| CAA55399.1 | X78822   | Chlamydomonas reinhardtii<br>thioredoxin h. Trx.  |
| CAA56850.1 | X80887   | Chlamydomonas reinhardtii<br>thioredoxin h. Trx h.  |
| AAD33596.1 | AF133127 | Hevea brasiliensis<br>thioredoxin h.  |
| CAA35826.1 | X51462   | Spinacia oleracea<br>thioredoxin M precursor (AA -67 to 114).   |

|            |          |  |
|------------|----------|--|
| CAA35827.1 | X51463   | Spinacia oleracea<br>thioredoxin M precursor (AA -67 to 114).                              |
| CAA45098.1 | X63537   | Pisum sativum<br>thioredoxin F. isoform.   |
| AAC49357.1 | U35830   | Pisum sativum<br>thioredoxin f.  |
| AAC19392.1 | AF069314 | Mesembryanthemum crystallinum<br>thioredoxin F precursor.                                  |
| AAC04671.1 | AF018174 | Brassica napus<br>thioredoxin-f. TRXF.   |
| AAB47556.1 | U87141   | Mesembryanthemum crystallinum<br>thioredoxin h.  |
| CAA53900.1 | X76269   | Pisum sativum<br>thioredoxin m.  |
| AAC49358.1 | U35831   | Pisum sativum<br>thioredoxin m. chloroplastic.   |
| CAA33082.1 | X14959   | Spinacia oleracea<br>pre-thioredoxin f (AA -77 to 113).                                    |
| CAA06736.1 | AJ005841 | Oryza sativa<br>thioredoxin M.   |
| CAA55398.1 | X78821   | Chlamydomonas reinhardtii<br>thioredoxin m. Trx.   |
| CAA56851.1 | X80888   | Chlamydomonas reinhardtii<br>thioredoxin m. Trx m.   |
| CAA44209.1 | X62335   | Chlamydomonas reinhardtii<br>thioredoxin Ch2. Trx.   |
| AAA92464.1 | L40957   | Zea mays<br>regulation of activities of photosynthetic enzymes. thioredoxin M. putative.   |
| CAA06735.1 | AJ005840 | Triticum aestivum<br>thioredoxin M.  |
| AAB52409.1 | U76831   | Brassica napus<br>thioredoxin-m.   |
| AAD45358.1 | AF160870 | Brassica napus<br>thioredoxin-m precursor.   |
| CAA71103.1 | Y09987   | Solanum tuberosum<br>CDSP32 protein (Chloroplast Drought-induced Stress Protein of 32kDa). |
| AAA32662.1 | M82973   | Medicago sativa<br>putative endomembrane protein; putative.                                |
| CAA77575.1 | Z11499   | Medicago sativa<br>protein disulfide isomerase.  |

SEQ ID NO: 148



|                |          |                           |   |
|----------------|----------|---------------------------|---|
| AAA33376.1     | L36129   | Helianthus annuus         | NADPH thioredoxin reductase.  |
| SEQ ID NO: 150 |          |                           |   |
| AAA92013.1     | U49454   | Prunus persica            | beta-1,3-glucanase. Gns1.   |
| CAA54952.1     | X77990   | Brassica rapa             | beta-1,3-glucanase. bgl.  |
| AAF33405.1     | AF230109 | Populus x canescens       | beta-1,3 glucanase. BGLUC.  |
| AAA33946.1     | M37753   | Glycine max               | beta-1,3-endoglucanase (EC 3.2.1.39).   |
| CAA03908.1     | AJ000081 | Citrus sinensis           | glucan hydrolase. beta-1,3-glucanase. gns1.   |
| CAB91554.1     | AJ277900 | Vitis vinifera            | beta 1-3 glucanase. gl.   |
| AAB03501.1     | U41323   | Glycine max               | beta-1,3-glucanase. SGN1.   |
| AAA34078.1     | M63634   | Nicotiana plumbaginifolia | regulator of beta(1,3)-glucanase. beta(1,3)-glucanase regulator.  |
| CAA30261.1     | X07280   | Nicotiana plumbaginifolia | beta-glucanase.   |
| AAA51643.1     | M23120   | Nicotiana plumbaginifolia | beta-glucanase precursor.   |
| AAA87456.1     | U22147   | Hevea brasiliensis        | beta-1,3-glucanase. HGN1. hydrolytic enzyme.  |
| CAB38443.1     | AJ133470 | Hevea brasiliensis        | beta-1,3-glucanase. hgn1.   |
| AAA03618.1     | M80608   | Lycopersicon esculentum   | beta-1,3-glucanase.   |
| AAA18928.1     | U01901   | Solanum tuberosum         | catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). glub2. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| AAC19114.1     | AF067863 | Solanum tuberosum         | 1,3-beta-glucan glucanohydrolase. glucanase.  |
| CAA92278.1     | Z68154   | Gossypium hirsutum        | 1,3-beta-glucanase.   |
| AAG24921.1     | AF311749 | Hevea brasiliensis        | beta-1,3-glucanase.   |
| AAA63539.1     | M60402   | Nicotiana tabacum         | glucan beta-1,3-glucanase. glucanase GLA.   |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAA63540.1     | M60403   | Nicotiana tabacum             | glucan-1,3-beta-glucosidase. glucanase GLB.   |
| AAA88794.1     | U01900   | Solanum tuberosum             | catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). gluB1. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| AAA63541.1     | M59442   | Nicotiana tabacum             | basic beta-1,3-glucanase. glucanase.  |
| AAB82772.2     | AF001523 | Musa acuminata                | beta-1, 3-glucanase. similar to beta-1, 3-glucanase.  |
| CAA37289.1     | X53129   | Phaseolus vulgaris            | 1,3,-beta-D-glucanase.  |
| AAF08679.1     | AF004838 | Musa acuminata                | beta-1,3-glucanase.   |
| AAD33881.1     | AF141654 | Nicotiana tabacum             | beta-1,3-glucanase. GGL4.   |
| AAD33880.1     | AF141653 | Nicotiana tabacum             | beta-1,3-glucanase. GGL1.   |
| AAA33648.1     | L02212   | Pisum sativum                 | beta-1,3-glucan hydrolysis. beta-1,3-glucanase. putative.   |
| AAA34082.1     | M20620   | Nicotiana tabacum             | prepro-beta-1,3-glucanase precursor.  |
| AAA19111.1     | U01902   | Solanum tuberosum             | catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). gluB3. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| CAA57255.1     | X81560   | Nicotiana tabacum             | (1-)-beta-glucanase. Sp41a.   |
| AAA34053.1     | M60464   | Nicotiana tabacum             | beta-1,3-glucanase.   |
| AAA63542.1     | M59443   | Nicotiana tabacum             | acidic beta-1,3-glucanase. glucanase.   |
| AAB24398.1     | S51479   | Pisum sativum                 | beta-1,3-glucanase. beta-1,3-glucanase. This sequence comes from Fig. 1B.   |
| AAB41551.1     | U27179   | Medicago sativa subsp. sativa | acidic glucanase.   |
| AAD10384.1     | U72253   | Oryza sativa                  | beta-1,3-glucanase precursor. Gns7.   |
| AAA03617.1     | M80604   | Lycopersicon esculentum       | beta-1,3-glucanase.   |
| SEQ ID NO: 151 |          |                               |   |

|            |          |  |  |
|------------|----------|--|--|
| BAA19102.1 | AB000408 | Populus kitakamiensis                  | o-methyltransferase. caffeoyl-CoA 3-O-methyltransferase.   |
| AAC28973.1 | U20736   | Medicago sativa subsp. sativa          | synthesis of feruloyl-CoA from caffeoyl-CoA and S-adenosyl-L-methionine. S-adenosyl-L-methionine:trans-caffeoyl-CoA 3-O-methyltransferase. CCOMT.  |
| CAA12198.1 | AJ224894 | Populus balsamifera subsp. trichocarpa | methylates lignin precursors. caffeoyl-CoA 3-O-methyltransferase.  |
| CAA11496.1 | AJ223621 | Populus balsamifera subsp. trichocarpa | caffeoyl CoA 3-O-methyltransferase. CCoAOMT1.  |
| AAB80931.1 | AF022775 | Nicotiana tabacum                      | caffeoyl-CoA 3-O-methyltransferase 5. CCoAOMT-5. implicated in lignification and defense reaction against pathogens.   |
| CAA83943.1 | Z33878   | Petroselinum crispum                   | caffeoyl-CoA 3-O-methyltransferase.  |
| AAA33851.1 | M69184   | Petroselinum crispum                   | caffeoyl-CoA 3-O-methyltransferase. CCoAMT.  |
| CAA90894.1 | Z54183   | Petroselinum crispum                   | trans-caffeoyl-CoA 3-O-methyltransferase. CCoAOMT.   |
| CAA90969.1 | Z54233   | Vitis vinifera                         | plant defense and lignification. caffeoyl-CoA O-methyltransferase.   |
| AAA59389.1 | U13151   | Zinnia elegans                         | S-adenosyl-L-methionine:trans-caffeoyl-CoA 3-O-methyltransferase. CCoAOMT.   |
| CAB05369.1 | Z82982   | Nicotiana tabacum                      | methylation of caffeoyl-CoA in lignin biosynthesis. caffeoyl-CoA O-methyltransferase 5. CCoAOMT-5.   |
| AAA80651.1 | U27116   | Populus tremuloides                    | caffeoyl-CoA 3-O-methyltransferase. S-adenosyl-L-methionine caffeoyl-CoA 3-O-methyltransferase; similar to Swiss-Prot Accession Number P28034; similar to proteins encoded by GenBank accession Numbers U20736, U13151, and L22203; Mr = 27.9 kDa and pI = 5.16. |
| AAC08395.1 | AF053553 | Mesembryanthemum crystallinum          | caffeoyl-CoA O-methyltransferase.  |
| CAA12200.1 | AJ224896 | Populus balsamifera subsp. trichocarpa | methylates lignin precursors. caffeoyl-CoA 3-O-methyltransferase.  |
| CAA12199.1 | AJ224895 | Populus balsamifera subsp. trichocarpa | methylates lignin precursors. caffeoyl-CoA 3-O-methyltransferase.  |
| CAA11495.1 | AJ223620 | Populus balsamifera subsp. trichocarpa | caffeoyl CoA 3-O-methyltransferase. CCoAOMT2.  |
| AAD50443.1 | AF168780 | Eucalyptus globulus                    | caffeoyl-CoA O-methyltransferase. CCoAOMT2.  |
| AAF44689.1 | AF240466 | Populus tomentosa                      | caffeoyl-CoA O-methyltransferase. CCoAOMT.   |

|            |          |                                   |  |
|------------|----------|-----------------------------------|--|
| AAC49913.1 | U38612   | Nicotiana tabacum                 | methylation of caffeoyl-CoA in feruloyl-coA in phenylpropanoid pathway. caffeoyl-coenzymeA O-methyltransferase. CCoAOMT-1.   |
| CAA72911.1 | Y12228   | Eucalyptus gunnii                 | caffeoyl-CoA O-methyltransferase. COOAMT.  |
| AAC49916.1 | U62736   | Nicotiana tabacum                 | methylation of caffeoyl-CoA in feruloyl-CoA in phenylpropanoid pathway (lignin biosynthesis). caffeoyl-CoA O-methyltransferase 4. CCoAOMT-4.   |
| BAA78733.1 | AB023482 | Oryza sativa                      | ESTs AU058067(E20733), AAU058070(E20873) correspond to a region of the predicted gene.; Similar to Populus tremuloides caffeoyl-CoA 3-O-methyltransferase mRNA, complete cds.(U27116). |
| CAA91228.1 | Z56282   | Nicotiana tabacum                 | plant defense and lignification. caffeoyl-CoA O-methyltransferase. NTCCOAOMT.  |
| AAC49914.1 | U62734   | Nicotiana tabacum                 | methylation of caffeoyl-CoA in feruloyl-CoA in phenylpropanoid pathway (lignin biosynthesis). caffeoyl-CoA O-methyltransferase 2. CCoAOMT-2.   |
| AAC26191.1 | AF046122 | Eucalyptus globulus               | catalyses the methylation of caffeoyl CoA in lignin biosynthesis. caffeoyl-CoA 3-O-methyltransferase. CCOMT. S-adenosyl-L-methionine:caffeoyl-CoA 3-O-methyltransferase.               |
| AAC49915.1 | U62735   | Nicotiana tabacum                 | methylation of caffeoyl-CoA in feruloyl-CoA in phenylpropanoid pathway (lignin biosynthesis). caffeoyl-CoA O-methyltransferase 3. CCoAOMT-3.   |
| AAD02050.1 | AF036095 | Pinus taeda                       | caffeoyl-CoA O-methyltransferase. CCoAOMT. lignin pathway O-methyltransferase.   |
| AAK16714.1 | AF327458 | Populus alba x Populus glandulosa | caffeoyl-CoA 3-O-methyltransferase. CCoAOMT.   |
| CAB45150.1 | AJ242981 | Zea mays                          | lignin synthesis. Caffeoyl CoA O-methyltransferase. ccoAOMT.   |
| CAB45149.1 | AJ242980 | Zea mays                          | lignin synthesis. Caffeoyl CoA O-methyltransferase. ccoAOMT.   |
| AAB61680.1 | L22203   | Stellaria longipes                | S-adenosyl-L-methionine:trans-caffeoyl-CoA 3-O-methyltransferase. 26.7-kDa; pI=5.3.  |
| BAA88234.1 | AB035144 | Citrus natsudaikai                | Methylation of caffeoyl-coA in feruloyl-coA in phenylpropanoid pathway. caffeoyl-CoA 3-O-methyltransferase. CCoAMT.  |
| BAA81776.1 | AP000364 | Oryza sativa                      | ESTs C98431(E0144),C71728(E0144) correspond to a region of the predicted gene.; Similar to Medicago sativa S-adenosyl-L-methionine. (U20736).  |
| BAA81774.1 | AP000364 | Oryza sativa                      | ESTs AU030740(E60171),AU030739(E60171) correspond to a region of the predicted gene.; Similar to Populus tremuloides caffeoyl-CoA 3-O-methyltransferase. (U27116).                     |
| BAA81777.1 | AP000364 | Oryza sativa                      | Similar to Petroselinum crispum caffeoyl-CoA 3-O-methyltransferase. (S49342).  |

---

CAA10217.1 AJ130841 *Populus balsamifera* subsp. *trichocarpa*  
methylates lignin precursors. caffeoyl-CoA 3-O-methyltransferase.

---

CAA04769.1 AJ001447 *Fragaria vesca*  
caffeoyl-CoA 3-O-methyltransferase. putative.

---

AAD50441.1 AF168778 *Eucalyptus globulus*  
caffeoyl-CoA O-methyltransferase. CCoAOMT1.

---

AAD50442.1 AF168779 *Eucalyptus globulus*  
caffeoyl-CoA O-methyltransferase. CCoAOMT2.

---

AAC15067.1 AF060180 *Nicotiana tabacum*  
plant lignification and defense. caffeoyl-coenzyme A trunc2. truncated caffeoyl-coenzyme A.

---

## SEQ ID NO: 152

---

AAK11255.1 AF329729 *Nicotiana tabacum*  
regulator of gene silencing. rgs-CaM; calmodulin-related protein.

---

AAD10245.1 AF030033 *Phaseolus vulgaris*  
calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal  
transduction pathways.

---

CAA62150.1 X90560 *Physcomitrella patens*  
Calmodulin. CaM.

---

BAA94696.1 AB041711 *Chara corallina*  
calmodulin. cccam1.

---

BAA96536.1 AB044286 *Chara corallina*  
calmodulin. ccam.

---

BAA94697.1 AB041712 *Chara corallina*  
calmodulin. cccam2.

---

BAA87825.1 AP000815 *Oryza sativa*  
ESTs AU030013(E50493),AU081341(E50493) correspond to a region of the predicted gene.  
Similar to O.sativa gene encoding calmodulin. (Z12828).

---

CAA61980.1 X89890 *Bidens pilosa*  
Calmodulin.

---

AAA19571.1 U10150 *Brassica napus*  
calcium binding. calmodulin. bcm1.

---

AAA87347.1 M88307 *Brassica juncea*  
calmodulin.

---

CAA74111.1 Y13784 *Mougeotia scalaris*  
Calmodulin.

---

AAA92677.1 U13736 *Pisum sativum*  
binds calcium. calmodulin-like protein.

---

AAA34015.1 L01433 *Glycine max*  
calcium-binding regulatory protein. calmodulin. SCaM-4. putative.

---

AAA33083.1 M20729 *Chlamydomonas reinhardtii*  
calmodulin.

---

AAK25753.1 AF334833 *Castanea sativa*  
calmodulin. CAM2.

---

|            |          |   |
|------------|----------|---|
| AAF73157.1 | AF150059 | Brassica napus<br>calmodulin. CaM1. involved in seed germination.   |
| CAA74307.1 | Y13974   | Zea mays<br>calmodulin.   |
| AAA34238.1 | L20507   | Vigna radiata<br>calmodulin.  |
| AAA34237.1 | L20691   | Vigna radiata<br>calmodulin.  |
| AAC49587.1 | U49105   | Triticum aestivum<br>calmodulin TaCaM4-1. calcium-binding protein.  |
| AAC49586.1 | U49104   | Triticum aestivum<br>calmodulin TaCaM3-3. calcium-binding protein.  |
| AAC49585.1 | U49103   | Triticum aestivum<br>calmodulin TaCaM3-2. calcium-binding protein.  |
| AAC49584.1 | U48693   | Triticum aestivum<br>calmodulin TaCaM3-1. calcium-binding protein.  |
| AAC49580.1 | U48689   | Triticum aestivum<br>calmodulin TaCaM1-3. calcium-binding protein.  |
| AAC49579.1 | U48688   | Triticum aestivum<br>calmodulin TaCaM1-2. calcium binding protein.  |
| AAC49578.1 | U48242   | Triticum aestivum<br>calmodulin TaCaM1-1. calcium-binding.  |
| AAA34014.1 | L01432   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-3. putative.  |
| AAA03580.1 | L01431   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-2. putative.  |
| AAA34013.1 | L01430   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-1. putative.  |
| AAB36130.1 | S81594   | Vigna radiata<br>auxin-regulated calmodulin. auxin-regulated calmodulin, arCaM. This sequence comes from Fig. 1; arCaM. |
| AAA33901.1 | L18913   | Oryza sativa<br>calcium binding protein, signal transduction. calmodulin. putative.                                     |
| AAA92681.1 | U13882   | Pisum sativum<br>calcium-binding protein. calmodulin.   |
| AAA33706.1 | M80836   | Petunia x hybrida<br>calmodulin. CAM81.   |
| AAA33705.1 | M80831   | Petunia x hybrida<br>calmodulin-related protein. CAM53.   |
| CAA78287.1 | Z12827   | Oryza sativa<br>calcium binding protein, signal transduction. calmodulin.   |
| CAA46150.1 | X65016   | Oryza sativa<br>calmodulin. cam.  |

|                |          |  |
|----------------|----------|--|
| CAA36644.1     | X52398   | Medicago sativa<br>calmodulin (AA 1-149).  |
| CAA43143.1     | X60738   | Malus x domestica<br>Calmodulin. CaM.  |
| CAA78301.1     | Z12839   | Lilium longiflorum<br>calcium binding protein, signal transduction. calmodulin.  |
| AAA33397.1     | L18912   | Lilium longiflorum<br>calcium binding protein, signal transduction. calmodulin. putative.  |
| AAB68399.1     | U79736   | Helianthus annuus<br>calmodulin. HaCaM.  |
| CAA42423.1     | X59751   | Daucus carota<br>calmodulin. Ccam-1.   |
| AAA32938.1     | M27303   | Hordeum vulgare<br>calmodulin.   |
| BAA88540.1     | AP000969 | Oryza sativa<br>ESTs AU081349(E61253), D41425(S3918) correspond to a region of the predicted gene.<br>Similar to calmodulin. (AF042840). |
| AAG27432.1     | AF295637 | Elaeis guineensis<br>calmodulin.   |
| AAG11418.1     | AF292108 | Prunus avium<br>calmodulin.  |
| AAC36059.1     | AF042840 | Oryza sativa<br>calmodulin. CaM1.  |
| SEQ ID NO: 153 |          |  |
| CAA40474.1     | X57187   | Phaseolus vulgaris<br>chitinase. Chi4.   |
| AAB65776.1     | U97521   | Vitis vinifera<br>class IV endochitinase. VvChi4A.   |
| CAA61281.1     | X88803   | Vigna unguiculata<br>chitinase class 4. CHI4.  |
| AAB65777.1     | U97522   | Vitis vinifera<br>class IV endochitinase. VvChi4B.   |
| BAA22966.1     | D45182   | Chenopodium amaranticolor<br>chitinase.  |
| BAA22968.1     | D45184   | Chenopodium amaranticolor<br>chitinase.  |
| BAA22965.1     | D45181   | Chenopodium amaranticolor<br>chitinase.  |
| CAA43708.1     | X61488   | Brassica napus<br>chitinase.   |
| BAA22967.1     | D45183   | Chenopodium amaranticolor<br>chitinase.  |

|            |          |   |
|------------|----------|---|
| CAA53544.1 | X75945   | Beta vulgaris<br>chitinase. Ch4.  |
| AAC49435.1 | U52845   | Daucus carota<br>class IV chitinase EP3-1/H5. EP3.                                    |
| AAB08468.1 | U52846   | Daucus carota<br>class IV chitinase EP3-2/H1. EP3.                                    |
| AAB08470.1 | U52848   | Daucus carota<br>class IV chitinase EP3B/E6. EP3.                                     |
| AAB08469.1 | U52847   | Daucus carota<br>class IV chitinase EP3-3/E7. EP3.                                    |
| AAA33445.1 | M84165   | Zea mays<br>chitinase B. seed chitinase.  |
| AAA33444.1 | M84164   | Zea mays<br>chitinase A. seed chitinase.  |
| AAA32916.1 | L25826   | Beta vulgaris<br>chitinase. SP2.  |
| AAD28733.1 | AF112966 | Triticum aestivum<br>chitinase IV precursor. Cht4.                                    |
| BAB21377.1 | AB054811 | Oryza sativa<br>PR-3 class IV chitinase. Cht4. Catalytic domain.                      |
| BAB21374.1 | AB054687 | Oryza sativa<br>PR-3 class IV chitinase. Cht4. catalytic domain.                      |
| BAA19793.1 | AB003194 | Oryza sativa<br>chitinase IIb.  |
| AAA85364.1 | L42467   | Picea glauca<br>chitinase. chi.   |
| AAB01665.1 | U21848   | Brassica napus<br>chitinase class IV. LSC222.   |
| AAC35981.1 | AF090336 | Citrus sinensis<br>chitin hydrolase. chitinase CHI1. chil.                            |
| AAD28730.1 | AF112963 | Triticum aestivum<br>chitinase II precursor. Cht2.                                    |
| AAF04454.1 | AF000966 | Poa pratensis<br>chitinase. Chi2.   |
| CAC17793.1 | AJ301671 | Nicotiana sylvestris<br>hydrolysis of chitin. endochitinase. chnb. class I chitinase. |
| AAF04453.1 | AF000964 | Poa pratensis<br>chitinase. Chi1.   |
| CAA34812.1 | X16938   | Nicotiana tabacum<br>chitinase precursor.   |
| CAA34813.1 | X16939   | Nicotiana tabacum<br>chitinase precursor (AA -23 to 306).                             |



|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA45822.1     | X64519   | Nicotiana tabacum       | chitinase B class I. CHN200.   |
| CAA35945.1     | X51599   | Nicotiana tabacum       | chitinase. CHN50.  |
| AAB23374.1     | S44869   | Nicotiana tabacum       | basic chitinase. basic chitinase. This sequence comes from Fig. 1.   |
| AAA34070.1     | M15173   | Nicotiana tabacum       | endochitinase precursor (EC 3.2.1.14).   |
| CAA30142.1     | X07130   | Solanum tuberosum       | endochitinase.   |
| CAA33517.1     | X15494   | Solanum tuberosum       | pre-chitinase (AA -26 to 302).   |
| AAG53609.1     | AF280437 | Secale cereale          | 31.7 kDa class I endochitinase-antifreeze protein precursor. cht9.   |
| CAB01591.1     | Z78202   | Persea americana        | hydrolysis of the 1,4-beta-linkages of chitin. endochitinase. chl1.  |
| CAA53626.1     | X76041   | Triticum aestivum       | endochitinase. CHI.  |
| CAA78845.1     | Z15140   | Lycopersicon esculentum | chitinase. Encodes 30 kD basic intracellular chitinase.  |
| AAC16010.1     | AF061805 | Elaeagnus umbellata     | acidic chitinase.  |
| AAA32640.1     | M94106   | Allium sativum          | chitinase. chitinase.  |
| AAA32641.1     | M94105   | Allium sativum          | chitinase. chitinase.  |
| AAA56787.1     | L34211   | Hordeum vulgare         | hydrolysis of chitin. chitinase. CHI33.  |
| AAA17409.1     | U02607   | Solanum tuberosum       | catalyzes the random hydrolysis of 1,4-beta-(2-acetamido-2-deoxy-D-glucoside) linkages in chitin; plant defense gene. chitinase. chtB3. induced expression in response to infection, elicitor, ethylene, wounding; preproprotein; gene product subunit is monomer.   |
| AAA18332.1     | U02605   | Solanum tuberosum       | catalyzes the random hydrolysis of 1,4-beta-(2-acetamido-2-deoxy-D-glucoside) linkages in chitin; plant defense gene. chitinase. chtB1. induced expression in response to infection, elicitor, ethylene, wounding; preproprotein; gene product subunit is a monomer. |
| CAA45821.1     | X64518   | Nicotiana tabacum       | chitinase C class I. CHN14.  |
| SEQ ID NO: 154 |          |                         |  |
| AAB35812.1     | S80554   | Arabidopsis             | chalcone synthase. chalcone synthase, CHS. This sequence comes from Fig. 5; CHS.   |
| AAF23570.1     | AF112095 | Arabidopsis halleri     | chalcone synthase. CHS.  |

|            |          |   |
|------------|----------|---|
| AAF23568.1 | AF112093 | <i>Arabidopsis griffithiana</i><br>chalcone synthase. CHS.                  |
| AAG43351.1 | AF144533 | <i>Arabidopsis korshinskyi</i><br>chalcone synthase. chs.                   |
| AAF23581.1 | AF112106 | <i>Capsella rubella</i><br>chalcone synthase. CHS.                          |
| AAF23569.1 | AF112094 | <i>Halimolobos perplexa</i> var. <i>perplexa</i><br>chalcone synthase. CHS. |
| AAG43349.1 | AF144531 | <i>Arabidopsis himalaica</i><br>chalcone synthase. chs.                     |
| AAF23575.1 | AF112100 | <i>Arabidopsis lyrata</i> subsp. <i>lyrata</i><br>chalcone synthase. CHS.   |
| AAF23567.1 | AF112092 | <i>Arabidopsis griffithiana</i><br>chalcone synthase. CHS.                  |
| AAF23578.1 | AF112103 | <i>Arabidopsis lyrata</i> subsp. <i>petraea</i><br>chalcone synthase. CHS.  |
| AAF23576.1 | AF112101 | <i>Arabis parishii</i><br>chalcone synthase. CHS.                           |
| AAF23574.1 | AF112099 | <i>Arabis lyallii</i><br>chalcone synthase. CHS.                            |
| AAF23566.1 | AF112091 | <i>Arabis glabra</i><br>chalcone synthase. CHS.                             |
| AAF23565.1 | AF112090 | <i>Arabis fendleri</i><br>chalcone synthase. CHS.                           |
| AAF23563.1 | AF112088 | <i>Arabis drummondii</i><br>chalcone synthase. CHS.                         |
| AAF23564.1 | AF112089 | <i>Arabis drummondii</i><br>chalcone synthase. CHS.                         |
| AAF23579.1 | AF112104 | <i>Arabidopsis lyrata</i> subsp. <i>petraea</i><br>chalcone synthase. CHS.  |
| AAF23573.1 | AF112098 | <i>Arabis lignifera</i><br>chalcone synthase. CHS.                          |
| AAF23560.1 | AF112085 | <i>Cardamine amara</i><br>chalcone synthase. CHS.                           |
| AAG43348.1 | AF144530 | <i>Rorippa amphibia</i><br>chalcone synthase. chs.                          |
| AAG43356.1 | AF144538 | <i>Cardamine penzesii</i><br>chalcone synthase. chs.                        |
| AAG43359.1 | AF144541 | <i>Sisymbrium irio</i><br>chalcone synthase. chs.                           |
| AAG43352.1 | AF144534 | <i>Lepidium campestre</i><br>chalcone synthase. chs.                        |

|            |          |  |
|------------|----------|--|
| CAA32495.1 | X14314   | <i>Sinapis alba</i><br>chalcone synthase (AA 1-395).       |
| AAG43357.1 | AF144539 | <i>Cardamine rivularis</i><br>chalcone synthase. chs.      |
| AAF23583.1 | AF112108 | <i>Barbarea vulgaris</i><br>chalcone synthase. CHS.        |
| AAC31914.1 | AF076336 | <i>Brassica napus</i><br>chalcone synthase B2. CHSB2.      |
| AAC31912.1 | AF076334 | <i>Brassica napus</i><br>chalcone synthase A2. CHSA2.      |
| AAF23577.1 | AF112102 | <i>Arabis pauciflora</i><br>chalcone synthase. CHS.        |
| AAG43350.1 | AF144532 | <i>Cochlearia danica</i><br>chalcone synthase. chs.        |
| CAA34460.1 | X16437   | <i>Sinapis alba</i><br>chalcone synthase.                  |
| CAA35600.1 | X17577   | <i>Matthiola incana</i><br>chalcone synthase (AA 1-394).   |
| AAG43358.1 | AF144540 | <i>Cardamine pratensis</i><br>chalcone synthase. chs.      |
| AAG43353.1 | AF144535 | <i>Thlaspi arvense</i><br>chalcone synthase. chs.          |
| AAC31913.1 | AF076335 | <i>Brassica napus</i><br>chalcone synthase B1. CHSB1.      |
| AAF23571.1 | AF112096 | <i>Arabis hirsuta</i><br>chalcone synthase. CHS.           |
| AAF23582.1 | AF112107 | <i>Arabis turrita</i><br>chalcone synthase. CHS.           |
| AAG43406.1 | AF174529 | <i>Aubrieta deltoidea</i><br>chalcone synthase. chs.       |
| AAG43355.1 | AF144537 | <i>Alliaria petiolata</i><br>chalcone synthase. chs.       |
| AAF23580.1 | AF112105 | <i>Arabis procurrens</i><br>chalcone synthase. CHS.        |
| AAF23572.1 | AF112097 | <i>Arabis jacquini</i><br>chalcone synthase. CHS.          |
| AAF23562.1 | AF112087 | <i>Arabis blepharophylla</i><br>chalcone synthase. CHS.    |
| AAF23584.1 | AF112109 | <i>Aubrieta deltoidea</i><br>chalcone synthase. CHS.       |
| AAG43354.1 | AF144536 | <i>Microthlaspi perfoliatum</i><br>chalcone synthase. chs. |

|                |          |                               |  |
|----------------|----------|-------------------------------|--|
| AAF23557.1     | AF112082 | Aethionema grandiflora        | chalcone synthase. CHS.  |
| AAF23558.1     | AF112083 | Arabis alpina                 | chalcone synthase. CHS.  |
| AAF23559.1     | AF112084 | Arabis alpina                 | chalcone synthase. CHS.  |
| AAB87072.1     | AF031922 | Raphanus sativus              | chalcone synthase. CHS.  |
| AAG43360.1     | AF144542 | Ionopsidium abulense          | chalcone synthase. chs.  |
| AAC31911.1     | AF076333 | Brassica napus                | chalcone synthase A1. CHSA1.   |
| SEQ ID NO: 156 |          |                               |  |
| AAD10327.1     | U63534   | Fragaria x ananassa           | catalyzes the reduction of cinnamylaldehydes leading to monolignols. cinnamyl alcohol dehydrogenase. CAD. involved with lignin biosynthesis.   |
| AAK28509.1     | AF320110 | Fragaria x ananassa           | cinnamyl alcohol dehydrogenase.  |
| AAB38503.1     | U79770   | Mesembryanthemum crystallinum | cinnamyl-alcohol dehydrogenase Eli3.   |
| CAA48028.1     | X67817   | Petroselinum crispum          | Eli3.  |
| AAC15467.1     | U24561   | Apium graveolens              | converts mannitol to mannose. mannitol dehydrogenase. Mtd. 1-oxidoreductase; induced with sodium salicylate; similar to the plant defense gene ELI3 in Arabidopsis thaliana, PIR Accession Number S28044; EC number unassigned; MTD. |
| AAC35846.1     | AF083333 | Medicago sativa               | cinnamyl-alcohol dehydrogenase. MsaCad1.   |
| AAC61854.1     | AF067082 | Apium graveolens              | oxidizes mannitol to mannose. mannitol dehydrogenase. Mtd. mannitol 1-oxidoreductase.  |
| AAA74882.1     | L36823   | Stylosanthes humilis          | cinnamyl-alcohol dehydrogenase. CAD1.  |
| AAA74883.1     | L36456   | Stylosanthes humilis          | cinnamyl-alcohol dehydrogenase. CAD3.  |
| AAF72100.1     | AF146691 | Lycopersicon esculentum       | ELI3. Eli3. similar] to cinnamyl alcohol dehydrogenase.  |
| CAA86072.1     | Z37991   | Pinus taeda                   | cinnamyl alcohol dehydrogenase.  |
| CAA05095.1     | AJ001924 | Picea abies                   | cinnamyl alcohol dehydrogenase. cad2.  |
| AAB38774.1     | U62394   | Pinus radiata                 | cinnamyl alcohol dehydrogenase. CAD.   |

|            |          |   |
|------------|----------|---|
| CAA51226.1 | X72675   | <i>Picea abies</i><br>cinnamyl-alcohol dehydrogenase.   |
| CAA05097.1 | AJ001926 | <i>Picea abies</i><br>cinnamyl alcohol dehydrogenase. cad8.   |
| CAA05096.1 | AJ001925 | <i>Picea abies</i><br>cinnamyl alcohol dehydrogenase. cad7.   |
| AAC31166.1 | AF060491 | <i>Pinus radiata</i><br>cinnamyl alcohol dehydrogenase. CAD.  |
| CAA86073.1 | Z37992   | <i>Pinus taeda</i><br>cinnamyl alcohol dehydrogenase.   |
| CAA44216.1 | X62343   | <i>Nicotiana tabacum</i><br>cinnamyl-alcohol dehydrogenase. CAD14.  |
| CAA44217.1 | X62344   | <i>Nicotiana tabacum</i><br>cinnamyl-alcohol dehydrogenase. CAD19.  |
| BAA03099.1 | D13991   | <i>Aralia cordata</i><br>cinnamyl alcohol dehydrogenase. cadac1.  |
| CAA79625.1 | Z19573   | <i>Medicago sativa</i><br>lignin biosynthesis. cinnamyl alcohol dehydrogenase.  |
| AAF43140.1 | AF217957 | <i>Populus tremuloides</i><br>cinnamyl alcohol dehydrogenase. CAD.  |
| AAC35845.1 | AF083332 | <i>Medicago sativa</i><br>cinnamyl-alcohol dehydrogenase. MsaCad2.  |
| CAC07423.1 | AJ295837 | <i>Populus balsamifera</i> subsp. <i>trichocarpa</i><br>lignin monomer biosynthesis. cinnamyl alcohol dehydrogenase. cad.   |
| CAA79622.1 | Z19568   | <i>Populus deltoides</i><br>lignin biosynthesis. cinnamyl alcohol dehydrogenase.  |
| AAC07987.1 | AF038561 | <i>Eucalyptus globulus</i><br>catalyses the reduction of cinnamaldehydes to the corresponding cinnamyl alcohols as the last step in the production of lignin monomers. cinnamyl alcohol dehydrogenase. CAD.                     |
| AAG15553.1 | AF294793 | <i>Eucalyptus saligna</i><br>cinnamyl alcohol dehydrogenase. cad. CAD.  |
| AAK00679.1 | AF229407 | <i>Brassica napus</i><br>Eli3 product. ELI3-BN-2. protein identity based on similarity to Arabidopsis sequence; intron/exon junctions deduced from alignments to DNA or mRNA sequences already present in the GenBank database. |
| CAA46585.1 | X65631   | <i>Eucalyptus gunnii</i><br>cinnamyl-alcohol dehydrogenase. cad.  |
| CAA53211.1 | X75480   | <i>Eucalyptus gunnii</i><br>cinnamyl-alcohol dehydrogenase. CAD.  |
| AAK00681.1 | AF229409 | <i>Brassica napus</i><br>Eli3 product. ELI3-BN-4. protein identity based on similarity to Arabidopsis sequence; intron/exon junctions deduced from alignments to DNA or mRNA sequences already present in the GenBank database. |

|            |          |  |
|------------|----------|--|
| AAB70908.1 | AF010290 | <i>Lolium perenne</i><br>cinnamyl alcohol dehydrogenase. CAD.  |
| AAK00682.1 | AF229410 | <i>Brassica oleracea</i><br>Eli3 product. ELI3-BO-1. protein identity based on similarity to Arabidopsis sequence;<br>intron/exon junctions deduced from alignments to DNA or mRNA sequences already present<br>in the GenBank database. |
| CAA74070.1 | Y13733   | <i>Zea mays</i><br>cinnamyl alcohol dehydrogenase. cad.  |
| CAA06687.1 | AJ005702 | <i>Zea mays</i><br>cinnamyl alcohol dehydrogenase. cad.  |
| AAK00678.1 | AF229406 | <i>Brassica napus</i><br>Eli3 product. ELI3-BN-1. protein identity based on similarity to Arabidopsis sequence;<br>intron/exon junctions deduced from alignments to DNA or mRNA sequences already present<br>in the GenBank database.    |
| CAA13177.1 | AJ231135 | <i>Saccharum officinarum</i><br>lignin biosynthesis. cinnamyl alcohol dehydrogenase. cad.  |
| AAK00684.1 | AF229412 | <i>Brassica rapa</i><br>Eli3 product. ELI3-BR-2. protein identity based on similarity to Arabidopsis sequence;<br>intron/exon junctions deduced from alignments to DNA or mRNA sequences already present<br>in the GenBank database.     |
| AAK00680.1 | AF229408 | <i>Brassica napus</i><br>Eli3 product. ELI3-BN-3. protein identity based on similarity to Arabidopsis sequence;<br>intron/exon junctions deduced from alignments to DNA or mRNA sequences already present<br>in the GenBank database.    |
| AAK00683.1 | AF229411 | <i>Brassica rapa</i><br>Eli3 product. ELI3-BR-1. protein identity based on similarity to Arabidopsis sequence;<br>intron/exon junctions deduced from alignments to DNA or mRNA sequences already present<br>in the GenBank database.     |
| BAA19487.1 | D86590   | <i>Zinnia elegans</i><br>cinnamyl alcohol dehydrogenase. ZCAD1.  |
| BAA04046.1 | D16624   | <i>Eucalyptus botryoides</i><br>cinnamyl alcohol dehydrogenase. Cad1:Eb:1.   |
| AAD18000.1 | AF109157 | <i>Eucalyptus globulus</i><br>cinnamyl alcohol dehydrogenase. CAD.   |
| AAF23409.1 | AF207552 | <i>Brassica napus</i><br>cinnamyl alcohol dehydrogenase. CADa-1.   |
| AAF23412.1 | AF207555 | <i>Brassica rapa</i><br>cinnamyl alcohol dehydrogenase. CADa.  |
| AAF23411.1 | AF207554 | <i>Brassica oleracea</i><br>cinnamyl alcohol dehydrogenase. CADa.  |
| AAF23410.1 | AF207553 | <i>Brassica napus</i><br>cinnamyl alcohol dehydrogenase. CADa-2.   |
| AAF23416.1 | AF207559 | <i>Brassica rapa</i><br>cinnamyl alcohol dehydrogenase. CADb.  |

|                |          |                      |   |
|----------------|----------|----------------------|---|
| AAF23415.1     | AF207558 | Brassica oleracea    | cinnamyl alcohol dehydrogenase. CADb.   |
| SEQ ID NO: 158 |          |                      |   |
| BAA87853.1     | AP000816 | Oryza sativa         | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAA78764.1     | AB023482 | Oryza sativa         | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAF43496.1     | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| AAK11674.1     | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| BAB16871.1     | AP002537 | Oryza sativa         | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).  |
| AAG16628.1     | AY007545 | Brassica napus       | protein serine/threonine kinase BNK1.   |
| AAC27894.1     | AF023164 | Zea mays             | leucine-rich repeat transmembrane protein kinase 1. ltk1.   |
| BAA94509.1     | AB041503 | Populus nigra        | protein kinase 1. PnPK1.  |
| BAA94510.1     | AB041504 | Populus nigra        | protein kinase 2. PnPK2.  |
| BAB03429.1     | AP002817 | Oryza sativa         | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).             |
| BAB07999.1     | AP002525 | Oryza sativa         | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB39409.1     | AP002901 | Oryza sativa         | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| AAC27895.1     | AF023165 | Zea mays             | leucine-rich repeat transmembrane protein kinase 2. ltk2.   |
| BAB21241.1     | AP002953 | Oryza sativa         | Putative Pto kinase interactor 1. P0426D06.21. contains ESTs AU108280(E0721),D48017(S13927).  |
| AAG59657.1     | AC084319 | Oryza sativa         | putative protein kinase. OSJNBa0004B24.20.  |
| AAF91337.1     | AF249318 | Glycine max          | Pt1l kinase-like protein. Pt1lb. protein kinase.  |
| AAF91336.1     | AF249317 | Glycine max          | Pt1l kinase-like protein. Pt1la. protein kinase.  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAB51834.1 | 00069    | Oryza sativa            | 11332.5. contains eukaryotic protein kinase domain PF.  |
| BAA87852.1 | AP000816 | Oryza sativa            | Similar to putative Ser/Thr protein kinase. (AC004218).   |
| BAA92221.1 | AP001278 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).        |
| AAK00425.1 | AC069324 | Oryza sativa            | Putative protein kinase. OSJNBa0071K19.11.  |
| AAB09771.1 | U67422   | Zea mays                | CRINKLY4 precursor. cr4. receptor kinase homolog.   |
| BAB18321.1 | AP002865 | Oryza sativa            | putative receptor protein kinase. P0034C11.11.  |
| BAB40081.1 | AP003074 | Oryza sativa            | putative receptor protein kinase. OSJNBa0004G10.30.   |
| AAC61805.1 | U28007   | Lycopersicon esculentum | serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.  |
| BAB39873.1 | AP002882 | Oryza sativa            | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).                                 |
| AAD38286.1 | AC007789 | Oryza sativa            | putative protein kinase. OSJNBa0049B20.13.  |
| BAB21240.1 | AP002953 | Oryza sativa            | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).                                    |
| AAK21965.1 | AY028699 | Brassica napus          | receptor protein kinase PERK1.  |
| AAG03090.1 | AC073405 | Oryza sativa            | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).                                      |
| BAB18292.1 | AP002860 | Oryza sativa            | putative receptor-like protein kinase. P0409B08.19.   |
| AAF34428.1 | AF172282 | Oryza sativa            | receptor-like protein kinase. DUPR11.18.  |
| AAG25966.1 | AF302082 | Nicotiana tabacum       | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment. |
| AAF66615.1 | AF142596 | Nicotiana tabacum       | LRR receptor-like protein kinase.   |
| CAB51836.1 | AJ243961 | Oryza sativa            | Putative Ser/Thr protein kinase. 11332.7.   |
| BAA90808.1 | AP001168 | Oryza sativa            | Similar to putative receptor-like protein kinase (AL035679).  |



|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA97692.1     | Z73295   | Catharanthus roseus     | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.     |
| BAA84787.1     | AP000559 | Oryza sativa            | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| BAA83373.1     | AP000391 | Oryza sativa            | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| AAF76313.1     | AF220603 | Lycopersicon esculentum | Pto kinase. LescPth5.  |
| AAB47421.1     | U59316   | Lycopersicon esculentum | serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.                             |
| CAA79355.1     | Z18921   | Brassica oleracea       | S-receptor kinase-like protein.  |
| AAA33915.1     | L27821   | Oryza sativa            | receptor type serine/threonine kinase. protein kinase.   |
| BAA95893.1     | AP002071 | Oryza sativa            | Similar to Arabidopsis thaliana wak4 gene; wall-associated kinase 4. (AJ009695).   |
| AAC36318.1     | AF053127 | Malus x domestica       | leucine-rich receptor-like protein kinase. LRPKm1.   |
| SEQ ID NO: 159 |          |                         |  |
| AAA86687.1     | U15777   | Lupinus albus           | farnesyl pyrophosphate synthase. fps1.   |
| AAA87729.1     | U20771   | Lupinus albus           | farnesyl pyrophosphate synthase. fps1.   |
| CAA72793.1     | Y12072   | Gossypium arboreum      | farnesyl pyrophosphate synthase. fps1.   |
| BAB40665.1     | AB053486 | Humulus lupulus         | farnesyl pyrophosphate synthase. fpps.   |
| BAB40666.1     | AB053487 | Humulus lupulus         | farnesyl pyrophosphate synthase. fpps.   |
| CAA57893.1     | X82543   | Parthenium argentatum   | farnesyl diphosphate synthase. fps2.   |
| AAC78557.1     | AF019892 | Helianthus annuus       | farnesyl pyrophosphate synthase. FPS.  |
| CAA57892.1     | X82542   | Parthenium argentatum   | farnesyl diphosphate synthase. fps1.   |
| AAC49452.1     | U36376   | Artemisia annua         | farnesyl diphosphate synthase. fps1.   |
| CAA59170.1     | X84695   | Capsicum annuum         | dimethylallyltransferase. Fps, farnesyl pyrophosphate synthase gene.   |

|            |          |  |
|------------|----------|--|
| AAC73051.1 | AF048747 | <i>Lycopersicon esculentum</i><br>synthesis of farnesyl pyrophosphate. farnesyl pyrophosphate synthase. FPS1. prenyl transferase; farnesyl diphosphate synthetase. |
| AAD17204.1 | AF112881 | <i>Artemisia annua</i><br>farnesyl diphosphate synthase.   |
| BAA19856.1 | D85317   | <i>Oryza sativa</i><br>farnesyl pyrophosphate synthase. dimethylallyltransferase; geranyltranstransferase.   |
| BAA36276.1 | AB021747 | <i>Oryza sativa</i><br>farnesyl diphosphate synthase. FPPS1.   |
| AAD32648.1 | AF136602 | <i>Artemisia annua</i><br>farnesyl diphosphate synthase. fps2.   |
| AAB39276.1 | L39789   | <i>Zea mays</i><br>farnesyl pyrophosphate synthetase. fps. putative.   |
| AAD27558.1 | AF111710 | <i>Oryza sativa</i> subsp. <i>indica</i><br>putative farnesyl pyrophosphate synthase. similar to <i>Oryza sativa</i> EST clones E10230_1A, C52647_1A, 232.         |
| BAA36347.1 | AB021979 | <i>Oryza sativa</i><br>farnesyl diphosphate synthase. FPPS2.   |
| AAD37789.1 | AF149257 | <i>Artemisia annua</i><br>farnesyl diphosphate synthase. FPP synthase.   |
| BAB20822.1 | AB045713 | <i>Taraxacum japonicum</i><br>putative FPP synthase. TJFPPS.   |
| BAB16688.1 | AB041627 | <i>Eucommia ulmoides</i><br>FPP synthase 2. EUFPPS2. putative.   |
| BAB21061.1 | AB046212 | <i>Sonchus oleraceus</i><br>putative FPP synthase. SoFPPS.   |
| AAD45122.1 | AF164026 | <i>Xanthoceras sorbifolium</i><br>synthesis of farnesyl pyrophosphate. farnesyl pyrophosphate synthase. FPS. prenyl transferase; farnesyl diphosphate synthetase.  |
| AAB93951.1 | U97330   | <i>Nicotiana tabacum</i><br>farnesylpyrophosphate synthase. FPPS.  |
| BAB16687.1 | AB041626 | <i>Eucommia ulmoides</i><br>FPP synthase 1. EUFPPS1. putative.   |
| BAB39479.1 | AB049086 | <i>Youngia japonica</i><br>putative FPP synthase 1. YjFPPS1.   |
| AAB93984.1 | AF005201 | <i>Parthenium argentatum</i><br>farnesyl pyrophosphate synthase. FPS3.   |

## SEQ ID NO: 160

|            |          |   |
|------------|----------|---|
| BAB03615.1 | AP002522 | <i>Oryza sativa</i><br>putative gamma-glutamyltransferase. P0009G03.15. contains ESTs AU056150(S20332), AU056151(S20332). |
| BAB03616.1 | AP002522 | <i>Oryza sativa</i><br>putative gamma-glutamyltransferase. P0009G03.16. contains EST AU056150(S20332).                    |

## SEQ ID NO: 162

CAB96145.1 AJ250951 *Mesembryanthemum crystallinum*  
phospholipid hydroperoxide glutathione peroxidase-like protein. gpxmc1.

CAA42780.1 X60219 *Nicotiana sylvestris*  
homologous to animal glutathione peroxidases.

BAB16430.1 AB041518 *Nicotiana tabacum*  
glutathione peroxidase Nt-SubC08. Nt-SubC08.

CAA75054.1 Y14762 *Lycopersicon esculentum*  
glutathione peroxidase. GPXle-1.

AAB94892.1 AF037051 *Gossypium hirsutum*  
glutathione peroxidase.

CAB59893.1 AJ238697 *Hordeum vulgare*  
GPX12Hv, glutathione peroxidase-like protein.

BAA22194.1 D63425 *Spinacia oleracea*  
phospholipid hydroperoxide glutathione peroxidase-like protein. similar to mammalian  
phospholipid hydroperoxide glutathione peroxidases.

CAB59895.1 AJ238745 *Hordeum vulgare*  
glutathione peroxidase-like protein GPX54Hv.

AAC78466.1 AF053311 *Zantedeschia aethiopica*  
glutathione peroxidase. gpx.

CAA04142.1 AJ000508 *Pisum sativum*  
phospholipid glutathione peroxidase. plastid-localised.

CAA75009.1 Y14707 *Helianthus annuus*  
glutathione peroxidase. GPxha-2.

CAB59894.1 AJ238744 *Hordeum vulgare*  
glutathione peroxidase-like protein GPX15Hv.

CAA74775.1 Y14429 *Helianthus annuus*  
glutathione peroxidase. GPxha-1.

CAC17628.1 AJ270955 *Oryza sativa*  
putative role in antioxidative systems. putative phospholipid hydroperoxide glutathione  
peroxidase. riPHGPX.

BAA83594.1 AB009083 *Chlamydomonas* sp. W80  
glutathione peroxidase.

AAB66330.1 AF014927 *Chlamydomonas reinhardtii*  
glutathione peroxidase homolog. gpxh.

CAA75055.1 Y14763 *Lycopersicon esculentum*  
glutathione peroxidase. GPXle-2.

CAA09194.1 AJ010455 *Triticum aestivum*  
glutathione peroxidase. PHGPX6.

CAB66331.1 AJ279689 *Betula pendula*  
glutathione peroxidase. gpx.

## SEQ ID NO: 163

|            |          |                                 |   |
|------------|----------|---------------------------------|---|
| AAF67753.1 | AF255651 | Brassica rapa subsp. pekinensis | conversion of oxidized glutathione to reduced glutathione. cytosolic glutathione reductase. GR1.                      |
| AAC49980.2 | AF008441 | Brassica rapa                   | glutathione reductase. BcGR1. cytosolic.  |
| BAA11214.1 | D78136   | Oryza sativa                    | Glutathione Reductase. putative.  |
| CAA66924.1 | X98274   | Pisum sativum                   | glutathione reductase. cytosolic.   |
| BAA36283.1 | D85751   | Oryza sativa                    | glutathione reductase.  |
| BAA37092.1 | AB009592 | Oryza sativa                    | conversion of oxidized glutathione to reduced glutathione. cytosolic glutathione reductase. RGRC2. Amino Acids 1-496. |
| BAA07108.1 | D37870   | Spinacia oleracea               | Glutathione Reductase precursor. Chloroplastic glutathione reductase.   |
| CAC13956.1 | AJ400816 | Mesembryanthemum crystallinum   | reduction of glutathione. glutathione reductase. gr1.   |
| CAB66332.1 | AJ279690 | Betula pendula                  | glutathione reductase. gr.  |
| CAA53925.1 | X76293   | Nicotiana tabacum               | glutathione reductase (NADPH). gor.   |
| CAA42921.1 | X60373   | Pisum sativum                   | glutathione reductase (NADPH). Protein sequence is in conflict with the conceptual translation.                       |
| AAK27157.1 | AF349449 | Brassica juncea                 | glutathione reductase. GR2.   |
| AAD28177.1 | AF109694 | Brassica juncea                 | glutathione reductase. GR1.   |
| CAA62482.1 | X90996   | Pisum sativum                   | glutathione reductase (NADPH). gr. alpha II subunit.  |
| AAF26175.1 | AF105199 | Glycine max                     | glutathione reductase. GR-5.  |
| AAB70837.1 | AF019907 | Vitis vinifera                  | glutathione reductase (NADPH). GOR. VvGR1.  |
| AAA33962.1 | L11632   | Glycine max                     | glutathione reductase. GR.  |
| CAA54043.1 | X76533   | Nicotiana tabacum               | glutathione reductase (NADPH). gor.   |
| CAA06835.1 | AJ006055 | Zea mays                        | NADPH-dependent reduction of glutathione disulphide. glutathione reductase. gor1.                                     |
| CAA53993.1 | X76455   | Nicotiana tabacum               | glutathione reductase. gor.   |

|                |          |                                       |  |
|----------------|----------|---------------------------------------|--|
| AAB30526.1     | S70187   | Glycine max                           | ferric leghemoglobin reductase. ferric leghemoglobin reductase, FLbR. Method: conceptual translation with partial peptide sequencing; This sequence comes from Fig. 3; FLbR. |
| AAC26053.1     | AF074940 | Glycine max                           | ferric leghemoglobin reductase-2 precursor. FLbR homolog;FLbR-2.   |
| AAD53185.1     | AF181096 | Vigna unguiculata                     | ferric leghemoglobin reductase. flbr.  |
| AAA60979.1     | U06461   | Pisum sativum                         | catalyzes the conversion of monodehydroascorbate to ascorbate, oxidizing NADH in the process, binds to flavin as a single subunit. monodehydroascorbate reductase.           |
| BAA05408.1     | D26392   | Cucumis sativus                       | monodehydroascorbate reductase.  |
| AAC41654.1     | L41345   | Lycopersicon esculentum               | ascorbate free radical reductase. AFRR.  |
| AAD53522.1     | AF158602 | Zantedeschia aethiopica               | monodehydroascorbate reductase. MDAR. putative.  |
| AAD28178.1     | AF109695 | Brassica juncea                       | monodehydroascorbate reductase. MDAR1.   |
| BAA77214.1     | D85764   | Oryza sativa                          | cytosolic monodehydroascorbate reductase.  |
| SEQ ID NO: 164 |          |                                       |  |
| CAA04391.1     | AJ000923 | Carica papaya                         | glutathione transferase. PGST1.  |
| AAC18566.1     | AF048978 | Glycine max                           | 2,4-D inducible glutathione S-transferase. GSTa.   |
| AAG34800.1     | AF243365 | Glycine max                           | glutathione S-transferase GST 10.  |
| AAF22647.1     | AF193439 | Lycopersicon esculentum               | glutathione S-transferase/peroxidase. BI-GST/GPX.  |
| CAA71784.1     | Y10820   | Glycine max                           | glutathione transferase.   |
| AAG34799.1     | AF243364 | Glycine max                           | glutathione S-transferase GST 9.   |
| AAG16760.1     | AY007562 | Lycopersicon esculentum               | putative glutathione S-transferase T5.   |
| CAA48717.1     | X68819   | Glycine max                           | lactoylglutathione lyase. glyoxalase I.  |
| CAC24549.1     | AJ296343 | Cichorium intybus x Cichorium endivia | glutathione S-transferase. chi-GST1. auxin-induced GST.  |
| AAC28101.1     | AF079511 | Mesembryanthemum crystallinum         | glutathione S-transferase.   |
| AAG34806.1     | AF243371 | Glycine max                           | glutathione S-transferase GST 16.  |

|            |          |   |
|------------|----------|---|
| AAF23357.1 | AF109194 | Hordeum vulgare<br>glutathione-S-transferase.   |
| CAA73369.1 | Y12862   | Zea mays<br>glutathione transferase. GST5.  |
| AAG32470.1 | AF309377 | Oryza sativa subsp. japonica<br>putative glutathione S-transferase OsGSTU5.   |
| CAA74197.1 | Y13898   | Brassica juncea<br>glutathione-S-transferase. gst.  |
| AAG34827.1 | AF244684 | Zea mays<br>glutathione S-transferase GST 19.   |
| AAC05216.1 | AF050102 | Oryza sativa<br>glutathione s-transferase. GST1.  |
| AAC32118.1 | AF051214 | Picea mariana<br>probable glutathione S-transferase. Sb18. similar to Nicotiana tabacum probable glutathione<br>S-transferase encoded by GenBank Accession Number X56266. |
| AAG16758.1 | AY007560 | Lycopersicon esculentum<br>putative glutathione S-transferase T3.   |
| AAG34798.1 | AF243363 | Glycine max<br>glutathione S-transferase GST 8.   |
| AAG34842.1 | AF244699 | Zea mays<br>glutathione S-transferase GST 34.   |
| AAG34807.1 | AF243372 | Glycine max<br>glutathione S-transferase GST 17.  |
| AAG34809.1 | AF243374 | Glycine max<br>glutathione S-transferase GST 19.  |
| AAG34839.1 | AF244696 | Zea mays<br>glutathione S-transferase GST 31.   |
| AAG34804.1 | AF243369 | Glycine max<br>glutathione S-transferase GST 14.  |
| AAG34844.1 | AF244701 | Zea mays<br>glutathione S-transferase GST 36.   |
| AAG34831.1 | AF244688 | Zea mays<br>glutathione S-transferase GST 23.   |
| AAG34797.1 | AF243362 | Glycine max<br>glutathione S-transferase GST 7.   |
| AAG34832.1 | AF244689 | Zea mays<br>glutathione S-transferase GST 24.   |
| AAG32471.1 | AF309378 | Oryza sativa subsp. japonica<br>putative glutathione S-transferase OsGSTU4.   |
| AAG34796.1 | AF243361 | Glycine max<br>glutathione S-transferase GST 6.   |
| AAG34810.1 | AF243375 | Glycine max<br>glutathione S-transferase GST 20.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAG34835.1     | AF244692 | Zea mays                | glutathione S-transferase GST 27.   |
| AAC32139.1     | AF051238 | Picea mariana           | probable glutathione S-transferase. Sb52. similar to Nicotiana tabacum probable glutathione S-transferase encoded by GenBank Accession Number X56266. |
| SEQ ID NO: 165 |          |                         |   |
| CAA71878.1     | Y10984   | Brassica juncea         | ATP-dependent addition of glycine to gamma-glutamylcysteine. glutathione synthetase. gshII.   |
| AAB71231.1     | AF017984 | Lycopersicon esculentum | glutathione synthetase. GSH2.   |
| AAF98157.1     | AF258320 | Phaseolus vulgaris      | homoglutathione synthetase. hgshs.  |
| AAF98156.1     | AF258319 | Pisum sativum           | putative homoglutathione synthetase. hgshs.   |
| CAB91078.1     | AJ272035 | Glycine max             | homoglutathione synthetase. hGS. putatively predicted to be targetted to the chloroplast.   |
| AAF98121.1     | AF231137 | Pisum sativum           | glutathione synthetase precursor. gshs. putative mitochondrial protein.   |
| AAD29848.1     | AF075699 | Medicago truncatula     | putative glutathione synthetase. GSHS1.   |
| AAD29849.1     | AF075700 | Medicago truncatula     | putative glutathione synthetase. GSHS2.   |
| SEQ ID NO: 166 |          |                         |   |
| BAA83711.1     | AB014484 | Nicotiana tabacum       | heat shock factor. NtHSF2.  |
| AAF37579.1     | AF235958 | Medicago sativa         | heat shock transcription factor. HSFA4-6. MsHSFA4-6.  |
| CAA58117.1     | X82943   | Zea mays                | heat shock factor. hsfb.  |
| CAA47868.1     | X67599   | Lycopersicon esculentum | heat stress transcription factor 8. hsf8.   |
| CAA47869.1     | X67600   | Lycopersicon peruvianum | heat shock transcription factor 8. hsf8.  |
| CAA47870.1     | X67601   | Lycopersicon peruvianum | heat stress transcription factor HSF30. hsf30.  |
| AAF74563.1     | AF208544 | Lycopersicon peruvianum | heat stress transcription factor A3. HSFA3.   |
| CAA87080.1     | Z46956   | Glycine max             | heat shock transcription factor 5. HSF.   |
| CAA87076.1     | Z46952   | Glycine max             | heat shock transcription factor 21. HSF.  |
| CAA39034.1     | X55347   | Lycopersicon peruvianum | heat stress transcription factor. Lp-HSF24.   |

---

BAA83710.1 AB014483 *Nicotiana tabacum*  
heat shock factor. NtHSF1.

---

CAA87077.1 Z46953 *Glycine max*  
heat shock transcription factor 34. HSF. corresponds to longest open reading frame; preceeded by four short open reading frames in the 5' leader sequence.

---

BAB19067.1 AP002744 *Oryza sativa*  
putative heat shock factor protein 1 (HSF 1). P0006C01.9.

---

CAA09301.1 AJ010644 *Pisum sativum*  
heat shock transcription factor (HSFA). hsfA.

---

CAA87079.1 Z46955 *Glycine max*  
heat shock transcription factor 31. HSF.

---

CAA87075.1 Z46951 *Glycine max*  
heat shock transcription factor 29. HSF.

---

CAA09300.1 AJ010643 *Pisum sativum*  
heat shock transcription factor (HSFA). hsfA.

---

SEQ ID NO: 168

---

BAA83710.1 AB014483 *Nicotiana tabacum*  
heat shock factor. NtHSF1.

---

CAA39034.1 X55347 *Lycopersicon peruvianum*  
heat stress transcription factor. Lp-HSF24.

---

CAA87077.1 Z46953 *Glycine max*  
heat shock transcription factor 34. HSF. corresponds to longest open reading frame; preceeded by four short open reading frames in the 5' leader sequence.

---

CAA87080.1 Z46956 *Glycine max*  
heat shock transcription factor 5. HSF.

---

CAA47869.1 X67600 *Lycopersicon peruvianum*  
heat shock transcription factor 8. hsf8.

---

CAA47868.1 X67599 *Lycopersicon esculentum*  
heat stress transcription factor 8. hsf8.

---

BAA83711.1 AB014484 *Nicotiana tabacum*  
heat shock factor. NtHSF2.

---

CAA58117.1 X82943 *Zea mays*  
heat shock factor. hsfB.

---

CAA87075.1 Z46951 *Glycine max*  
heat shock transcription factor 29. HSF.

---

CAA87076.1 Z46952 *Glycine max*  
heat shock transcription factor 21. HSF.

---

CAA47870.1 X67601 *Lycopersicon peruvianum*  
heat stress transcription factor HSF30. hsf30.

---

AAF74563.1 AF208544 *Lycopersicon peruvianum*  
heat stress transcription factor A3. HSFA3.

---

AAF37579.1 AF235958 *Medicago sativa*  
heat shock transcription factor. HSFA4-6. MsHSFA4-6.

---



|                |          |                     |   |
|----------------|----------|---------------------|---|
| CAA87079.1     | Z46955   | Glycine max         | heat shock transcription factor 31. HSF.  |
| BAB19067.1     | AP002744 | Oryza sativa        | putative heat shock factor protein 1 (HSF 1). P0006C01.9.                         |
| CAA09301.1     | AJ010644 | Pisum sativum       | heat shock transcription factor (HSFA). hsfA.                                     |
| CAA09300.1     | AJ010643 | Pisum sativum       | heat shock transcription factor (HSFA). hsfA.                                     |
| SEQ ID NO: 169 |          |                     |   |
| AAB72109.1     | AF022217 | Brassica rapa       | low molecular weight heat-shock protein. BcHSP17.6. 17.6 kDa; cytosolic class I.  |
| CAB93512.1     | AJ243565 | Brassica oleracea   | putative class I small heat shock protein. HSP17.7-a protein. hsp17.7-a.          |
| CAA37847.1     | X53851   | Daucus carota       | heat shock protein.   |
| AAD49336.1     | AF166277 | Nicotiana tabacum   | low molecular weight heat-shock protein. LHS-1. TLHS-1.                           |
| BAA33062.1     | AB017273 | Cuscuta japonica    | low-molecular-weight heat shock protein. CJHSP17.                                 |
| CAB36910.1     | AJ000691 | Quercus suber       | stress protein chaperone. heat shock protein 17.4. hsp17.                         |
| CAA08908.1     | AJ009880 | Castanea sativa     | molecular chaperone. cytosolic class I small heat-shock protein HSP17.5. hsp17.5. |
| AAA33975.1     | M11395   | Glycine max         | small heat shock protein.   |
| CAA25578.1     | X01104   | Glycine max         | heat shock protein 6871 (aa 1-153).   |
| AAB03893.1     | M11318   | Glycine max         | 17.5 kd heat shock protein Gmhsp17.6L.  |
| CAA41547.1     | X58711   | Medicago sativa     | heat shock protein.   |
| AAB63310.1     | U46544   | Helianthus annuus   | 18.6 kDa heat-shock protein. Class I low-molecular-weight heat-shock protein.     |
| CAB08441.1     | Z95153   | Helianthus annuus   | 17.6 kD class I small heat-shock protein HSP17.6. Ha hsp17.6.                     |
| CAA42222.1     | X59701   | Helianthus annuus   | 17.6 kDa heat shock protein.  |
| CAA37848.1     | X53852   | Daucus carota       | heat shock protein.   |
| AAC39360.1     | U63631   | Fragaria x ananassa | LMW heat shock protein.   |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAA33672.1 | M33899   | Pisum sativum           | 18.1 kDa heat shock protein (hsp18.1).  |
| AAB63311.1 | U46545   | Helianthus annuus       | 17.7 kDa heat shock protein. Class I low-molecular-weight heat-shock protein. |
| AAA33974.1 | M11317   | Glycine max             | 17.6 kd heat shock protein Gmhsp17.6L.  |
| CAA63903.1 | X94193   | Pennisetum glaucum      | heat shock protein 17.9. hsp17.9.   |
| AAA61632.1 | U08601   | Papaver somniferum      | low molecular weight heat-shock protein.                                      |
| CAB55634.2 | AJ237596 | Helianthus annuus       | 17.9 kDa heat-shock protein. hsp17.9.   |
| AAC78392.1 | U83669   | Oryza sativa            | low molecular mass heat shock protein Oshsp17.3. OSHSP17.3. class I LMMHSP.   |
| AAA33910.1 | M80939   | Oryza sativa            | 16.9 kDa heat shock protein.  |
| BAA02160.1 | DI2635   | Oryza sativa            | 'low molecular weight heat shock protein'.                                    |
| CAA43210.1 | X60820   | Oryza sativa            | 16.9 KD low molecular weight heat shock protein.                              |
| CAA37864.1 | X53870   | Chenopodium rubrum      | heat-shock protein.   |
| AAA33909.1 | M80938   | Oryza sativa            | 16.9 kDa heat shock protein.  |
| AAC78393.1 | U83670   | Oryza sativa            | low molecular mass heat shock protein Oshsp18.0. OSHSP18.0. class I LMMHSP.   |
| AAB39856.1 | U81385   | Oryza sativa            | heat shock protein. Oshsp16.9C. class I, low molecular mass.                  |
| AAD30454.1 | AF123257 | Lycopersicon esculentum | 17.6 kD class I small heat shock protein. HSP17.6.                            |
| AAA33671.1 | M33900   | Pisum sativum           | 17.9 kDa heat shock protein (hsp17.9).  |
| AAC78394.1 | U83671   | Oryza sativa            | low molecular mass heat shock protein Oshsp17.7. OSHSP17.7. class I LMMHSP.   |
| CAB93514.1 | AJ243567 | Brassica oleracea       | Putative class I small heat shock protein. HSP17.x protein. hsp17.x.          |
| CAA63901.1 | X94191   | Pennisetum glaucum      | heat shock protein 17.0. hsp17.0.   |
| CAA63902.1 | X94192   | Pennisetum glaucum      | heat shock protein 16.9. hsp16.9.   |
| AAD30452.1 | AF123255 | Lycopersicon esculentum | 17.7 kD class I small heat shock protein. HSP17.7.                            |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA46641.1     | X65725   | Zea mays                | heat shock protein 17.2. Zmhsp 17.2. Class I low molecular weight heat shock protein.                              |
| CAA39603.1     | X56138   | Lycopersicon esculentum | small heat shock protein (class I).  |
| CAA63570.1     | X92983   | Pseudotsuga menziesii   | low molecular weight heat-shock protein.   |
| AAD30453.1     | AF123256 | Lycopersicon esculentum | 17.8 kD class I small heat shock protein. HSP17.8.   |
| CAA63571.1     | X92984   | Pseudotsuga menziesii   | low molecular weight heat-shock protein.   |
| CAA31785.1     | X13431   | Triticum aestivum       | put. heat shock protein (AA 1 -151).   |
| CAA53286.1     | X75616   | Oryza sativa            | heat shock protein 17.8.   |
| SEQ ID NO: 170 |          |                         |  |
| AAC14577.1     | U72396   | Lycopersicon esculentum | class II small heat shock protein Le-HSP17.6. heat treatment/chilling tolerance related protein from tomato fruit. |
| AAA33670.1     | M33901   | Pisum sativum           | 17.7 kDa heat shock protein (hsp17.7).   |
| CAA82653.1     | Z29554   | Helianthus annuus       | 17.9 kDa heat-shock protein.   |
| AAD41409.1     | AF159562 | Prunus dulcis           | cytosolic class II low molecular weight heat shock protein. hsp17.5.   |
| CAA65020.1     | X95716   | Petroselinum crispum    | small heat shock protein. cytoplasmic class II HSP.  |
| AAC36312.1     | AF090115 | Lycopersicon esculentum | cytosolic class II small heat shock protein HCT2. HSP17.4.   |
| AAB01561.1     | L47717   | Picea glauca            | heat shock protein 17.0. EMB27.  |
| AAB39336.1     | M99430   | Ipomoea nil             | small heat shock protein.  |
| AAB01562.1     | L47740   | Picea glauca            | class II cytoplasmic small molecular weight heat shock protein 17.1. EMB29, SMW HSP17.1.                           |
| CAA67206.1     | X98617   | Medicago sativa         | 17kD heat shock protein.   |
| BAA99529.1     | AP002484 | Oryza sativa            | putative heat shock protein, 18K - maize. P0489A01.20. contains ESTs C99035(E4351),AU093460(E3974).                |
| CAA41218.1     | X58279   | Triticum aestivum       | heat shock protein 17.3. Tahsp17.3.  |
| CAA67726.1     | X99346   | Picea abies             | small heat shock protein.  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAA38012.1 | X54075   | Zea mays                | 18kDa heat shock protein.   |
| CAA38013.1 | X54076   | Zea mays                | 18kDa heat shock protein.   |
| AAB26481.1 | S59777   | Zea mays                | HSP18. HSP18. 18 kda heat shock protein; This sequence comes from Fig. 2B.    |
| AAB39335.1 | M99429   | Ipomoea nil             | small heat shock protein.   |
| AAD09184.1 | AF089845 | Funaria hygrometrica    | cytosolic II small heat shock protein HSP16.4II. HSP16.4II.                   |
| BAA04841.1 | D21817   | Lilium longiflorum      | small heat shock protein. LIM11.  |
| AAD09185.1 | AF089846 | Funaria hygrometrica    | cytosolic II small heat shock protein HSP18.3II. HSP18.3II.                   |
| BAA04842.1 | D21818   | Lilium longiflorum      | small heat shock protein. LIM12.  |
| CAA63570.1 | X92983   | Pseudotsuga menziesii   | low molecular weight heat-shock protein.                                      |
| CAA63571.1 | X92984   | Pseudotsuga menziesii   | low molecular weight heat-shock protein.                                      |
| AAC39360.1 | U63631   | Fragaria x ananassa     | LMW heat shock protein.   |
| AAD09178.1 | AF087640 | Funaria hygrometrica    | cytosolic I small heat shock protein HSP17.2IA. HSP17.2IA.                    |
| AAD09182.1 | AF089843 | Funaria hygrometrica    | cytosolic I small heat shock protein HSP17.2IC. HSP17.2IC.                    |
| BAA04840.1 | D21816   | Lilium longiflorum      | small heat shock protein. LIM10.  |
| CAB93514.1 | AJ243567 | Brassica oleracea       | Putative class I small heat shock protein. HSP17.x protein. hsp17.x.          |
| AAD30452.1 | AF123255 | Lycopersicon esculentum | 17.7 kD class I small heat shock protein. HSP17.7.                            |
| AAB63311.1 | U46545   | Helianthus annuus       | 17.7 kDa heat shock protein. Class I low-molecular-weight heat-shock protein. |
| CAA31785.1 | X13431   | Triticum aestivum       | put. heat shock protein (AA 1 -151).  |
| CAA63901.1 | X94191   | Pennisetum glaucum      | heat shock protein 17.0. hsp17.0.   |
| AAA33672.1 | M33899   | Pisum sativum           | 18.1 kDa heat shock protein (hsp18.1).  |
| CAA41547.1 | X58711   | Medicago sativa         | heat shock protein.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA46641.1     | X65725   | Zea mays                | heat shock protein 17.2. Zmhspl7.2. Class I low molecular weight heat shock protein.                               |
| AAB39856.1     | U81385   | Oryza sativa            | heat shock protein. Oshsp16.9C. class I, low molecular mass.   |
| AAC78394.1     | U83671   | Oryza sativa            | low molecular mass heat shock protein Oshsp17.7. OSHSP17.7. class I LMMHSP.  |
| AAC78392.1     | U83669   | Oryza sativa            | low molecular mass heat shock protein Oshsp17.3. OSHSP17.3. class I LMMHSP.  |
| CAA39603.1     | X56138   | Lycopersicon esculentum | small heat shock protein (class I).  |
| AAA33910.1     | M80939   | Oryza sativa            | 16.9 kDa heat shock protein.   |
| AAA33909.1     | M80938   | Oryza sativa            | 16.9 kDa heat shock protein.   |
| CAA43210.1     | X60820   | Oryza sativa            | 16.9 KD low molecular weight heat shock protein.   |
| AAD30453.1     | AF123256 | Lycopersicon esculentum | 17.8 kD class I small heat shock protein. HSP17.8.   |
| AAA34294.1     | L14444   | Triticum aestivum       | heat shock protein 16.9C. hsp16.9C.  |
| CAA69172.1     | Y07844   | Hordeum vulgare         | 17 kDa class I small heat shock protein. hsp17. putative.  |
| AAD30454.1     | AF123257 | Lycopersicon esculentum | 17.6 kD class I small heat shock protein. HSP17.6.   |
| SEQ ID NO: 171 |          |                         |  |
| CAA82653.1     | Z29554   | Helianthus annuus       | 17.9 kDa heat-shock protein.   |
| AAC14577.1     | U72396   | Lycopersicon esculentum | class II small heat shock protein Le-HSP17.6. heat treatment/chilling tolerance related protein from tomato fruit. |
| CAA65020.1     | X95716   | Petroselinum crispum    | small heat shock protein. cytoplasmic class II HSP.  |
| AAB39336.1     | M99430   | Ipomoea nil             | small heat shock protein.  |
| AAA33670.1     | M33901   | Pisum sativum           | 17.7 kDa heat shock protein (hsp17.7).   |
| AAD41409.1     | AF159562 | Prunus dulcis           | cytosolic class II low molecular weight heat shock protein. hsp17.5.   |
| CAA67206.1     | X98617   | Medicago sativa         | 17kD heat shock protein.   |
| BAA99529.1     | AP002484 | Oryza sativa            | putative heat shock protein, 18K - maize. P0489A01.20. contains ESTs C99035(E4351),AU093460(E3974).                |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAC36312.1 | AF090115 | Lycopersicon esculentum | cytosolic class II small heat shock protein HCT2. HSP17.4.                               |
| CAA38012.1 | X54075   | Zea mays                | 18kDa heat shock protein.  |
| CAA38013.1 | X54076   | Zea mays                | 18kDa heat shock protein.  |
| CAA41218.1 | X58279   | Triticum aestivum       | heat shock protein 17.3. Tahsp17.3.  |
| AAB26481.1 | S59777   | Zea mays                | HSP18. HSP18. 18 kda heat shock protein; This sequence comes from Fig. 2B.               |
| AAB01561.1 | L47717   | Picea glauca            | heat shock protein 17.0. EMB27.  |
| AAB39335.1 | M99429   | Ipomoea nil             | small heat shock protein.  |
| AAB01562.1 | L47740   | Picea glauca            | class II cytoplasmic small molecular weight heat shock protein 17.1. EMB29, SMW HSP17.1. |
| CAA67726.1 | X99346   | Picea abies             | small heat shock protein.  |
| AAD09184.1 | AF089845 | Funaria hygrometrica    | cytosolic II small heat shock protein HSP16.4II. HSP16.4II.                              |
| BAA04841.1 | D21817   | Lilium longiflorum      | small heat shock protein. LIM11.   |
| BAA04842.1 | D21818   | Lilium longiflorum      | small heat shock protein. LIM12.   |
| AAD09185.1 | AF089846 | Funaria hygrometrica    | cytosolic II small heat shock protein HSP18.3II. HSP18.3II.                              |
| BAA04840.1 | D21816   | Lilium longiflorum      | small heat shock protein. LIM10.   |
| CAA63570.1 | X92983   | Pseudotsuga menziesii   | low molecular weight heat-shock protein.   |
| CAA63571.1 | X92984   | Pseudotsuga menziesii   | low molecular weight heat-shock protein.   |
| AAD30452.1 | AF123255 | Lycopersicon esculentum | 17.7 kD class I small heat shock protein. HSP17.7.                                       |
| AAD09178.1 | AF087640 | Funaria hygrometrica    | cytosolic I small heat shock protein HSP17.2IA. HSP17.2IA.                               |
| AAB63311.1 | U46545   | Helianthus annuus       | 17.7 kDa heat shock protein. Class I low-molecular-weight heat-shock protein.            |
| CAA39603.1 | X56138   | Lycopersicon esculentum | small heat shock protein (class I).  |
| AAC39360.1 | U63631   | Fragaria x ananassa     | LMW heat shock protein.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAA33672.1     | M33899   | Pisum sativum           | 18.1 kDa heat shock protein (hsp18.1).  |
| AAD30453.1     | AF123256 | Lycopersicon esculentum | 17.8 kD class I small heat shock protein. HSP17.8.  |
| AAD30454.1     | AF123257 | Lycopersicon esculentum | 17.6 kD class I small heat shock protein. HSP17.6.  |
| CAA41547.1     | X58711   | Medicago sativa         | heat shock protein.   |
| CAA31785.1     | X13431   | Triticum aestivum       | put. heat shock protein (AA 1 -151).  |
| AAB39856.1     | U81385   | Oryza sativa            | heat shock protein. Oshsp16.9C. class I, low molecular mass.  |
| CAA41546.1     | X58710   | Medicago sativa         | heat shock protein.   |
| CAB93514.1     | AJ243567 | Brassica oleracea       | Putative class I small heat shock protein. HSP17.x protein. hsp17.x.  |
| CAA63901.1     | X94191   | Pennisetum glaucum      | heat shock protein 17.0. hsp17.0.   |
| AAA33910.1     | M80939   | Oryza sativa            | 16.9 kDa heat shock protein.  |
| AAA33909.1     | M80938   | Oryza sativa            | 16.9 kDa heat shock protein.  |
| CAA43210.1     | X60820   | Oryza sativa            | 16.9 KD low molecular weight heat shock protein.  |
| AAB03097.1     | U21723   | Glycine max             | Hsp22.3. Gmhsp22.3. low molecular weight heat shock protein.  |
| AAC78394.1     | U83671   | Oryza sativa            | low molecular mass heat shock protein Oshsp17.7. OSHSP17.7. class I LMMHSP.   |
| AAC78392.1     | U83669   | Oryza sativa            | low molecular mass heat shock protein Oshsp17.3. OSHSP17.3. class I LMMHSP.   |
| AAD09182.1     | AF089843 | Funaria hygrometrica    | cytosolic I small heat shock protein HSP17.2IC. HSP17.2IC.  |
| SEQ ID NO: 173 |          |                         |   |
| AAD30456.1     | AF123259 | Lycopersicon esculentum | heat shock protein 90. HSP90.   |
| AAF31705.1     | AF221856 | Euphorbia esula         | heat-shock protein 80.  |
| AAC32131.1     | AF051230 | Picea mariana           | heat shock protein. Sb40. similar to Oryza sativa heat shock protein 82 encoded by GenBank Accession Number Z11920.   |
| AAF64453.1     | AF239931 | Euphorbia esula         | putative heat-shock protein 90. GRP94; similar to endoplasmic homolog precursor; contains the endoplasmic reticulum targeting sequence KDEL at the 3'-tail. |

|                |          |                     |   |
|----------------|----------|---------------------|---|
| CAA78738.1     | Z15018   | Oryza sativa        | heat shock protein. heat shock protein hsp82. hsp82.  |
| SEQ ID NO: 174 |          |                     |   |
| AAG43546.1     | AF211528 | Nicotiana tabacum   | Avr9/Cf-9 rapidly elicited protein 4. ACRE4. similar to Nicotiana glutinosa Ntr truncated N resistance protein encoded by GenBank Accession Number U15605.              |
| AAA50763.1     | U15605   | Nicotiana glutinosa | virus resistance. N.  |
| CAA08797.1     | AJ009719 | Solanum tuberosum   | disease resistance. NL25. nl25.   |
| CAA08798.1     | AJ009720 | Solanum tuberosum   | disease resistance. NL27. nl27.   |
| AAG09951.1     | AF175388 | Glycine max         | resistance protein LM6.   |
| AAG09954.1     | AF175399 | Glycine max         | resistance protein MG13.  |
| AAD25974.1     | AF093647 | Linum usitatissimum | flax rust resistance protein. L.  |
| AAD25966.1     | AF093639 | Linum usitatissimum | flax rust resistance protein. L.  |
| AAD25969.1     | AF093642 | Linum usitatissimum | flax rust resistance protein. L.  |
| AAD25965.1     | AF093638 | Linum usitatissimum | flax rust resistance protein. L.  |
| AAD25968.1     | AF093641 | Linum usitatissimum | flax rust resistance protein. L.  |
| AAA91021.1     | U27081   | Linum usitatissimum | rust resistance. L6tr. L6. The shorter of two alternate protein products of The L6 gene that results from retention of intron 3 in the mRNA; truncated L6 gene product. |
| AAA91022.1     | U27081   | Linum usitatissimum | rust resistance. L6. L6. The longer of two alternate proteins encoded by the L6 gene.   |
| AAD25967.1     | AF093640 | Linum usitatissimum | flax rust resistance protein. L.  |
| AAK28803.1     | AF310958 | Linum usitatissimum | resistance-like protein P1-A. p1-A.   |
| AAK28808.1     | AF310961 | Linum usitatissimum | resistance-like protein P3-A. p3-A.   |
| AAK28805.1     | AF310960 | Linum usitatissimum | resistance-like protein P2-A. p2-A.   |
| AAD25976.1     | AF093649 | Linum usitatissimum | flax rust resistance protein. L.  |
| AAD25973.1     | AF093646 | Linum usitatissimum | flax rust resistance protein. L.  |



|            |          |   |
|------------|----------|---|
| AAD25972.1 | AF093645 | Linum usitatissimum<br>flax rust resistance protein. L.   |
| AAD25971.1 | AF093644 | Linum usitatissimum<br>flax rust resistance protein. L.   |
| AAD25970.1 | AF093643 | Linum usitatissimum<br>flax rust resistance protein. L.   |
| AAD25975.1 | AF093648 | Linum usitatissimum<br>flax rust resistance protein. L.   |
| AAG48132.1 | AF322632 | Glycine max<br>putative resistance protein. L20a.   |
| AAG01052.1 | AF175395 | Glycine max<br>resistance protein MG23.   |
| CAC35330.1 | AJ310155 | Linum usitatissimum<br>N1-D protein. N1-D. N locus resistance gene homolog: TIR-NBS-LRR protein.        |
| CAC35333.1 | AJ310158 | Linum usitatissimum<br>N2-C protein. N2-C. N locus rust resistance gene homolog: TIR-NBS-LRR protein.   |
| AAG48133.1 | AF322633 | Glycine max<br>putative resistance protein. L33.  |
| AAF61452.1 | AF139523 | Tagetes erecta<br>disease-resistance protein NRSA1.   |
| CAC35334.1 | AJ310159 | Linum usitatissimum<br>N2-D protein. N2-D. N locus rust resistance gene homolog: TIR-NBS-LRR protein.   |
| CAC35337.1 | AJ310162 | Linum usitatissimum<br>Nbi-C protein. Nbi-C. N locus rust resistance gene homolog: TIR-NBS-LRR protein. |
| CAC35339.1 | AJ310164 | Linum usitatissimum<br>Nho-C protein. Nho-C. N locus rust resistance gene homolog: TIR-NBS-LRR protein. |
| CAC35326.1 | AJ310151 | Linum usitatissimum<br>Ngc-C protein. Ngc-C. N rust resistance gene homolog: TIR-NBS-LRR protein.       |
| AAG01051.1 | AF175394 | Glycine max<br>resistance protein LM12.   |
| CAC35338.1 | AJ310163 | Linum usitatissimum<br>Nbi-D protein. Nbi-D. N locus rust resistance gene homolog: TIR-NBS-LRR protein. |
| CAC35329.1 | AJ310154 | Linum usitatissimum<br>N1-C protein. N1-C. N locus resistance gene homolog: TIR-NBS-LRR protein.        |
| AAK28804.1 | AF310959 | Linum usitatissimum<br>resistance-like protein P1-B. p1-B.  |
| AAK28809.1 | AF310962 | Linum usitatissimum<br>resistance-like protein P3-B. p3-B.  |
| CAC35332.1 | AJ310157 | Linum usitatissimum<br>N2-B protein. N2-B. N locus rust resistance gene homolog: TIR-NBS-LRR protein.   |
| CAC35328.1 | AJ310153 | Linum usitatissimum<br>N1-B protein. N1-B. N locus resistance gene homolog: TIR-NBS-LRR protein.        |

|  |          |                     |
|--|----------|---------------------|
| CAC35325.1   | AJ310150 | Linum usitatissimum |
| Ngc-B protein. Ngc-B. N rust resistance gene homolog: TIR-NBS-LRR protein.   |          |                     |
| CAC35336.1   | AJ310161 | Linum usitatissimum |
| Nbi-B protein. Nbi-B. N locus rust resistance gene homolog: TIR-NBS-LRR protein.                                   |          |                     |
| CAC35321.1   | AJ310150 | Linum usitatissimum |
| Ngc-D protein. Ngc-D. N rust resistance gene: TIR-NBS-LRR protein.   |          |                     |
| CAC35331.1   | AJ310156 | Linum usitatissimum |
| N2-A protein. N2-A. N locus rust resistance gene homolog: TIR-NBS-LRR protein.                                     |          |                     |
| CAC35323.1   | AJ310150 | Linum usitatissimum |
| Ngc-A protein. Ngc-A. N rust resistance gene homolog: TIR-NBS-LRR protein.   |          |                     |
| AAB47618.1   | U73916   | Linum usitatissimum |
| rust resistance protein M. nucleotide-binding site, leucine-rich repeat class of plant disease resistance protein. |          |                     |
| AAG01053.1   | AF175396 | Glycine max         |
| resistance protein MG55.   |          |                     |
| CAC35327.1   | AJ310152 | Linum usitatissimum |
| N1-A protein. N1-A. N locus resistance gene homolog: TIR-NBS-LRR protein.  |          |                     |
| AAK28811.1   | AF310966 | Linum usitatissimum |
| resistance-like protein P-B. p-B.  |          |                     |
| SEQ ID NO: 175   |          |                     |
| CAB52796.1   | AJ245861 | Solanum tuberosum   |
| respiratory chain. putative internal rotenone-insensitive NADH dehydrogenase. nda1.                                |          |                     |
| CAB52797.1   | AJ245862 | Solanum tuberosum   |
| respiratory chain. putative external rotenone-insensitive NADH dehydrogenase. ndb1.                                |          |                     |
| SEQ ID NO: 176   |          |                     |
| BAA05648.1   | D26601   | Nicotiana tabacum   |
| protein kinase.  |          |                     |
| CAA08997.1   | AJ010093 | Brassica napus      |
| MAP3K beta 1 protein kinase. MAP3K beta 1.   |          |                     |
| CAA08995.1   | AJ010091 | Brassica napus      |
| MAP3K alpha 1 protein kinase. MAP3K alpha 1.   |          |                     |
| AAF34436.1   | AF172282 | Oryza sativa        |
| similar to mitogen-activated protein kinases. DUPR11.32.   |          |                     |
| CAB54520.1   | AJ238845 | Brassica napus      |
| putative role in cell cycle control. MAP3K epsilon 1 protein kinase. MAP3Ke1.                                      |          |                     |
| CAA08758.1   | AJ009609 | Brassica napus      |
| BnMAP4K alpha2.  |          |                     |
| CAA08757.1   | AJ009608 | Brassica napus      |
| BnMAP4K alpha1.  |          |                     |
| AAC83393.1   | U83625   | Zea mays            |
| protein kinase ZmMEK1. mitogen-activated; ERK-activating protein kinase (MEK) homolog.                             |          |                     |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAG53979.1 | AF325168 | Nicotiana tabacum       | mitogen-activated protein kinase 2. MEK2. upstream kinase for SIPK and WIPK, two tobacco MAP kinases.  |
| AAG40578.1 | AF216314 | Oryza sativa            | MAP kinase kinase 1. protein kinase; MEK1.   |
| CAA04261.2 | AJ000728 | Lycopersicon esculentum | MAP kinase kinase. mek1.   |
| AAF67262.1 | AF165186 | Nicotiana tabacum       | MAP kinase kinase.   |
| AAG45491.1 | AY013245 | Oryza sativa            | 36I5.3. putative serine/threonine kinase.  |
| BAB32405.1 | AB055514 | Nicotiana tabacum       | NQK1 MAPKK. nqk1.  |
| AAG49001.1 | AY013246 | Hordeum vulgare         | putative serine/threonine kinase. 635P2.3.   |
| AAF19403.1 | AF203481 | Lycopersicon esculentum | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.   |
| AAF19402.1 | AF203480 | Lycopersicon esculentum | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.   |
| CAC24705.1 | AJ302651 | Nicotiana tabacum       | protein kinase. MAP kinase. mek1.  |
| BAA06731.1 | D31964   | Nicotiana tabacum       | NPK2. protein kinase.  |
| AAG31141.1 | AF305911 | Oryza sativa            | EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.  |
| AAF19401.1 | AF203479 | Glycine max             | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.   |
| AAD46406.1 | AF096250 | Lycopersicon esculentum | ethylene-responsive protein kinase TCTR1. ER50. serine/threonine kinase; similar to Arabidopsis thaliana negative regulator of the ethylene response pathway encoded by GenBank Accession Number L08789. |
| CAA73722.1 | Y13273   | Lycopersicon esculentum | putative protein kinase.   |
| BAA05649.1 | D26602   | Nicotiana tabacum       | protein kinase.  |
| AAD23582.1 | AF128443 | Glycine max             | probably involved in plant stress responses possibly regulates gene expression. SNF-1-like serine/threonine protein kinase. expressed in nodules, roots and leaves.                                      |
| CAA06334.1 | AJ005077 | Lycopersicon esculentum | protein kinase. TCTR2 protein. TCTR2.  |

|            |          |   |
|------------|----------|---|
| AAD10057.1 | AF110519 | <i>Lycopersicon esculentum</i><br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1v.  |
| CAB89082.1 | AJ277534 | <i>Asparagus officinalis</i><br>S6 ribosomal protein kinase. pk1. putative.   |
| AAC78558.1 | AF030879 | <i>Solanum tuberosum</i><br>protein kinase CPK1.  |
| AAD10056.1 | AF110518 | <i>Lycopersicon esculentum</i><br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1.   |
| AAK18832.1 | AC082645 | <i>Oryza sativa</i><br>putative protein kinase. OSJNBb0033N16.9.  |
| AAG31142.1 | AF305912 | <i>Hordeum vulgare</i><br>EDR1. EDR1. MAP kinase kinase kinase; similar to <i>Arabidopsis thaliana</i> EDR1.  |
| AAC24961.1 | AF009337 | <i>Tradescantia virginiana</i><br>CDPK-related protein kinase. CRK1.  |
| BAB21278.1 | AP002844 | <i>Oryza sativa</i><br>putative MAP kinase. P0410E03.9.   |
| AAA61682.1 | L27484   | <i>Zea mays</i><br>calcium-dependent protein kinase. CDPK.  |
| CAA56313.1 | X79992   | <i>Avena sativa</i><br>putative pp70 ribosomal protein S6 kinase. Aspk11.   |
| AAD31900.1 | AF145482 | <i>Mesembryanthemum crystallinum</i><br>putative serine/threonine protein kinase.   |
| AAC25423.1 | AF072908 | <i>Nicotiana tabacum</i><br>calcium-dependent protein kinase. CDPK1.  |
| CAA71142.1 | Y10036   | <i>Cucumis sativus</i><br>SNF1-related protein kinase.  |
| BAA83689.1 | AB011968 | <i>Oryza sativa</i><br>OsPK7. OsPK7. protein kinase.  |
| CAA46554.1 | X65604   | <i>Hordeum vulgare</i><br>protein kinase. BKIN12.   |
| CAA46556.1 | X65606   | <i>Hordeum vulgare</i><br>protein kinase. BKIN12.   |
| CAA65500.1 | X96723   | <i>Medicago sativa</i><br>protein kinase. CDPK.   |
| AAF05112.1 | AF158091 | <i>Mesembryanthemum crystallinum</i><br>day/night regulation of carbon fixation by crassulacean acid metabolism pathway.<br>phosphoenolpyruvate carboxylase-kinase. SNIK. protein kinase; salt/night induced kinase;<br>phosphorylase; member of Ca <sup>2+</sup> /Cam protein kinase family; lacks both CAM domain and<br>autoinhibitory domain; Ser/Thr kinase. |
| AAD17800.1 | AF090835 | <i>Mesembryanthemum crystallinum</i><br>Ca <sup>2+</sup> -dependent protein kinase. CPK1. serine/threonine protein kinase.  |

CAA07813.1 AJ007990 *Hordeum vulgare*  
SnRK1-type protein kinase. kin12a.

SEQ ID NO: 177

BAB16335.1 AP002818 *Oryza sativa*  
putative NAM protein. P0436E04.18. contains ESTs E10793(C19698),E10793(C99379).

BAA84803.1 AP000559 *Oryza sativa*  
Similar to NAM like protein (AC005310).

BAB19365.1 AP002542 *Oryza sativa*  
putative NAM (no apical meristem) protein. P0679C08.4.

BAB16328.1 AP002818 *Oryza sativa*  
putative NAM protein. P0436E04.11. contains ESTs R4069(AU032425),R4069(AU082730).

CAA63102.2 X92205 *Petunia x hybrida*  
apical meristem formation. NAM.

CAA63101.1 X92204 *Petunia x hybrida*  
apical meristem formation. NAM.

AAK13151.1 AC078829 *Oryza sativa*  
putative NAM (no apical meristem) protein. OSJNBa0026O12.6.

BAB03447.1 AP002817 *Oryza sativa*  
ESTs C96615(C10106),C26336(C12127),D21959(C10106) correspond to a region of the predicted gene. Similar to *Arabidopsis thaliana* DNA chromosome 4, BAC clone T6K21; NAM (no apical meristem) - like protein (AL021889).

BAA92400.1 AP001366 *Oryza sativa*  
ESTs C96615(C10106),C26336(C12127),D21959(C10106) correspond to a region of the predicted gene. Similar to NAM (AL021889).

SEQ ID NO: 178

BAA25434.1 AB000708 *Raphanus sativus*  
SAUR.

AAG14454.1 AF283706 *Tulipa gesneriana*  
auxin-induced protein TGSAUR12. SAUR12. small auxin upregulated RNA.

AAG14455.1 AF283707 *Tulipa gesneriana*  
auxin-induced protein TGSAUR21. SAUR21. small auxin upregulated RNA.

AAG14456.1 AF283708 *Tulipa gesneriana*  
auxin-induced protein TGSAUR22. SAUR22. small auxin upregulated RNA.

SEQ ID NO: 179

BAA78738.1 AB023482 *Oryza sativa*  
EST AU055776(S20048) corresponds to a region of the predicted gene.; Similar to *Arabidopsis thaliana* AP2 domain containing protein RAP2.10 mRNA, partial cds.(AF003103).

CAB96899.1 AJ251249 *Catharanthus roseus*  
transcription factor. AP2-domain DNA-binding protein. orca3.

CAB96900.1 AJ251250 *Catharanthus roseus*  
transcription factor. AP2-domain DNA-binding protein. orca3.

|            |          |  |
|------------|----------|--|
| AAF76898.1 | AF274033 | <i>Atriplex hortensis</i><br>apetala2 domain-containing protein.   |
| CAC12822.1 | AJ299252 | <i>Nicotiana tabacum</i><br>AP2 domain-containing transcription factor. ap2.   |
| AAC14323.1 | AF058827 | <i>Nicotiana tabacum</i><br>TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.   |
| BAA97123.1 | AB016265 | <i>Nicotiana sylvestris</i><br>ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3. |
| CAB93940.1 | AJ238740 | <i>Catharanthus roseus</i><br>putative transcription factor. AP2-domain DNA-binding protein. orca2.  |
| AAC24587.1 | AF071893 | <i>Prunus armeniaca</i><br>AP2 domain containing protein. AP2DCP.  |
| BAB03248.1 | AB037183 | <i>Oryza sativa</i><br>ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.   |
| BAB16083.1 | AB036883 | <i>Oryza sativa</i><br>transcriptional repressor. osERF3. osERF3. ERF protein family ERF3 associated repression domain.  |
| BAA76734.1 | AB024575 | <i>Nicotiana tabacum</i><br>ethylene responsive element binding factor.  |
| CAB93939.1 | AJ238739 | <i>Catharanthus roseus</i><br>putative transcription factor. AP2-domain DNA-binding protein. orca1.  |
| AAG43545.1 | AF211527 | <i>Nicotiana tabacum</i><br>Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.   |
| AAG43548.1 | AF211530 | <i>Nicotiana tabacum</i><br>Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.   |
| AAG43549.1 | AF211531 | <i>Nicotiana tabacum</i><br>Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.   |
| AAF23899.1 | AF193803 | <i>Oryza sativa</i><br>transcription factor EREBP1. EREBP/AP2-like transcription factor.   |
| AAK31271.1 | AC079890 | <i>Oryza sativa</i><br>putative transcriptional factor. OSJNBb0089A17.22.  |
| AAF63205.1 | AF245119 | <i>Mesembryanthemum crystallinum</i><br>AP2-related transcription factor. CDBP. stress induced transcription factor.   |
| BAA07321.1 | D38123   | <i>Nicotiana tabacum</i><br>ERF1. ethylene-responsive transcription factor.  |
| BAA97122.1 | AB016264 | <i>Nicotiana sylvestris</i><br>ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2. |
| BAA97124.1 | AB016266 | <i>Nicotiana sylvestris</i><br>ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4. |

|                |          |                   |  |
|----------------|----------|-------------------|--|
| BAA94514.2     | AP001800 | Oryza sativa      | Similar to Arabidopsis thaliana chromosome 4, BAC clone F9D16; putative Ap2 domain protein (AL035394).   |
| AAC62619.1     | AF057373 | Nicotiana tabacum | transcription factor. ethylene response element binding protein 1. EREBP1.   |
| AAK01088.1     | AF298230 | Hordeum vulgare   | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.   |
| BAA99376.1     | AP002526 | Oryza sativa      | ESTs AU093391(E60370), AU091593(C60458), AU093392(E60370) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana BAC F21J9; AP2 domain protein. (AC000103). |
| AAK01089.1     | AF298231 | Hordeum vulgare   | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.  |
| AAG59618.1     | AF239616 | Hordeum vulgare   | CRT/DRE-binding factor. CBF.   |
| SEQ ID NO: 181 |          |                   |  |
| CAB71134.1     | AJ271667 | Cicer arietinum   | putative proteasome regulatory subunit.  |
| SEQ ID NO: 185 |          |                   |  |
| BAA85440.1     | AP000616 | Oryza sativa      | ESTs AU055729(S20023), AU055730(S20023) correspond to a region of the predicted gene.; similar to Medicago nodulin N21-like protein (AC004218).                                    |
| CAB53493.1     | AJ245900 | Oryza sativa      | CAA303720.1 protein. q3037.20. Similar to Medicago nodulin N21 (MtN21).  |
| BAB17350.1     | AP002747 | Oryza sativa      | putative nodulin. P0698G03.34. contains ESTs D39891(S1543), D41717(S4395), AU033037(S1543).  |
| SEQ ID NO: 186 |          |                   |  |
| BAA02724.1     | D13506   | Glycine max       | early nodulin.   |
| BAA33816.1     | AB018378 | Glycine max       | early nodulin. GmENOD93.   |
| BAA83560.1     | AP000399 | Oryza sativa      | EST AU077941(C12908) corresponds to a region of the predicted gene. Similar to OsENOD93a gene for early nodulin (AB018375).  |
| BAA33815.1     | AB018377 | Oryza sativa      | early nodulin. OsENOD93b.  |
| BAA83566.1     | AP000399 | Oryza sativa      | ESTs C98280(C1391), D15843(C1391) correspond to a region of the predicted gene. Similar to OsENOD93a gene for early nodulin (AB018375).  |
| BAA83568.1     | AP000399 | Oryza sativa      | EST AU077972(C53511) corresponds to a region of the predicted gene. Similar to OsENOD93a gene for early nodulin (AB018375).  |

|                |          |                    |  |
|----------------|----------|--------------------|--|
| BAA83565.1     | AP000399 | Oryza sativa       | ESTs C98096(C0688),C98097(C0688) correspond to a region of the predicted gene. Similar to OsENOD93a gene for early nodulin (AB018375).               |
| AAD30134.1     | AF140229 | Oryza sativa       | early nodulin.   |
| BAA33814.1     | AB018376 | Oryza sativa       | early nodulin. OsENOD93a.  |
| BAA33813.1     | AB018375 | Oryza sativa       | early nodulin. OsENOD93a.  |
| BAA83567.1     | AP000399 | Oryza sativa       | EST C97982(C0324) corresponds to a region of the predicted gene. Similar to OsENOD93a gene for early nodulin (AB018375).                             |
| BAA83559.1     | AP000399 | Oryza sativa       | EST C97982(C0324) corresponds to a region of the predicted gene. Similar to OsENOD93a gene for early nodulin (AB018375).                             |
| BAA83557.1     | AP000399 | Oryza sativa       | Similar to OsENOD93a gene for early nodulin (AB018375).  |
| SEQ ID NO: 188 |          |                    |  |
| AAA33811.1     | L02830   | Solanum tuberosum  | calcium-binding protein.   |
| AAG43547.1     | AF211529 | Nicotiana tabacum  | Avr9/Cf-9 rapidly elicited protein 31. ACRE31. similar to Solanum tuberosum CAST calcium binding protein encoded by GenBank Accession Number L02830. |
| CAB63264.1     | AJ251808 | Lotus japonicus    | calcium-binding protein. cbp1.   |
| AAA34015.1     | L01433   | Glycine max        | calcium-binding regulatory protein. calmodulin. SCaM-4. putative.  |
| AAA92677.1     | U13736   | Pisum sativum      | binds calcium. calmodulin-like protein.  |
| AAF31152.1     | AF078680 | Olea europaea      | calcium-binding protein. PCA23. Pca23.   |
| AAA33948.1     | L19359   | Glycine max        | calcium-binding regulatory protein. calmodulin. SCaM-5. putative.  |
| AAD10245.1     | AF030033 | Phaseolus vulgaris | calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal transduction pathways.                                      |
| AAF31151.1     | AF078679 | Olea europaea      | calcium-binding protein. PCA18. Pca18.   |
| AAA19571.1     | U10150   | Brassica napus     | calcium binding. calmodulin. bcm1.   |
| AAC49587.1     | U49105   | Triticum aestivum  | calmodulin TaCaM4-1. calcium-binding protein.  |



|                |          |   |
|----------------|----------|---|
| AAC49586.1     | U49104   | Triticum aestivum<br>calmodulin TaCaM3-3. calcium-binding protein.                  |
| AAC49585.1     | U49103   | Triticum aestivum<br>calmodulin TaCaM3-2. calcium-binding protein.                  |
| AAC49584.1     | U48693   | Triticum aestivum<br>calmodulin TaCaM3-1. calcium-binding protein.                  |
| AAC49580.1     | U48689   | Triticum aestivum<br>calmodulin TaCaM1-3. calcium-binding protein.                  |
| AAC49579.1     | U48688   | Triticum aestivum<br>calmodulin TaCaM1-2. calcium binding protein.                  |
| AAC49578.1     | U48242   | Triticum aestivum<br>calmodulin TaCaM1-1. calcium-binding.                          |
| AAA85157.1     | U20297   | Solanum tuberosum<br>calcium-binding protein. calmodulin.                           |
| AAA85156.1     | U20296   | Solanum tuberosum<br>calcium-binding protein. calmodulin.                           |
| AAA62351.1     | U20295   | Solanum tuberosum<br>calcium-binding protein. calmodulin.                           |
| AAA85155.1     | U20294   | Solanum tuberosum<br>calcium-binding protein. calmodulin.                           |
| AAA34014.1     | L01432   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-3. putative.    |
| AAA03580.1     | L01431   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-2. putative.    |
| AAA34013.1     | L01430   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-1. putative.    |
| AAA33901.1     | L18913   | Oryza sativa<br>calcium binding protein, signal transduction. calmodulin. putative. |
| AAA33900.1     | L18914   | Oryza sativa<br>calcium binding protein, signal transduction. calmodulin.           |
| AAA92681.1     | U13882   | Pisum sativum<br>calcium-binding protein. calmodulin.                               |
| CAA78288.1     | Z12828   | Oryza sativa<br>calcium binding protein, signal transduction. calmodulin.           |
| SEQ ID NO: 190 |          |   |
| AAF72197.1     | AF263737 | Euphorbia esula<br>glutathione S-transferase. theta class GST.                      |
| AAG34815.1     | AF243380 | Glycine max<br>glutathione S-transferase GST 25.                                    |
| AAG34825.1     | AF244682 | Zea mays<br>glutathione S-transferase GST 17.                                       |

|            |          |   |
|------------|----------|---|
| AAG32474.1 | AF309381 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTZ1.   |
| AAG34826.1 | AF244683 | <i>Zea mays</i><br>glutathione S-transferase GST 18.  |
| AAA33277.1 | M64268   | <i>Dianthus caryophyllus</i><br>glutathione transferase. CARSR8.  |
| CAA41279.1 | X58390   | <i>Dianthus caryophyllus</i><br>glutathione s-transferase. CARSR8.  |
| AAD09190.1 | AF109714 | <i>Triticum aestivum</i><br>glutathione S-transferase. GST.   |
| AAB60886.1 | AF002211 | <i>Triticum aestivum</i><br>glutathione-S-transferase.  |
| AAA51450.1 | L05916   | <i>Dianthus caryophyllus</i><br>glutathione s-transferase. GST2.  |
| AAC50036.1 | U42463   | <i>Coccothryxa</i> sp. PA<br>glutathione S-transferase. GST.  |
| AAG34801.1 | AF243366 | <i>Glycine max</i><br>glutathione S-transferase GST 11.   |
| AAG32469.1 | AF309376 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTU6.   |
| AAG34850.1 | AF244707 | <i>Zea mays</i><br>glutathione S-transferase GST 42.  |
| AAG34807.1 | AF243372 | <i>Glycine max</i><br>glutathione S-transferase GST 17.   |
| CAB38119.1 | AJ010296 | <i>Zea mays</i><br>Glutathione transferase III(b). gst3b.   |
| AAF23357.1 | AF109194 | <i>Hordeum vulgare</i><br>glutathione-S-transferase.  |
| AAG34849.1 | AF244706 | <i>Zea mays</i><br>glutathione S-transferase GST 41.  |
| CAB38118.1 | AJ010295 | <i>Zea mays</i><br>Glutathione transferase III(a). gst3a.   |
| AAG32475.1 | AF309382 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTF5.   |
| AAB65163.1 | AF002692 | <i>Solanum commersonii</i><br>glutathione S-transferase, class-phi. GST1. low temperature induced.  |
| AAF29773.1 | AF159229 | <i>Gossypium hirsutum</i><br>glutathione S-transferase. GST.  |
| BAB39927.1 | AP002914 | <i>Oryza sativa</i><br>putative glutathione S-transferase. P0493G01.1. contains ESTs<br>AU031696(R0596), C97559(C60386), C28218(C60386), D28287(R0596). |
| AAG32473.1 | AF309380 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTU2.   |

|            |          |   |
|------------|----------|---|
| CAA09188.1 | AJ010449 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1b.  |
| AAG34846.1 | AF244703 | <i>Zea mays</i><br>glutathione S-transferase GST 38.  |
| AAG34812.1 | AF243377 | <i>Glycine max</i><br>glutathione S-transferase GST 22.   |
| CAA55039.1 | X78203   | <i>Hyoscyamus muticus</i><br>glutathione transferase.   |
| AAG34803.1 | AF243368 | <i>Glycine max</i><br>glutathione S-transferase GST 13.   |
| CAA09189.1 | AJ010450 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1c.  |
| AAG34808.1 | AF243373 | <i>Glycine max</i><br>glutathione S-transferase GST 18.   |
| AAG34813.1 | AF243378 | <i>Glycine max</i><br>glutathione S-transferase GST 23.   |
| AAD10129.1 | AF004358 | <i>Aegilops tauschii</i><br>chloroacetamide herbicide metabolism. glutathione S-transferase TSI-1. GST isozyme. |
| AAG41204.1 | AF321437 | <i>Suaeda maritima</i><br>glutathione transferase.  |
| AAG34810.1 | AF243375 | <i>Glycine max</i><br>glutathione S-transferase GST 20.   |
| AAG34804.1 | AF243369 | <i>Glycine max</i><br>glutathione S-transferase GST 14.   |
| AAG34814.1 | AF243379 | <i>Glycine max</i><br>glutathione S-transferase GST 24.   |
| AAG32472.1 | AF309379 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTU3.                       |
| AAG32471.1 | AF309378 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTU4.                       |
| CAA09187.1 | AJ010448 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1a.  |
| AAG34835.1 | AF244692 | <i>Zea mays</i><br>glutathione S-transferase GST 27.  |
| AAF64449.1 | AF239927 | <i>Euphorbia esula</i><br>glutathione S-transferase. theta class GST.   |
| AAG16758.1 | AY007560 | <i>Lycopersicon esculentum</i><br>putative glutathione S-transferase T3.  |
| AAG34848.1 | AF244705 | <i>Zea mays</i><br>glutathione S-transferase GST 40.  |

SEQ ID NO: 191

|                |          |                                 |  |
|----------------|----------|---------------------------------|--|
| AAD32141.1     | AF123503 | Nicotiana tabacum               | Nt-gh3 deduced protein.  |
| CAA42636.1     | X60033   | Glycine max                     | auxin-responsive GH3 product. GH3.   |
| BAA96221.1     | AP002094 | Oryza sativa                    | ESTs C19814(E10971),AU090481(E10971) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, sequence from clones T3F17, F11C10, F13A10; auxin-responsive GH3-like protein (AC006526). |
| SEQ ID NO: 192 |          |                                 |  |
| AAG13589.1     | AC051633 | Oryza sativa                    | putative ubiquitin protein. OSJNBb0015I11.23.  |
| SEQ ID NO: 193 |          |                                 |  |
| CAC09348.1     | AL442007 | Oryza sativa                    | putative phosphoglycerate dehydrogenase. H0212B02.4.   |
| CAA79702.2     | Z21493   | Solanum tuberosum               | mitochondrial formate dehydrogenase precursor.   |
| BAA77337.1     | AB019533 | Oryza sativa                    | Nad-dependent formate dehydrogenase.   |
| BAA36181.1     | D88272   | Hordeum vulgare                 | formate dehydrogenase.   |
| SEQ ID NO: 194 |          |                                 |  |
| AAD46412.1     | AF096262 | Lycopersicon esculentum         | ER6 protein. ethylene-inducible; similar to sequence F21M12 from Arabidopsis thaliana encoded by GenBank Accession Number AC000132.  |
| SEQ ID NO: 195 |          |                                 |  |
| AAK13154.1     | AC078829 | Oryza sativa                    | putative casein kinase. OSJNBa0026O12.5.   |
| BAA92986.1     | AP001550 | Oryza sativa                    | ESTs D41826(S4655),C22685(S4655) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana chromosome 4, BAC clone F16A16; protein kinase-like protein (AL035353).                                    |
| AAF19807.1     | AF180356 | Brassica oleracea               | casein kinase I-like protein. CK1b. strong similarity to Arabidopsis thaliana casein kinase 1.   |
| AAD20819.1     | AF107592 | Dendrobium grex Madame Thong-In | putative casein kinase I. otg16.   |
| AAF19403.1     | AF203481 | Lycopersicon esculentum         | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.   |
| AAF19402.1     | AF203480 | Lycopersicon esculentum         | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.   |
| BAA05648.1     | D26601   | Nicotiana tabacum               | protein kinase.  |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAF23901.2 | AF194414 | Oryza sativa                  | calcium-dependent protein kinase. CDPK5. OsCDPK5.   |
| AAC04324.1 | U73937   | Nicotiana tabacum             | ethylene signal transduction. PK12 protein kinase. PK12. component of the LAMMER family of protein kinases; dual-specificity protein kinase.                        |
| AAF23900.1 | AF194413 | Oryza sativa                  | calcium-dependent protein kinase. CDPK1. OsCDPK1.   |
| BAA34675.1 | AB011670 | Triticum aestivum             | wpk4 protein kinase. wpk4.  |
| BAA13440.1 | D87707   | Ipomoea batatas               | calcium dependent protein kinase. CDPK.   |
| AAD17800.1 | AF090835 | Mesembryanthemum crystallinum | Ca <sup>2+</sup> -dependent protein kinase. CPK1. serine/threonine protein kinase.  |
| CAA57157.1 | X81394   | Oryza sativa                  | calcium-dependent protein kinase. OSCPK2.   |
| AAD23582.1 | AF128443 | Glycine max                   | probably involved in plant stress responses possibly regulates gene expression. SNF-1-like serine/threonine protein kinase. expressed in nodules, roots and leaves. |
| CAA39936.1 | X56599   | Daucus carota                 | calcium- dependent protein kinase. DcPK431.   |
| BAA12715.1 | D85039   | Zea mays                      | calcium-dependent protein kinase.   |
| CAA58750.1 | X83869   | Daucus carota                 | CDPK-related protein kinase. CRK (or PK421).  |
| AAB80693.1 | U69174   | Glycine max                   | calmodulin-like domain protein kinase isoenzyme gamma. CDPK gamma.  |
| AAF21062.1 | AF216527 | Dunaliella tertiolecta        | calcium-dependent protein kinase. CPK1; CDPK.   |
| CAA74646.1 | Y14274   | Sorghum bicolor               | putative serine/threonine protein kinase. SNFL3.  |
| AAD28192.2 | AF115406 | Solanum tuberosum             | calcium-dependent protein kinase. CDPK; catalytic domain.   |
| BAA05649.1 | D26602   | Nicotiana tabacum             | protein kinase.   |
| AAA69507.1 | U28376   | Zea mays                      | calcium-dependent protein kinase. MZECDPK2.   |
| AAB05457.1 | U55768   | Oryza sativa                  | SNF1-related protein kinase. RSk1. Ser/Thr protein kinase homolog.  |
| AAG36872.1 | AF239819 | Zea mays                      | protein kinase CK2 catalytic subunit CK2 alpha-3.   |
| CAA89202.1 | Z49233   | Chlamydomonas eugametos       | calcium-stimulated protein kinase.  |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| CAA72362.1     | Y11649   | Zea mays                      | protein kinase CK2, alpha subunit.  |
| CAA72290.1     | Y11526   | Zea mays                      | casein kinase II alpha subunit. CK2.  |
| CAA43659.1     | X61387   | Zea mays                      | casein kinase II alpha subunit. ZMACK2.   |
| CAA65244.1     | X95997   | Solanum tuberosum             | SNF1-related protein kinase. PKIN1.   |
| AAF76187.1     | AF271237 | Zea mays                      | casein kinase II alpha subunit.   |
| BAB21591.1     | AB036788 | Oryza sativa                  | casein kinase II alpha subunit. OSCKA2.   |
| BAB21589.1     | AB036786 | Oryza sativa                  | casein kinase II alpha subunit. OSCKA2.   |
| CAB89082.1     | AJ277534 | Asparagus officinalis         | S6 ribosomal protein kinase. pk1. putative.   |
| AAF06970.1     | AF162662 | Kalanchoe fedtschenkoi        | phosphoenolpyruvate carboxylase kinase. calcium-independent protein kinase.             |
| AAF06969.1     | AF162661 | Kalanchoe fedtschenkoi        | phosphoenolpyruvate carboxylase kinase. calcium-independent protein kinase.             |
| CAA41172.1     | X58194   | Oryza sativa                  | cdc2+/CDC28-related protein kinase.   |
| CAA65500.1     | X96723   | Medicago sativa               | protein kinase. CDPK.   |
| BAA12691.1     | D84507   | Zea mays                      | CDPK-related protein kinase. Does not require calcium for its activity (by similarity). |
| AAB47181.1     | S82324   | Zea mays                      | /gene="calcium/calmodulin-dependent protein kinase. This sequence comes from Fig. 1.    |
| CAA07481.1     | AJ007366 | Zea mays                      | calcium-dependent protein kinase.   |
| BAA99439.1     | AP002743 | Oryza sativa                  | putative protein kinase. P0710E05.26. contains ESTs C22394(C30013),C22393(C30013).      |
| BAB12687.1     | AP002746 | Oryza sativa                  | putative protein kinase. P0671B11.2. contains ESTs C22394(C30013),C22393(C30013).       |
| AAF40430.1     | AF234652 | Mesembryanthemum crystallinum | protein kinase MK5.   |
| BAB21081.1     | AP002819 | Oryza sativa                  | putative calcium-dependent protein kinase. P0501G01.10.                                 |
| SEQ ID NO: 196 |          |                               |   |
| AAF37267.1     | AF220406 | Vitis riparia                 | 26S proteasome regulatory ATPase subunit S10b. Rev136-3.                                |
| SEQ ID NO: 197 |          |                               |   |

|            |          |  |
|------------|----------|--|
| BAA08104.1 | D45074   | <i>Panicum miliaceum</i><br>2-oxoglutarate/malate translocator. mitochondrial 2-oxoglutarate/malate translocator.  |
| BAA08103.1 | D45073   | <i>Panicum miliaceum</i><br>2-oxoglutarate/malate translocator. mitochondrial 2-oxoglutarate/malate translocator.  |
| BAA08105.1 | D45075   | <i>Panicum miliaceum</i><br>2-oxoglutarate/malate translocator. mitochondrial 2-oxoglutarate/malate translocator.  |
| CAA72107.1 | Y11220   | <i>Solanum tuberosum</i><br>mitochondrial uncoupling protein.  |
| CAC12820.1 | AJ299250 | <i>Nicotiana tabacum</i><br>mitochondrial 2-oxoglutarate/malate carrier protein. momc1.  |
| AAB71744.1 | U75346   | <i>Chlamydomonas reinhardtii</i><br>envelope protein. LIP-36G2. low CO <sub>2</sub> inducible carrier protein LIP-36 with a molecular weight of 36 kDa.                                    |
| AAB71743.1 | U75345   | <i>Chlamydomonas reinhardtii</i><br>envelope protein. LIP-36G1. low CO <sub>2</sub> inducible carrier protein LIP-36 with a molecular weight of 36 kDa.                                    |
| CAA07568.1 | AJ007580 | <i>Ribes nigrum</i><br>Mitochondrial carrier protein. prib7.   |
| BAB40117.1 | AP003311 | <i>Oryza sativa</i><br>putative peroxisomal Ca-dependent solute carrier protein. P0024G09.9.   |
| BAB16462.1 | AP002483 | <i>Oryza sativa</i><br>putative peroxisomal Ca-dependent solute carrier protein. P0019D06.21.  |
| CAA56325.1 | X80023   | <i>Triticum turgidum</i><br>ATP/ADP carrier protein.   |
| CAA46311.1 | X65194   | <i>Chlamydomonas reinhardtii</i><br>mitochondrial ADP/ATP translocator protein. CRANT.   |
| CAA67107.1 | X98474   | <i>Solanum tuberosum</i><br>mitochondrial energy transfer protein. brittle1.   |
| CAC27140.1 | AJ132535 | <i>Picea abies</i><br>ADP, ATP carrier protein precursor.  |
| CAA69726.1 | Y08499   | <i>Betula pendula</i><br>mitochondrial phosphate translocator.   |
| BAA31583.1 | AB016064 | <i>Zea mays</i><br>mitochondrial phosphate transporter.  |
| CAB61741.1 | AJ275306 | <i>Cicer arietinum</i><br>mitochondrial phosphate transporter.   |
| AAG45489.1 | AY013245 | <i>Oryza sativa</i><br>36I5.1. putative mitochondrial carrier protein.   |
| BAA31584.1 | AB016065 | <i>Oryza sativa</i><br>mitochondrial phosphate transporter.  |
| BAA92520.1 | AP001383 | <i>Oryza sativa</i><br>ESTs AU068633(C30614),AU068634(C30614) correspond to a region of the predicted gene. Similar to <i>Bos taurus</i> mitochondrial solute carrier protein. (AF049236). |

## SEQ ID NO: 199

BAA03455.1 D14605 Daucus carota  
AX110P. AX110.

## SEQ ID NO: 204

AAD21872.1 AF078082 Phaseolus vulgaris  
receptor-like protein kinase homolog RK20-1.

AAC23542.1 U20948 Ipomoea trifida  
receptor protein kinase. IRK1.

CAA73133.1 Y12530 Brassica oleracea  
serine /threonine kinase. ARLK.

CAA67145.1 X98520 Brassica oleracea  
receptor-like kinase. SFR2.

CAB41879.1 Y18260 Brassica oleracea  
SRK15 protein. SRK15. receptor-like kinase.

BAA92836.1 AB032473 Brassica oleracea  
S18 S-locus receptor kinase. SRK18.

CAA74662.1 Y14286 Brassica oleracea  
SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular  
kinase domain: from 1413.

AAB93834.1 U82481 Zea mays  
KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein  
kinase.

CAB41878.1 Y18259 Brassica oleracea  
SRK5 protein. SRK5. receptor-like kinase.

CAA73134.1 Y12531 Brassica oleracea  
serine/threonine kinase. BRLK.

CAA74661.1 Y14285 Brassica oleracea  
SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular  
kinase domain: 1412-2554.

BAA23676.1 AB000970 Brassica rapa  
receptor kinase 1. BcRK1.

AAA33000.1 M76647 Brassica oleracea  
receptor protein kinase. SKR6.

AAA62232.1 U00443 Brassica napus  
S-receptor kinase. protein contains an immunoglobulin-like domain.

CAB89179.1 AJ245479 Brassica napus subsp. napus  
ser /thr kinase. S-locus receptor kinase. srk.

AAA33008.1 M97667 Brassica napus  
serine/threonine kinase receptor.

BAA21132.1 D88193 Brassica rapa  
S-receptor kinase. SRK9 (B.c).

BAA06285.1 D30049 Brassica rapa  
S-receptor kinase SRK9.



|            |          |                     |  |
|------------|----------|---------------------|--|
| AAK02023.1 | AC074283 | Oryza sativa        | Putative protein kinase-like. OSJNBa0087H07.5.   |
| CAA79355.1 | Z18921   | Brassica oleracea   | S-receptor kinase-like protein.  |
| BAA92837.1 | AB032474 | Brassica oleracea   | S60 S-locus receptor kinase. SRK60.  |
| BAB21001.1 | AB054061 | Brassica rapa       | S locus receptor kinase. SRK22.  |
| BAA07577.2 | D38564   | Brassica rapa       | receptor protein kinase SRK12.   |
| BAA07576.1 | D38563   | Brassica rapa       | receptor protein kinase SRK8.  |
| AAD52097.1 | AF088885 | Nicotiana tabacum   | receptor-like kinase CHRK1. Chrkl.   |
| BAB18292.1 | AP002860 | Oryza sativa        | putative receptor-like protein kinase. P0409B08.19.  |
| BAB21240.1 | AP002953 | Oryza sativa        | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).   |
| BAA87853.1 | AP000816 | Oryza sativa        | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308). |
| BAB16871.1 | AP002537 | Oryza sativa        | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).                           |
| BAB39873.1 | AP002882 | Oryza sativa        | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| AAA33915.1 | L27821   | Oryza sativa        | receptor type serine/threonine kinase. protein kinase.   |
| CAB51836.1 | AJ243961 | Oryza sativa        | Putitive Ser/Thr protein kinase. 11332.7.  |
| CAA97692.1 | Z73295   | Catharanthus roseus | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.   |
| AAK21965.1 | AY028699 | Brassica napus      | receptor protein kinase PERK1.   |
| BAA94529.2 | AP001800 | Oryza sativa        | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAA94510.1 | AB041504 | Populus nigra       | protein kinase 2. PnPK2.   |
| BAA92954.1 | AP001551 | Oryza sativa        | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).  |

BAA94509.1 AB041503 *Populus nigra*  
protein kinase 1. PnPK1.

SEQ ID NO: 211

AAG13478.1 AC026758 *Oryza sativa*  
putative trehalose-6-phosphate phosphatase. OSJNBa0015J15.3.

SEQ ID NO: 212

AAC27894.1 AF023164 *Zea mays*  
leucine-rich repeat transmembrane protein kinase 1. ltk1.

AAC27895.1 AF023165 *Zea mays*  
leucine-rich repeat transmembrane protein kinase 2. ltk2.

AAK21965.1 AY028699 *Brassica napus*  
receptor protein kinase PERK1.

AAC61805.1 U28007 *Lycopersicon esculentum*  
serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.

BAB21241.1 AP002953 *Oryza sativa*  
Putative Pto kinase interactor 1. P0426D06.21. contains ESTs  
AU108280(E0721),D48017(S13927).

AAF91336.1 AF249317 *Glycine max*  
Pti1 kinase-like protein. Pti1a. protein kinase.

AAF91337.1 AF249318 *Glycine max*  
Pti1 kinase-like protein. Pti1b. protein kinase.

BAA82394.1 AP000367 *Oryza sativa*  
ESTs D23521(C2939),C22481(C2939) correspond to a region of the predicted gene.; Similar  
to serine/threonine protein kinase like protein. (AL022140).

BAA87853.1 AP000816 *Oryza sativa*  
EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative  
NAK-like Ser/Thr protein kinase. (AF001308).

AAG16628.1 AY007545 *Brassica napus*  
protein serine/threonine kinase BNK1.

BAB07999.1 AP002525 *Oryza sativa*  
putative protein kinase. P0462H08.22. contains EST C22619(S11214).

BAB03429.1 AP002817 *Oryza sativa*  
EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis  
thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).

BAB16871.1 AP002537 *Oryza sativa*  
putative protein kinase APK1A Arabidopsis thaliana. P0001B06.24. contains ESTs  
C22608(R3192),D25110(R3192).

BAA94510.1 AB041504 *Populus nigra*  
protein kinase 2. PnPK2.

BAB39409.1 AP002901 *Oryza sativa*  
putative protein kinase. P0456F08.9. contains EST C23560(R0290).

BAA94509.1 AB041503 *Populus nigra*  
protein kinase 1. PnPK1.

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| AAK11674.1 | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| AAF43496.1 | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| BAB21240.1 | AP002953 | Oryza sativa         | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| BAA78764.1 | AB023482 | Oryza sativa         | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAB09771.1 | U67422   | Zea mays             | CRINKLY4 precursor. cr4. receptor kinase homolog.   |
| AAK00425.1 | AC069324 | Oryza sativa         | Putative protein kinase. OSJNBa0071K19.11.  |
| AAG59657.1 | AC084319 | Oryza sativa         | putative protein kinase. OSJNBa0004B24.20.  |
| AAF66615.1 | AF142596 | Nicotiana tabacum    | LRR receptor-like protein kinase.   |
| AAG03090.1 | AC073405 | Oryza sativa         | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| CAB51834.1 | 00069    | Oryza sativa         | 1132.5. contains eukaryotic protein kinase domain PF.   |
| AAG25966.1 | AF302082 | Nicotiana tabacum    | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.   |
| CAA97692.1 | Z73295   | Catharanthus roseus  | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.  |
| BAA90808.1 | AP001168 | Oryza sativa         | Similar to putative receptor-like protein kinase (AL035679).  |
| BAB40081.1 | AP003074 | Oryza sativa         | putative receptor protein kinase. OSJNBa0004G10.30.   |
| BAB18321.1 | AP002865 | Oryza sativa         | putative receptor protein kinase. P0034C11.11.  |
| AAD38286.1 | AC007789 | Oryza sativa         | putative protein kinase. OSJNBa0049B20.13.  |
| CAB51480.1 | Y14600   | Sorghum bicolor      | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |

|                |          |                      |  |
|----------------|----------|----------------------|--|
| BAB19337.1     | AP003044 | Oryza sativa         | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).                 |
| AAF91324.1     | AF244890 | Glycine max          | receptor-like protein kinase 3. RLK3. GmRLK3.  |
| AAF91323.1     | AF244889 | Glycine max          | receptor-like protein kinase 2. RLK2. GmRLK2.  |
| BAB16326.1     | AP002818 | Oryza sativa         | putative receptor ser/thr protein kinase. P0436E04.9. contains ESTs S10111(AU070304),S10111(AU083519). |
| BAA82556.1     | AB030083 | Populus nigra        | lectin-like protein kinase. PnLPK.   |
| BAA94516.1     | AP001800 | Oryza sativa         | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).                                  |
| BAB07906.1     | AP002835 | Oryza sativa         | putative S-receptor kinase. P0417G05.14.   |
| AAF59906.1     | AF197947 | Glycine max          | receptor protein kinase-like protein. CLV1B.   |
| AAF59905.1     | AF197946 | Glycine max          | receptor protein kinase-like protein. CLV1A.   |
| BAB07904.1     | AP002835 | Oryza sativa         | putative S-receptor kinase. P0417G05.12.   |
| SEQ ID NO: 214 |          |                      |  |
| AAB51442.1     | U63012   | Sophora japonica     | lectin precursor.  |
| CAA93829.1     | Z69999   | Phaseolus lunatus    | lectin 3.  |
| BAA36415.1     | AB012634 | Robinia pseudoacacia | lectin.  |
| CAA93828.1     | Z69998   | Phaseolus lunatus    | lectin 2.  |
| AAC49137.1     | U21959   | Cladrastis kentukea  | lectin precursor.  |
| AAC49150.1     | U21940   | Cladrastis kentukea  | storage protein precursor. lectin.   |
| CAB96391.1     | AJ271873 | Phaseolus lunatus    | carbohydrate-binding. lectin. lb15.  |
| CAB96392.1     | AJ271874 | Phaseolus lunatus    | carbohydrate-binding. lectin. lb16.  |
| CAA76366.1     | Y16754   | Medicago sativa      | lectin. lec2.  |
| BAA82556.1     | AB030083 | Populus nigra        | lectin-like protein kinase. PnLPK.   |

|            |          |   |
|------------|----------|---|
| CAA93830.1 | Z70000   | Phaseolus lunatus<br>lectin 4.                      |
| AAG16779.1 | AF190633 | Ulex europaeus<br>lectin II.                        |
| AAC49136.1 | U21958   | Cladrastis kentukea<br>lectin precursor.            |
| AAB39933.1 | U65009   | Maackia amurensis<br>lectin precursor.              |
| AAB39934.1 | U65010   | Maackia amurensis<br>lectin precursor.              |
| AAA33766.1 | L26237   | Phaseolus lunatus<br>lectin II.                     |
| AAA33143.1 | M34270   | Dolichos biflorus<br>seed lectin.                   |
| CAA57697.1 | X82216   | Medicago truncatula<br>lectin. lec3.                |
| BAA36413.1 | AB012632 | Robinia pseudoacacia<br>lectin.                     |
| AAA80182.1 | U12783   | Robinia pseudoacacia<br>lectin.                     |
| BAA04604.1 | D17757   | Robinia pseudoacacia<br>lectin precursor.           |
| CAA68497.1 | Y00440   | Pisum sativum<br>lectin-precursor (AA -30 to 245).  |
| AAC49271.1 | U24249   | Robinia pseudoacacia<br>lectin precursor.           |
| AAA80181.1 | U12782   | Robinia pseudoacacia<br>lectin.                     |
| BAA36416.1 | AB012635 | Robinia pseudoacacia<br>lectin-related polypeptide. |
| AAA33676.1 | M18160   | Pisum sativum<br>lectin.                            |
| CAA47011.1 | X66368   | Pisum sativum<br>Psl lectin. psl.                   |
| AAA33141.1 | J02721   | Dolichos biflorus<br>lectin subunit I precursor.    |
| BAA36414.1 | AB012633 | Robinia pseudoacacia<br>lectin.                     |
| BAA02049.1 | D12481   | Bauhinia purpurea<br>lectin.                        |
| AAA80183.1 | U12784   | Robinia pseudoacacia<br>lectin.                     |

|                |          |                      |  |
|----------------|----------|----------------------|--|
| AAC49272.1     | U24250   | Robinia pseudoacacia | lectin precursor.  |
| AAA82737.1     | U18296   | Medicago sativa      | lectin. Mslec1.  |
| AAA74571.1     | U22468   | Arachis hypogaea     | agglutinin. galactose-binding lectin precursor. lec. lectin.   |
| AAB51441.1     | U63011   | Sophora japonica     | lectin precursor.  |
| AAA74574.1     | U22471   | Arachis hypogaea     | agglutinin. galactose-binding lectin precursor. lec. lectin.   |
| AAG00508.1     | AF285121 | Sophora flavescens   | lectin.  |
| AAB39932.1     | U65008   | Maackia amurensis    | lectin precursor.  |
| SEQ ID NO: 215 |          |                      |  |
| CAA64327.1     | X94624   | Brassica napus       | acyl-CoA synthetase.   |
| CAA96523.1     | Z72153   | Brassica napus       | acyl CoA synthetase.   |
| CAC19877.1     | AJ401089 | Brassica napus       | activation of free fatty acids. long chain acyl-CoA synthetase. acs6. activity confirmed by expression in E. coli. |
| CAA06820.1     | AJ006025 | Cicer arietinum      | acyl-coA synthetase.   |
| AAC39365.1     | AF008183 | Populus x generosa   | 4-coumarate:CoA ligase 2. 4CL2.  |
| CAA31697.1     | X13325   | Petroselinum crispum | 4-coumarate:CoA ligase Pc4Cl-2 (AA 1-544).   |
| CAA31696.1     | X13324   | Petroselinum crispum | 4-coumarate:CoA ligase Pc4Cl-1 (AA 1-544).   |
| AAC39366.1     | AF008184 | Populus x generosa   | 4-coumarate:CoA ligase 1. 4CL1.  |
| AAF37734.1     | AF052223 | Lolium perenne       | 4-coumarate--CoA ligase 4CL3.  |
| AAF91309.1     | AF239686 | Rubus idaeus         | 4-coumarate:coA ligase 2. adenylate-forming enzyme; 4CL2.  |
| AAF37733.1     | AF052222 | Lolium perenne       | 4-coumarate--CoA ligase 4CL2.  |
| CAA36850.1     | X52623   | Oryza sativa         | 4-coumarate-CoA ligase.  |
| BAA07828.1     | D43773   | Nicotiana tabacum    | 4-coumarate:coenzyme A ligase.   |

|   |          |                            |
|---|----------|----------------------------|
| BAA08365.1  | D49366   | Lithospermum erythrorhizon |
| 4-coumarate:CoA ligase.                                   |          |                            |
| AAB18637.1  | U50845   | Nicotiana tabacum          |
| 4-coumarate:coenzyme A ligase. 4CL1. Nt4CL-1.             |          |                            |
| AAD40664.1  | AF150686 | Solanum tuberosum          |
| 4-coumarate:coenzyme A ligase. 4CL-2a.                    |          |                            |
| AAA33842.1  | M62755   | Solanum tuberosum          |
| 4-coumarate--CoA ligase. St4C1-1.                         |          |                            |
| AAB18638.1  | U50846   | Nicotiana tabacum          |
| 4-coumarate:coenzyme A ligase. 4CL2. Nt4CL-19.            |          |                            |
| AAF91310.1  | AF239687 | Rubus idaeus               |
| 4-coumarate:coA ligase 1. adenylate-forming enzyme; 4CL1. |          |                            |
| AAC24503.1  | AF041049 | Populus tremuloides        |
| 4-coumarate:CoA ligase.                                   |          |                            |
| AAF37732.1  | AF052221 | Lolium perenne             |
| 4-coumarate--CoA ligase 4CL1.                             |          |                            |
| AAA92669.1  | U12013   | Pinus taeda                |
| 4-coumarate-CoA ligase enzyme.                            |          |                            |
| AAB42382.1  | U39404   | Pinus taeda                |
| 4-coumarate:CoA ligase. lp4CL-2.                          |          |                            |
| AAB42383.1  | U39405   | Pinus taeda                |
| 4-coumarate:CoA ligase. lp4CL-1.                          |          |                            |
| AAF91308.1  | AF239685 | Rubus idaeus               |
| 4-coumarate:coA ligase 3. adenylate-forming enzyme; 4CL3. |          |                            |
| AAC24504.1  | AF041050 | Populus tremuloides        |
| 4-coumarate:CoA ligase.                                   |          |                            |
| AAA92668.1  | U12012   | Pinus taeda                |
| 4-coumarate-CoA ligase enzyme.                            |          |                            |
| CAC36095.1  | X69955   | Glycine max                |
| 4-coumarate:Coenzyme A ligase isoenzyme 4. 4CL4.          |          |                            |
| AAG43823.1  | AF212317 | Capsicum annuum            |
| 4-coumarate:coenzyme A ligase. 4CL.                       |          |                            |
| AAA69580.1  | L43362   | Oryza sativa               |
| 4-coumarate:CoA ligase isoform 2. 4cl.2. putative.        |          |                            |
| BAA08366.2  | D49367   | Lithospermum erythrorhizon |
| 4-coumarate:CoA ligase.                                   |          |                            |
| CAA49575.1  | X69954   | Glycine max                |
| 4-coumarate--CoA ligase.                                  |          |                            |
| CAB97359.1  | AJ278455 | Juglans nigra              |
| 4-coumarate-CoA ligase. 4CL.                              |          |                            |
| AAF73995.2  | AF144502 | Pinus armandii             |
| 4-coumarate:CoA ligase. 4CL.                              |          |                            |

|                              |          |                                  |
|------------------------------|----------|----------------------------------|
| AAF73994.2                   | AF144501 | <i>Pinus armandii</i>            |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74018.2                   | AF144525 | <i>Tsuga canadensis</i>          |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74010.2                   | AF144517 | <i>Abies holophylla</i>          |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74020.2                   | AF144527 | <i>Pseudolarix amabilis</i>      |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74016.2                   | AF144523 | <i>Nothotsuga longibracteata</i> |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF73997.2                   | AF144504 | <i>Picea smithiana</i>           |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74013.2                   | AF144520 | <i>Abies beshanzuensis</i>       |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74008.2                   | AF144515 | <i>Abies firma</i>               |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74022.2                   | AF144529 | <i>Cedrus atlantica</i>          |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74005.2                   | AF144512 | <i>Larix gmelini</i>             |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74021.2                   | AF144528 | <i>Pseudolarix amabilis</i>      |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74003.2                   | AF144510 | <i>Pseudotsuga sinensis</i>      |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |
| AAF74019.2                   | AF144526 | <i>Tsuga canadensis</i>          |
| 4-coumarate:CoA ligase. 4CL. |          |                                  |

SEQ ID NO: 221

|                                    |        |                          |
|------------------------------------|--------|--------------------------|
| AAB37246.1                         | U58971 | <i>Nicotiana tabacum</i> |
| calmodulin-binding protein. TCB60. |        |                          |

SEQ ID NO: 224

|  |          |                           |
|--|----------|---------------------------|
| BAB19413.1                                       | AP002870 | <i>Oryza sativa</i>       |
| putative acetone-cyanohydrin lyase. P0458A05.22. |          |                           |
| AAC49184.1                                       | U40402   | <i>Hevea brasiliensis</i> |
| hydroxynitrile lyase. hnl.                       |          |                           |
| CAA11219.1                                       | AJ223281 | <i>Manihot esculenta</i>  |
| alpha-hydroxynitrile lyase. HNL4.                |          |                           |
| CAA82334.1                                       | Z29091   | <i>Manihot esculenta</i>  |
| alpha-hydroxynitrile lyase.                      |          |                           |

SEQ ID NO: 225

|   |          |                     |
|---|----------|---------------------|
| BAB16335.1  | AP002818 | <i>Oryza sativa</i> |
| putative NAM protein. P0436E04.18. contains ESTs E10793(C19698),E10793(C99379). |          |                     |



|                |          |                           |  |
|----------------|----------|---------------------------|--|
| AAK13151.1     | AC078829 | Oryza sativa              | putative NAM (no apical meristem) protein. OSJNBa0026O12.6.  |
| CAA63102.2     | X92205   | Petunia x hybrida         | apical meristem formation. NAM.  |
| CAA63101.1     | X92204   | Petunia x hybrida         | apical meristem formation. NAM.  |
| BAB16328.1     | AP002818 | Oryza sativa              | putative NAM protein. P0436E04.11. contains ESTs R4069(AU032425),R4069(AU082730).  |
| BAB19365.1     | AP002542 | Oryza sativa              | putative NAM (no apical meristem) protein. P0679C08.4.   |
| BAA84803.1     | AP000559 | Oryza sativa              | Similar to NAM like protein (AC005310).  |
| BAB03447.1     | AP002817 | Oryza sativa              | ESTs C96615(C10106),C26336(C12127),D21959(C10106) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana DNA chromosome 4, BAC clone T6K21; NAM (no apical meristem) - like protein (AL021889).     |
| BAA92400.1     | AP001366 | Oryza sativa              | ESTs C96615(C10106),C26336(C12127),D21959(C10106) correspond to a region of the predicted gene. Similar to NAM (AL021889).   |
| SEQ ID NO: 226 |          |                           |  |
| BAA96221.1     | AP002094 | Oryza sativa              | ESTs C19814(E10971),AU090481(E10971) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, sequence from clones T3F17, F11C10, F13A10; auxin-responsive GH3-like protein (AC006526). |
| AAD32141.1     | AF123503 | Nicotiana tabacum         | Nt-gh3 deduced protein.  |
| CAA42636.1     | X60033   | Glycine max               | auxin-responsive GH3 product. GH3.   |
| SEQ ID NO: 227 |          |                           |  |
| CAA70403.1     | Y09204   | Nicotiana tabacum         | histidinol-phosphate aminotransferase. hpa.  |
| CAC20728.1     | AJ278767 | Nicotiana plumbaginifolia | essential for histidine biosynthesis. histidinol phosphate aminotransferase. hpa.  |
| SEQ ID NO: 229 |          |                           |  |
| AAF33670.1     | AF079872 | Nicotiana tabacum         | cyclic nucleotide-gated calmodulin-binding ion channel. CBP4.  |
| AAF33669.1     | AF079871 | Nicotiana tabacum         | cyclic nucleotide-gated calmodulin-binding ion channel. CBP7.  |
| AAK16188.1     | AC079887 | Oryza sativa              | putative cyclic nucleotide and calmodulin-regulated ion channel protein. OSJNBa0040E01.13.   |
| AAB53255.1     | U65390   | Nicotiana tabacum         | cyclic nucleotide gated channel protein. CaMB-channel protein. channel protein homolog.  |

|            |          |                                       |  |
|------------|----------|---------------------------------------|--|
| CAB54856.1 | AJ132686 | Zea mays                              | potassium channel protein ZMK2. ZMK2.  |
| AAD16278.1 | AF099095 | Samanea saman                         | pulvinus inward-rectifying channel for potassium SPICK1. similar to Arabidopsis potassium channel AKT3.  |
| CAA71598.1 | Y10579   | Vicia faba                            | potassium channel.   |
| CAA56175.1 | X79779   | Solanum tuberosum                     | K+ channel inward rectifying. KST1.  |
| CAC05489.1 | AJ271447 | Populus tremula x Populus tremuloides | potassium channel. potassium channel 2. ptk2.  |
| AAD39492.1 | AF145272 | Samanea saman                         | pulvinus inward-rectifying channel SPICK2. potassium channel; similar to AKT2/3.   |
| CAA70870.1 | Y09699   | Solanum tuberosum                     | putative inward rectifying potassium channel. SKT2.  |
| CAB62555.1 | AJ249962 | Daucus carota                         | potassium channel. kdc1.   |
| BAA96192.1 | AP002093 | Oryza sativa                          | Similar to Arabidopsis thaliana potassium channel protein (M86990).  |
| BAA96150.1 | AP002092 | Oryza sativa                          | Similar to Arabidopsis thaliana potassium channel protein (M86990).  |
| BAA84085.1 | AB032074 | Nicotiana paniculata                  | potassium channel. NpKT1.  |
| AAF81251.1 | AF267755 | Mesembryanthemum crystallinum         | potassium channel protein Mkt2p.   |
| CAA68912.1 | Y07632   | Zea mays                              | potassium channel. ZMK1.   |
| CAA60016.1 | X86021   | Solanum tuberosum                     | potassium channel. SKT1 gene. putative start codon.  |
| CAA65254.1 | X96390   | Lycopersicon esculentum               | potassium channel. LKT1.   |
| CAA12645.1 | AJ225805 | Egeria densa                          | inward potassium channel alpha subunit. homologous to the sequences of the family of inwardly rectifying potassium channels in plants which is structurally related to the shaker family of outwardly rectifying channels in Drosophila. |
| AAF36832.1 | AF207745 | Triticum aestivum                     | AKT1-like potassium channel. TaAKT1.   |
| CAC10514.1 | AJ299019 | Samanea saman                         | potassium release. outwardly rectifying potassium channel. spork1.   |
| AAF81249.1 | AF267753 | Mesembryanthemum crystallinum         | putative potassium channel protein Mkt1p.  |
| CAC05488.1 | AJ271446 | Populus tremula x Populus tremuloides | potassium channel. outward rectifying potassium channel. ptork.  |

## SEQ ID NO: 232

AAA80575.1 U13148 Pennisetum ciliare  
possible apospory-associated protein.

AAF34174.1 AF195243 Chlamydomonas reinhardtii  
apospory-associated protein C. APOC.

## SEQ ID NO: 233

AAB97366.1 AF039531 Oryza sativa  
lysophospholipase homolog. LPL1.

## SEQ ID NO: 235

AAC61839.1 AF025430 Papaver somniferum  
berberine bridge enzyme. bbe1. (S)-reticuline:oxygen oxidoreductase (methylene bridge forming).

AAC39358.1 AF005655 Eschscholzia californica  
oxidizes the N-methyl group of (S)-reticuline to the berberine bridge carbon C-8 of (S)-scoulerine in berberine and benzophenanthridine alkaloid biosynthesis. berberine bridge enzyme. bbe1. covalently-bound FAD-dependent oxidase; elicitor-inducible.

AAB20352.1 S65550 Eschscholzia californica  
(S)-reticuline:oxygen oxidoreductase (methylene-bridge-forming). /gene="(S)-reticuline:oxygen oxidoreductase. This sequence comes from Fig 2; berberine bridge enzyme.

AAD17487.1 AF049347 Berberis stolonifera  
Catalyzes the oxidative cyclization of the N-methyl group of (S)-reticuline into the berberine bridge carbon (C-8) of (S)-scoulerine. berberine bridge enzyme. bbe1. Covalently flavinylated oxidase of isoquinoline alkaloid biosynthesis in plants.

## SEQ ID NO: 244

AAD17487.1 AF049347 Berberis stolonifera  
Catalyzes the oxidative cyclization of the N-methyl group of (S)-reticuline into the berberine bridge carbon (C-8) of (S)-scoulerine. berberine bridge enzyme. bbe1. Covalently flavinylated oxidase of isoquinoline alkaloid biosynthesis in plants.

AAB20352.1 S65550 Eschscholzia californica  
(S)-reticuline:oxygen oxidoreductase (methylene-bridge-forming). /gene="(S)-reticuline:oxygen oxidoreductase. This sequence comes from Fig 2; berberine bridge enzyme.

AAC39358.1 AF005655 Eschscholzia californica  
oxidizes the N-methyl group of (S)-reticuline to the berberine bridge carbon C-8 of (S)-scoulerine in berberine and benzophenanthridine alkaloid biosynthesis. berberine bridge enzyme. bbe1. covalently-bound FAD-dependent oxidase; elicitor-inducible.

AAC61839.1 AF025430 Papaver somniferum  
berberine bridge enzyme. bbe1. (S)-reticuline:oxygen oxidoreductase (methylene bridge forming).

## SEQ ID NO: 247

AAF37267.1 AF220406 Vitis riparia  
26S proteasome regulatory ATPase subunit S10b. Rev136-3.

## SEQ ID NO: 248

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAA78764.1 | AB023482 | Oryza sativa         | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| BAA94509.1 | AB041503 | Populus nigra        | protein kinase 1. PnPK1.  |
| AAG16628.1 | AY007545 | Brassica napus       | protein serine/threonine kinase BNK1.   |
| BAB03429.1 | AP002817 | Oryza sativa         | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).             |
| BAB07999.1 | AP002525 | Oryza sativa         | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAA94510.1 | AB041504 | Populus nigra        | protein kinase 2. PnPK2.  |
| AAK11674.1 | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| AAF43496.1 | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| BAB16871.1 | AP002537 | Oryza sativa         | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).  |
| BAA87853.1 | AP000816 | Oryza sativa         | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB39409.1 | AP002901 | Oryza sativa         | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| BAB21240.1 | AP002953 | Oryza sativa         | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| AAK21965.1 | AY028699 | Brassica napus       | receptor protein kinase PERK1.  |
| AAK00425.1 | AC069324 | Oryza sativa         | Putative protein kinase. OSJNBa0071K19.11.  |
| AAG03090.1 | AC073405 | Oryza sativa         | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| AAG59657.1 | AC084319 | Oryza sativa         | putative protein kinase. OSJNBa0004B24.20.  |
| AAF91336.1 | AF249317 | Glycine max          | Ptil kinase-like protein. Ptila. protein kinase.  |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAF91337.1 | AF249318 | Glycine max                   | Pti1 kinase-like protein. Pti1b. protein kinase.  |
| BAB21241.1 | AP002953 | Oryza sativa                  | Putative Pto kinase interactor 1. P0426D06.21. contains ESTs AU108280(E0721),D48017(S13927).  |
| CAB51834.1 | 00069    | Oryza sativa                  | 11332.5. contains eukaryotic protein kinase domain PF.  |
| AAC61805.1 | U28007   | Lycopersicon esculentum       | serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.  |
| AAG33377.1 | AF290411 | Oryza meyeriana               | serine/threonine protein kinase. R1.  |
| AAC27894.1 | AF023164 | Zea mays                      | leucine-rich repeat transmembrane protein kinase 1. ltk1.   |
| BAA90808.1 | AP001168 | Oryza sativa                  | Similar to putative receptor-like protein kinase (AL035679).  |
| AAC27895.1 | AF023165 | Zea mays                      | leucine-rich repeat transmembrane protein kinase 2. ltk2.   |
| AAB61708.1 | U93048   | Daucus carota                 | somatic embryogenesis receptor-like kinase. SERK.   |
| CAA97692.1 | Z73295   | Catharanthus roseus           | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.  |
| AAK11566.1 | AF318490 | Lycopersicon hirsutum         | Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.   |
| BAB18292.1 | AP002860 | Oryza sativa                  | putative receptor-like protein kinase. P0409B08.19.   |
| CAB51480.1 | Y14600   | Sorghum bicolor               | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| AAG25966.1 | AF302082 | Nicotiana tabacum             | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.   |
| AAG00510.1 | AF285172 | Phaseolus vulgaris            | leaf senescence-associated receptor-like protein kinase. SARK.  |
| AAC48932.1 | U13923   | Lycopersicon pimpinellifolium | Fen. putative serine/threonine protein kinase; similar to product encoded by Lycopersicon pimpinellifolium Pto gene, GenBank Accession Number U02271; Fen is a member of the Pto gene family. |
| AAB47424.1 | U59317   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Fen. Fen. fenthion sensitivity gene from tomato.  |
| AAF76307.1 | AF220602 | Lycopersicon pimpinellifolium | Fen kinase.   |
| AAF66615.1 | AF142596 | Nicotiana tabacum             | LRR receptor-like protein kinase.   |

|                |          |                              |  |
|----------------|----------|------------------------------|--|
| BAA92221.1     | AP001278 | Oryza sativa                 | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).                   |
| BAA87852.1     | AP000816 | Oryza sativa                 | Similar to putative Ser/Thr protein kinase. (AC004218).  |
| AAK11567.1     | AF318491 | Lycopersicon hirsutum        | Pto-like protein kinase F. LhirPtoF.   |
| CAA73134.1     | Y12531   | Brassica oleracea            | serine/threonine kinase. BRLK.   |
| BAB19337.1     | AP003044 | Oryza sativa                 | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).   |
| SEQ ID NO: 249 |          |                              |  |
| BAA22422.1     | AB001379 | Glycyrrhiza echinata         | cytochrome P450. CYP81E1.  |
| BAA74465.1     | AB022732 | Glycyrrhiza echinata         | cytochrome P450. CYP Ge-31.  |
| BAA93634.1     | AB025016 | Lotus japonicus              | cytochrome P450.   |
| CAB43505.1     | AJ239051 | Cicer arietinum              | cytochrome P450. cyp81E2.  |
| CAB41490.1     | AJ238439 | Cicer arietinum              | cytochrome P450 monooxygenase. cyp81E3v2.  |
| CAA10067.1     | AJ012581 | Cicer arietinum              | cytochrome P450. cyp81E3.  |
| CAA04117.1     | AJ000478 | Helianthus tuberosus         | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-race).                                    |
| CAA04116.1     | AJ000477 | Helianthus tuberosus         | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.  |
| AAK38079.1     | AF321855 | Lolium rigidum               | putative cytochrome P450.  |
| AAK38080.1     | AF321856 | Lolium rigidum               | putative cytochrome P450.  |
| AAK38081.1     | AF321857 | Lolium rigidum               | putative cytochrome P450.  |
| AAC34853.1     | AF082028 | Hemerocallis hybrid cultivar | putative cyt P450-containing fatty acid hydroxylase. senescence-associated protein 3. SA3. mRNA accumulates in senescing petals. |
| AAB94590.1     | AF022461 | Glycine max                  | CYP82C1p. CYP82C1. cytochrome P450 monooxygenase.  |
| CAB56742.1     | AJ249800 | Cicer arietinum              | cytochrome P450 monooxygenase. cyp81E5.  |

|            |          |  |
|------------|----------|--|
| AAD56282.1 | AF155332 | <i>Petunia x hybrida</i><br>flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.   |
| AAG09208.1 | AF175278 | <i>Pisum sativum</i><br>wound-inducible P450 hydroxylase. CYP82A1.   |
| CAA71515.1 | Y10491   | <i>Glycine max</i><br>putative cytochrome P450.  |
| CAA71516.1 | Y10492   | <i>Glycine max</i><br>putative cytochrome P450.  |
| AAC49188.2 | U29333   | <i>Pisum sativum</i><br>cytochrome P450 monooxygenase. CYP82. new cytochrome P450 family.  |
| CAA71876.1 | Y10982   | <i>Glycine max</i><br>putative cytochrome P450.  |
| BAA12159.1 | D83968   | <i>Glycine max</i><br>Cytochrome P-450 (CYP93A1).  |
| AAC39454.1 | AF014802 | <i>Eschscholzia californica</i><br>(S)-N-methylcoclaurine 3'-hydroxylase. CYP82B1. cytochrome P-450-dependent monooxygenase; methyl jasmonate-inducible cytochrome P-450-dependent, homologous to wound-inducible CYP82A1 of <i>Pisum sativum</i> GenBank Accession Number U29333. |
| CAA65580.1 | X96784   | <i>Nicotiana tabacum</i><br>cytochrome P450. hsr515.   |
| BAA13076.1 | D86351   | <i>Glycine max</i><br>cytochrome P-450 (CYP93A2).  |
| AAA32913.1 | M32885   | <i>Persea americana</i><br>cytochrome P-450LXXIA1 (cyp71A1).   |
| CAA71877.1 | Y10983   | <i>Glycine max</i><br>putative cytochrome P450.  |
| AAG44132.1 | AF218296 | <i>Pisum sativum</i><br>cytochrome P450. P450 isolog.  |
| CAA64635.1 | X95342   | <i>Nicotiana tabacum</i><br>cytochrome P450. hsr515. hypersensitivity-related gene.  |
| AAB94587.1 | AF022458 | <i>Glycine max</i><br>CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.  |
| AAD38930.1 | AF135485 | <i>Glycine max</i><br>cytochrome P450 monooxygenase CYP93D1. CYP93E1.  |
| BAA84072.1 | AB028152 | <i>Torenia hybrida</i><br>flavone synthase II. cytochrome P450. TFNS5.   |
| BAA74466.1 | AB022733 | <i>Glycyrrhiza echinata</i><br>cytochrome P450. CYP Ge-51.   |
| BAA22423.1 | AB001380 | <i>Glycyrrhiza echinata</i><br>cytochrome P450. CYP93B1.   |
| BAA35080.1 | AB015762 | <i>Nicotiana tabacum</i><br>putative cytochrome P450. CYP82E1.   |

|                |          |                                |   |
|----------------|----------|--------------------------------|---|
| BAA92894.1     | AB006790 | Petunia x hybrida              | cytochrome P450. IMT-2.   |
| AAC32274.1     | AF081575 | Petunia x hybrida              | flavonoid 3',5'-hydroxylase. Hfl. P450 enzyme.  |
| CAB56743.1     | AJ249801 | Cicer arietinum                | cytochrome P450 monooxygenase. cyp81E4.   |
| BAA84071.1     | AB028151 | Antirrhinum majus              | flavone synthase II. cytochrome P450. AFNS2.  |
| SEQ ID NO: 250 |          |                                |   |
| AAD55566.1     | AF110784 | Volvox carteri f. nagariensis  | protein disulfide isomerase precursor. pdi.   |
| AAD02069.1     | AF036939 | Chlamydomonas reinhardtii      | redox-regulator of 5'UTR psbA mRNA binding complex and translation. protein disulfide isomerase. localized to ER and chloroplast.   |
| AAC49896.1     | AF027727 | Chlamydomonas reinhardtii      | involved in the redox-regulated binding of chloroplast poly(A)-binding protein to the 5'-UTR of psbA mRNA; regulates chloroplast translational activation. protein disulfide isomerase RB60. PDI. |
| AAD28260.1     | AF131223 | Datisca glomerata              | protein disulfide isomerase homolog. PDI.   |
| AAB08519.1     | L39014   | Zea mays                       | protein disulfide isomerase. pdi. putative.   |
| AAA19660.1     | U11496   | Triticum aestivum              | protein disulfide isomerase. PDI.   |
| CAC21230.1     | AJ277379 | Triticum turgidum subsp. durum | catalyzes formation of disulfide bonds. protein disulfide isomerase. Pdi.   |
| CAC21228.1     | AJ277377 | Triticum turgidum subsp. durum | catalyzes the formation of disulfide bonds. protein disulfide isomerase. Pdi.   |
| AAA70345.1     | L33251   | Hordeum vulgare                | catalyze the formation of disulfide bonds. disulfide isomerase. PDI. putative.  |
| AAA70344.1     | L33250   | Hordeum vulgare                | catalyze the formation of disulfide bonds. disulfide isomerase. PDI. putative.  |
| AAB05641.1     | U41385   | Ricinus communis               | protein disulphide isomerase PDI. molecular chaperone.  |
| CAA77575.1     | Z11499   | Medicago sativa                | protein disulfide isomerase.  |
| CAC21231.1     | AJ277380 | Triticum turgidum subsp. durum | catalyzes formation of disulfide bonds. protein disulfide isomerase. Pdi.   |
| CAC21229.1     | AJ277378 | Triticum turgidum subsp. durum | catalyzes formation of disulfide bonds. protein disulfide isomerase. Pdi.   |
| BAB18780.1     | AB047268 | Cucumis sativus                | disulfide isomerase.  |



|                |          |                         |   |
|----------------|----------|-------------------------|---|
| BAA92322.1     | AB039278 | Oryza sativa            | protein disulfide isomerase. Pdi.   |
| AAA70346.1     | L33252   | Hordeum vulgare         | catalyze the formation of disulfide bonds. disulfide isomerase. PDI. putative.  |
| CAA72092.1     | Y11209   | Nicotiana tabacum       | protein disulfide-isomerase precursor. PDI.   |
| AAG13988.1     | AF298829 | Prunus avium            | putative protein disulfide-isomerase. PDI.  |
| SEQ ID NO: 251 |          |                         |   |
| CAA61275.1     | X88797   | Eucalyptus gunnii       | cinnamyl alcohol dehydrogenase. CAD1.   |
| AAC06319.1     | AF053084 | Malus x domestica       | putative cinnamyl alcohol dehydrogenase. CAD.   |
| SEQ ID NO: 253 |          |                         |   |
| CAB61745.1     | AJ275311 | Cicer arietinum         | farnesylated protein.   |
| AAD09515.1     | U64917   | Glycine max             | putative metal-binding protein. GMFP7. farnesylated protein.  |
| SEQ ID NO: 254 |          |                         |   |
| BAB19757.1     | AB052785 | Glycine max             | nitrate transporter NRT1-2. NRT1-2.   |
| BAB19756.1     | AB052784 | Glycine max             | nitrate transporter NRT1-1. NRT1-1.   |
| BAB19760.1     | AB052788 | Glycine max             | nitrate transporter NRT1-5. NRT1-5.   |
| AAC32034.1     | AF023472 | Hordeum vulgare         | peptide transporter. ptr1. PTR1; integral membrane protein.   |
| AAD01600.1     | AF016713 | Lycopersicon esculentum | LeOPT1. LeOPT1. oligopeptide transporter.   |
| BAB40113.1     | AP003311 | Oryza sativa            | putative peptide transport protein. P0024G09.4. contains ESTs D40448(S2437),C71800(E0368),AU102190(E2393), AU055921(S20154),AU102191(E2393),AU055922(S20154), C98524(E0368),AU097146(S2437).  |
| BAB16458.1     | AP002483 | Oryza sativa            | putative peptide transport protein. P0019D06.16. contains ESTs D40448(S2437),C71800(E0368),AU102190(E2393), AU055921(S20154),AU102191(E2393),AU055922(S20154), C98524(E0368),AU097146(S2437). |
| CAA93316.1     | Z69370   | Cucumis sativus         | nitrite transporter. NiTR1.   |
| AAK15441.1     | AC037426 | Oryza sativa            | putative nitrate transporter. OSJNBb0014I11.9.  |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| AAG21898.1     | AC026815 | Oryza sativa              | putative peptide transport protein. OSJNBa0079L16.13.   |
| CAC00544.1     | AJ277084 | Nicotiana plumbaginifolia | ion transport. putative low-affinity nitrate transporter. nrt1.1.                                       |
| AAG46153.1     | AC018727 | Oryza sativa              | putative peptide transporter. OSJNBa0056G17.8.  |
| CAC00545.1     | AJ277085 | Nicotiana plumbaginifolia | ion transport. putative low-affinity nitrate transporter. nrt1.2.                                       |
| AAG21906.1     | AC026815 | Oryza sativa              | putative peptide transport protein. OSJNBa0079L16.9.  |
| AAF20002.1     | AF213936 | Prunus dulcis             | amino acid/peptide transporter. PTR2. similar to transporters of nitrogenous compounds.                 |
| AAF07875.1     | AF140606 | Oryza sativa              | nitrate transporter. NRT1.  |
| BAB16322.1     | AP002818 | Oryza sativa              | putative peptide transporter-like protein. P0436E04.4.  |
| BAB19758.1     | AB052786 | Glycine max               | putative nitrate transporter NRT1-3. NRT1-3.  |
| AAA80582.1     | U17987   | Brassica napus            | putative nitrate transporter. RCH2 protein.   |
| CAC07206.1     | AJ278966 | Brassica napus            | Low-affinity nitrate transporter. nitrate transporter. nrt1.  |
| AAG46154.1     | AC018727 | Oryza sativa              | putative peptide transporter. OSJNBa0056G17.27.   |
| AAB69642.1     | AF000392 | Lotus japonicus           | peptide transporter. LjNOD65.   |
| BAB19759.1     | AB052787 | Glycine max               | putative nitrate transporter NRT1-4. NRT1-4.  |
| AAD16016.1     | AF080545 | Nepenthes alata           | peptide transporter. PTR1.  |
| AAD42860.1     | AF154930 | Prunus dulcis             | transporter-like protein. TLP1.   |
| AAG13513.1     | AC068924 | Oryza sativa              | putative peptide transporter. OSJNBa0026L12.7.  |
| SEQ ID NO: 255 |          |                           |   |
| AAB01567.1     | L47672   | Picea glauca              | EMB34. embryo-abundant protein.   |
| SEQ ID NO: 257 |          |                           |   |
| AAB71743.1     | U75345   | Chlamydomonas reinhardtii | envelope protein. LIP-36G1. low CO2 inducible carrier protein LIP-36 with a molecular weight of 36 kDa. |

|                |          |   |
|----------------|----------|---|
| AAB71744.1     | U75346   | <i>Chlamydomonas reinhardtii</i><br>envelope protein. LIP-36G2. low CO2 inducible carrier protein LIP-36 with a molecular weight of 36 kDa.   |
| BAA92520.1     | AP001383 | <i>Oryza sativa</i><br>ESTs AU068633(C30614),AU068634(C30614) correspond to a region of the predicted gene. Similar to <i>Bos taurus</i> mitochondrial solute carrier protein. (AF049236).                  |
| BAB16462.1     | AP002483 | <i>Oryza sativa</i><br>putative peroxisomal Ca-dependent solute carrier protein. P0019D06.21.   |
| BAB40117.1     | AP003311 | <i>Oryza sativa</i><br>putative peroxisomal Ca-dependent solute carrier protein. P0024G09.9.  |
| CAA07568.1     | AJ007580 | <i>Ribes nigrum</i><br>Mitochondrial carrier protein. prib7.  |
| CAC27140.1     | AJ132535 | <i>Picea abies</i><br>ADP, ATP carrier protein precursor.   |
| CAA56325.1     | X80023   | <i>Triticum turgidum</i><br>ATP/ADP carrier protein.  |
| CAC12820.1     | AJ299250 | <i>Nicotiana tabacum</i><br>mitochondrial 2-oxoglutarate/malate carrier protein. momc1.   |
| AAG48999.1     | AY013246 | <i>Hordeum vulgare</i><br>putative mitochondrial carrier protein. 635P2.1.  |
| SEQ ID NO: 258 |          |   |
| CAA05276.1     | AJ002236 | <i>Lycopersicon pimpinellifolium</i><br>resistance gene. Hcr9-9E. Hcr9-9E.  |
| AAC78591.1     | AF053993 | <i>Lycopersicon esculentum</i><br>disease resistance protein. Cf-5.   |
| AAC78596.1     | AF053998 | <i>Lycopersicon esculentum</i><br>Hcr2-5D. Hcr2-5D. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |
| CAA05279.1     | AJ002237 | <i>Lycopersicon esculentum</i><br>Hcr9-0. Hcr9-0. homologue of <i>Cladosporium fulvum</i> disease resistance gene Cf-9.   |
| AAC78593.1     | AF053995 | <i>Lycopersicon esculentum</i><br>Hcr2-0B. Hcr2-0B. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |
| AAA65235.1     | U15936   | <i>Lycopersicon pimpinellifolium</i><br>Cf-9 precursor. Cf-9. this is the ninth resistance gene to disease caused by <i>Cladosporium fulvum</i> to be isolated.   |
| CAA05274.1     | AJ002236 | <i>Lycopersicon pimpinellifolium</i><br>resistance gene. Cf-9. Cf-9.  |
| AAC78592.1     | AF053994 | <i>Lycopersicon esculentum</i><br>Hcr2-0A. Hcr2-0A. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |

|                |          |   |
|----------------|----------|---|
| AAC78595.1     | AF053997 | <i>Lycopersicon esculentum</i><br>Hcr2-5B. Hcr2-5B. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445.       |
| AAC78594.1     | AF053996 | <i>Lycopersicon pimpinellifolium</i><br>Hcr2-2A. Hcr2-2A. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |
| BAA96776.1     | AP002521 | <i>Oryza sativa</i><br>Similar to <i>Lycopersicon esculentum</i> disease resistance protein (AF053993).   |
| BAB08215.1     | AP002539 | <i>Oryza sativa</i><br>Similar to <i>Lycopersicon esculentum</i> disease resistance protein (AF053993).   |
| CAA05268.1     | AJ002235 | <i>Lycopersicon hirsutum</i><br>Resistance gene. Cf-4. Cf-4.  |
| AAG21897.1     | AC026815 | <i>Oryza sativa</i><br>putative disease resistance protein (3' partial). OSJNBa0079L16.21.  |
| AAD50430.1     | AF166121 | <i>Hordeum vulgare</i><br>Cf2/Cf5 disease resistance protein homolog. Big1. leucine rich repeat protein.  |
| AAG21917.1     | AC026815 | <i>Oryza sativa</i><br>putative disease resistance protein. OSJNBa0079L16.5.  |
| AAG21909.1     | AC026815 | <i>Oryza sativa</i><br>putative disease resistance protein. OSJNBa0079L16.3.  |
| CAB55409.1     | AL117265 | <i>Oryza sativa</i><br>zhb0001.1. Incomplete at 5'end, Similar to disease resistance protein; Method: conceptual translation with partial peptide sequencing.   |
| AAC49123.1     | U37133   | <i>Oryza sativa</i><br>receptor kinase-like protein. Xa21. Xa21 disease resistance gene.  |
| AAC80225.1     | U72723   | <i>Oryza longistaminata</i><br>receptor kinase-like protein. Xa21. disease resistance gene.   |
| SEQ ID NO: 259 |          |   |
| CAB52689.1     | AJ132224 | <i>Lycopersicon esculentum</i><br>hexose transporter. ht2.  |
| CAA09419.1     | AJ010942 | <i>Lycopersicon esculentum</i><br>hexose transporter protein.   |
| BAB19864.1     | AB052885 | <i>Oryza sativa</i><br>monosaccharide transporter 3. OsMST3.  |
| AAA18534.1     | L21753   | <i>Saccharum hybrid cultivar H65-7052</i><br>glucose transporter. putative.   |
| CAA47324.1     | X66856   | <i>Nicotiana tabacum</i><br>monosaccharid transporter. MST1.  |
| AAB06594.1     | U38651   | <i>Medicago truncatula</i><br>sugar transporter.  |
| AAA79761.1     | L08196   | <i>Ricinus communis</i><br>hexose transport. sugar carrier protein. RCSTC.  |

|            |          |                                    |  |
|------------|----------|------------------------------------|--|
| CAA04511.1 | AJ001061 | Vitis vinifera                     | hexose uptake. hexose transporter.   |
| CAA70777.1 | Y09590   | Vitis vinifera                     | hexose transporter.  |
| AAC61852.1 | AF061106 | Petunia x hybrida                  | putative monosaccharide transporter 1. pmt1. similar to hexose transporter protein; PMT1.                    |
| AAA79857.1 | L08188   | Ricinus communis                   | hexose transport. hexose carrier protein. HEX6.  |
| CAB06079.1 | Z83829   | Picea abies                        | monosaccharide transporter. PaMst-1. PaMst-1.  |
| BAB19863.1 | AB052884 | Oryza sativa                       | monosaccharide transporter 2. OsMST2.  |
| BAA83554.1 | AP000399 | Oryza sativa                       | Similar to hexose carrier protein HEX6 & RCCHCP_1 (Q07423).  |
| AAK31286.1 | AC079890 | Oryza sativa                       | putative hexose carrier protein. OSJNBb0089A17.11.   |
| CAA53192.1 | X75440   | Chlorella kessleri                 | hexose transporter like protein. HUP3.   |
| CAA68813.1 | Y07520   | Chlorella kessleri                 | H(+)/hexose cotransporter (AA 1-533).  |
| CAA39036.1 | X55349   | Chlorella kessleri                 | H(+)/hexose-cotransporter. HUP1.   |
| BAB19862.1 | AB052883 | Oryza sativa                       | monosaccharide transporter 1. OsMST1.  |
| AAA18533.1 | L21752   | Saccharum hybrid cultivar H65-7052 | glucose transporter. putative.   |
| CAB52688.1 | AJ132223 | Lycopersicon esculentum            | hexose transporter. ht1.   |
| AAD55054.1 | AF173655 | Beta vulgaris                      | glucose transporter. Gt.   |
| CAB52690.1 | AJ132225 | Lycopersicon esculentum            | hexose transporter. ht3.   |
| AAA33875.1 | L31352   | Ricinus communis                   | hexose transport. hexose carrier. Hex9. putative.  |
| AAK13147.1 | AC083945 | Oryza sativa                       | Putative sugar transporter. OSJNBa0058E19.22.  |
| AAB68029.1 | U64903   | Beta vulgaris                      | BvcDNA-397. putative sugar transporter; member of major facilitative superfamily; integral membrane protein. |
| AAB68028.1 | U64902   | Beta vulgaris                      | BvcDNA-205. putative sugar transporter; member of major facilitative superfamily; integral membrane protein. |

|                |          |                                      |   |
|----------------|----------|--------------------------------------|---|
| AAG43998.1     | AF215837 | <i>Apium graveolens</i> var. dulce   | mannitol transporter. Mat1.   |
| AAF74568.1     | AF215854 | <i>Zea mays</i>                      | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                       |
| AAF74567.1     | AF215853 | <i>Solanum tuberosum</i>             | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                       |
| AAF74566.1     | AF215852 | <i>Nicotiana tabacum</i>             | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                       |
| AAF74565.1     | AF215851 | <i>Spinacia oleracea</i>             | transport of hexoses across the plastid inner envelope membrane. hexose transporter. pGlcT.                       |
| AAG46179.1     | AC018727 | <i>Oryza sativa</i>                  | putative sugar transporter protein. OSJNBa0056G17.3.  |
| AAG00995.1     | AF286906 | <i>Mesembryanthemum crystallinum</i> | putative glucose translocator. metabolite transporter; targeted to plastid inner envelope membrane.               |
| AAB53155.1     | U43629   | <i>Beta vulgaris</i>                 | putative sugar transporter. integral membrane protein. member of major facilitator superfamily.                   |
| AAB88879.1     | AF000952 | <i>Prunus armeniaca</i>              | putative sugar transporter.   |
| AAA33874.1     | L31353   | <i>Ricinus communis</i>              | hexose transport. hexose carrier. Hex10. putative.  |
| AAD37424.1     | AF149282 | <i>Phaseolus vulgaris</i>            | hexose carrier protein 1. HCP1.   |
| AAD45934.1     | AF168773 | <i>Betula pendula</i>                | hexose transport protein. HEX2.   |
| SEQ ID NO: 260 |          |                                      |   |
| AAA33915.1     | L27821   | <i>Oryza sativa</i>                  | receptor type serine/threonine kinase. protein kinase.  |
| AAF34428.1     | AF172282 | <i>Oryza sativa</i>                  | receptor-like protein kinase. DUPR11.18.  |
| BAA92954.1     | AP001551 | <i>Oryza sativa</i>                  | Similar to <i>Oryza sativa</i> protein kinase (OSPK10) mRNA. (L27821).  |
| BAB07906.1     | AP002835 | <i>Oryza sativa</i>                  | putative S-receptor kinase. P0417G05.14.  |
| BAA94516.1     | AP001800 | <i>Oryza sativa</i>                  | Similar to <i>Zea mays</i> S-domain receptor-like protein kinase (AJ010166).                                      |
| BAA94528.1     | AP001800 | <i>Oryza sativa</i>                  | Similar to <i>Arabidopsis thaliana</i> chromosome 2 BAC T20K24; putative receptor-like protein kinase (AC002392). |
| BAA94517.1     | AP001800 | <i>Oryza sativa</i>                  | Similar to <i>Zea mays</i> S-domain receptor-like protein kinase (AJ010166).                                      |

|            |          |                   |  |
|------------|----------|-------------------|--|
| BAB07905.1 | AP002835 | Oryza sativa      | putative S-receptor kinase. P0417G05.13.   |
| BAA94529.2 | AP001800 | Oryza sativa      | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB07904.1 | AP002835 | Oryza sativa      | putative S-receptor kinase. P0417G05.12.   |
| BAA94518.1 | AP001800 | Oryza sativa      | Similar to Arabidopsis thaliana chromosome 2 section 111 of 255; putative receptor-like protein kinase (AC002392). |
| CAA47962.1 | X67733   | Zea mays          | receptor-like protein kinase. PK1.   |
| BAA83573.1 | AP000399 | Oryza sativa      | Similar to serine/threonine-specific protein kinase PK10 precursor (AL021811).                                     |
| CAA73134.1 | Y12531   | Brassica oleracea | serine/threonine kinase. BRLK.   |
| CAA74662.1 | Y14286   | Brassica oleracea | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.   |
| BAB17345.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.29.   |
| BAB17348.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.32.   |
| BAB17342.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.26.   |
| BAB39451.1 | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.24.   |
| BAB17126.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.16.   |
| BAB19337.1 | AP003044 | Oryza sativa      | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).                             |
| CAB51480.1 | Y14600   | Sorghum bicolor   | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.                                   |
| BAB17339.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.23.   |
| AAB61708.1 | U93048   | Daucus carota     | somatic embryogenesis receptor-like kinase. SERK.  |
| AAF78016.1 | AF238472 | Oryza sativa      | receptor-like kinase. RLG15. protein kinase.   |
| AAD46420.1 | AF100771 | Hordeum vulgare   | receptor-like kinase. Hv3ARK. similar to wheat ARK1AS.   |
| AAC49629.1 | U51330   | Triticum aestivum | rust resistance kinase Lr10. LRK10.  |

|                |          |                   |   |
|----------------|----------|-------------------|---|
| BAB17139.1     | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.31.  |
| BAB17331.1     | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.12.  |
| AAC01746.1     | AF044489 | Oryza sativa      | receptor-like protein kinase. drpk1.  |
| AAC27489.1     | AF077130 | Oryza sativa      | receptor-like protein kinase.   |
| AAC02535.1     | AF044260 | Oryza sativa      | receptor serine/threonine kinase. protein kinase.   |
| BAB39434.1     | AP003338 | Oryza sativa      | receptor-like kinase. OJ1212_B09.1.   |
| AAF78020.1     | AF238476 | Oryza sativa      | receptor-like kinase. RLG11. protein kinase.  |
| BAA92953.1     | AP001551 | Oryza sativa      | Similar to Arabidopsis thaliana chromosome 4 BAC clone F10M6 ; S-receptor kinase -like protein. (AL021811). |
| AAF78018.1     | AF238474 | Oryza sativa      | receptor-like kinase. RLG16. protein kinase.  |
| AAD46917.1     | AF164021 | Oryza sativa      | receptor kinase.  |
| BAB39438.1     | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.7.   |
| BAB39435.1     | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.2.   |
| BAB17129.1     | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.20.  |
| BAB17321.1     | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.1.   |
| AAF68398.1     | AF237568 | Oryza sativa      | receptor-like protein kinase. RLG2.   |
| BAB39437.1     | AP003338 | Oryza sativa      | receptor-like kinase. OJ1212_B09.6.   |
| AAF78019.1     | AF238475 | Oryza sativa      | receptor-like kinase. RLG17. protein kinase.  |
| SEQ ID NO: 261 |          |                   |   |
| AAD09343.1     | AF026538 | Hordeum vulgare   | ABA-responsive protein.   |
| SEQ ID NO: 263 |          |                   |   |
| BAA22813.1     | D26015   | Nicotiana tabacum | aspartic protease activity. CND41, chloroplast nucleoid DNA binding protein. cnd41.                         |



|                |          |                      |  |
|----------------|----------|----------------------|--|
| BAB21205.1     | AP002913 | Oryza sativa         | nucleoid DNA-binding protein cnd41-like protein. P0480E02.11. contains ESTs AU166073(E31027),AU029516(E31027).       |
| SEQ ID NO: 270 |          |                      |  |
| BAB12719.1     | AP002746 | Oryza sativa         | putative regulatory protein NPR1. P0671B11.35.   |
| BAB16860.1     | AP002537 | Oryza sativa         | Arabidopsis thaliana regulatory protein NPR1 like protein. P0001B06.13.  |
| SEQ ID NO: 271 |          |                      |  |
| AAG35658.1     | AF204925 | Petroselinum crispum | transcription factor WRKY4. WRKY4. binds to W box (TTGACC) elements.   |
| BAA87058.1     | AB028022 | Nicotiana tabacum    | WIZZ. wizz. wound-induced transcription factor.  |
| BAA86031.1     | AB026890 | Nicotiana tabacum    | transcription factor NtWRKY4.  |
| AAD16139.1     | AF096299 | Nicotiana tabacum    | DNA-binding protein 2. WRKY2. transcription factor.  |
| BAA77383.1     | AB020590 | Nicotiana tabacum    | transcription factor NtWRKY2.  |
| AAF23898.1     | AF193802 | Oryza sativa         | zinc finger transcription factor WRKY1.  |
| AAD55974.1     | AF121353 | Petroselinum crispum | zinc-finger type transcription factor WRKY1. WRKY1.  |
| CAB97004.1     | AJ278507 | Solanum tuberosum    | putative transcription factor. WRKY DNA binding protein. WRKY1.  |
| BAA82107.1     | AB022693 | Nicotiana tabacum    | transcription factor. NtWRKY1.   |
| AAD32677.1     | AF140554 | Avena sativa         | DNA-binding protein WRKY1. wrky1. putative transcription factor.   |
| BAB16432.1     | AB041520 | Nicotiana tabacum    | WRKY transcription factor Nt-SubD48. Nt-SubD48.  |
| AAD16138.1     | AF096298 | Nicotiana tabacum    | DNA-binding protein 1. WRKY1. transcription factor.  |
| AAD32676.1     | AF140553 | Avena sativa         | DNA-binding protein WRKY3. wrky3. putative transcription factor.   |
| AAD27591.1     | AF121354 | Petroselinum crispum | binds sequence specifically to W Boxes (TTGACC). transcription factor. WRKY3. sequence specific DNA-binding protein. |
| AAF61864.1     | AF193771 | Nicotiana tabacum    | DNA-binding protein 4. WRKY4. transcription factor.  |
| AAG35659.1     | AF204926 | Petroselinum crispum | transcription factor WRKY5. WRKY5. binds to W box (TTGACC) elements.   |

|   |          |                         |
|---|----------|-------------------------|
| AAF61863.1  | AF193770 | Nicotiana tabacum       |
| DNA-binding protein 3. WRKY3. transcription factor. |          |                         |
| SEQ ID NO: 272                                      |          |                         |
| BAA07395.1  | D38220   | Brassica napus          |
| nitrate reductase.                                  |          |                         |
| BAA07394.1  | D38219   | Brassica napus          |
| nitrate reductase.                                  |          |                         |
| AAG30576.1  | AF314093 | Ricinus communis        |
| nitrate reductase. NIA.                             |          |                         |
| CAA32217.1  | X14059   | Nicotiana tabacum       |
| nitrate reductase.                                  |          |                         |
| AAA33713.1  | L13691   | Petunia x hybrida       |
| nitrate reductase. putative.                        |          |                         |
| CAA32218.1  | X14060   | Lycopersicon esculentum |
| nitrate reductase.                                  |          |                         |
| AAA33712.1  | L11563   | Petunia x hybrida       |
| nitrate reductase apoenzyme. nia.                   |          |                         |
| CAA32216.1  | X14058   | Nicotiana tabacum       |
| nitrate reductase.                                  |          |                         |
| CAA56696.1  | X80670   | Lotus japonicus         |
| nitrate reductase (NADH). NIA.                      |          |                         |
| AAB52786.1  | U95317   | Solanum tuberosum       |
| NADH nitrate reductase. StNR3.                      |          |                         |
| AAB18985.1  | U76701   | Solanum tuberosum       |
| NADH nitrate reductase. StNR2.                      |          |                         |
| AAA95940.1  | U01029   | Phaseolus vulgaris      |
| nitrate reductase. PVNR2.                           |          |                         |
| AAA34033.1  | M32600   | Spinacia oleracea       |
| NADH nitrate reductase.                             |          |                         |
| CAA38031.1  | X54097   | Betula pendula          |
| nitrate reductase (NADH). nia1.                     |          |                         |
| BAA13047.1  | D86226   | Spinacia oleracea       |
| nitrate reductase.                                  |          |                         |
| AAA33114.1  | M33154   | Cucurbita maxima        |
| nitrate reductase.                                  |          |                         |
| AAD19790.1  | AF055369 | Glycine max             |
| nitrate reductase. nr2.                             |          |                         |
| CAA58909.1  | X84103   | Cichorium intybus       |
| nitrate reductase (NADH). nia.                      |          |                         |
| AAA96813.1  | U13987   | Glycine max             |
| inducible nitrate reductase 2. INR2.                |          |                         |

|            |          |                           |  |
|------------|----------|---------------------------|--|
| CAA37672.1 | X53603   | Phaseolus vulgaris        | nitrate reductase.   |
| AAA96727.1 | L23854   | Glycine max               | nitrate reductase. INR1.   |
| AAA62316.1 | U20450   | Zea mays                  | nitrate reductase.   |
| AAD38068.1 | AF153448 | Zea mays                  | nitrate reductase. NR1.  |
| CAA40975.1 | X57844   | Hordeum vulgare           | nitrate reductase. cDNA is 9bp short of atg.   |
| CAA40976.1 | X57845   | Hordeum vulgare           | nitrate reductase.   |
| CAA42739.1 | X60173   | Hordeum vulgare           | nitrate reductase (NAD(P)H). nar7.   |
| AAB93560.1 | AF022780 | Glycine max               | nitrate reductase. BCNR-A.   |
| AAFI7595.1 | AF203033 | Chlamydomonas reinhardtii | nitrate reductase. NIT1.   |
| CAA45497.1 | X64136   | Volvox carteri            | nitrate reductase (NADH). nitA.  |
| AAC49460.1 | U39931   | Chlorella vulgaris        | nitrate reductase.   |
| AAC49459.1 | U39930   | Chlorella vulgaris        | nitrate reductase.   |
| CAA29497.1 | X06134   | Nicotiana tabacum         | nitrate reductase.   |
| AAA18377.1 | U08029   | Spinacia oleracea         | reduces nitrate to nitrite with NADH. NADH:nitrate reductase.  |
| AAB39553.1 | U64308   | Agrostemma githago        | nitrate reductase. agnr1. NADH; similar to agnr2 product encoded by GenBank Accession Number U64309 and to agnr3 product encoded by GenBank Accession Number U64310. |
| AAA03202.1 | M27821   | Zea mays                  | NADH:nitrate reductase; (EC 1.6.6.1).  |
| AAA33483.1 | M77792   | Zea mays                  | enzyme. nitrate reductase. NAR1S.  |
| AAB39555.1 | U64310   | Agrostemma githago        | nitrate reductase. agnr3. NADH; similar to agnr1 product encoded by GenBank Accession Number U64308 and agnr2 product encoded by GenBank Accession Number U64309.    |
| AAB39554.1 | U64309   | Agrostemma githago        | nitrate reductase. agnr2. NADH; similar to agnr1 product encoded by GenBank Accession Number U64308 and agnr3 product encoded by GenBank Accession Number U64310.    |

|                |          |                           |  |
|----------------|----------|---------------------------|--|
| CAA33819.1     | X15820   | Oryza sativa              | nitrate reductase apoenzyme (AA 472-916); Protein sequence is in conflict with the conceptual translation. |
| CAA33817.1     | X15819   | Oryza sativa              | nitrate reductase apoenzyme.   |
| AAA33998.1     | L23853   | Glycine max               | nitrate reductase. mutant.   |
| CAA58908.1     | X84102   | Cichorium intybus         | nitrate reductase (NADH). nia.   |
| CAA40090.1     | X56771   | Chlorella vulgaris        | nitrate reductase (NADH).  |
| CAA45776.1     | X64446   | Zea mays                  | nitrate reductase (NAD(P)H). nar.  |
| AAD17694.1     | AF077372 | Zea mays                  | possible reduction of Fe <sup>3+</sup> -chelates. cytochrome b5 reductase. NFR.                            |
| AAA96242.1     | L40147   | Avena strigosa            | nitrate reductase.   |
| AAB20155.1     | S61885   | Nicotiana plumbaginifolia | nitrate reductase heme domain. nitrate reductase heme domain, NR. This sequence comes from fig3; NR.       |
| AAA96245.1     | L40151   | Hordeum pusillum          | nitrate reductase.   |
| AAA96247.1     | L40153   | Hordeum stenostachys      | nitrate reductase.   |
| SEQ ID NO: 273 |          |                           |  |
| BAA07395.1     | D38220   | Brassica napus            | nitrate reductase.   |
| BAA07394.1     | D38219   | Brassica napus            | nitrate reductase.   |
| AAA33713.1     | L13691   | Petunia x hybrida         | nitrate reductase. putative.   |
| CAA32218.1     | X14060   | Lycopersicon esculentum   | nitrate reductase.   |
| AAA33712.1     | L11563   | Petunia x hybrida         | nitrate reductase apoenzyme. nia.  |
| AAG30576.1     | AF314093 | Ricinus communis          | nitrate reductase. NIA.  |
| CAA32217.1     | X14059   | Nicotiana tabacum         | nitrate reductase.   |
| CAA32216.1     | X14058   | Nicotiana tabacum         | nitrate reductase.   |
| AAA33114.1     | M33154   | Cucurbita maxima          | nitrate reductase.   |

|            |          |                           |  |
|------------|----------|---------------------------|--|
| AAB52786.1 | U95317   | Solanum tuberosum         | NADH nitrate reductase. StNR3.   |
| AAB18985.1 | U76701   | Solanum tuberosum         | NADH nitrate reductase. StNR2.   |
| AAA34033.1 | M32600   | Spinacia oleracea         | NADH nitrate reductase.  |
| BAA13047.1 | D86226   | Spinacia oleracea         | nitrate reductase.   |
| CAA38031.1 | X54097   | Betula pendula            | nitrate reductase (NADH). nia1.  |
| CAA56696.1 | X80670   | Lotus japonicus           | nitrate reductase (NADH). NIA.   |
| AAA95940.1 | U01029   | Phaseolus vulgaris        | nitrate reductase. PVNR2.  |
| CAA58909.1 | X84103   | Cichorium intybus         | nitrate reductase (NADH). nia.   |
| AAD19790.1 | AF055369 | Glycine max               | nitrate reductase. nr2.  |
| AAA96813.1 | U13987   | Glycine max               | inducible nitrate reductase 2. INR2.   |
| CAA40976.1 | X57845   | Hordeum vulgare           | nitrate reductase.   |
| AAA96727.1 | L23854   | Glycine max               | nitrate reductase. INR1.   |
| CAA37672.1 | X53603   | Phaseolus vulgaris        | nitrate reductase.   |
| CAA33819.1 | X15820   | Oryza sativa              | nitrate reductase apoenzyme (AA 472-916); Protein sequence is in conflict with the conceptual translation. |
| AAD38068.1 | AF153448 | Zea mays                  | nitrate reductase. NR1.  |
| CAA40975.1 | X57844   | Hordeum vulgare           | nitrate reductase. cDNA is 9bp short of atg.   |
| AAB93560.1 | AF022780 | Glycine max               | nitrate reductase. BCNR-A.   |
| AAA62316.1 | U20450   | Zea mays                  | nitrate reductase.   |
| CAA42739.1 | X60173   | Hordeum vulgare           | nitrate reductase (NAD(P)H). nar7.   |
| AAF17595.1 | AF203033 | Chlamydomonas reinhardtii | nitrate reductase. NIT1.   |
| CAA45497.1 | X64136   | Volvox carteri            | nitrate reductase (NADH). nitA.  |

|                |          |                      |  |
|----------------|----------|----------------------|--|
| AAC49460.1     | U39931   | Chlorella vulgaris   | nitrate reductase.   |
| AAC49459.1     | U39930   | Chlorella vulgaris   | nitrate reductase.   |
| CAA29497.1     | X06134   | Nicotiana tabacum    | nitrate reductase.   |
| AAA18377.1     | U08029   | Spinacia oleracea    | reduces nitrate to nitrite with NADH. NADH:nitrate reductase.  |
| AAB39553.1     | U64308   | Agrostemma githago   | nitrate reductase. agnr1. NADH; similar to agnr2 product encoded by GenBank Accession Number U64309 and to agnr3 product encoded by GenBank Accession Number U64310. |
| AAA03202.1     | M27821   | Zea mays             | NADH:nitrate reductase; (EC 1.6.6.1).  |
| AAA33483.1     | M77792   | Zea mays             | enzyme. nitrate reductase. NAR1S.  |
| AAB39555.1     | U64310   | Agrostemma githago   | nitrate reductase. agnr3. NADH; similar to agnr1 product encoded by GenBank Accession Number U64308 and agnr2 product encoded by GenBank Accession Number U64309.    |
| AAB39554.1     | U64309   | Agrostemma githago   | nitrate reductase. agnr2. NADH; similar to agnr1 product encoded by GenBank Accession Number U64308 and agnr3 product encoded by GenBank Accession Number U64310.    |
| CAA33817.1     | X15819   | Oryza sativa         | nitrate reductase apoenzyme.   |
| CAA58908.1     | X84102   | Cichorium intybus    | nitrate reductase (NADH). nia.   |
| AAA33998.1     | L23853   | Glycine max          | nitrate reductase. mutant.   |
| CAA40090.1     | X56771   | Chlorella vulgaris   | nitrate reductase (NADH).  |
| CAA45776.1     | X64446   | Zea mays             | nitrate reductase (NAD(P)H). nar.  |
| AAD17694.1     | AF077372 | Zea mays             | possible reduction of Fe <sup>3+</sup> -chelates. cytochrome b5 reductase. NFR.  |
| AAA96242.1     | L40147   | Avena strigosa       | nitrate reductase.   |
| AAA96250.1     | L40149   | Hordeum chilense     | nitrate reductase.   |
| AAA96245.1     | L40151   | Hordeum pusillum     | nitrate reductase.   |
| AAA96247.1     | L40153   | Hordeum stenostachys | nitrate reductase.   |
| SEQ ID NO: 274 |          |                      |  |

|            |          |  |
|------------|----------|--|
| AAC39318.1 | AF029858 | <i>Sorghum bicolor</i><br>second multifunctional cytochrome P450 in the biosynthetic pathway of the cyanogenic glucoside dhurrin. Catalyzes the conversion of p-hydroxyphenylacetaldoxime to p-hydroxymandelonitrile. cytochrome P450 CYP71E1. CYP71E1. No EST#s identified. |
| BAB40323.1 | AB037244 | <i>Asparagus officinalis</i><br>cytochrome P450. ASPI-1.   |
| AAA32913.1 | M32885   | <i>Persea americana</i><br>cytochrome P-450LXXIA1 (cyp71A1).   |
| BAB40324.1 | AB037245 | <i>Asparagus officinalis</i><br>cytochrome P450. ASPI-2.   |
| AAA19701.1 | L24438   | <i>Thlaspi arvense</i><br>cytochrome P450.   |
| CAA71513.1 | Y10489   | <i>Glycine max</i><br>putative cytochrome P450.  |
| AAB94589.1 | AF022460 | <i>Glycine max</i><br>CYP83D1p. CYP83D1. cytochrome P450 monooxygenase.  |
| AAB94588.1 | AF022459 | <i>Glycine max</i><br>CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.  |
| AAB61965.1 | U48435   | <i>Solanum chacoense</i><br>putative cytochrome P450.  |
| CAA70575.1 | Y09423   | <i>Nepeta racemosa</i><br>cytochrome P450. CYP71A5.  |
| CAA71517.1 | Y10493   | <i>Glycine max</i><br>putative cytochrome P450.  |
| AAD47832.1 | AF166332 | <i>Nicotiana tabacum</i><br>cytochrome P450.   |
| AAK38084.1 | AF321860 | <i>Lolium rigidum</i><br>putative cytochrome P450.   |
| AAB94584.1 | AF022157 | <i>Glycine max</i><br>capable of catalyzing the metabolism of phenylurea herbicides. CYP71A10. CYP71A10. cytochrome P450 monooxygenase.  |
| AAF27282.1 | AF122821 | <i>Capsicum annuum</i><br>cytochrome P450. PepCYP.   |
| AAK38083.1 | AF321859 | <i>Lolium rigidum</i><br>putative cytochrome P450.   |
| CAA50312.1 | X70981   | <i>Solanum melongena</i><br>P450 hydroxylase. CYPEG2.  |
| CAB56503.1 | AJ238612 | <i>Catharanthus roseus</i><br>cytochrome P450.   |
| AAB61964.1 | U48434   | <i>Solanum chacoense</i><br>putative cytochrome P450.  |
| AAK38087.1 | AF321863 | <i>Lolium rigidum</i><br>putative cytochrome P450.   |

|                |          |  |
|----------------|----------|--|
| CAA71514.1     | Y10490   | Glycine max<br>putative cytochrome P450.   |
| CAA50645.1     | X71654   | Solanum melongena<br>P450 hydroxylase.   |
| BAA03635.1     | D14990   | Solanum melongena<br>Cytochrome P-450EG4.  |
| AAD44151.1     | AF124816 | Mentha x piperita<br>cytochrome p450 isoform PM17.   |
| AAD44152.1     | AF124817 | Mentha x piperita<br>cytochrome p450 isoform PM2.  |
| CAA83941.1     | Z33875   | Mentha x piperita<br>cytochrome P-450 oxidase.   |
| AAD44150.1     | AF124815 | Mentha spicata<br>cytochrome p450.   |
| AAB69644.1     | AF000403 | Lotus japonicus<br>putative cytochrome P450. LjNP450.  |
| BAB40322.1     | AB036772 | Triticum aestivum<br>cytochrome P450. N-1.   |
| CAC27827.1     | AJ295719 | Catharanthus roseus<br>geraniol hydroxylase. cytochrome P450. cyp71.   |
| AAG44132.1     | AF218296 | Pisum sativum<br>cytochrome P450. P450 isolog.   |
| AAD56282.1     | AF155332 | Petunia x hybrida<br>flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.  |
| AAG14963.1     | AF214009 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H3.   |
| AAD37433.1     | AF150881 | Lycopersicon esculentum x Lycopersicon peruvianum<br>catalyzes the hydroxylation of ferulic acid to 5-hydroxyferulic acid. ferulate-5-hydroxylase.<br>CYP84. cytochrome P450-dependent monooxygenase; F5H; FAH1. |
| BAB39252.1     | AP002968 | Oryza sativa<br>putative cytochrome P450. P0416G11.1.  |
| AAK38088.1     | AF321864 | Lolium rigidum<br>putative cytochrome P450.  |
| CAA65580.1     | X96784   | Nicotiana tabacum<br>cytochrome P450. hsr515.  |
| AAG14961.1     | AF214007 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H1.   |
| AAG14962.1     | AF214008 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H2.   |
| AAB17562.1     | U72654   | Eustoma grandiflorum<br>flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.  |
| SEQ ID NO: 279 |          |  |



|            |          |                            |  |
|------------|----------|----------------------------|--|
| AAA34122.1 | M84466   | Nicotiana tabacum          | phenylalanine ammonia lyase. tpal.   |
| BAA22948.1 | AB008200 | Nicotiana tabacum          | phenylalanine ammonia-lyase. palB.   |
| AAA34176.1 | M90692   | Lycopersicon esculentum    | phenylalanine ammonia-lyase. PAL5.   |
| AAF40224.1 | AF237955 | Rubus idaeus               | phenylalanine ammonia-lyase 2. PAL2. PAL; phenylpropanoid; multigene; flavonoid. |
| CAA37129.1 | X52953   | Glycine max                | phenylalanine ammonia-lyase. PAL1.   |
| CAA68036.1 | X99705   | Triticum aestivum          | phenylalanine ammonia-lyase. PAL.  |
| AAA33389.1 | M29232   | Ipomoea batatas            | phenylalanine ammonia-lyase.   |
| AAA34179.2 | M83314   | Lycopersicon esculentum    | deamination of phenylalanine to coumarate. phenylalanine ammonia lyase. pal.     |
| BAA21643.1 | D30656   | Populus kitakamiensis      | phenylalanine ammonia-lyase.   |
| AAB67733.1 | U43338   | Citrus limon               | phenylalanine ammonia-lyase. pal6.   |
| BAA95629.1 | AB042520 | Catharanthus roseus        | phenylalanine ammonia lyase.   |
| BAA05643.1 | D26596   | Camellia sinensis          | phenylalanine ammonia-lyase.   |
| CAA73065.1 | Y12461   | Helianthus annuus          | phenylalanine ammonia lyase. PAL.  |
| BAA24929.1 | D83076   | Lithospermum erythrorhizon | phenylalanine ammonia-lyase.   |
| BAA24928.1 | D83075   | Lithospermum erythrorhizon | phenylalanine ammonia-lyase.   |
| BAA00885.1 | D10001   | Pisum sativum              | phenylalanine ammonia-lyase.   |
| AAA84889.1 | U39792   | Pinus taeda                | phenylalanine ammonia-lyase. lpPAL.  |
| CAA61198.1 | X87946   | Oryza sativa               | phenylalanine ammonia-lyase. ZB8.  |
| CAA41169.1 | X58180   | Medicago sativa            | phenylalanine ammonia-lyase. PAL.  |
| BAA00887.1 | D10003   | Pisum sativum              | phenylalanine ammonia-lyase. PAL2.   |
| BAA00886.1 | D10002   | Pisum sativum              | phenylalanine ammonia-lyase. PAL1.   |

|            |          |                                       |  |
|------------|----------|---------------------------------------|--|
| AAA17993.1 | M91192   | Trifolium subterraneum                | phenylalanine ammonia-lyase. PAL1.   |
| AAA33805.1 | L11747   | Populus x generosa                    | phenylalanine ammonia lyase. PAL.  |
| AAC78457.1 | AF036948 | Prunus avium                          | phenylalanine ammonia-lyase. PAL1.   |
| BAA23367.1 | D85850   | Daucus carota                         | phenylalanine ammonia-lyase. gDcPAL1.  |
| CAB42793.1 | AJ238753 | Citrus clementina x Citrus reticulata | phenylalanine-ammonia lyase. pal1.   |
| AAA99500.1 | L36822   | Stylosanthes humilis                  | phenylalanine ammonia lyase. PAL17.1.  |
| CAA55075.1 | X78269   | Nicotiana tabacum                     | phenylalanine ammonia-lyase.   |
| BAA22963.1 | D17467   | Nicotiana tabacum                     | phenylalanine ammonia-lyase. TOBPAL1.  |
| BAA22947.1 | AB008199 | Nicotiana tabacum                     | phenylalanine ammonia-lyase. palA.   |
| CAA57057.1 | X81159   | Petroselinum crispum                  | phenylalanine ammonia-lyase 3. PAL3. tetramere subunit.                          |
| AAG49585.1 | AF325496 | Ipomoea nil                           | phenylalanine ammonia-lyase.   |
| CAA57056.1 | X81158   | Petroselinum crispum                  | phenylalanine ammonia-lyase 2. PAL2. deaminase subunit.                          |
| CAB42794.1 | AJ238754 | Citrus clementina x Citrus reticulata | phenylalanine-ammonia lyase. pal2.   |
| CAA05251.1 | AJ002221 | Digitalis lanata                      | phenylalanine ammonia lyase.   |
| BAA07860.1 | D43802   | Populus kitakamiensis                 | phenylalanine ammonia-lyase.   |
| CAB60719.1 | AJ250836 | Cicer arietinum                       | phenylpropanoid pathway. phenylalanine ammonia-lyase. pal.                       |
| CAA68256.1 | X99997   | Bromheadia finlaysoniana              | phenylalanine ammonia-lyase. pal.  |
| AAK15640.1 | AF326116 | Agastache rugosa                      | phenylalanine ammonia-lyase. PAL.  |
| CAA34226.1 | X16099   | Oryza sativa subsp. japonica          | phenylalanine ammonia-lyase.   |
| AAF40223.1 | AF237954 | Rubus idaeus                          | phenylalanine ammonia-lyase 1. PAL1. PAL; phenylpropanoid; multigene; flavonoid. |
| BAA11459.1 | D78640   | Ipomoea batatas                       | Phenylalanine Ammonia-Lyase.   |

BAA06337.1 D30657 *Populus kitakamiensis*  
phenylalanine ammonia-lyase.

AAD45384.1 AF165998 *Vigna unguiculata*  
phenylalanine ammonia-lyase.

CAA53733.1 X76130 *Cucumis melo*  
phenylalanine ammonia-lyase. pal.

AAA51873.1 U16130 *Persea americana*  
phenylalanine ammonia lyase. PAL.

BAB19128.1 AB041361 *Dianthus caryophyllus*  
phenylalanine ammonia-lyase. Dcpal1.

CAA34715.1 X16772 *Petroselinum crispum*  
phenylalanine ammonia-lyase (AA 137 - 716) (1 is 3rd base in codon).

BAA07861.1 D43803 *Populus kitakamiensis*  
phenylalanine ammonia-lyase.

SEQ ID NO: 280

AAG43550.1 AF211532 *Nicotiana tabacum*  
Avr9/Cf-9 rapidly elicited protein 132. ACRE132. similar to RING finger proteins.

AAK00436.1 AC060755 *Oryza sativa*  
putative zinc finger protein. OSJNBa0003O19.23.

BAA78746.1 AB023482 *Oryza sativa*  
Similar to *Arabidopsis thaliana* RING-H2 finger protein RHX1a mRNA, partial cds.(AF079184).

CAA74911.1 Y14573 *Hordeum vulgare*  
ring finger protein. putative.

AAG46117.1 AC073166 *Oryza sativa*  
putative ring finger protein. OSJNBb0064P21.7.

BAA96875.1 AB045121 *Oryza sativa*  
RING finger 1. RRF1.

BAA90357.1 AP001080 *Oryza sativa*  
EST AU070319(S10788) corresponds to a region of the predicted gene. Similar to RING-H2 finger protein RHA2b (AC006200).

BAA90806.1 AP001168 *Oryza sativa*  
ESTs C26000(C11448),AU082130(C11448) correspond to a region of the predicted gene.;  
Similar to mRNA for zinc-finger protein (Z36749).

SEQ ID NO: 286

AAG14454.1 AF283706 *Tulipa gesneriana*  
auxin-induced protein TGSAUR12. SAUR12. small auxin upregulated RNA.

AAG14456.1 AF283708 *Tulipa gesneriana*  
auxin-induced protein TGSAUR22. SAUR22. small auxin upregulated RNA.

AAG14455.1 AF283707 *Tulipa gesneriana*  
auxin-induced protein TGSAUR21. SAUR21. small auxin upregulated RNA.

AAC08401.1 AF053564 *Mesembryanthemum crystallinum*  
auxin-induced protein. similar to auxin-induced proteins from soybean.

## SEQ ID NO: 290

AAB65498.1 U73856 *Chlamydomonas reinhardtii*  
carbonic anhydrase, alpha type. CAH3.

AAC49983.1 U40871 *Chlamydomonas reinhardtii*  
intracellular carbonic anhydrase, alpha type. CAH3.

AAF04292.2 AF190735 *Dunaliella salina*  
carbonic anhydrase. CA.

AAC49378.1 U53811 *Dunaliella salina*  
carbonic anhydrase. dca.

AAF22644.1 AF183939 *Dunaliella salina*  
duplicated carbonic anhydrase. DCA1. DCA; carbonic anhydrase gene family member; salt-inducible; intra-duplicated.

AAD51633.1 AF170173 *Acetabularia acetabulum*  
putative carbonic anhydrase 2. CA2. AaCA2.

AAD51634.1 AF170174 *Acetabularia acetabulum*  
putative carbonic anhydrase 1. CA1. AaCA1.

AAD51635.1 AF170175 *Acetabularia acetabulum*  
putative carbonic anhydrase 1. CA1. AaCA1.

BAA14232.1 D90206 *Chlamydomonas reinhardtii*  
carbonic anhydrase.

BAA28217.1 AB013804 *Chlorella sorokiniana*  
soluble carbonic anhydrase precursor. CAH1.

## SEQ ID NO: 301

AAG03089.2 AC073405 *Oryza sativa*  
similar to an Arabidopsis putative P-type transporting ATPase (AC010926).

BAA89544.1 AP001072 *Oryza sativa*  
Similar to chromaffin granule ATPase II homolog. (U75321).

BAA88191.1 AP000836 *Oryza sativa*  
Similar to chromaffin granule ATPase II homolog (U75321).

BAA90510.2 AP001111 *Oryza sativa*  
rice EST AU030811, similar to rice Ca<sup>2+</sup>-ATPase (U82966).

AAD11618.1 AF050496 *Lycopersicon esculentum*  
Ca<sup>2+</sup>-ATPase. LCA1B; alternative transcript.

AAA34138.1 M96324 *Lycopersicon esculentum*  
The calcium ATPase is a calcium ion pump. Ca<sup>2+</sup>-ATPase. LCA1.

AAD11617.1 AF050495 *Lycopersicon esculentum*  
Ca<sup>2+</sup>-ATPase. LCA1A; alternative transcript.

AAF73985.1 AF096871 *Zea mays*  
calcium pump. calcium ATPase. cap1.

AAD31896.1 AF145478 *Mesembryanthemum crystallinum*  
calcium ATPase.

AAG28436.1 AF195029 *Glycine max*  
plasma membrane Ca<sup>2+</sup>-ATPase. SCA2.

|            |          |                           |  |
|------------|----------|---------------------------|--|
| CAA63790.1 | X93592   | Dunaliella bioculata      | P-type ATPase. cal. calcium pumping; CA1.                                |
| AAG28435.1 | AF195028 | Glycine max               | plasma membrane Ca <sup>2+</sup> -ATPase. SCA1.                          |
| CAA68234.1 | X99972   | Brassica oleracea         | calmodulin-stimulated calcium-ATPase.                                    |
| AAB58910.1 | U82966   | Oryza sativa              | Ca <sup>2+</sup> -ATPase.  |
| CAB69824.1 | AJ271439 | Prunus persica            | plasma membrane H <sup>+</sup> ATPase. PPA1.                             |
| AAB60276.1 | U09989   | Zea mays                  | H(+)-transporting ATPase. Mha1.  |
| BAA01058.1 | D10207   | Oryza sativa              | H-ATPase. OSA1.  |
| CAC29435.1 | AJ310523 | Vicia faba                | P-type H <sup>+</sup> -ATPase. vha4. predominantly expressed in flowers. |
| AAD20330.1 | AF110268 | Oryza sativa              | plasma membrane proton-ATPase gene OSA3.                                 |
| AAA34098.1 | M80490   | Nicotiana plumbaginifolia | plasma membrane H <sup>+</sup> ATPase. pma3.                             |
| AAB84203.1 | AF029257 | Kosteletzkya virginica    | plasma membrane H <sup>+</sup> -ATPase.                                  |
| CAC28224.1 | AJ286749 | Sesbania rostrata         | p-type H <sup>+</sup> -ATPase. ha5.                                      |
| AAD46188.1 | AF156691 | Nicotiana plumbaginifolia | plasma membrane proton ATPase. pma9.                                     |
| AAA34173.1 | M60166   | Lycopersicon esculentum   | H <sup>+</sup> -ATPase. LHA1.  |
| AAA34094.1 | M80489   | Nicotiana plumbaginifolia | plasma membrane H <sup>+</sup> ATPase. pma1.                             |
| AAA34052.1 | M27888   | Nicotiana plumbaginifolia | H <sup>+</sup> -translocating ATPase.                                    |
| CAC28221.1 | AJ286746 | Sesbania rostrata         | p-type H <sup>+</sup> -ATPase. ha2.                                      |
| CAA54045.1 | X76535   | Solanum tuberosum         | H(+)-transporting ATPase. PHA2.  |
| BAA06629.1 | D31843   | Oryza sativa              | plasma membrane H <sup>+</sup> -ATPase. OSA2.                            |
| CAA64406.1 | X94936   | Phaseolus vulgaris        | H(+)-transporting ATPase. BHA-2.   |
| AAF98344.1 | AF275745 | Lycopersicon esculentum   | plasma membrane H <sup>+</sup> -ATPase. LHA2. P-type ion pump.           |

AAD55399.1 AF179442 *Lycopersicon esculentum*  
plasma membrane H<sup>+</sup>-ATPase isoform LHA2. LHA2.

CAA54046.1 X76536 *Solanum tuberosum*  
H(+)-transporting ATPase. PHA1.

SEQ ID NO: 302

CAA70575.1 Y09423 *Nepeta racemosa*  
cytochrome P450. CYP71A5.

CAA70576.1 Y09424 *Nepeta racemosa*  
cytochrome P450. CYP71A6.

CAA50312.1 X70981 *Solanum melongena*  
P450 hydroxylase. CYPEG2.

AAA32913.1 M32885 *Persea americana*  
cytochrome P-450LXXIA1 (cyp71A1).

BAA03635.1 D14990 *Solanum melongena*  
Cytochrome P-450EG4.

CAA50645.1 X71654 *Solanum melongena*  
P450 hydroxylase.

CAA83941.1 Z33875 *Mentha x piperita*  
cytochrome P-450 oxidase.

CAA71513.1 Y10489 *Glycine max*  
putative cytochrome P450.

AAB94584.1 AF022157 *Glycine max*  
capable of catalyzing the metabolism of phenylurea herbicides. CYP71A10. CYP71A10.  
cytochrome P450 monooxygenase.

AAB61965.1 U48435 *Solanum chacoense*  
putative cytochrome P450.

AAB69644.1 AF000403 *Lotus japonicus*  
putative cytochrome P450. LjNP450.

BAB40324.1 AB037245 *Asparagus officinalis*  
cytochrome P450. ASPI-2.

AAD47832.1 AF166332 *Nicotiana tabacum*  
cytochrome P450.

AAB61964.1 U48434 *Solanum chacoense*  
putative cytochrome P450.

BAB40323.1 AB037244 *Asparagus officinalis*  
cytochrome P450. ASPI-1.

CAA71514.1 Y10490 *Glycine max*  
putative cytochrome P450.

AAB94588.1 AF022459 *Glycine max*  
CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.

AAF27282.1 AF122821 *Capsicum annuum*  
cytochrome P450. PepCYP.

|            |          |                     |  |
|------------|----------|---------------------|--|
| CAA50313.1 | X70982   | Solanum melongena   | P450 hydroxylase. CYPEG3.  |
| AAA19701.1 | L24438   | Thlaspi arvense     | cytochrome P450.   |
| BAA12159.1 | D83968   | Glycine max         | Cytochrome P-450 (CYP93A1).  |
| AAK38082.1 | AF321858 | Lolium rigidum      | putative cytochrome P450.  |
| BAB40322.1 | AB036772 | Triticum aestivum   | cytochrome P450. N-1.  |
| CAA71516.1 | Y10492   | Glycine max         | putative cytochrome P450.  |
| CAA65580.1 | X96784   | Nicotiana tabacum   | cytochrome P450. hsr515.   |
| AAK38083.1 | AF321859 | Lolium rigidum      | putative cytochrome P450.  |
| AAK38084.1 | AF321860 | Lolium rigidum      | putative cytochrome P450.  |
| CAA71517.1 | Y10493   | Glycine max         | putative cytochrome P450.  |
| CAA72196.1 | Y11368   | Zea mays            | cytochrome p450. cyp71c4.  |
| CAA57425.1 | X81831   | Zea mays            | cytochrome P450. CYP71C4. family CYP71, subfamily CYP71C.  |
| AAC39318.1 | AF029858 | Sorghum bicolor     | second multifunctional cytochrome P450 in the biosynthetic pathway of the cyanogenic glucoside dhurrin. Catalyzes the conversion of p-hydroxyphenylacetaldoxime to p-hydroxymandelonitrile. cytochrome P450 CYP71E1. CYP71E1. No EST#s identified. |
| AAD56282.1 | AF155332 | Petunia x hybrida   | flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.   |
| CAB56503.1 | AJ238612 | Catharanthus roseus | cytochrome P450.   |
| AAK38087.1 | AF321863 | Lolium rigidum      | putative cytochrome P450.  |
| CAA57421.1 | X81827   | Zea mays            | cytochrome P450. CYP71C1. family CYP71, subfamily CYP71C.  |
| CAA57422.1 | X81828   | Zea mays            | cytochrome P450. CYP71C1. family CYP71, subfamily CYP71C.  |
| BAB39252.1 | AP002968 | Oryza sativa        | putative cytochrome P450. P0416G11.1.  |
| BAA13076.1 | D86351   | Glycine max         | cytochrome P-450 (CYP93A2).  |

|                |          |                     |   |
|----------------|----------|---------------------|---|
| AAB94589.1     | AF022460 | Glycine max         | CYP83D1p. CYP83D1. cytochrome P450 monooxygenase.   |
| AAD38930.1     | AF135485 | Glycine max         | cytochrome P450 monooxygenase CYP93D1. CYP93E1.   |
| CAA57423.1     | X81829   | Zea mays            | cytochrome P450. CYP71C2. family CYP71, subfamily CYP71C.   |
| CAA72208.1     | Y11404   | Zea mays            | cytochrome p450. cyp71c2.   |
| AAG44132.1     | AF218296 | Pisum sativum       | cytochrome P450. P450 isolog.   |
| AAK38088.1     | AF321864 | Lolium rigidum      | putative cytochrome P450.   |
| CAA64635.1     | X95342   | Nicotiana tabacum   | cytochrome P450. hsr515. hypersensitivity-related gene.   |
| SEQ ID NO: 303 |          |                     |   |
| AAC49826.1     | U71604   | Catharanthus roseus | involved in the second to last step in vindoline biosynthesis. desacetoxyvindoline 4-hydroxylase. CRSD4H. 2-oxoglutarate dependent dioxygenase.                   |
| AAB97311.1     | AF008597 | Catharanthus roseus | desacetoxyvindoline-4-hydroxylase. CRSD4H. 2-oxoglutarate dependent dioxygenase; involved in the second to last step in vindoline biosynthesis.                   |
| AAC49827.1     | U71605   | Catharanthus roseus | involved in the second to last step in vindoline biosynthesis. desacetoxyvindoline 4-hydroxylase. CRSD4H. 2-oxoglutarate dependent dioxygenase.                   |
| BAA95828.1     | AP002069 | Oryza sativa        | ESTs D47168(S12332), D46350(S10967) correspond to a region of the predicted gene. Similar to Prunus armeniaca ethylene-forming-enzyme-like dioxygenase. (U97530). |
| BAA37127.1     | AB012203 | Lactuca sativa      | 2-oxoglutarate-dependent dioxygenase. gibberelin 20-oxidase. Ls20ox1.   |
| CAA54557.1     | X77368   | Solanum melongena   | dioxygenase. DIOX.  |
| BAA81862.1     | AB026295 | Oryza sativa        | Similar to leucoanthocyanidin dioxygenase.(AI440611).   |
| SEQ ID NO: 304 |          |                     |   |
| AAF61647.1     | AF190634 | Nicotiana tabacum   | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAA89009.1     | AB027455 | Petunia x hybrida   | anthocyanin 5-O-glucosyltransferase. PH1.   |
| AAF17077.1     | AF199453 | Sorghum bicolor     | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |



|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAF98390.1 | AF287143 | Brassica napus          | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT. |
| BAA36423.1 | AB013598 | Verbena x hybrida       | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| AAK16181.1 | AC079887 | Oryza sativa            | putative glucosyltransferase. OSJNBa0040E01.16.   |
| BAA36421.1 | AB013596 | Perilla frutescens      | UDP-glucose:anthocynsin 5-O-glucosyltransferase. PF3R4.   |
| BAB07962.1 | AP002524 | Oryza sativa            | putative anthocyanin 5-O-glucosyltransferase. P0406H10.16. contains ESTs AU067881(C10481),AU067882(C10481).   |
| AAK16178.1 | AC079887 | Oryza sativa            | putative glucosyltransferase. OSJNBa0040E01.5.  |
| BAA93039.1 | AB033758 | Citrus unshiu           | limonoid UDP-glucosyltransferase. LGTase.   |
| AAK16172.1 | AC079887 | Oryza sativa            | putative glucosyltransferase. OSJNBa0040E01.14.   |
| BAA36422.1 | AB013597 | Perilla frutescens      | UDP-glucose:anthocyanin 5-O-glucosyltransferase homologue. PF3R6.   |
| BAB17182.1 | AP002843 | Oryza sativa            | putative UTP-glucose glucosyltransferase. P0407B12.19.  |
| BAA12737.1 | D85186   | Gentiana triflora       | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| AAG25643.1 | AF303396 | Phaseolus vulgaris      | UDP-glucosyltransferase HRA25. putative; defense associated.  |
| AAB48444.1 | U82367   | Solanum tuberosum       | UDP-glucose glucosyltransferase.  |
| BAA19659.1 | AB002818 | Perilla frutescens      | flavonoid 3-O-glucosyltransferase. UDP glucose.   |
| CAA54612.1 | X77462   | Manihot esculenta       | UTP-glucose glucosyltransferase. CGT5.  |
| BAA89008.1 | AB027454 | Petunia x hybrida       | anthocyanidin 3-O-glucosyltransferase. PGT8.  |
| AAK16175.1 | AC079887 | Oryza sativa            | putative glucosyltransferase. OSJNBa0040E01.15.   |
| BAA83484.1 | AB031274 | Scutellaria baicalensis | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.   |
| AAB36652.1 | U32643   | Nicotiana tabacum       | immediate-early salicylate-induced glucosyltransferase. IS10a.  |
| AAK28304.1 | AF346432 | Nicotiana tabacum       | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.  |

|                |          |                                 |   |
|----------------|----------|---------------------------------|---|
| AAB36653.1     | U32644   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS5a.   |
| AAD04166.1     | AF101972 | Phaseolus lunatus               | catalyzes formation of O-glucosylzeatin from zeatin and UDPG; catalyzes formation of O-xylosylzeatin from zeatin and UDPX. zeatin O-glucosyltransferase. ZOG1. cytokinin O-glucosyltransferase. |
| AAD21086.1     | AF127218 | Forsythia x intermedia          | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.   |
| CAA54611.1     | X77461   | Manihot esculenta               | UTP-glucose glucosyltransferase. CGT2.  |
| AAK28303.1     | AF346431 | Nicotiana tabacum               | phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.  |
| CAA54609.1     | X77459   | Manihot esculenta               | UTP-glucose glucosyltransferase. CGT1.  |
| CAA54613.1     | X77463   | Manihot esculenta               | UTP-glucose glucosyltransferase. CGT6.  |
| BAB41026.1     | AB047099 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT2.   |
| BAB41024.1     | AB047097 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT2.   |
| BAB41022.1     | AB047095 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT2.   |
| BAB41017.1     | AB047090 | Vitis labrusca x Vitis vinifera | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT1. The gene was derived from one of the parents V. labruscana cv. Ishiharawase.  |
| BAB41020.1     | AB047093 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT2.   |
| CAA59450.1     | X85138   | Lycopersicon esculentum         | twil. homologous to glucosyltransferases.   |
| BAB41025.1     | AB047098 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.   |
| BAB41023.1     | AB047096 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.   |
| BAB41018.1     | AB047091 | Vitis labrusca x Vitis vinifera | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT2. The gene originated in one of the parents V. vinifera cv. Centennial.   |
| SEQ ID NO: 306 |          |                                 |   |
| AAB06458.1     | U64806   | Brassica napus                  | pathogenesis-related protein PR1. Ypr1.   |
| AAB01666.1     | U21849   | Brassica napus                  | PR-1a. LSC94.   |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAB09587.1 | U70666   | Brassica napus          | pathogenesis-related protein PR1. Ypr1.   |
| CAA47374.1 | X66942   | Nicotiana tabacum       | prb-1b. PRB-1B.   |
| AAK30143.1 | AF348141 | Capsicum annuum         | pathogenesis-related protein PR-1 precursor.                                      |
| CAA36790.1 | X52555   | Nicotiana tabacum       | PR-1 protein (AA 1-184).  |
| CAA35666.1 | X17681   | Nicotiana tabacum       | pathogenesis-related protein 1b (AA 1-168).                                       |
| CAA31010.1 | X12487   | Nicotiana tabacum       | PR1c preprotein.  |
| CAA29023.1 | X05454   | Nicotiana tabacum       | PR-1c protein.  |
| CAA32228.1 | X14065   | Nicotiana tabacum       | PRP 1 precursor (AA -23 to 154).  |
| BAA14220.1 | D90196   | Nicotiana tabacum       | PR1a protein precursor.   |
| CAA31233.1 | X12737   | Nicotiana tabacum       | PR-1a protein (AA 1 - 168).   |
| CAA29392.1 | X05959   | Nicotiana tabacum       | PR-1a precursor (AA -30 to 138).  |
| CAA29660.1 | X06361   | Nicotiana tabacum       | PR1a precursor (AA -30 to -1).  |
| CAA09671.1 | AJ011520 | Lycopersicon esculentum | pathogenesis-related protein PR1a (P4). pr1a (P4).                                |
| AAA03615.1 | M69247   | Lycopersicon esculentum | pathogenesis-related protein P4. P4.  |
| CAA30017.1 | X06930   | Nicotiana tabacum       | PR-1a protein (AA 1 - 168).   |
| CAA31008.1 | X12485   | Nicotiana tabacum       | PR1a preprotein.  |
| CAA52893.1 | X74939   | Hordeum vulgare         | PR-1a pathogenesis related protein (Hv-1a).                                       |
| CAB58263.1 | AJ250136 | Solanum tuberosum       | pathogenesis related protein PR-1. pr1-1.   |
| AAB49685.1 | U89895   | Oryza sativa            | pathogenesis-related protein class 1. PR-1. induced by pathogen attack in plants. |
| CAA27183.1 | X03465   | Nicotiana tabacum       | PR-1b precursor; (aa -30-138).  |
| CAA35665.1 | X17680   | Nicotiana tabacum       | pathogenesis-related protein 1b (AA 1-168).                                       |

|   |          |                         |
|---|----------|-------------------------|
| BAA14221.1  | D90197   | Nicotiana tabacum       |
| PR1b protein precursor.                                     |          |                         |
| CAA48672.1  | X68738   | Lycopersicon esculentum |
| P1(p14) protein. pTE28.1.                                   |          |                         |
| CAA81229.1  | Z26320   | Hordeum vulgare         |
| pathogenesis-related protein. pathogenesis-related protein. |          |                         |
| AAA03616.1  | M69248   | Lycopersicon esculentum |
| pathogenesis-related protein P6. P6.                        |          |                         |
| CAA70042.1  | Y08804   | Lycopersicon esculentum |
| PR protein. PR1b1.  |          |                         |
| AAB05225.1  | U49241   | Nicotiana glutinosa     |
| pathogenesis-related protein-1.                             |          |                         |
| CAA31009.1  | X12486   | Nicotiana tabacum       |
| PR1b preprotein.  |          |                         |
| AAC25629.1  | U82200   | Zea mays                |
| pathogenesis related protein-1. PR-1.                       |          |                         |
| AAF78528.1  | AF195237 | Pyrus pyrifolia         |
| pathogenesis-related protein. PR-1b.                        |          |                         |
| AAD33696.1  | AF136636 | Glycine max             |
| PR1a precursor. PR1a.                                       |          |                         |
| CAA79703.1  | Z21494   | Hordeum vulgare         |
| Pathogenesis-related protein 1.                             |          |                         |
| CAA52894.1  | X74940   | Hordeum vulgare         |
| PR-1b pathogenesis related protein (Hv-8).                  |          |                         |
| CAA81234.1  | Z26333   | Hordeum vulgare         |
| pathogenesis-related protein. pathogenesis-related protein. |          |                         |
| CAA81230.1  | Z26321   | Hordeum vulgare         |
| pathogenesis-related protein. pathogenesis-related protein. |          |                         |
| CAA04881.1  | AJ001627 | Lycopersicon esculentum |
| pathogenesis-related protein. PR1d.                         |          |                         |
| CAA07473.1  | AJ007348 | Triticum aestivum       |
| pathogenesis-related protein 1.1. PR-1.1.                   |          |                         |
| CAA07474.1  | AJ007349 | Triticum aestivum       |
| pathogenesis-related protein 1.2. PR-1.2.                   |          |                         |
| CAA50596.1  | X71592   | Lycopersicon esculentum |
| PR-1a1.   |          |                         |
| CAA70070.1  | Y08844   | Lycopersicon esculentum |
| PR protein. PR1a2.  |          |                         |
| CAA38223.1  | X54325   | Zea mays                |
| pathogenesis-related protein. PRms.                         |          |                         |
| AAF78527.1  | AF195236 | Pyrus pyrifolia         |
| pathogenesis-related proteins. PR-1a.                       |          |                         |

|                |          |                              |   |
|----------------|----------|------------------------------|---|
| CAC03571.1     | AJ278436 | Oryza sativa                 | defence response. PR1a protein. Pr1a.   |
| AAG44566.1     | AF251277 | Oryza sativa subsp. japonica | acidic PR-1 type pathogenesis-related protein PR-1a. PR-1a. induced by pathogen attack. |
| AAC06244.1     | AF053343 | Capsicum annuum              | PR-1 protein precursor. pathogen-induced PR1 protein.                                   |
| CAA56174.1     | X79778   | Medicago truncatula          | PR-1.   |
| SEQ ID NO: 307 |          |                              |   |
| AAF06347.1     | AF195654 | Vitis vinifera               | SCUTL2. thaumatin-like protein.   |
| BAA28872.1     | AB006009 | Pyrus pyrifolia              | thaumatin-like protein precursor. PsTL1.  |
| AAB38064.1     | U32440   | Prunus avium                 | thaumatin-like protein precursor.   |
| BAA95017.1     | AB031870 | Cestrum elegans              | thaumatin-like protein. CETLP.  |
| BAA74546.2     | AB000834 | Nicotiana tabacum            | thaumatin-like protein SE39b.   |
| AAC36740.1     | AF090143 | Malus x domestica            | thaumatin-like protein precursor Mdtl1. MDTL1. pathogenesis-related.                    |
| CAC10270.1     | AJ243427 | Malus x domestica            | thaumatin-like protein. tl. allergen, pathogenesis-related.                             |
| AAB95118.1     | U71244   | Brassica rapa                | pathogenesis-related group 5 protein. BFTP. thaumatin-like protein; PR-5.               |
| CAC09477.1     | AL442113 | Oryza sativa                 | thaumatin-like protein. H0806H05.10.  |
| CAB62167.1     | AJ242828 | Castanea sativa              | antifungal. thaumatin-like protein. tl1.  |
| CAA06927.1     | AJ006233 | Nicotiana tabacum            | putative thaumatin-like protein precursor.  |
| AAF06346.1     | AF195653 | Vitis vinifera               | SCUTL1. thaumatin-like protein.   |
| AAB02259.1     | U57787   | Avena sativa                 | permatin precursor. thaumatin-like protein.   |
| AAD55090.1     | AF178653 | Vitis riparia                | thaumatin. osmotin; pathogenesis-related protein.                                       |
| CAA10492.1     | AJ131731 | Pseudotsuga menziesii        | Thaumatin-like protein. 5A1A.16.  |
| CAA09228.1     | AJ010501 | Cicer arietinum              | thaumatin-like protein PR-5b.   |

|                |          |  |  |
|----------------|----------|--|--|
| BAA95165.1     | AB029918 | Nicotiana tabacum                      | pistil transmitting tissue specific thaumatin (SE39b)-like protein. SE39b.           |
| AAF82264.1     | AF227324 | Vitis vinifera                         | thaumatin-like protein.  |
| CAB85636.1     | AJ237998 | Vitis vinifera                         | putative thaumatin-like protein. TL2.  |
| AAB53368.1     | U77657   | Oryza sativa                           | pathogenesis-related thaumatin-like protein.   |
| AAB61590.1     | AF003007 | Vitis vinifera                         | VVTL1. osmotin-like protein, PR-5 protein; thaumatin-like protein.                   |
| AAB53367.1     | U77656   | Oryza sativa                           | pathogenesis-related thaumatin-like protein.   |
| CAB85637.1     | AJ237999 | Vitis vinifera                         | putative thaumatin-like protein. TL1. alternative name grip 51.                      |
| SEQ ID NO: 308 |          |  |  |
| BAA14143.1     | D90115   | Armoracia rusticana                    | peroxidase isozyme.  |
| BAA14144.1     | D90116   | Armoracia rusticana                    | peroxidase isozyme.  |
| BAA11853.1     | D83225   | Populus nigra                          | peroxidase.  |
| CAA66035.1     | X97349   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.   |
| CAA66036.1     | X97350   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.   |
| BAA11852.1     | D83224   | Populus nigra                          | peroxidase.  |
| CAA66034.1     | X97348   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.   |
| CAA66037.1     | X97351   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.   |
| BAA07241.1     | D38051   | Populus kitakamiensis                  | peroxidase. prxA4a.  |
| AAB47602.1     | L07554   | Linum usitatissimum                    | peroxidase. FLXPER1.   |
| BAA06334.1     | D30652   | Populus kitakamiensis                  | peroxidase.  |
| AAC98519.1     | AF007211 | Glycine max                            | peroxidase precursor. GMIPER1. pathogen-induced.                                     |
| AAB97734.1     | AF014502 | Glycine max                            | seed coat peroxidase precursor. Ep. H2O2 oxidoreductase; class III plant peroxidase. |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| BAA06335.1 | D30653   | Populus kitakamiensis   | peroxidase.   |
| AAD37427.1 | AF149277 | Phaseolus vulgaris      | peroxidase 1 precursor. FBP1. secretory peroxidase.   |
| CAA62226.1 | X90693   | Medicago sativa         | peroxidase1B. prx1B.  |
| CAA62227.1 | X90694   | Medicago sativa         | peroxidase1C. prx1C.  |
| AAB41811.1 | L36157   | Medicago sativa         | peroxidase. pxdC. amino acid feature: conserved domains, aa 123 .. 129, 191 .. 198; amino acid feature: heme-binding domain, aa 68 .. 73. |
| CAA62225.1 | X90692   | Medicago sativa         | peroxidase1A. prx1A.  |
| AAB41810.1 | L36156   | Medicago sativa         | peroxidase. pxdA. amino acid feature: conserved motifs, aa 181 .. 188; amino acid feature: heme-binding domain, aa 60 .. 65.              |
| AAD37430.1 | AF149280 | Phaseolus vulgaris      | peroxidase 5 precursor. FBP5. secretory peroxidase.   |
| CAB94692.1 | AJ242742 | Ipomoea batatas         | Removal of H2O2, oxidation of toxic reductants, defence response toward wounding. peroxidase. pod.  |
| AAA34108.1 | J02979   | Nicotiana tabacum       | lignin-forming peroxidase precursor (EC 1.11.1.7).  |
| CAA40796.1 | X57564   | Armoracia rusticana     | peroxidase. peroxidase precursor.   |
| BAA01877.1 | D11102   | Populus kitakamiensis   | peroxidase. prxA1.  |
| BAA01992.1 | D11396   | Nicotiana tabacum       | 'peroxidase'.   |
| CAB67121.1 | Y19023   | Lycopersicon esculentum | peroxidase. cevi-1.   |
| CAA50597.1 | X71593   | Lycopersicon esculentum | peroxidase. CEVI-1.   |
| AAD43561.1 | AF155124 | Gossypium hirsutum      | bacterial-induced peroxidase precursor. Perx_Goshiko.   |
| BAA92500.1 | AP001383 | Oryza sativa            | ESTs D39300(R3292),AU030751(E60187) correspond to a region of the predicted gene. Similar to peroxidase ATP6a. (X98774).                  |
| AAF63027.1 | AF244924 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.  |
| AAB02554.1 | L37790   | Stylosanthes humilis    | cationic peroxidase.  |

|                |          |                       |  |
|----------------|----------|-----------------------|--|
| AAB06183.1     | M37636   | Arachis hypogaea      | cationic peroxidase. PNC1.   |
| CAA59487.1     | X85230   | Triticum aestivum     | peroxidase. pox4.  |
| BAA94962.1     | AB042103 | Asparagus officinalis | peroxidase. AspPOX1.   |
| AAF63026.1     | AF244923 | Spinacia oleracea     | hydrogen peroxide catabolism. peroxidase prx14 precursor. type III peroxidase.   |
| CAB99487.1     | AJ276227 | Hordeum vulgare       | defence against plant pathogens. peroxidase. prx8.   |
| CAA71492.1     | Y10466   | Spinacia oleracea     | peroxidase. prxr5.   |
| CAB65334.1     | AJ250121 | Picea abies           | peroxidase. SPI2 protein. spi2.  |
| AAA33121.1     | M32742   | Cucumis sativus       | peroxidase (CuPer2).   |
| CAA39486.1     | X56011   | Triticum aestivum     | peroxidase.  |
| BAA92422.1     | AP001366 | Oryza sativa          | ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene. Similar to A.thaliana mRNA for peroxidase ATP18a. (X98804). |
| BAA92497.1     | AP001383 | Oryza sativa          | ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene. Similar to peroxidase ATP18a. (X98804).                     |
| CAA59485.1     | X85228   | Triticum aestivum     | peroxidase. POX2.  |
| CAA76680.1     | Y17192   | Cucurbita pepo        | peroxidase. aprx. type III peroxidase.   |
| SEQ ID NO: 309 |          |                       |  |
| BAA85400.1     | AP000615 | Oryza sativa          | similar to OsMlo-h1. (Z95353).   |
| CAB06083.1     | Z83834   | Hordeum vulgare       | Mlo. Mlo.  |
| CAA74909.1     | Y14573   | Hordeum vulgare       | Mlo protein. Mlo.  |
| AAG46114.1     | AC073166 | Oryza sativa          | putative Mlo (pathogen resistance) protein. OSJNBb0064P21.5.   |
| CAA06487.1     | AJ005341 | Linum usitatissimum   | MLO. homolog.  |
| SEQ ID NO: 310 |          |                       |  |
| AAC63113.1     | AF000307 | Brassica napus        | steroid sulfotransferase 3. BnST3.   |



|                |          |                           |   |
|----------------|----------|---------------------------|---|
| AAC63112.1     | AF000306 | Brassica napus            | steroid sulfotransferase 2. BnST2.  |
| AAC63111.1     | AF000305 | Brassica napus            | steroid sulfotransferase 1. BnST1.  |
| AAA61638.1     | U10275   | Flaveria bidentis         | O-sulfation of position 3 of flavonols. flavonol 3-sulfotransferase.  |
| AAA33342.2     | M84135   | Flaveria chloraefolia     | flavonol 3-sulfotransferase.  |
| AAA87399.1     | U10277   | Flaveria bidentis         | transfers sulfate group into flavonol. sulfotransferase-like flavonol.  |
| AAA33343.1     | M84136   | Flaveria chloraefolia     | O-sulfation of position 4' of flavonol. flavonol 4'-sulfotransferase.   |
| SEQ ID NO: 312 |          |                           |   |
| AAD22970.1     | AF124148 | Glycine max               | trehalase 1 GMTRE1. expressed constitutively in many tissues of soybean at a low level; similar to the Arabidopsis thaliana trehalase precursor encoded by GenBank Accession Number AC002343. |
| AAG13442.1     | AC051634 | Oryza sativa              | putative trehalase. OSJNBb0018B10.19.   |
| CAB50901.1     | AJ238651 | Medicago truncatula       | trehalase. TRE1 protein. tre1.  |
| SEQ ID NO: 313 |          |                           |   |
| BAA19928.1     | AB003491 | Oryza sativa              | tryptophan synthase B. trpB.  |
| AAA33491.1     | M76685   | Zea mays                  | tryptophan synthase beta-subunit. TSB2.   |
| AAB97526.1     | AF042321 | Camptotheca acuminata     | tryptophan synthase beta. TSB.  |
| AAB97087.1     | AF042320 | Camptotheca acuminata     | tryptophan synthase beta subunit.   |
| AAA33490.1     | M76684   | Zea mays                  | tryptophan synthase beta-subunit. TSB1.   |
| AAC25986.1     | AF047024 | Chlamydomonas reinhardtii | tryptophan synthase beta. MAA7.   |
| SEQ ID NO: 314 |          |                           |   |
| AAG42689.1     | AF271384 | Zea mays                  | putative tryptophan synthase alpha. TSAlike.  |
| CAA54131.1     | X76713   | Zea mays                  | tryptophan synthase, alpha subunit. trpA.   |
| AAG42688.1     | AF271383 | Zea mays                  | putative tryptophan synthase alpha. TSAlike.  |
| SEQ ID NO: 315 |          |                           |   |

AAB97526.1 AF042321 *Camptotheca acuminata*  
tryptophan synthase beta. TSB.

AAB97087.1 AF042320 *Camptotheca acuminata*  
tryptophan synthase beta subunit.

BAA19928.1 AB003491 *Oryza sativa*  
tryptophan synthase B. trpB.

AAA33491.1 M76685 *Zea mays*  
tryptophan synthase beta-subunit. TSB2.

AAA33490.1 M76684 *Zea mays*  
tryptophan synthase beta-subunit. TSB1.

AAC25986.1 AF047024 *Chlamydomonas reinhardtii*  
tryptophan synthase beta. MAA7.

## SEQ ID NO: 316

AAA33967.1 M76981 *Glycine max*  
vegetative storage protein. vspA.

BAA23563.1 D50094 *Phaseolus vulgaris*  
pod storage protein.

BAA19152.1 AB000585 *Phaseolus vulgaris*  
pod storage protein. PSP.

AAA34020.1 M20037 *Glycine max*  
vegetative storage protein.

AAA34022.1 M76980 *Glycine max*  
vegetative storage protein. vspB.

AAA34021.1 M20038 *Glycine max*  
vegetative storage protein.

## SEQ ID NO: 321

BAA87043.1 AB035183 *Ipomoea batatas*  
N-hydroxycinnamoyl/benzoyltransferase. hcbt.

CAB06427.1 Z84383 *Dianthus caryophyllus*  
phytoalexin biosynthesis. anthranilate N-hydroxycinnamoyl/benzoyltransferase.

CAB06429.1 Z84385 *Dianthus caryophyllus*  
phytoalexin biosynthesis. anthranilate N-hydroxycinnamoyl/benzoyltransferase.

CAB06430.1 Z84386 *Dianthus caryophyllus*  
phytoalexin biosynthesis. anthranilate N-hydroxycinnamoyl/benzoyltransferase.

CAB11466.1 Z98758 *Dianthus caryophyllus*  
carnation phytoalexin biosynthesis. anthranilate N-hydroxycinnamoyl/benzoyltransferase.

CAB06538.1 Z84571 *Dianthus caryophyllus*  
phytoalexin biosynthesis. anthranilate N-hydroxycinnamoyl/benzoyltransferase.

CAB06428.1 Z84384 *Dianthus caryophyllus*  
phytoalexin biosynthesis. anthranilate N-hydroxycinnamoyl/benzoyltransferase.

## SEQ ID NO: 323

BAB12694.1 AP002746 *Oryza sativa*  
putative zinc finger transcription factor. P0671B11.10. contains ESTs  
AU098331(E31537),C91783(E31537).

## SEQ ID NO: 325

BAB03447.1 AP002817 *Oryza sativa*  
ESTs C96615(C10106),C26336(C12127),D21959(C10106) correspond to a region of the  
predicted gene. Similar to *Arabidopsis thaliana* DNA chromosome 4, BAC clone T6K21;  
NAM (no apical meristem) - like protein (AL021889).

BAA92400.1 AP001366 *Oryza sativa*  
ESTs C96615(C10106),C26336(C12127),D21959(C10106) correspond to a region of the  
predicted gene. Similar to NAM (AL021889).

BAB16335.1 AP002818 *Oryza sativa*  
putative NAM protein. P0436E04.18. contains ESTs E10793(C19698),E10793(C99379).

BAB16328.1 AP002818 *Oryza sativa*  
putative NAM protein. P0436E04.11. contains ESTs R4069(AU032425),R4069(AU082730).

CAA63102.2 X92205 *Petunia x hybrida*  
apical meristem formation. NAM.

CAA63101.1 X92204 *Petunia x hybrida*  
apical meristem formation. NAM.

AAK13151.1 AC078829 *Oryza sativa*  
putative NAM (no apical meristem) protein. OSJNBa0026O12.6.

BAB19365.1 AP002542 *Oryza sativa*  
putative NAM (no apical meristem) protein. P0679C08.4.

BAA84803.1 AP000559 *Oryza sativa*  
Similar to NAM like protein (AC005310).

## SEQ ID NO: 326

CAB51836.1 AJ243961 *Oryza sativa*  
Putative Ser/Thr protein kinase. I1332.7.

BAB18292.1 AP002860 *Oryza sativa*  
putative receptor-like protein kinase. P0409B08.19.

BAB39873.1 AP002882 *Oryza sativa*  
putative protein kinase. P0439B06.8. contains ESTs  
AU056701(S20808),AU056702(S20808).

BAB21240.1 AP002953 *Oryza sativa*  
Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).

AAA33000.1 M76647 *Brassica oleracea*  
receptor protein kinase. SKR6.

CAA74661.1 Y14285 *Brassica oleracea*  
SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular  
kinase domain: 1412-2554.

CAB89179.1 AJ245479 *Brassica napus* subsp. *napus*  
ser /thr kinase. S-locus receptor kinase. srk.

|            |          |                    |  |
|------------|----------|--------------------|--|
| AAA33008.1 | M97667   | Brassica napus     | serine/threonine kinase receptor.  |
| AAA62232.1 | U00443   | Brassica napus     | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| BAA87853.1 | AP000816 | Oryza sativa       | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308). |
| CAA73134.1 | Y12531   | Brassica oleracea  | serine/threonine kinase. BRLK.   |
| BAA92836.1 | AB032473 | Brassica oleracea  | S18 S-locus receptor kinase. SRK18.  |
| AAK21965.1 | AY028699 | Brassica napus     | receptor protein kinase PERK1.   |
| BAA23676.1 | AB000970 | Brassica rapa      | receptor kinase 1. BcRK1.  |
| CAA73133.1 | Y12530   | Brassica oleracea  | serine /threonine kinase. ARLK.  |
| CAA67145.1 | X98520   | Brassica oleracea  | receptor-like kinase. SFR2.  |
| AAK00425.1 | AC069324 | Oryza sativa       | Putative protein kinase. OSJNBa0071K19.11.   |
| AAG03090.1 | AC073405 | Oryza sativa       | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).   |
| CAB41878.1 | Y18259   | Brassica oleracea  | SRK5 protein. SRK5. receptor-like kinase.  |
| AAD21872.1 | AF078082 | Phaseolus vulgaris | receptor-like protein kinase homolog RK20-1.   |
| BAA21132.1 | D88193   | Brassica rapa      | S-receptor kinase. SRK9 (B.c).   |
| BAA06285.1 | D30049   | Brassica rapa      | S-receptor kinase SRK9.  |
| AAB61708.1 | U93048   | Daucus carota      | somatic embryogenesis receptor-like kinase. SERK.  |
| BAB16871.1 | AP002537 | Oryza sativa       | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).                           |
| AAC23542.1 | U20948   | Ipomoea trifida    | receptor protein kinase. IRK1.   |
| CAB41879.1 | Y18260   | Brassica oleracea  | SRK15 protein. SRK15. receptor-like kinase.  |
| AAB93834.1 | U82481   | Zea mays           | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.                                 |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| BAB39409.1     | AP002901 | Oryza sativa            | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| CAA74662.1     | Y14286   | Brassica oleracea       | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.  |
| CAA79355.1     | Z18921   | Brassica oleracea       | S-receptor kinase-like protein.   |
| BAA92837.1     | AB032474 | Brassica oleracea       | S60 S-locus receptor kinase. SRK60.   |
| BAA07577.2     | D38564   | Brassica rapa           | receptor protein kinase SRK12.  |
| BAA07576.1     | D38563   | Brassica rapa           | receptor protein kinase SRK8.   |
| AAG59657.1     | AC084319 | Oryza sativa            | putative protein kinase. OSJNBa0004B24.20.  |
| BAB03429.1     | AP002817 | Oryza sativa            | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587). |
| BAB07999.1     | AP002525 | Oryza sativa            | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| SEQ ID NO: 327 |          |                         |   |
| AAC04717.1     | AF034131 | Gossypium hirsutum      | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-G. similar to MYB A encoded by GenBank Accession Number L04497.                      |
| BAA23340.1     | D88620   | Oryza sativa            | transfactor. OSMYB4. OsmYb4.  |
| AAK19611.1     | AF336278 | Gossypium hirsutum      | BNLGH1233. bnlghi6233. similar to myb.  |
| AAA33482.1     | M37153   | Zea mays                | c1 locus myb homologue; putative.   |
| AAK09326.1     | AF320613 | Zea mays                | activates anthocyanin transcription. anthocyanin regulatory C1. c1. transcription factor.   |
| AAK09327.1     | AF320614 | Zea mays                | activates anthocyanin transcription. anthocyanin regulatory C1. c1. transcription factor.   |
| AAA82943.1     | U39448   | Picea mariana           | MYB-like transcriptional factor MBF1. putative DNA binding region highly similar to the maize C1.   |
| AAK19618.1     | AF336285 | Gossypium hirsutum      | GHMYB38. ghmyb38. similar to myb.   |
| CAA64614.1     | X95296   | Lycopersicon esculentum | transcription factor. THM27. myb-related.   |
| AAK19619.1     | AF336286 | Gossypium hirsutum      | GHMYB9. ghmyb9. similar to myb.   |

---

AAC04720.1 AF034134 *Gossypium hirsutum*  
putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-O.  
similar to MYB A encoded by GenBank Accession Number L04497.

---

AAC04718.1 AF034132 *Gossypium hirsutum*  
putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-J.  
similar to MYB A encoded by GenBank Accession Number L04497.

---

SEQ ID NO: 329

---

CAB93939.1 AJ238739 *Catharanthus roseus*  
putative transcription factor. AP2-domain DNA-binding protein. orca1.

---

AAF76898.1 AF274033 *Atriplex hortensis*  
apetala2 domain-containing protein.

---

AAG43545.1 AF211527 *Nicotiana tabacum*  
Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.

---

CAC12822.1 AJ299252 *Nicotiana tabacum*  
AP2 domain-containing transcription factor. ap2.

---

BAA78738.1 AB023482 *Oryza sativa*  
EST AU055776(S20048) corresponds to a region of the predicted gene.; Similar to  
*Arabidopsis thaliana* AP2 domain containing protein RAP2.10 mRNA, partial  
cds.(AF003103).

---

AAC14323.1 AF058827 *Nicotiana tabacum*  
TS1. Ts1. contains putative AP2 DNA-binding domain; similar to Pti6.

---

BAA97122.1 AB016264 *Nicotiana sylvestris*  
ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-  
related) gene of higher plant. ethylene-responsive element binding factor. nserf2.

---

CAB93940.1 AJ238740 *Catharanthus roseus*  
putative transcription factor. AP2-domain DNA-binding protein. orca2.

---

BAA76734.1 AB024575 *Nicotiana tabacum*  
ethylene responsive element binding factor.

---

CAB96900.1 AJ251250 *Catharanthus roseus*  
transcription factor. AP2-domain DNA-binding protein. orca3.

---

CAB96899.1 AJ251249 *Catharanthus roseus*  
transcription factor. AP2-domain DNA-binding protein. orca3.

---

AAF63205.1 AF245119 *Mesembryanthemum crystallinum*  
AP2-related transcription factor. CDBP. stress induced transcription factor.

---

AAC24587.1 AF071893 *Prunus armeniaca*  
AP2 domain containing protein. AP2DCP.

---

BAA94514.2 AP001800 *Oryza sativa*  
Similar to *Arabidopsis thaliana* chromosome 4, BAC clone F9D16; putative Ap2 domain  
protein (AL035394).

---

BAA97124.1 AB016266 *Nicotiana sylvestris*  
ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-  
related) gene of higher plant. ethylene-responsive element binding factor. nserf4.

---

|                |          |                      |   |
|----------------|----------|----------------------|---|
| BAA07321.1     | D38123   | Nicotiana tabacum    | ERF1. ethylene-responsive transcription factor.   |
| BAB16083.1     | AB036883 | Oryza sativa         | transcriptional repressor. osERF3. osERF3. ERF protein family ERF3 associated repression domain.  |
| BAB03248.1     | AB037183 | Oryza sativa         | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.   |
| BAA97123.1     | AB016265 | Nicotiana sylvestris | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.   |
| AAF23899.1     | AF193803 | Oryza sativa         | transcription factor EREBP1. EREBP/AP2-like transcription factor.   |
| AAK31271.1     | AC079890 | Oryza sativa         | putative transcriptional factor. OSJNBb0089A17.22.  |
| AAC62619.1     | AF057373 | Nicotiana tabacum    | transcription factor. ethylene response element binding protein 1. EREBP1.  |
| AAG43548.1     | AF211530 | Nicotiana tabacum    | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.  |
| BAA90812.1     | AP001168 | Oryza sativa         | Similar to mRNA for DREB1A (AB007787).  |
| AAG43549.1     | AF211531 | Nicotiana tabacum    | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.  |
| BAA99376.1     | AP002526 | Oryza sativa         | ESTs AU093391(E60370),AU091593(C60458), AU093392(E60370) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana BAC F21J9; AP2 domain protein. (AC000103).   |
| AAK01089.1     | AF298231 | Hordeum vulgare      | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.   |
| AAG59618.1     | AF239616 | Hordeum vulgare      | CRT/DRE-binding factor. CBF.  |
| AAG59619.1     | AF243384 | Oryza sativa         | CRT/DRE binding factor. CBF. DREB.  |
| AAK01088.1     | AF298230 | Hordeum vulgare      | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.  |
| AAC49567.1     | U41466   | Zea mays             | Glossy15. Glossy15. AP2 DNA-binding domain protein; similar to the Arabidopsis floral homeotic gene APETALA2; maize AP2-domain regulator of leaf epidermal traits; homeotic regulator of leaf epidermal cell identity; allele: W64A; putative transcription factor. |
| SEQ ID NO: 330 |          |                      |   |
| AAC67571.1     | AF060884 | Brassica napus       | desiccation protein. Cdes. induced by dehydration.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAA61564.1     | U08108   | Glycine max             | putative desiccation protectant protein, homolog of Lea14, GenBank Accession Number M88321.   |
| AAA18543.1     | M88322   | Gossypium hirsutum      | probable desiccation protectant. Group 4 late embryogenesis-abundant protein. Lea14-A. This CDS is colinear and 66% identical with that in cDNA clone pcC27-45 from <i>Craterostigma plantagineum</i> desiccated leaves (Piatkowski et al., 1990, Plant Physiol. 94: 1682-1688).; putative. |
| AAA18542.1     | M88321   | Gossypium hirsutum      | probable desiccation protectant. Group 4 late embryogenesis-abundant protein. Lea14-A. putative.  |
| AAD25354.1     | AF115314 | Glycine max             | possible desiccation protectant. seed maturation protein PM22. PM22. similar to desiccation protectant protein encoded by GenBank Accession Number U08108; late embryogenesis abundant protein; LEA protein.  |
| AAF64451.1     | AF239929 | Euphorbia esula         | late-embryogenesis abundant protein. similar to desiccation protectant protein and late-embryogenesis abundant protein LEA14.   |
| AAB96796.1     | U77719   | Lycopersicon esculentum | ethylene-responsive late embryogenesis-like protein. ER5. LEA-like protein; drought-inducible; ABA-inducible; putative desiccation protectant protein; similar to cotton Lea14A product encoded by GenBank Accession Number M88321.   |
| SEQ ID NO: 345 |          |                         |   |
| AAF61647.1     | AF190634 | Nicotiana tabacum       | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAA36423.1     | AB013598 | Verbena x hybrida       | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| BAA89009.1     | AB027455 | Petunia x hybrida       | anthocyanin 5-O-glucosyltransferase. PH1.   |
| BAA36421.1     | AB013596 | Perilla frutescens      | UDP-glucose:anthocyanin 5-O-glucosyltransferase. PF3R4.   |
| BAA93039.1     | AB033758 | Citrus unshiu           | limonoid UDP-glucosyltransferase. LGTase.   |
| BAA36422.1     | AB013597 | Perilla frutescens      | UDP-glucose:anthocyanin 5-O-glucosyltransferase homologue. PF3R6.   |
| AAF98390.1     | AF287143 | Brassica napus          | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT.   |
| BAB07962.1     | AP002524 | Oryza sativa            | putative anthocyanin 5-O-glucosyltransferase. P0406H10.16. contains ESTs AU067881(C10481),AU067882(C10481).   |



|            |          |   |   |
|------------|----------|---|---|
| AAF17077.1 | AF199453 | <i>Sorghum bicolor</i>                        | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |
| AAK16172.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.14.   |
| AAB36653.1 | U32644   | <i>Nicotiana tabacum</i>                      | immediate-early salicylate-induced glucosyltransferase. IS5a.   |
| AAK28303.1 | AF346431 | <i>Nicotiana tabacum</i>                      | phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.  |
| BAA83484.1 | AB031274 | <i>Scutellaria baicalensis</i>                | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.   |
| AAK16181.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.16.   |
| AAD21086.1 | AF127218 | <i>Forsythia x intermedia</i>                 | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.   |
| AAB36652.1 | U32643   | <i>Nicotiana tabacum</i>                      | immediate-early salicylate-induced glucosyltransferase. IS10a.  |
| AAK28304.1 | AF346432 | <i>Nicotiana tabacum</i>                      | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.  |
| CAA59450.1 | X85138   | <i>Lycopersicon esculentum</i>                | twil. homologous to glucosyltransferases.   |
| AAK16178.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.5.  |
| AAK16175.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.15.   |
| AAK16180.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.21.   |
| CAB56231.1 | Y18871   | <i>Dorotheanthus bellidiformis</i>            | betanidin-5-O-glucosyltransferase.  |
| BAA12737.1 | D85186   | <i>Gentiana triflora</i>                      | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| AAG25643.1 | AF303396 | <i>Phaseolus vulgaris</i>                     | UDP-glucosyltransferase HRA25, putative; defense associated.  |
| BAB41017.1 | AB047090 | <i>Vitis labrusca</i> x <i>Vitis vinifera</i> | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT1. The gene was derived from one of the parents <i>V. labruscana</i> cv. Ishiharawase. |
| AAB81683.1 | AF000372 | <i>Vitis vinifera</i>                         | UDP glucose:flavonoid 3-o-glucosyltransferase.  |
| BAB41020.1 | AB047093 | <i>Vitis vinifera</i>                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT2.   |
| BAB41022.1 | AB047095 | <i>Vitis vinifera</i>                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT2.   |

|                |          |  |   |
|----------------|----------|--|---|
| BAB41019.1     | AB047092 | Vitis vinifera                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT1.   |
| AAB81682.1     | AF000371 | Vitis vinifera                         | UDP glucose:flavonoid 3-o-glucosyltransferase.  |
| BAB41025.1     | AB047098 | Vitis vinifera                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.   |
| BAB41023.1     | AB047096 | Vitis vinifera                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.   |
| BAB41021.1     | AB047094 | Vitis vinifera                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT1.   |
| BAB41026.1     | AB047099 | Vitis vinifera                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT2.   |
| BAB41024.1     | AB047097 | Vitis vinifera                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT2.   |
| BAB41018.1     | AB047091 | Vitis labrusca x Vitis vinifera        | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT2. The gene originated in one of the parents V. vinifera cv. Centennial. |
| BAB17182.1     | AP002843 | Oryza sativa                           | putative UTP-glucose glucosyltransferase. P0407B12.19.  |
| BAA19659.1     | AB002818 | Perilla frutescens                     | flavonoid 3-O-glucosyltransferase. UDP glucose.   |
| BAA89008.1     | AB027454 | Petunia x hybrida                      | anthocyanidin 3-O-glucosyltransferase. PGT8.  |
| BAB17176.1     | AP002843 | Oryza sativa                           | putative UTP-glucose glucosyltransferase. P0407B12.13.  |
| AAB86473.1     | AF028237 | Ipomoea purpurea                       | UDP glucose: flavonoid 3-O-glucosyltransferase. UF3GT-FL1.  |
| SEQ ID NO: 350 |          |  |   |
| CAA44216.1     | X62343   | Nicotiana tabacum                      | cinnamyl-alcohol dehydrogenase. CAD14.  |
| BAA03099.1     | D13991   | Aralia cordata                         | cinnamyl alcohol dehydrogenase. cadacl.   |
| CAA79622.1     | Z19568   | Populus deltoides                      | lignin biosynthesis. cinnamyl alcohol dehydrogenase.  |
| CAC07423.1     | AJ295837 | Populus balsamifera subsp. trichocarpa | lignin monomer biosynthesis. cinnamyl alcohol dehydrogenase. cad.   |
| AAF43140.1     | AF217957 | Populus tremuloides                    | cinnamyl alcohol dehydrogenase. CAD.  |
| CAA44217.1     | X62344   | Nicotiana tabacum                      | cinnamyl-alcohol dehydrogenase. CAD19.  |
| CAA79625.1     | Z19573   | Medicago sativa                        | lignin biosynthesis. cinnamyl alcohol dehydrogenase.  |

|            |          |   |
|------------|----------|---|
| AAC35845.1 | AF083332 | <i>Medicago sativa</i><br>cinnamyl-alcohol dehydrogenase. MsaCad2.  |
| AAC07987.1 | AF038561 | <i>Eucalyptus globulus</i><br>catalyses the reduction of cinnamaldehydes to the corresponding cinnamyl alcohols as the last step in the production of lignin monomers. cinnamyl alcohol dehydrogenase. CAD. |
| CAA46585.1 | X65631   | <i>Eucalyptus gunnii</i><br>cinnamyl-alcohol dehydrogenase. cad.  |
| AAG15553.1 | AF294793 | <i>Eucalyptus saligna</i><br>cinnamyl alcohol dehydrogenase. cad. CAD.  |
| CAA53211.1 | X75480   | <i>Eucalyptus gunnii</i><br>cinnamyl-alcohol dehydrogenase. CAD.  |
| AAB70908.1 | AF010290 | <i>Lolium perenne</i><br>cinnamyl alcohol dehydrogenase. CAD.   |
| CAA74070.1 | Y13733   | <i>Zea mays</i><br>cinnamyl alcohol dehydrogenase. cad.   |
| CAA06687.1 | AJ005702 | <i>Zea mays</i><br>cinnamyl alcohol dehydrogenase. cad.   |
| CAA13177.1 | AJ231135 | <i>Saccharum officinarum</i><br>lignin biosynthesis. cinnamyl alcohol dehydrogenase. cad.   |
| BAA19487.1 | D86590   | <i>Zinnia elegans</i><br>cinnamyl alcohol dehydrogenase. ZCAD1.   |
| CAA51226.1 | X72675   | <i>Picea abies</i><br>cinnamyl-alcohol dehydrogenase.   |
| CAA05097.1 | AJ001926 | <i>Picea abies</i><br>cinnamyl alcohol dehydrogenase. cad8.   |
| CAA05096.1 | AJ001925 | <i>Picea abies</i><br>cinnamyl alcohol dehydrogenase. cad7.   |
| CAA05095.1 | AJ001924 | <i>Picea abies</i><br>cinnamyl alcohol dehydrogenase. cad2.   |
| AAB38774.1 | U62394   | <i>Pinus radiata</i><br>cinnamyl alcohol dehydrogenase. CAD.  |
| AAC31166.1 | AF060491 | <i>Pinus radiata</i><br>cinnamyl alcohol dehydrogenase. CAD.  |
| CAA86073.1 | Z37992   | <i>Pinus taeda</i><br>cinnamyl alcohol dehydrogenase.   |
| CAA86072.1 | Z37991   | <i>Pinus taeda</i><br>cinnamyl alcohol dehydrogenase.   |
| BAA04046.1 | D16624   | <i>Eucalyptus botryoides</i><br>cinnamyl alcohol dehydrogenase. Cad1:Eb:1.  |
| AAD10327.1 | U63534   | <i>Fragaria x ananassa</i><br>catalyzes the reduction of cinnamylaldehydes leading to monolignols. cinnamyl alcohol dehydrogenase. CAD. involved with lignin biosynthesis.                                  |

|                |          |   |
|----------------|----------|---|
| AAK28509.1     | AF320110 | <i>Fragaria x ananassa</i><br>cinnamyl alcohol dehydrogenase.   |
| AAB38503.1     | U79770   | <i>Mesembryanthemum crystallinum</i><br>cinnamyl-alcohol dehydrogenase Eli3.  |
| AAC35846.1     | AF083333 | <i>Medicago sativa</i><br>cinnamyl-alcohol dehydrogenase. MsaCad1.  |
| AAA74882.1     | L36823   | <i>Stylosanthes humilis</i><br>cinnamyl-alcohol dehydrogenase. CAD1.  |
| AAF23409.1     | AF207552 | <i>Brassica napus</i><br>cinnamyl alcohol dehydrogenase. CADA-1.  |
| AAC15467.1     | U24561   | <i>Apium graveolens</i><br>converts mannitol to mannose. mannitol dehydrogenase. Mtd. 1-oxidoreductase; induced with sodium salicylate; similar to the plant defense gene ELI3 in <i>Arabidopsis thaliana</i> , PIR Accession Number S28044; EC number unassigned; MTD. |
| AAF23411.1     | AF207554 | <i>Brassica oleracea</i><br>cinnamyl alcohol dehydrogenase. CADA.   |
| AAF23412.1     | AF207555 | <i>Brassica rapa</i><br>cinnamyl alcohol dehydrogenase. CADA.   |
| AAC61854.1     | AF067082 | <i>Apium graveolens</i><br>oxidizes mannitol to mannose. mannitol dehydrogenase. Mtd. mannitol 1-oxidoreductase.  |
| AAF23410.1     | AF207553 | <i>Brassica napus</i><br>cinnamyl alcohol dehydrogenase. CADA-2.  |
| AAA74883.1     | L36456   | <i>Stylosanthes humilis</i><br>cinnamyl-alcohol dehydrogenase. CAD3.  |
| AAD18000.1     | AF109157 | <i>Eucalyptus globulus</i><br>cinnamyl alcohol dehydrogenase. CAD.  |
| AAF72100.1     | AF146691 | <i>Lycopersicon esculentum</i><br>ELI3. Eli3. similar] to cinnamyl alcohol dehydrogenase.   |
| CAA63410.1     | X92754   | <i>Hordeum vulgare</i><br>cinnamyl alcohol dehydrogenase. CAD.  |
| AAF23416.1     | AF207559 | <i>Brassica rapa</i><br>cinnamyl alcohol dehydrogenase. CADb.   |
| SEQ ID NO: 351 |          |   |
| AAB71227.1     | AF004809 | <i>Glycine max</i><br>Ca <sup>2+</sup> -binding EF hand protein. GmPM13. encodes EF-hand motifs.  |
| AAF13743.1     | AF109921 | <i>Sesamum indicum</i><br>caleosin. 27 kDa calcium-binding protein.   |
| CAA61981.1     | X89891   | <i>Oryza sativa</i><br>EFA27 for EF hand, abscisic acid, 27kD. efa27.   |
| CAB71337.1     | AJ250283 | <i>Hordeum vulgare</i><br>putative calcium binding EF-hand protein. bci-4.  |
| CAB42585.1     | AJ238627 | <i>Chlorella protothecoides</i><br>putative Ca <sup>++</sup> binding protein. dee112.   |

## SEQ ID NO: 353

AAF60316.1 AF236108 Glycine max  
putative purple acid phosphatase precursor. PAP. type 5 acid phosphatase.

AAF60315.1 AF236107 Ipomoea batatas  
putative purple acid phosphatase precursor. PAP. type 5 acid phosphatase.

AAF60317.1 AF236109 Phaseolus vulgaris  
putative purple acid phosphatase precursor. PAP. type 5 acid phosphatase.

## SEQ ID NO: 359

CAA55039.1 X78203 Hyoscyamus muticus  
glutathione transferase.

AAB65163.1 AF002692 Solanum commersonii  
glutathione S-transferase, class-phi. GST1. low temperature induced.

CAA96431.1 Z71749 Nicotiana plumbaginifolia  
glutathione S-transferase.

BAA01394.1 D10524 Nicotiana tabacum  
glutathione S-transferase. parB.

AAA33930.1 M84968 Silene vulgaris  
glutathione-S-transferase.

AAA33931.1 M84969 Silene vulgaris  
glutathione-S-transferase.

AAF65767.1 AF242309 Euphorbia esula  
glutathione S-transferase. putative auxin-binding GST.

AAF61392.1 AF133894 Persea americana  
glutathione S-transferase. GTH.

CAB38119.1 AJ010296 Zea mays  
Glutathione transferase III(b). gst3b.

CAB38118.1 AJ010295 Zea mays  
Glutathione transferase III(a). gst3a.

BAB39935.1 AP002914 Oryza sativa  
putative glutathione S-transferase. P0493G01.17.

AAG34811.1 AF243376 Glycine max  
glutathione S-transferase GST 21.

BAB39941.1 AP002914 Oryza sativa  
putative glutathione S-transferase. P0493G01.23.

CAA09190.1 AJ010451 Alopecurus myosuroides  
glutathione transferase. GST2a.

CAA09192.1 AJ010453 Alopecurus myosuroides  
glutathione transferase. GST2c.

CAA09193.1 AJ010454 Alopecurus myosuroides  
glutathione transferase. GST2d.

AAG34814.1 AF243379 Glycine max  
glutathione S-transferase GST 24.

|            |          |   |
|------------|----------|---|
| CAA09191.1 | AJ010452 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST2b.  |
| BAB39939.1 | AP002914 | <i>Oryza sativa</i><br>putative glutathione S-transferase. P0493G01.21.   |
| AAG32476.1 | AF309383 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTF4.   |
| AAG34812.1 | AF243377 | <i>Glycine max</i><br>glutathione S-transferase GST 22.   |
| BAB39929.1 | AP002914 | <i>Oryza sativa</i><br>putative glutathione transferase. P0493G01.7.  |
| CAA39487.1 | X56012   | <i>Triticum aestivum</i><br>glutathione transferase. gstA1.   |
| AAD56395.1 | AF184059 | <i>Triticum aestivum</i><br>glutathione S-transferase. GST1.  |
| BAB39940.1 | AP002914 | <i>Oryza sativa</i><br>putative glutathione S-transferase. P0493G01.22.   |
| CAA68993.1 | Y07721   | <i>Petunia x hybrida</i><br>conjugates glutathione to anthocyanin to facilitate transport to the vacuole. glutathione S-transferase. an9 locus.         |
| AAA33469.1 | M16902   | <i>Zea mays</i><br>glutathione S-transferase I.   |
| AAG32477.1 | AF309384 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTF3.   |
| AAA33470.1 | M16901   | <i>Zea mays</i><br>glutathione S-transferase I.   |
| AAA20585.1 | U12679   | <i>Zea mays</i><br>glutathione S-transferase IV. GSTIV.   |
| CAA56047.1 | X79515   | <i>Zea mays</i><br>glutathione transferase. GST27.  |
| CAA39480.1 | X56004   | <i>Triticum aestivum</i><br>glutathione transferase. gstA2.   |
| AAC64007.1 | AF062403 | <i>Oryza sativa</i><br>glutathione S-transferase II.  |
| BAB39927.1 | AP002914 | <i>Oryza sativa</i><br>putative glutathione S-transferase. P0493G01.1. contains ESTs<br>AU031696(R0596), C97559(C60386), C28218(C60386), D28287(R0596). |
| AAG32475.1 | AF309382 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTF5.   |
| AAG34823.1 | AF244680 | <i>Zea mays</i><br>glutathione S-transferase GST 15.  |
| AAG34817.1 | AF244674 | <i>Zea mays</i><br>glutathione S-transferase GST 9.   |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| CAA05354.1     | AJ002380 | Oryza sativa            | glutathione S-transferase. Rgst I.  |
| AAG34820.1     | AF244677 | Zea mays                | glutathione S-transferase GST 11.   |
| AAG34821.1     | AF244678 | Zea mays                | glutathione S-transferase GST 13.   |
| CAB66333.1     | AJ279691 | Betula pendula          | glutathione-S-transferase. gst.   |
| AAG34818.1     | AF244675 | Zea mays                | glutathione S-transferase GST 10.   |
| AAG34816.1     | AF244673 | Zea mays                | glutathione S-transferase GST 8.  |
| AAG34822.1     | AF244679 | Zea mays                | glutathione S-transferase GST 14.   |
| CAA05355.1     | AJ002381 | Oryza sativa            | glutathione S-transferase. Rgst II.   |
| SEQ ID NO: 360 |          |                         |   |
| AAA33710.1     | L16977   | Petunia x hybrida       | glutamate decarboxylase. gad.   |
| AAA33709.1     | L16797   | Petunia x hybrida       | glutamate decarboxylase. gad.   |
| AAC24195.1     | AF020425 | Nicotiana tabacum       | calmodulin binding protein. glutamate decarboxylase isozyme 1. NtGAD1. calcium-calmodulin-dependent enzyme.                               |
| AAB40608.1     | U54774   | Nicotiana tabacum       | glutamate decarboxylase. NtGAD1. calmodulin regulated enzyme; calmodulin-binding protein.   |
| AAK18620.1     | AF352732 | Nicotiana tabacum       | converts glutamate to gamma-aminobutyric acid. glutamate decarboxylase isozyme 3. GAD; GAD3; NtGAD3; calcium/calmodulin-dependent enzyme. |
| AAC39483.1     | AF020424 | Nicotiana tabacum       | glutamate decarboxylase isozyme 2. NtGAD2. calcium-calmodulin-dependent enzyme.   |
| BAB32870.1     | AB056062 | Oryza sativa            | glutamate decarboxylase. GAD.   |
| BAB32868.1     | AB056060 | Oryza sativa            | glutamate decarboxylase. GAD.   |
| BAB32869.1     | AB056061 | Oryza sativa            | glutamate decarboxylase. GAD.   |
| BAB32871.1     | AB056063 | Oryza sativa            | glutamate decarboxylase. GAD.   |
| CAA56812.1     | X80840   | Lycopersicon esculentum | homology to pyrooxidal-5'-phosphate-dependant glutamate decarboxylases; putative start codon.   |

|   |          |                         |
|---|----------|-------------------------|
| CAA50719.1  | X71900   | Lycopersicon esculentum |
| histidine decarboxylase. hdc. pyridoxal 5'-phosphate dependant.   |          |                         |
| SEQ ID NO: 362  |          |                         |
| AAG13467.1  | AC026758 | Oryza sativa            |
| putative proline oxidase. OSJNBa0015J15.31.   |          |                         |
| AAD48490.1  | AF171226 | Brassica napus          |
| proline dehydrogenase. pdh.   |          |                         |
| SEQ ID NO: 363  |          |                         |
| AAA33967.1  | M76981   | Glycine max             |
| vegetative storage protein. vspA.   |          |                         |
| AAA34022.1  | M76980   | Glycine max             |
| vegetative storage protein. vspB.   |          |                         |
| AAA34021.1  | M20038   | Glycine max             |
| vegetative storage protein.   |          |                         |
| BAA23563.1  | D50094   | Phaseolus vulgaris      |
| pod storage protein.  |          |                         |
| BAA19152.1  | AB000585 | Phaseolus vulgaris      |
| pod storage protein. PSP.   |          |                         |
| AAA34020.1  | M20037   | Glycine max             |
| vegetative storage protein.   |          |                         |
| SEQ ID NO: 364  |          |                         |
| AAB86939.1  | AF030387 | Oryza sativa            |
| NOI protein.  |          |                         |
| AAC03022.1  | AF045033 | Zea mays                |
| nitrate-induced NOI protein.  |          |                         |
| AAB86937.1  | AF030385 | Zea mays                |
| nitrate-induced NOI protein.  |          |                         |
| SEQ ID NO: 366  |          |                         |
| AAF75824.1  | AF101788 | Pinus taeda             |
| phytoecyanin homolog.   |          |                         |
| AAC32448.1  | U76296   | Spinacia oleracea       |
| plantacyanin. member of a strictly plant-specific family of mononuclear blue copper proteins known as phytoecyanins, which are further classified into three distinct subfamilies: Uclacyanins, Stellacyanins, and Plantacyanins; member of the subfamily of Plantacyanins; non-glycosylated. |          |                         |



---

AAC32421.1 U65511 *Cucumis sativus*  
 putative oxygen activation and/or lignin formation. stellacyanin. member of a strictly plant-specific family of mononuclear blue copper proteins known as phytocyanins, which are further classified into three distinct subfamilies: Uclacyanins, Stellacyanins, and Plantacyanins; member of the subfamily of Stellacyanins; similar to *Rhus vernicifera* stellacyanin: SwissProt Accession Number P00302; similar to umecyanin: SwissProt Accession Number P42849; similar to mavicyanin; SwissProt Accession Number P80728; similar to BCB encoded by GenBank Accession Number Z15058; similar to F18A8.9, encoded by GenBank Accession Number AC003105; similar to F7F1.27 encoded by GenBank Accession Number AC004669; similar to F9D12.16 encoded by GenBank Accession Number AF077407; glycoprotein.

---

AAF66243.1 AF243181 *Lycopersicon esculentum*  
 plantacyanin. naturally occurring cupredoxin with a Val residue in the position of the axial ligand Met; member of the plantacyanin subfamily of a strictly plant-specific family of mononuclear blue copper proteins known as phytocyanins.

---

BAA90481.1 AB035146 *Ipomoea nil*  
 phytocyanin-related protein.

---

SEQ ID NO: 367

---

AAD11617.1 AF050495 *Lycopersicon esculentum*  
 Ca<sup>2+</sup>-ATPase. LCA1A; alternative transcript.

---

AAA34138.1 M96324 *Lycopersicon esculentum*  
 The calcium ATPase is a calcium ion pump. Ca<sup>2+</sup>-ATPase. LCA1.

---

AAD11618.1 AF050496 *Lycopersicon esculentum*  
 Ca<sup>2+</sup>-ATPase. LCA1B; alternative transcript.

---

CAA63790.1 X93592 *Dunaliella bioculata*  
 P-type ATPase. cal. calcium pumping; CA1.

---

AAF73985.1 AF096871 *Zea mays*  
 calcium pump. calcium ATPase. cap1.

---

AAB58910.1 U82966 *Oryza sativa*  
 Ca<sup>2+</sup>-ATPase.

---

AAG28435.1 AF195028 *Glycine max*  
 plasma membrane Ca<sup>2+</sup>-ATPase. SCA1.

---

AAG28436.1 AF195029 *Glycine max*  
 plasma membrane Ca<sup>2+</sup>-ATPase. SCA2.

---

AAB49042.1 U54690 *Dunaliella acidophila*  
 plasma membrane proton ATPase. dha1. DaDHA1; proton pump.

---

AAB35314.2 S79323 *Vicia faba*  
 plasma membrane H<sup>(+)</sup>-ATPase precursor. plasma membrane H<sup>(+)</sup>-ATPase. This sequence comes from Fig. 1; conceptual translation presented here differs from translation in publication.

---

CAB85495.1 AJ132892 *Medicago truncatula*  
 proton pump. H<sup>+</sup>-ATPase. ha1.

---

CAB85494.1 AJ132891 *Medicago truncatula*  
 proton pump. H<sup>+</sup>-ATPase. ha1.

---

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAB17186.1     | U72148   | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase. LHA4. plasma membrane proton pumping ATPase.  |
| CAC29436.1     | AJ310524 | Vicia faba                    | P-type H <sup>+</sup> -ATPase. ha5. predominantly expressed in guard cells and flowers.   |
| AAB41898.1     | U84891   | Mesembryanthemum crystallinum | plasma membrane proton pump. H <sup>+</sup> -transporting ATPase. PMA.  |
| CAC29435.1     | AJ310523 | Vicia faba                    | P-type H <sup>+</sup> -ATPase. vha4. predominantly expressed in flowers.  |
| AAF98344.1     | AF275745 | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase. LHA2. P-type ion pump.  |
| SEQ ID NO: 369 |          |                               |   |
| BAB17726.1     | AB050900 | Raphanus sativus              | asparagine synthetase. Asn1.  |
| CAA59138.1     | X84448   | Brassica oleracea             | asparagine synthase (glutamine-hydrolysing).  |
| AAC16325.1     | AF061740 | Elaeagnus umbellata           | asparagine synthetase. AS.  |
| CAA08913.1     | AJ009952 | Phaseolus vulgaris            | asparagine synthesis. asparagine synthetase type II. as2.   |
| AAF02775.1     | AF190728 | Helianthus annuus             | asparagine synthetase. HAS1.  |
| AAC49613.1     | U77678   | Glycine max                   | catalyzes the ATP-dependent transfer of the amide group of glutamine to aspartate producing asparagine and glutamate. asparagine synthetase 2. AS2. |
| AAB81011.1     | U89923   | Medicago sativa               | asparagine synthetase.  |
| AAC09952.1     | U55874   | Glycine max                   | asparagine synthetase.  |
| AAB48058.1     | L40327   | Medicago sativa               | asparagine synthetase.  |
| CAA61589.1     | X89409   | Lotus japonicus               | asparagine synthase (glutamine-hydrolysing). AS.  |
| CAA67889.1     | X99552   | Asparagus officinalis         | asparagine synthetase.  |
| AAF74755.1     | AF263432 | Helianthus annuus             | asparagine synthetase. HAS1.1.  |
| AAD05035.1     | AF014057 | Triphysaria versicolor        | asparagine synthetase. AS. glutamine-hydrolyzing.   |
| AAD05034.1     | AF014056 | Triphysaria versicolor        | asparagine synthetase. AS. glutamine-hydrolyzing.   |
| AAD05033.1     | AF014055 | Triphysaria versicolor        | asparagine synthetase. AS. glutamine-hydrolyzing.   |

|            |          |                        |  |
|------------|----------|------------------------|--|
| CAA96526.1 | Z72354   | Vicia faba             | synthesis of asparagine from aspartate and glutamine. asparagine synthetase. VfAS1.  |
| CAA48141.1 | X67958   | Asparagus officinalis  | asparagine synthase (glutamine-hydrolysing).   |
| CAA61590.1 | X89410   | Lotus japonicus        | asparagine synthase (glutamine-hydrolysing). AS.   |
| CAA36429.1 | X52179   | Pisum sativum          | asparagine synthase (glutamine-hydrolysing).   |
| BAA96252.1 | AB035248 | Astragalus sinicus     | asparagine synthetase. AsAS2.  |
| CAA36430.1 | X52180   | Pisum sativum          | asparagine synthase (glutamine-hydrolysing).   |
| BAA96251.1 | AB035247 | Astragalus sinicus     | asparagine synthetase. AsAS1.  |
| CAB57292.1 | AJ133522 | Phaseolus vulgaris     | asparagine synthetase (type-I). as1.   |
| AAC49614.1 | U77679   | Glycine max            | catalyzes the ATP-dependent transfer of the amide group of glutamine to aspartate producing asparagine and glutamate. asparagine synthetase 1. AS1.      |
| BAA18951.1 | D83378   | Oryza sativa           | asparagine synthetase.   |
| AAB03991.1 | U55873   | Oryza sativa           | asparagine synthetase.   |
| AAF02776.1 | AF190729 | Helianthus annuus      | asparagine synthetase. HAS2.   |
| AAB71532.1 | AF005724 | Sandersonia aurantiaca | role in flower senescence. asparagine synthetase. SAND1.   |
| CAA58052.1 | X82849   | Zea mays               | asparagine synthetase. AS.   |
| AAB91481.1 | AF037363 | Helianthus annuus      | asparagine synthetase.   |
| CAA73762.1 | Y13321   | Pisum sativum          | asparagine synthetase 1. AS1.  |
| CAA73763.1 | Y13322   | Pisum sativum          | asparagine synthetase 2. AS2.  |
| BAA96452.1 | AB021793 | Pyrus pyrifolia        | asparagine synthetase. PPFRU32.  |
| AAA73943.1 | L23833   | Glycine max            | production of phosphoribosylamine using glutamine and phosphoribosylpyrophosphate as substrates. glutamine phosphoribosylpyrophosphate amidotransferase. |

SEQ ID NO: 370

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAG21985.1     | AF271636 | Zea mays                | lysine ketoglutarate reductase/saccharopine dehydrogenase. LKRSDH. bifunctional enzyme; LKR/SDH; lysine 2-oxoglutarate reductase/saccharopine dehydrogenase. |
| AAC18622.2     | AF003551 | Zea mays                | lysine-ketoglutarate reductase/saccharopine dehydrogenase bifunctional enzyme.   |
| AAG28387.1     | AF191667 | Brassica oleracea       | lysine-ketoglutarate reductase/saccharopine dehydrogenase.   |
| AAG28386.1     | AF191666 | Brassica napus          | lysine-ketoglutarate reductase/saccharopine dehydrogenase.   |
| AAB97685.1     | AF042184 | Brassica napus          | lysine-ketoglutarate reductase/saccharopine dehydrogenase.   |
| AAG14462.1     | AF293461 | Brassica napus          | lysine-ketoglutarate reductase. LKR.   |
| SEQ ID NO: 371 |          |                         |  |
| CAB62537.1     | AJ012583 | Hevea brasiliensis      | pseudo-hevein.   |
| AAA33357.1     | M36986   | Hevea brasiliensis      | hevein (HEV1) precursor.   |
| CAA05978.1     | AJ003196 | Hevea brasiliensis      | N-acetyl-D-glucosamine/N-acetyl-D-neuraminic acid binding lectin. prohevein.   |
| AAF61435.1     | AF137352 | Pisum sativum           | pre-hevein-like protein. PHLP. stress-induced; the coding region is putative in the 5' end.  |
| SEQ ID NO: 372 |          |                         |  |
| AAG28503.1     | AF196966 | Citrus sinensis         | hexokinase.  |
| AAF18584.1     | AF118132 | Spinacia oleracea       | chloroplast outer envelope hexokinase 1. Hxk1.   |
| AAF18585.1     | AF118133 | Nicotiana tabacum       | chloroplast outer envelope hexokinase 1. Hxk1.   |
| AAF14186.1     | AF106068 | Solanum tuberosum       | hexokinase 2.  |
| AAG35735.1     | AF208543 | Lycopersicon esculentum | hexokinase. Hxk2.  |
| CAA63966.1     | X94302   | Solanum tuberosum       | hexokinase. hxk.   |
| BAA99425.1     | AP002743 | Oryza sativa            | putative chloroplast outer envelope hexokinase 1. P0710E05.10.   |
| SEQ ID NO: 374 |          |                         |  |
| AAC83688.2     | AF083343 | Nicotiana tabacum       | 101 kDa heat shock protein. HSP101.  |
| AAF01280.1     | AF174433 | Triticum aestivum       | heat shock protein 101. HSP101. ClpB family member.  |

|                |          |                                  |   |
|----------------|----------|----------------------------------|---|
| AAD33606.1     | AF133840 | <i>Zea mays</i>                  | heat shock protein HSP101. HSP101. 101 kDa protein.   |
| AAD25223.1     | AF077337 | <i>Zea mays</i>                  | heat shock protein 101. HSP101. ClpB/Hsp100 protein homolog; 101 kDa heat shock protein.  |
| AAF91178.1     | AF203700 | <i>Phaseolus lunatus</i>         | ClpB. clpB. heat shock protein HSP100.  |
| AAD22629.1     | AF097363 | <i>Triticum aestivum</i>         | heat shock protein 101. Hsp101a.  |
| AAC83689.2     | AF083344 | <i>Triticum aestivum</i>         | 101 kDa heat shock protein. HSP101.   |
| AAD26530.1     | AF083327 | <i>Zea mays</i>                  | 101 kDa heat shock protein. HSP101. similar to HSP100/ClpB; HSP104.   |
| SEQ ID NO: 375 |          |                                  |   |
| CAA04611.1     | AJ001208 | <i>Brassica juncea</i>           | APS reductase. apsr8. putative mitochondrial or plastidic transit peptide.  |
| CAA04610.1     | AJ001207 | <i>Brassica juncea</i>           | APS reductase. apsr2. putative mitochondrial or plastidic transit peptide.  |
| AAB05871.2     | U63784   | <i>Catharanthus roseus</i>       | reduction of adenylyl sulfate (APS). PAPS-reductase-like protein. par2neu.  |
| CAB65911.1     | AJ249831 | <i>Lemna minor</i>               | APR reducing enzyme, sulphur assimilation. adenosine 5'-phosphosulphate reductase. lapr.  |
| AAF18999.1     | AF212155 | <i>Allium cepa</i>               | APS-reductase.  |
| AAC26855.1     | AF069951 | <i>Enteromorpha intestinalis</i> | catalyzes the formation of sulfite and 5'-AMP from APS and reduced glutathione. 5'-adenylylsulfate reductase. EAPR1; sulfate assimilation enzyme; similar to <i>Escherichia coli</i> 3'-phosphoadenosine, 5'-phosphosulfate (PAPS) reductase encoded by cysH. |
| AAD02069.1     | AF036939 | <i>Chlamydomonas reinhardtii</i> | redox-regulator of 5'UTR psbA mRNA binding complex and translation. protein disulfide isomerase. localized to ER and chloroplast.   |
| AAC49896.1     | AF027727 | <i>Chlamydomonas reinhardtii</i> | involved in the redox-regulated binding of chloroplast poly(A)-binding protein to the 5'-UTR of psbA mRNA; regulates chloroplast translational activation. protein disulfide isomerase RB60. PDI.   |
| SEQ ID NO: 376 |          |                                  |   |
| AAC62017.1     | AF077547 | <i>Brassica juncea</i>           | arginine decarboxylase.   |
| AAF26434.1     | AF220097 | <i>Brassica juncea</i>           | arginine decarboxylase.   |
| AAF26435.1     | AF220098 | <i>Brassica juncea</i>           | arginine decarboxylase.   |
| AAB60880.1     | AF002017 | <i>Dianthus caryophyllus</i>     | arginine decarboxylase. gCARADC8.   |

|            |          |                          |  |
|------------|----------|--------------------------|--|
| AAF42972.1 | AF127241 | Nicotiana tabacum        | arginine decarboxylase 2. ADC2.  |
| BAA25685.1 | AB012873 | Nicotiana sylvestris     | arginine decarboxylase. NsADC-1.   |
| CAA85773.1 | Z37540   | Pisum sativum            | arginine decarboxylase.  |
| CAB64599.1 | AJ251898 | Datura stramonium        | polyamine biosynthesis. arginine decarboxylase 1. adc1.  |
| AAD09204.1 | U35367   | Glycine max              | arginine decarboxylase.  |
| BAA84799.1 | AP000559 | Oryza sativa             | ESTs C99670(E21043),C99671(E21043),<br>AU078262(R10938),AU078261(R10938),D15282(C0402) correspond to a region of the<br>predicted gene.; Similar to arginine decarboxylase (U52851). |
| AAB67887.1 | U63832   | Dianthus caryophyllus    | arginine decarboxylase. ADC.   |
| AAF42971.1 | AF127240 | Nicotiana tabacum        | arginine decarboxylase 1. ADC1.  |
| AAF42970.1 | AF127239 | Nicotiana tabacum        | arginine decarboxylase 1. ADC1.  |
| AAC68511.1 | AF045666 | Theobroma cacao          | arginine decarboxylase. spe2.  |
| CAA65585.1 | X96791   | Vitis vinifera           | arginine decarboxylase. ADC.   |
| AAA61347.1 | L16582   | Lycopersicon esculentum  | decarboxylation of L-arginine. arginine decarboxylase.   |
| AAC68530.1 | AF045685 | Arabidopsis arenosa      | arginine decarboxylase. spe2.  |
| AAC68529.1 | AF045684 | Capsella bursa-pastoris  | arginine decarboxylase. spe2.  |
| AAC68525.1 | AF045680 | Arabis drummondii        | arginine decarboxylase. spe2.  |
| AAC68526.1 | AF045681 | Barbarea vulgaris        | arginine decarboxylase. spe2.  |
| AAC68535.1 | AF045690 | Nasturtium officinale    | arginine decarboxylase. spe2.  |
| AAC68534.1 | AF045689 | Thellungiella salsuginea | arginine decarboxylase. spe2.  |
| AAC68533.1 | AF045688 | Thlaspi arvense          | arginine decarboxylase. spe2.  |
| AAC68532.1 | AF045687 | Stanleya pinnata         | arginine decarboxylase. spe2.  |

|                               |          |                                 |
|-------------------------------|----------|---------------------------------|
| AAC68531.1                    | AF045686 | <i>Sisymbrium altissimum</i>    |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68510.1                    | AF045665 | <i>Aethionema grandiflora</i>   |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68528.1                    | AF045683 | <i>Brassica oleracea</i>        |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68519.1                    | AF045674 | <i>Arabidopsis arenosa</i>      |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68527.1                    | AF045682 | <i>Brassica nigra</i>           |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68523.1                    | AF045678 | <i>Thellungiella salsuginea</i> |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68514.1                    | AF045669 | <i>Arabis drummondii</i>        |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68524.1                    | AF045679 | <i>Nasturtium officinale</i>    |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68522.1                    | AF045677 | <i>Thlaspi arvense</i>          |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68513.1                    | AF045668 | <i>Polanisia dodecandra</i>     |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68518.1                    | AF045673 | <i>Capsella bursa-pastoris</i>  |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68515.1                    | AF045670 | <i>Barbarea vulgaris</i>        |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68521.1                    | AF045676 | <i>Stanleya pinnata</i>         |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68520.1                    | AF045675 | <i>Sisymbrium altissimum</i>    |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68517.1                    | AF045672 | <i>Brassica oleracea</i>        |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68516.1                    | AF045671 | <i>Brassica nigra</i>           |
| arginine decarboxylase. spe2. |          |                                 |
| AAC68512.1                    | AF045667 | <i>Carica papaya</i>            |
| arginine decarboxylase. spe2. |          |                                 |
| CAA40137.1                    | X56802   | <i>Avena sativa</i>             |
| arginine decarboxylase. spe1. |          |                                 |
| AAD24801.1                    | AF132498 | <i>Brassica napus</i>           |
| arginine decarboxylase. ADC.  |          |                                 |
| BAA21617.1                    | AB005880 | <i>Nicotiana tabacum</i>        |
| arginine decarboxylase.       |          |                                 |
| AAB82607.1                    | AF026809 | <i>Ipomoea nil</i>              |
| arginine decarboxylase. adc.  |          |                                 |

## SEQ ID NO: 377

- AAG22606.1 AF258809 *Lycopersicon esculentum*  
aldehyde oxidase. AO2. molybdenum cofactor-binding enzyme.
- BAA23226.1 D88451 *Zea mays*  
aldehyde oxidase. zmAO-1.
- AAG22607.1 AF258810 *Lycopersicon esculentum*  
aldehyde oxidase. AO3. molybdenum cofactor-binding enzyme.
- AAG22605.1 AF258808 *Lycopersicon esculentum*  
aldehyde oxidase. AO1. molybdenum cofactor-binding enzyme.
- BAA23227.1 D88452 *Zea mays*  
aldehyde oxidase-2. zmAO-2. putative.
- AAB41742.1 U82559 *Lycopersicon esculentum*  
aldehyde oxidase 1 homolog. TAO1. has sequence properties of the molybdenum cofactor containing the hydroxylase genes aldehyde oxidase and xanthine dehydrogenase; belongs to a multigene family.
- AAG22608.1 AF259793 *Lycopersicon esculentum*  
aldehyde oxidase. AO1. molybdenum cofactor-binding enzyme.
- AAB41741.1 U82558 *Lycopersicon esculentum*  
aldehyde oxidase 1 homolog. TAO1. has sequence properties of the molybdenum cofactor containing the hydroxylase genes aldehyde oxidase and xanthine dehydrogenase; belongs to a multigene family.

## SEQ ID NO: 378

- BAB21211.1 AP002913 *Oryza sativa*  
putative DNA binding protein RAV2. P0480E02.17.
- BAB21218.1 AP002913 *Oryza sativa*  
putative DNA binding protein RAV2. P0480E02.24.
- BAA90643.1 AP001129 *Oryza sativa*  
Similar to *Arabidopsis thaliana* chromosome II BAC F11F19 genomic sequence, putative DNA-binding protein RAV2. (AC007017).
- BAA85426.1 AP000616 *Oryza sativa*  
similar to putative DNA-binding protein RAV2 (AC007017).

## SEQ ID NO: 380

- CAB65369.1 AJ250832 *Pisum sativum*  
germin-like protein. ger1.
- AAF03355.1 AF132671 *Nicotiana plumbaginifolia*  
nectarin I precursor. NEC1. germin-like protein.
- AAD38298.1 AC007789 *Oryza sativa*  
putative oxalate oxidase (germin protein). OSJNBa0049B20.25.
- BAB18339.1 AP002865 *Oryza sativa*  
putative germin protein. P0034C11.30. contains EST C97263(C53484).
- BAA25197.1 AB012138 *Lycopersicon esculentum*  
adaptation to Mn-deficiency. germin-like protein. Mdip1.



|            |          |                          |   |
|------------|----------|--------------------------|---|
| AAC78470.1 | AF067731 | <i>Solanum tuberosum</i> | germin-like protein. OXAOXA. similar to oxalate oxidase.  |
| AAC04835.1 | AF032974 | <i>Oryza sativa</i>      | germin-like protein 4. GER4. similar to wheat and barley oxalate oxidase.   |
| BAB39980.1 | AP003020 | <i>Oryza sativa</i>      | probable germin protein 4. P0498A12.8. contains ESTs AU101991(S4037),AU070167(R0031).   |
| BAB39965.1 | AP003018 | <i>Oryza sativa</i>      | probable germin protein 4. OSJNBa0004B13.19. contains ESTs AU101991(S4037),AU070167(R0031).   |
| AAC04833.1 | AF032972 | <i>Oryza sativa</i>      | germin-like protein 2. GER2. similar to wheat and barley oxalate oxidase.   |
| AAG00425.1 | AF250933 | <i>Hordeum vulgare</i>   | germin A. GerA. apoplastic protein; contains prepeptide for targeting into the cell wall.   |
| AAD43972.1 | AF141879 | <i>Oryza sativa</i>      | germin-like protein 2 precursor. RGLP2. similar to barley epidermis/papilla-specific oxalate oxidase-like protein.                  |
| AAD43973.1 | AF141880 | <i>Oryza sativa</i>      | germin-like protein 1 precursor. RGLP1. similar to barley epidermis/papilla-specific oxalate oxidase-like protein.                  |
| AAD43971.1 | AF141878 | <i>Oryza sativa</i>      | germin-like protein 1 precursor. RGLP1. similar to barley epidermis/papilla-specific oxalate oxidase-like protein.                  |
| CAB55559.1 | AJ237943 | <i>Triticum aestivum</i> | germin-like protein. glp2b.   |
| CAB55558.1 | AJ237942 | <i>Triticum aestivum</i> | germin-like protein. glp2a.   |
| AAC04837.1 | AF032976 | <i>Oryza sativa</i>      | germin-like protein 6. GER6. similar to wheat and barley oxalate oxidase.   |
| AAC04832.1 | AF032971 | <i>Oryza sativa</i>      | germin-like protein 1. GER1. similar to wheat and barley oxalate oxidase.   |
| CAA63659.1 | X93171   | <i>Hordeum vulgare</i>   | oxalate oxidase-like protein or germin-like protein.  |
| CAB55394.1 | AL117264 | <i>Oryza sativa</i>      | zwh0010.1. similar to Arabidopsis germin-like protein 6 (AF032976); Method: conceptual translation with partial peptide sequencing. |
| AAB97470.1 | AF042489 | <i>Oryza sativa</i>      | germin-like protein 16. glp16.  |
| AAC25777.1 | AF072694 | <i>Oryza sativa</i>      | germin-like protein 7. GER7. similar to wheat and barley oxalate oxidase.   |
| AAG00427.1 | AF250935 | <i>Hordeum vulgare</i>   | germin F. GerF. apoplastic protein; contains prepeptide for targeting into the cell wall.   |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAG00426.1 | AF250934 | Hordeum vulgare               | germin B. GerB. apoplastic protein; contains prepeptide for targeting into the cell wall. |
| BAA78563.1 | AB024338 | Atriplex lentiformis          | germin-like protein.  |
| AAA20245.1 | U01963   | Hordeum vulgare               | germin subunit.   |
| AAG00428.1 | AF250936 | Hordeum vulgare               | germin D. GerD. apoplastic protein; contains prepeptide for targeting into the cell wall. |
| AAC99473.1 | AF039201 | Pinus caribaea                | germin-like protein. PcGER1.  |
| AAC04834.1 | AF032973 | Oryza sativa                  | germin-like protein 3. GER3. similar to wheat and barley oxalate oxidase.                 |
| AAA34271.1 | M63224   | Triticum aestivum             | germin. germin 9f-3.8.  |
| AAC05146.1 | AF049065 | Pinus radiata                 | germin-like protein. PRGer1.  |
| AAA34268.1 | M21962   | Triticum aestivum             | germin protein precursor.   |
| AAA34270.1 | M63223   | Triticum aestivum             | germin. germin 9f-2.8.  |
| CAA71052.1 | Y09917   | Triticum aestivum             | germin homolog. pSBGer3.  |
| AAG00429.1 | AF250937 | Hordeum vulgare               | germin E. GerE. apoplastic protein.   |
| BAA86880.1 | AB028454 | Barbula unguiculata           | germin-like protein.  |
| CAB65370.1 | AJ250833 | Pisum sativum                 | germin-like protein. ger2a. 1st variant of this clone.                                    |
| AAA33030.1 | M93041   | Mesembryanthemum crystallinum | germin-like protein. germin-like protein.   |
| CAB65371.1 | AJ250834 | Pisum sativum                 | germin-like protein. ger2b. 2nd variant of the clone PsGER2.                              |
| CAA71050.1 | Y09915   | Triticum aestivum             | germin homolog. pSBGer1.  |
| CAA71051.1 | Y09916   | Triticum aestivum             | germin homolog. pSBGer2.  |
| AAA86365.1 | U21743   | Brassica napus                | germin-like protein. similar to product encoded by GenBank Accession Number X84786.       |
| CAC34417.1 | AJ311624 | Pisum sativum                 | Germin-like protein. glp3.  |

SEQ ID NO: 389

|                |          |  |
|----------------|----------|--|
| CAA04703.1     | AJ001370 | <i>Olea europaea</i><br>cytochrome b5. cytb5-2.  |
| AAA32990.1     | M87514   | <i>Brassica oleracea</i><br>cytochrome b-5. cytochrome b-5.  |
| CAA50575.1     | X71441   | <i>Nicotiana tabacum</i><br>cytochrome b5.   |
| CAA53366.1     | X75670   | <i>Oryza sativa</i><br>cytochrome b5.  |
| CAA04702.1     | AJ001369 | <i>Olea europaea</i><br>cytochrome b5. cytb5-1.  |
| AAA62621.1     | L22209   | <i>Cuscuta reflexa</i><br>associated with cytokinin-induced haustoria formation in <i>Cuscuta reflexa</i> . cytochrome b5.   |
| AAC49701.1     | U79011   | <i>Borago officinalis</i><br>haem-binding protein. cytochrome b5.  |
| CAA56318.1     | X80008   | <i>Nicotiana tabacum</i><br>cytochrome b5.   |
| CAA48240.1     | X68140   | <i>Nicotiana tabacum</i><br>cytochrome b5.   |
| AAD10774.1     | AF098510 | <i>Petunia x hybrida</i><br>involved in anthocyanin biosynthesis. cytochrome b5 DIF-F. diff. required for full activity of flavonoid 3',5' hydroxylase.                                    |
| AAF60299.1     | AF233640 | <i>Petunia x hybrida</i><br>involved in anthocyanin biosynthesis. cytochrome b5 DIF-F. diff.   |
| CAA11033.1     | AJ222981 | <i>Physcomitrella patens</i><br>delta6-acyl-lipid desaturase. des6. des6 represents a fusion between a C-terminal desaturase with a cytochrome b5-related part and a N-terminal extension. |
| SEQ ID NO: 392 |          |  |
| AAK28303.1     | AF346431 | <i>Nicotiana tabacum</i><br>phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.   |
| AAB36653.1     | U32644   | <i>Nicotiana tabacum</i><br>immediate-early salicylate-induced glucosyltransferase. IS5a.  |
| BAB17061.1     | AP002523 | <i>Oryza sativa</i><br>putative glucosyl transferase. P0013F10.7. contains EST C73149(E2992).  |
| AAB36652.1     | U32643   | <i>Nicotiana tabacum</i><br>immediate-early salicylate-induced glucosyltransferase. IS10a.   |
| AAK28304.1     | AF346432 | <i>Nicotiana tabacum</i><br>phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.   |
| BAB17059.1     | AP002523 | <i>Oryza sativa</i><br>putative glucosyl transferase. P0013F10.5.  |
| BAB17060.1     | AP002523 | <i>Oryza sativa</i><br>putative glucosyl transferase. P0013F10.6.  |
| CAA59450.1     | X85138   | <i>Lycopersicon esculentum</i><br>twil. homologous to glucosyltransferases.  |

|            |          |   |
|------------|----------|---|
| CAB56231.1 | Y18871   | <i>Dorotheanthus bellidiformis</i><br>betanidin-5-O-glucosyltransferase.  |
| BAA83484.1 | AB031274 | <i>Scutellaria baicalensis</i><br>UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.   |
| AAB48444.1 | U82367   | <i>Solanum tuberosum</i><br>UDP-glucose glucosyltransferase.  |
| BAA36410.1 | AB012114 | <i>Vigna mungo</i><br>UDP-glycose:flavonoid glycosyltransferase. UFGlyT.  |
| CAA54610.1 | X77460   | <i>Manihot esculenta</i><br>UTP-glucose glucosyltransferase. CGT4.  |
| CAB88666.1 | AJ400861 | <i>Cicer arietinum</i><br>flavonoid glycosyltransferase. putative UDP-glycose.  |
| AAK16180.1 | AC079887 | <i>Oryza sativa</i><br>putative glucosyltransferase. OSJNBa0040E01.21.  |
| AAD51778.1 | AF116858 | <i>Phaseolus vulgaris</i><br>utilizes UDPX as the sugar donor and catalyzes the formation of O-xylosylzeatin from zeatin.<br>zeatin O-xylosyltransferase. ZOX1.   |
| AAD04166.1 | AF101972 | <i>Phaseolus lunatus</i><br>catalyzes formation of O-glucosylzeatin from zeatin and UDPG; catalyzes formation of O-xylosylzeatin from zeatin and UDPX. zeatin O-glucosyltransferase. ZOG1. cytokinin O-glucosyltransferase. |
| BAA89009.1 | AB027455 | <i>Petunia x hybrida</i><br>anthocyanin 5-O-glucosyltransferase. PH1.   |
| CAC09351.1 | AL442007 | <i>Oryza sativa</i><br>putative glucosyltransferase. H0212B02.7.  |
| AAB62270.1 | AF006081 | <i>Solanum berthaultii</i><br>UDPG glucosyltransferase. PLGT.   |
| AAK16172.1 | AC079887 | <i>Oryza sativa</i><br>putative glucosyltransferase. OSJNBa0040E01.14.  |
| AAF61647.1 | AF190634 | <i>Nicotiana tabacum</i><br>UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| CAA54612.1 | X77462   | <i>Manihot esculenta</i><br>UTP-glucose glucosyltransferase. CGT5.  |
| AAK16181.1 | AC079887 | <i>Oryza sativa</i><br>putative glucosyltransferase. OSJNBa0040E01.16.  |
| AAK16178.1 | AC079887 | <i>Oryza sativa</i><br>putative glucosyltransferase. OSJNBa0040E01.5.   |
| BAA36412.1 | AB012116 | <i>Vigna mungo</i><br>UDP-glycose:flavonoid glycosyltransferase. UFGlyT.  |
| AAK16175.1 | AC079887 | <i>Oryza sativa</i><br>putative glucosyltransferase. OSJNBa0040E01.15.  |

|            |          |                        |   |
|------------|----------|------------------------|---|
| AAF98390.1 | AF287143 | Brassica napus         | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT.               |
| BAA36423.1 | AB013598 | Verbena x hybrida      | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| AAF17077.1 | AF199453 | Sorghum bicolor        | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |
| BAA93039.1 | AB033758 | Citrus unshiu          | limonoid UDP-glucosyltransferase. LGTase.   |
| BAB17176.1 | AP002843 | Oryza sativa           | putative UTP-glucose glucosyltransferase. P0407B12.13.  |
| BAB17182.1 | AP002843 | Oryza sativa           | putative UTP-glucose glucosyltransferase. P0407B12.19.  |
| BAA89008.1 | AB027454 | Petunia x hybrida      | anthocyanidin 3-O-glucosyltransferase. PGT8.  |
| CAA54558.1 | X77369   | Solanum melongena      | glycosyl transferase. GT.   |
| CAA54611.1 | X77461   | Manihot esculenta      | UTP-glucose glucosyltransferase. CGT2.  |
| CAA54609.1 | X77459   | Manihot esculenta      | UTP-glucose glucosyltransferase. CGT1.  |
| BAA12737.1 | D85186   | Gentiana triflora      | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| AAD21086.1 | AF127218 | Forsythia x intermedia | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.   |
| AAG25643.1 | AF303396 | Phaseolus vulgaris     | UDP-glucosyltransferase HRA25. putative; defense associated.  |
| BAA36411.1 | AB012115 | Vigna mungo            | UDP-glycosyl:flavonoid glycosyltransferase. UFGlyT.   |
| BAA19155.1 | AB000623 | Nicotiana tabacum      | glucosyl transferase. JIGT.   |
| AAD55985.1 | AF165148 | Petunia x hybrida      | catalyzes the penultimate step of flavonol glucosylgalactoside biosynthesis from UDP-galactose and flavonol aglycones in petunia pollen. UDP-galactose:flavonol 3-O-galactosyltransferase. F3galtase. |
| CAA54613.1 | X77463   | Manihot esculenta      | UTP-glucose glucosyltransferase. CGT6.  |
| BAA36421.1 | AB013596 | Perilla frutescens     | UDP-glucose:anthocyanin 5-O-glucosyltransferase. PF3R4.   |

|                |          |  |
|----------------|----------|--|
| BAA19659.1     | AB002818 | <i>Perilla frutescens</i><br>flavonoid 3-O-glucosyltransferase. UDP glucose.   |
| AAB81683.1     | AF000372 | <i>Vitis vinifera</i><br>UDP glucose:flavonoid 3-o-glucosyltransferase.  |
| BAB41025.1     | AB047098 | <i>Vitis vinifera</i><br>UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.   |
| BAB41023.1     | AB047096 | <i>Vitis vinifera</i><br>UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.   |
| SEQ ID NO: 393 |          |  |
| CAB56756.1     | AJ011589 | <i>Pisum sativum</i><br>5,10-methylenetetrahydrofolate dehydrogenase: 5,10-methenyltetrahydrofolate<br>cyclohydrolase. bifunctional enzyme.  |
| AAD01907.1     | AF030516 | <i>Pisum sativum</i><br>5,10-methylenetetrahydrofolate dehydrogenase-5,10-methenyltetrahydrofolate cyclohydrolase.<br>bifunctional protein; 31.3 kDa protein.                      |
| AAG48834.1     | AC084218 | <i>Oryza sativa</i><br>similar to <i>Pisum sativum</i> methylenetetrahydrofolate dehydrogenase (NADP+) (EC 1.5.1.5)<br>(AF030516).   |
| SEQ ID NO: 406 |          |  |
| AAD46491.1     | AF135014 | <i>Zea mays</i><br>dihydrolipoamide S-acetyltransferase. pyruvate dehydrogenase complex E2 subunit.  |
| BAA90623.1     | AP001129 | <i>Oryza sativa</i><br>ESTs AU033004(S0924),C74754(E50863) correspond to a region of the predicted gene.;<br>Similar to Rat mRNA for dihydrolipoamide acetyltransferase. (D10655). |
| BAA77024.1     | AB026124 | <i>Lithospermum erythrorhizon</i><br>dihydrolipoamide acetyltransferase.   |
| SEQ ID NO: 407 |          |  |
| AAF69017.1     | AF261654 | <i>Dianthus caryophyllus</i><br>ethylene-insensitive 3-like protein 1. EIL1. EIN3-like protein.  |
| CAC09582.1     | AJ298994 | <i>Fagus sylvatica</i><br>gibberellic acid (GA3)-induced. ethylene insensitive (EIN3/EIL)-like transcription regulator.<br>ein11.  |
| AAG00419.1     | AF247568 | <i>Nicotiana tabacum</i><br>EIN3. component in ethylene signal transduction pathway.   |
| SEQ ID NO: 408 |          |  |
| AAC15870.1     | AF002016 | <i>Cucurbita</i> sp.<br>acyl CoA oxidase homolog.  |
| AAF14635.1     | AF202987 | <i>Petroselinum crispum</i><br>acyl-CoA oxidase. ACO. peroxisomal acyl-CoA oxidase.  |
| AAB67883.1     | U66299   | <i>Phalaenopsis</i> sp. 'True Lady'<br>acyl-CoA oxidase homolog.   |

|                |          |  |
|----------------|----------|--|
| AAC32108.1     | AF051203 | <i>Picea mariana</i><br>acyl-CoA oxidase homolog. Sb06. similar to Phalaenopsis sp. acyl-CoA oxidase homolog<br>encoded by GenBank Accession Number U66299.  |
| CAA04688.1     | AJ001341 | <i>Hordeum vulgare</i><br>putative acyl-CoA oxidase.   |
| BAB08201.1     | AP002539 | <i>Oryza sativa</i><br>ESTs AU056822(S20908), C26441(C12328), C28477(C61243) correspond to a region of the<br>predicted gene. Similar to <i>Arabidopsis thaliana</i> putative acyl-coA dehydrogenase (AF049236). |
| BAA96762.1     | AP002521 | <i>Oryza sativa</i><br>ESTs AU056822(S20908), C26441(C12328), C28477(C61243) correspond to a region of the<br>predicted gene. Similar to <i>Arabidopsis thaliana</i> putative acyl-coA dehydrogenase (AF049236). |
| CAB55555.1     | AJ010946 | <i>Pisum sativum</i><br>auxin binding protein (ABP44). abp44/ivdh. has both auxin binding capability and enzymatic<br>activity; isovaleryl-CoA Dehydrogenase.  |
| CAB55554.1     | AJ010945 | <i>Pisum sativum</i><br>auxin binding protein (ABP44). abp44/ivdh. has both auxin binding capability and enzymatic<br>activity; Isovaleryl-CoA Dehydrogenase.  |
| CAC08233.1     | AJ278987 | <i>Solanum tuberosum</i><br>leucine catabolism. isovaleryl-CoA dehydrogenase. ivd1.  |
| CAC08234.1     | AJ278988 | <i>Solanum tuberosum</i><br>leucine catabolism. isovaleryl-CoA dehydrogenase. ivd2.  |
| AAF02449.1     | AF127432 | <i>Picea abies</i><br>acyl-CoA oxidase homolog. Sb06.  |
| AAF02451.1     | AF127434 | <i>Picea abies</i><br>acyl-CoA oxidase homolog. Sb06.  |
| AAF02450.1     | AF127433 | <i>Picea abies</i><br>acyl-CoA oxidase homolog. Sb06.  |
| AAC32152.1     | AF051733 | <i>Picea mariana</i><br>acyl-CoA oxidase homolog. Sb06.  |
| AAC32153.1     | AF051734 | <i>Picea mariana</i><br>acyl-CoA oxidase homolog. Sb06.  |
| CAA58874.1     | X84055   | <i>Hordeum vulgare</i><br>mammalian acyl CoA oxidase homologous. cdr29. putative.  |
| SEQ ID NO: 409 |          |  |
| CAB64356.1     | AJ251511 | <i>Populus tremula</i> x <i>Populus tremuloides</i><br>mitochondrial oxidase. alternative oxidase. aox1.   |
| AAC60576.1     | S71335   | Mitochondrion <i>Nicotiana tabacum</i><br>alternative oxidase. Aox1. This sequence comes from Fig. 1; AOX.   |
| CAA56163.1     | X79768   | <i>Nicotiana tabacum</i><br>alternative oxidase. aox.  |
| CAA48653.1     | X68702   | <i>Glycine max</i><br>alternative oxidase. aox1. putative.   |

|            |          |                                       |  |
|------------|----------|---------------------------------------|--|
| AAC35354.1 | AF083880 | Glycine max                           | alternative oxidase precursor. Aox1. AOX1; terminal oxidase of mitochondrial electron transport chain. |
| BAB21500.1 | AB055060 | Catharanthus roseus                   | alternative oxidase.   |
| BAA23803.1 | AB009395 | Catharanthus roseus                   | alternative oxidase.   |
| BAA86963.1 | AB007452 | Oryza sativa                          | alternative oxidase. Aox1(Ao1-1).  |
| BAA28773.1 | AB004864 | Oryza sativa                          | alternative oxidase. AOX1a.  |
| BAA28772.1 | AB004813 | Oryza sativa                          | alternative oxidase. AOX1a.  |
| CAA78823.1 | Z15117   | Sauromatum guttatum                   | salicylic acid-inducible alternative oxidase. aox1.  |
| AAA34048.1 | M60330   | Sauromatum guttatum                   | alternative oxidase protein. aox1.   |
| AAD51707.1 | AF174004 | Triticum aestivum                     | alternative oxidase.   |
| CAA55892.1 | X79329   | Mangifera indica                      | alternative oxidase. AOMI 1.   |
| CAB72441.1 | AJ271889 | Populus tremula x Populus tremuloides | mitochondrial oxidase. alternative oxidase. aox1b.   |
| AAB97285.1 | U87906   | Glycine max                           | alternative oxidase. Aox2.   |
| BAA28771.1 | AB004813 | Oryza sativa                          | alternative oxidase. AOX1b.  |
| BAA28774.1 | AB004865 | Oryza sativa                          | alternative oxidase. AOX1b.  |
| AAB97286.1 | U87907   | Glycine max                           | alternative oxidase. Aox3.   |
| AAB97839.1 | AF040566 | Zea mays                              | alternative oxidase. Aox.  |
| AAG33634.1 | AF314255 | Chlamydomonas reinhardtii             | alternative oxidase 2. AOX2.   |
| AAG02081.1 | AF285187 | Chlamydomonas reinhardtii             | alternative oxidase. AOX2.   |
| AAG33633.1 | AF314254 | Chlamydomonas reinhardtii             | alternative oxidase 1. AOX1.   |
| AAC05743.2 | AF047832 | Chlamydomonas reinhardtii             | alternative oxidase. AOX1.   |
| BAA23725.1 | AB009087 | Chlamydomonas sp. W80                 | alternative oxidase.   |



|                |          |   |
|----------------|----------|---|
| AAC34192.1     | S81466   | Glycine max<br>alternative oxidase Aox1 precursor. Aox1. This sequence comes from Fig. 1. |
| AAG18450.1     | AF302932 | Lycopersicon esculentum<br>plastid quinol oxidase.  |
| AAG18449.1     | AF302931 | Lycopersicon esculentum<br>plastid quinol oxidase.  |
| AAG02287.1     | AF177980 | Lycopersicon esculentum<br>plastid terminal oxidase. PTOX.                                |
| AAG02286.1     | AF177979 | Lycopersicon esculentum<br>plastid terminal oxidase. PTOX.                                |
| AAG02288.1     | AF177981 | Capsicum annuum<br>plastid terminal oxidase. PTOX.  |
| AAB36072.1     | S81470   | Glycine max<br>Aox2. Aox2. alternative oxidase 2; This sequence comes from Fig. 1.        |
| AAC35554.2     | AF085174 | Oryza sativa<br>oxidase. IM1.   |
| AAG00450.1     | AF274001 | Triticum aestivum<br>oxidase. IM1.  |
| SEQ ID NO: 410 |          |   |
| CAA69600.1     | Y08292   | Nicotiana plumbaginifolia<br>NADH glutamate dehydrogenase. GDH A.                         |
| CAB94836.1     | AJ277949 | Nicotiana plumbaginifolia<br>NADH glutamate dehydrogenase. gdhA.                          |
| CAC18730.1     | AJ303070 | Vitis vinifera<br>NADH glutamate dehydrogenase. gdhA.                                     |
| CAA60507.1     | X86924   | Vitis vinifera<br>glutamate dehydrogenase. GDH.   |
| BAA08445.1     | D49475   | Zea mays<br>glutamate dehydrogenase.  |
| AAB51596.1     | U93561   | Zea mays<br>glutamate dehydrogenase. GDH1. wild type.                                     |
| AAB51595.1     | U93560   | Zea mays<br>glutamate dehydrogenase mutant. GDH1. mutant allele.                          |
| CAB94837.1     | AJ277950 | Nicotiana plumbaginifolia<br>NADH-glutamate dehydrogenase. gdhB.                          |
| CAA09478.1     | AJ011096 | Asparagus officinalis<br>glutamate dehydrogenase. gdhb.                                   |
| CAA69601.2     | Y08293   | Nicotiana plumbaginifolia<br>NADH glutamate dehydrogenase. GDH B.                         |
| AAB39508.1     | U48695   | Lycopersicon esculentum<br>glutamate dehydrogenase. legdh1.                               |

|                |          |  |
|----------------|----------|--|
| CAA09456.1     | AJ011006 | <i>Asparagus officinalis</i><br>NADH glutamate dehydrogenase. gdhA.  |
| CAA41635.1     | X58831   | <i>Chlorella sorokiniana</i><br>glutamate dehydrogenase (NADP+). gdhANC. There are other coding exons before the first exon given for this gene.   |
| CAA41636.1     | X58832   | <i>Chlorella sorokiniana</i><br>glutamate dehydrogenase (NADP+). gdhANC.   |
| SEQ ID NO: 412 |          |  |
| CAA65456.2     | X96681   | <i>Oryza sativa</i><br>transcription factor. DNA-binding protein. Oshox1. homeodomain leucine zipper gene.   |
| AAK00416.1     | AC069324 | <i>Oryza sativa</i><br>Putative DNA-binding protein. OSJNBa0071K19.2.  |
| BAA05622.1     | D26573   | <i>Daucus carota</i><br>transcriptional regulator. DNA-binding protein. homeodomain at nt 585-764; leucine zipper at nt 765-851.   |
| BAA05625.1     | D26576   | <i>Daucus carota</i><br>transcriptional regulator. DNA-binding protein. homeodomain at nt 300-479; leucine zipper at nt 480-587.   |
| BAA21017.1     | D26578   | <i>Daucus carota</i><br>transcriptional regulator. DNA-binding protein. homeodomain at nt 761-940; leucine zipper at nt 941-1048.  |
| BAA05624.1     | D26575   | <i>Daucus carota</i><br>transcriptional regulator. DNA-binding protein. homeodomain at nt 520-699; leucine zipper at nt 700-805.   |
| BAA05623.1     | D26574   | <i>Daucus carota</i><br>transcriptional regulator. DNA-binding protein. homeodomain at nt 498-677; leucine zipper at nt 678-785.   |
| SEQ ID NO: 414 |          |  |
| BAA95814.1     | AP002069 | <i>Oryza sativa</i><br>ESTs AU082579(S2069),D40238(S2069) correspond to a region of the predicted gene. Similar to <i>Arabidopsis thaliana</i> vesicle-associated membrane protein 7C; synaptobrevin 7C. (AF025332). |
| SEQ ID NO: 415 |          |  |
| CAA45025.1     | X63431   | <i>Rauvolfia mannii</i><br>strictosidine synthase. str1.   |
| CAA68725.1     | Y00756   | <i>Rauvolfia serpentina</i><br>strictosidine synthase.   |
| CAA44208.1     | X62334   | <i>Rauvolfia serpentina</i><br>strictosidine synthase. str1.   |
| CAA37671.1     | X53602   | <i>Catharanthus roseus</i><br>strictosidine synthase precursor.  |
| CAA71255.1     | Y10182   | <i>Catharanthus roseus</i><br>strictosidine synthase. str1.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| CAA43936.1     | X61932   | Catharanthus roseus     | strictosidine synthase. sss.  |
| AAF75751.1     | AF261141 | Lycopersicon esculentum | putative strictosidine synthase.  |
| CAB53484.1     | AJ245900 | Oryza sativa            |   |
| CAA303711.1    |          | protein. q3037.11.      | Similar to strictosidine synthase 3 precursor.  |
| SEQ ID NO: 418 |          |                         |   |
| AAG22607.1     | AF258810 | Lycopersicon esculentum | aldehyde oxidase. AO3. molybdenum cofactor-binding enzyme.  |
| AAG22606.1     | AF258809 | Lycopersicon esculentum | aldehyde oxidase. AO2. molybdenum cofactor-binding enzyme.  |
| AAG22605.1     | AF258808 | Lycopersicon esculentum | aldehyde oxidase. AO1. molybdenum cofactor-binding enzyme.  |
| BAA23227.1     | D88452   | Zea mays                | aldehyde oxidase-2. zmAO-2. putative.   |
| BAA23226.1     | D88451   | Zea mays                | aldehyde oxidase. zmAO-1.   |
| AAB41742.1     | U82559   | Lycopersicon esculentum | aldehyde oxidase 1 homolog. TAO1. has sequence properties of the molybdenum cofactor containing the hydroxylase genes aldehyde oxidase and xanthine dehydrogenase; belongs to a multigene family. |
| AAG22608.1     | AF259793 | Lycopersicon esculentum | aldehyde oxidase. AO1. molybdenum cofactor-binding enzyme.  |
| AAB41741.1     | U82558   | Lycopersicon esculentum | aldehyde oxidase 1 homolog. TAO1. has sequence properties of the molybdenum cofactor containing the hydroxylase genes aldehyde oxidase and xanthine dehydrogenase; belongs to a multigene family. |
| SEQ ID NO: 421 |          |                         |   |
| BAA13635.1     | D88530   | Spinacia oleracea       | serine acetyltransferase.   |
| BAA13634.1     | D88529   | Spinacia oleracea       | serine acetyltransferase.   |
| BAA93050.1     | AB040502 | Allium tuberosum        | serine acetyltransferase. ASAT5.  |
| BAA08479.1     | D49535   | Citrullus lanatus       | serine acetyltransferase.   |
| BAA12843.1     | D85624   | Citrullus lanatus       | serine acetyltransferase. Sat.  |
| BAA21827.1     | AB006530 | Citrullus lanatus       | serine acetyltransferase. Sat.  |
| AAF19000.1     | AF212156 | Allium cepa             | serine acetyltransferase.   |
| SEQ ID NO: 422 |          |                         |   |

|            |          |                   |  |
|------------|----------|-------------------|--|
| BAA05079.1 | D26086   | Petunia x hybrida | zinc-finger protein.   |
| CAB77055.1 | Y18788   | Medicago sativa   | putative TFIIIA (or kruppel)-like zinc finger protein.                 |
| AAD26942.1 | AF119050 | Datisca glomerata | zinc-finger protein 1. zfp1. DgZFP1.                                   |
| AAB39638.1 | U68763   | Glycine max       | putative transcription factor. SCOF-1. scof-1. zinc-finger protein.    |
| AAC06243.1 | AF053077 | Nicotiana tabacum | transcription factor. osmotic stress-induced zinc-finger protein. zfp. |
| BAA05077.1 | D26084   | Petunia x hybrida | zinc-finger DNA binding protein.                                       |
| BAA05076.1 | D26083   | Petunia x hybrida | zinc-finger DNA binding protein.                                       |
| BAA05078.1 | D26085   | Petunia x hybrida | zinc-finger DNA binding protein.                                       |
| AAK01713.1 | AF332876 | Oryza sativa      | zinc finger transcription factor ZF1.                                  |
| AAB53260.1 | U76554   | Brassica rapa     | transcription factor. zinc-finger protein-1. BR42.                     |
| AAB53261.1 | U76555   | Brassica rapa     | zinc-finger protein BcZFP1. BcZFP1(3-2z).                              |
| BAA96070.1 | AB035132 | Petunia x hybrida | C2H2 zinc-finger protein ZPT2-10. PETHy;ZPT2-10.                       |
| BAA21919.1 | AB006597 | Petunia x hybrida | ZPT2-10. C2H2 zinc finger protein, 2 finger.                           |
| BAA96071.1 | AB035133 | Petunia x hybrida | C2H2 zinc-finger protein ZPT3-3. ZPT3-3.                               |
| BAA21927.1 | AB006605 | Petunia x hybrida | ZPT3-3. C2H2 zinc finger protein, 3 finger.                            |
| BAA19112.1 | AB000453 | Petunia x hybrida | PETHy;ZPT3-1. Cys(2) His(2) zinc finger protein, 3 fingers.            |
| BAA21928.1 | AB006606 | Petunia x hybrida | ZPT4-4. C2H2 zinc finger protein, 4 finger.                            |
| BAA21922.1 | AB006600 | Petunia x hybrida | ZPT2-13. C2H2 zinc finger protein, 2finger.                            |
| BAA19114.1 | AB000455 | Petunia x hybrida | PETHy;ZPT4-1. Cys(2) His(2) zinc finger protein, 4 fingers.            |
| BAA21920.1 | AB006598 | Petunia x hybrida | ZPT2-11. C2H2 zinc finger protein, 2finger.                            |
| BAA21921.1 | AB006599 | Petunia x hybrida | ZPT2-12. C2H2 zinc finger protein, 2 finger.                           |

---

BAA19110.1 AB000451 *Petunia x hybrida*  
 PETHy;ZPT2-5. Cys(2) His(2) zinc finger protein, 2 fingers.

---

BAA21926.1 AB006604 *Petunia x hybrida*  
 ZPT2-9. C2H2 zinc finger protein, 2 finger.

---

BAA21925.1 AB006603 *Petunia x hybrida*  
 ZPT2-8. C2H2 zinc finger protein, 2 finger.

---

CAA60828.1 X87374 *Pisum sativum*  
 putative zinc finger protein.

---

BAA19111.1 AB000452 *Petunia x hybrida*  
 PETHy;ZPT2-6. Cys(2) His(2) zinc finger protein, 2 fingers.

---

BAA21923.1 AB006601 *Petunia x hybrida*  
 ZPT2-14. C2H2 zinc finger protein, 2 finger.

---

BAA21924.1 AB006602 *Petunia x hybrida*  
 ZPT2-7. C2H2 zinc finger protein, 2finger.

---

BAA19113.1 AB000454 *Petunia x hybrida*  
 PETHy;ZPT3-2. Cys(2) His(2) zinc finger protein, 3 fingers.

---

BAA19926.1 AB000456 *Petunia x hybrida*  
 PETHy; ZPT4-2. C2H2 zinc finger protein, 4 finger.

---

SEQ ID NO: 423

---

AAC49815.1 U87257 *Daucus carota*  
 4-hydroxyphenylpyruvate dioxygenase.

---

CAA04245.1 AJ000693 *Hordeum vulgare*  
 4-hydroxyphenylpyruvate dioxygenase.

---

SEQ ID NO: 424

---

AAC72193.1 AF069909 *Zea mays*  
 pyruvate dehydrogenase E1 beta subunit isoform 2.

---

AAC72192.1 AF069908 *Zea mays*  
 pyruvate dehydrogenase E1 beta subunit isoform 1.

---

AAC72194.1 AF069910 *Zea mays*  
 pyruvate dehydrogenase E1 beta subunit isoform 3.

---

AAB01223.1 U56697 *Pisum sativum*  
 pyruvate dehydrogenase E1beta.

---

AAC32149.1 AF051249 *Picea mariana*  
 pyruvate dehydrogenase E1 beta subunit. Sb68.

---

AAF43837.1 AF166114 *Chloroplast Mesostigma viride*  
 beta subunit of pyruvate dehydrogenase E1 component. odpB.

---

AAD22077.1 AF124755 *Pinus banksiana*  
 pyruvate dehydrogenase E1 beta subunit. Sb68.

---

CAA75778.1 Y15782 *Capsicum annuum*  
 transketolase 2.

---

AAB88295.1 AF024512 *Oryza sativa*  
 CLA1 transketolase-like protein. CLA1. similar to Arabidopsis CLA1 product; required for chloroplast development.

---

## SEQ ID NO: 426

AAB53764.1 U96713 Brassica rapa  
aminoalcoholphosphotransferase. AAPT1.

AAD56040.1 AF183933 Brassica rapa  
aminoalcoholphosphotransferase. AAPT3.

AAC79507.1 U96439 Pimpinella brachycarpa  
aminoalcoholphosphotransferase. AAPTase.

AAA67719.1 U12735 Glycine max  
CDP-choline:diacylglycerol cholinephosphotransferase activity and possibly CDP-ethanolamine: diacylglycerol ethanolaminephosphotransferase activity.  
aminoalcoholphosphotransferase. AAPT1.

## SEQ ID NO: 427

CAA56313.1 X79992 Avena sativa  
putative pp70 ribosomal protein S6 kinase. Aspk11.

CAB89082.1 AJ277534 Asparagus officinalis  
S6 ribosomal protein kinase. pk1. putative.

AAK18843.1 AC082645 Oryza sativa  
putative protein kinase. OSJNBb0033N16.3.

AAC05084.1 AF033097 Avena sativa  
NPH1-2. NPH1-2. putative serine/threonine protein kinase.

AAC05083.1 AF033096 Avena sativa  
NPH1-1. NPH1-1. putative serine/threonine protein kinase.

AAK13156.1 AC078829 Oryza sativa  
putative protein kinase. OSJNBa0026O12.14.

AAB88817.1 AF033263 Zea mays  
signal transduction for phototropism. nonphototropic hypocotyl 1. nph1. NPH1; putative serine/threonine kinase; similar to oat NPH1 proteins.

CAB82852.1 Z30329 Mesembryanthemum crystallinum  
protein kinase MK6.

BAB18104.1 AB042714 Chlamydomonas reinhardtii  
cyclic nucleotide dependent protein kinase. CL-PK1.

BAB18105.1 AB042715 Chlamydomonas reinhardtii  
cyclic nucleotide dependent protein kinase II. CL-PK2.

BAA83689.1 AB011968 Oryza sativa  
OsPK7. OsPK7. protein kinase.

AAD31900.1 AF145482 Mesembryanthemum crystallinum  
putative serine/threonine protein kinase.

BAA92970.1 AP001551 Oryza sativa  
Similar to Arabidopsis thaliana chromosome 4 BAC clone F28A21 ; putative protein kinase. (AL035526).

|            |          |                         |  |
|------------|----------|-------------------------|--|
| BAA92972.1 | AP001551 | Oryza sativa            | ESTs AU056183(S20356),AU056881(S20950) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 4 BAC clone F6I18 ; putative protein kinase. (AL022198). |
| BAA83688.1 | AB011967 | Oryza sativa            | OsPK4. OsPK4. protein kinase.  |
| AAF22219.1 | AF141378 | Zea mays                | protein kinase PK4. ZmPK4.   |
| CAA73067.1 | Y12464   | Sorghum bicolor         | serine/threonine kinase. SNFL1.  |
| BAA96628.1 | AP002482 | Oryza sativa            | ESTs D41739(S4522),AU055999(S20214), AU057588(S21592) correspond to a region of the predicted gene. Similar to Sorghum bicolor serine/threonine kinase (Y12465).                         |
| CAA89202.1 | Z49233   | Chlamydomonas eugametos | calcium-stimulated protein kinase.   |
| CAA73068.1 | Y12465   | Sorghum bicolor         | serine/threonine kinase. SNFL2.  |
| BAA34675.1 | AB011670 | Triticum aestivum       | wpk4 protein kinase. wpk4.   |
| CAA74646.1 | Y14274   | Sorghum bicolor         | putative serine/threonine protein kinase. SNFL3.   |
| AAF06970.1 | AF162662 | Kalanchoe fedtschenkoi  | phosphoenolpyruvate carboxylase kinase. calcium-independent protein kinase.  |
| AAF06969.1 | AF162661 | Kalanchoe fedtschenkoi  | phosphoenolpyruvate carboxylase kinase. calcium-independent protein kinase.  |
| CAA39936.1 | X56599   | Daucus carota           | calcium- dependent protein kinase. DcPK431.  |
| BAA99439.1 | AP002743 | Oryza sativa            | putative protein kinase. P0710E05.26. contains ESTs C22394(C30013),C22393(C30013).   |
| BAA90814.1 | AP001168 | Oryza sativa            | ESTs AU030197(E50746),AU030196(E50746) correspond to a region of the predicted gene.; Similar to calcium-dependent calmodulin-independent protein kinase CDPK (U90262).                  |
| BAB12687.1 | AP002746 | Oryza sativa            | putative protein kinase. P0671B11.2. contains ESTs C22394(C30013),C22393(C30013).  |
| AAB62693.1 | AF004947 | Oryza sativa            | protein kinase.  |
| CAA71142.1 | Y10036   | Cucumis sativus         | SNF1-related protein kinase.   |
| BAA05649.1 | D26602   | Nicotiana tabacum       | protein kinase.  |
| AAC25423.1 | AF072908 | Nicotiana tabacum       | calcium-dependent-protein kinase. CDPK1.   |

|                |          |   |
|----------------|----------|---|
| AAF19403.1     | AF203481 | <i>Lycopersicon esculentum</i><br>phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.    |
| AAF21062.1     | AF216527 | <i>Dunaliella tertiolecta</i><br>calcium-dependent protein kinase. CPK1; CDPK.  |
| AAF19402.1     | AF203480 | <i>Lycopersicon esculentum</i><br>phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.    |
| CAA65244.1     | X95997   | <i>Solanum tuberosum</i><br>SNF1-related protein kinase. PKIN1.   |
| CAA57898.1     | X82548   | <i>Hordeum vulgare</i><br>SNF1-related protein kinase. BKin2.   |
| BAA13608.1     | D88399   | <i>Oryza sativa</i><br>serine-threonine kinase. endosperm kinase. REK.  |
| CAA08995.1     | AJ010091 | <i>Brassica napus</i><br>MAP3K alpha 1 protein kinase. MAP3K alpha 1.   |
| AAF19401.1     | AF203479 | <i>Glycine max</i><br>phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                |
| BAA05648.1     | D26601   | <i>Nicotiana tabacum</i><br>protein kinase.   |
| AAC69450.1     | AF032465 | <i>Nicotiana tabacum</i><br>putative serine/threonine protein kinase. WAPK.   |
| AAD23582.1     | AF128443 | <i>Glycine max</i><br>probably involved in plant stress responses possibly regulates gene expression. SNF-1-like serine/threonine protein kinase. expressed in nodules, roots and leaves. |
| AAD00239.1     | U73938   | <i>Nicotiana tabacum</i><br>protein kinase. PK11-C1. PK11-C1. induced at the transcriptional level by the abscisic acid plant hormone; similar to serine/threonine protein kinase.        |
| AAG60195.1     | AC084763 | <i>Oryza sativa</i><br>protein kinase REK. OSJNBa0027P10.6.   |
| SEQ ID NO: 428 |          |   |
| CAC17753.1     | AJ294543 | <i>Dendrobium 'Sonia'</i><br>metabolic enzyme of cytokinins. cytokinin oxidase. cko1.   |
| CAC17752.1     | AJ294542 | <i>Dendrobium 'Sonia'</i><br>metabolic enzyme of cytokinins. cytokinin oxidase. cko1.   |
| CAA77151.1     | Y18377   | <i>Zea mays</i><br>cytokinin oxidase. cko.  |
| AAC27500.1     | AF044603 | <i>Zea mays</i><br>cytokinin oxidase. ckl1.   |
| BAB07927.1     | AP002836 | <i>Oryza sativa</i><br>putative cytokinin oxidase. P0512G09.9.  |



|  |          |                   |
|--|----------|-------------------|
| BAB03420.1   | AP002816 | Oryza sativa      |
| Similar to Zea mays mRNA for cytokinin oxidase. (Y18377).              |          |                   |
| SEQ ID NO: 429   |          |                   |
| BAA21922.1   | AB006600 | Petunia x hybrida |
| ZPT2-13. C2H2 zinc finger protein, 2finger.                            |          |                   |
| BAA21923.1   | AB006601 | Petunia x hybrida |
| ZPT2-14. C2H2 zinc finger protein, 2 finger.                           |          |                   |
| BAA21921.1   | AB006599 | Petunia x hybrida |
| ZPT2-12. C2H2 zinc finger protein, 2 finger.                           |          |                   |
| BAA19110.1   | AB000451 | Petunia x hybrida |
| PETHy;ZPT2-5. Cys(2) His(2) zinc finger protein, 2 fingers.            |          |                   |
| BAA21925.1   | AB006603 | Petunia x hybrida |
| ZPT2-8. C2H2 zinc finger protein, 2 finger.                            |          |                   |
| BAA21926.1   | AB006604 | Petunia x hybrida |
| ZPT2-9. C2H2 zinc finger protein, 2 finger.                            |          |                   |
| BAA21924.1   | AB006602 | Petunia x hybrida |
| ZPT2-7. C2H2 zinc finger protein, 2finger.                             |          |                   |
| BAA19111.1   | AB000452 | Petunia x hybrida |
| PETHy;ZPT2-6. Cys(2) His(2) zinc finger protein, 2 fingers.            |          |                   |
| BAA05077.1   | D26084   | Petunia x hybrida |
| zinc-finger DNA binding protein.                                       |          |                   |
| AAD26942.1   | AF119050 | Datisca glomerata |
| zinc-finger protein 1. zfp1. DgZFP1.                                   |          |                   |
| BAA05076.1   | D26083   | Petunia x hybrida |
| zinc-finger DNA binding protein.                                       |          |                   |
| CAB77055.1   | Y18788   | Medicago sativa   |
| putative TFIIIA (or kruppel)-like zinc finger protein.                 |          |                   |
| AAC06243.1   | AF053077 | Nicotiana tabacum |
| transcription factor. osmotic stress-induced zinc-finger protein. zfp. |          |                   |
| BAA21927.1   | AB006605 | Petunia x hybrida |
| ZPT3-3. C2H2 zinc finger protein, 3 finger.                            |          |                   |
| BAA96071.1   | AB035133 | Petunia x hybrida |
| C2H2 zinc-finger protein ZPT3-3. ZPT3-3.                               |          |                   |
| BAA21920.1   | AB006598 | Petunia x hybrida |
| ZPT2-11. C2H2 zinc finger protein, 2finger.                            |          |                   |
| AAB53261.1   | U76555   | Brassica rapa     |
| zinc-finger protein BcZFP1. BcZFP1(3-2z).                              |          |                   |
| AAB53260.1   | U76554   | Brassica rapa     |
| transcription factor. zinc-finger protein-1. BR42.                     |          |                   |
| CAA60828.1   | X87374   | Pisum sativum     |
| putative zinc finger protein.  |          |                   |

AAB39638.1 U68763 Glycine max  
putative transcription factor. SCOF-1. scof-1. zinc-finger protein.

AAK01713.1 AF332876 Oryza sativa  
zinc finger transcription factor ZF1.

BAA05079.1 D26086 Petunia x hybrida  
zinc-finger protein.

BAA96070.1 AB035132 Petunia x hybrida  
C2H2 zinc-finger protein ZPT2-10. PETHy;ZPT2-10.

BAA21919.1 AB006597 Petunia x hybrida  
ZPT2-10. C2H2 zinc finger protein, 2 finger.

BAA05078.1 D26085 Petunia x hybrida  
zinc-finger DNA binding protein.

BAA19112.1 AB000453 Petunia x hybrida  
PETHy;ZPT3-1: Cys(2) His(2) zinc finger protein, 3 fingers.

BAA19114.1 AB000455 Petunia x hybrida  
PETHy;ZPT4-1. Cys(2) His(2) zinc finger protein, 4 fingers.

BAA21928.1 AB006606 Petunia x hybrida  
ZPT4-4. C2H2 zinc finger protein, 4 finger.

BAA19926.1 AB000456 Petunia x hybrida  
PETHy; ZPT4-2. C2H2 zinc finger protein, 4 finger.

SEQ ID NO: 432

AAC05084.1 AF033097 Avena sativa  
NPH1-2. NPH1-2. putative serine/threonine protein kinase.

AAC05083.1 AF033096 Avena sativa  
NPH1-1. NPH1-1. putative serine/threonine protein kinase.

CAA82993.1 Z30332 Spinacia oleracea  
protein kinase.

AAB88817.1 AF033263 Zea mays  
signal transduction for phototropism. nonphototropic hypocotyl 1. nph1. NPH1; putative serine/threonine kinase; similar to oat NPH1 proteins.

CAB65325.1 AJ252142 Oryza sativa  
putative blue light receptor phototropin. non-phototropic hypocotyl NPH1. nph1.

CAA82994.1 Z30333 Mesembryanthemum crystallinum  
protein kinase.

BAA36192.1 AB012082 Adiantum capillus-veneris  
PHY3. PHY3. Ser/Thr protein kinase; chimeric structure of red/far-red light photoreceptive region (phytochrome) and putative blue light photoreceptor in phototropism of hypocotyl (NPH1).

SEQ ID NO: 446

AAB72047.1 AF006489 Gossypium hirsutum  
adenine nucleotide translocator 1. CANT1.

CAA05979.1 AJ003197 Lupinus albus  
transfer of ATP from mitochondria to cytosol. adenine nucleotide translocator. ant1.

|                |          |   |
|----------------|----------|---|
| CAA44054.1     | X62123   | <i>Solanum tuberosum</i><br>ADP /ATP translocator. ant. product has dimeric subunit structure.                    |
| AAB49700.1     | U89839   | <i>Lycopersicon esculentum</i><br>ADP/ATP translocator.   |
| CAA40782.1     | X57557   | <i>Solanum tuberosum</i><br>adenine nucleotide translocator. AAC.   |
| BAA02161.1     | D12637   | <i>Oryza sativa</i><br>ATP/ADP translocator.  |
| CAA41812.1     | X59086   | <i>Zea mays</i><br>adenine nucleotide translocator. MANT2.  |
| CAA40781.1     | X57556   | <i>Zea mays</i><br>adenine nucleotide translocator. MANT1.  |
| CAA33743.1     | X15712   | <i>Zea mays</i><br>adenine nucleotide translocator.   |
| CAA33742.1     | X15711   | <i>Zea mays</i><br>adenine nucleotide translocator.   |
| CAA65119.1     | X95863   | <i>Triticum turgidum</i><br>adenine nucleotide translocator.  |
| CAA26600.1     | X02842   | <i>Zea mays</i><br>put. ATP/ADP translocator.   |
| CAA65120.1     | X95864   | <i>Triticum turgidum</i><br>adenine nucleotide translocator.  |
| CAA46311.1     | X65194   | <i>Chlamydomonas reinhardtii</i><br>mitochondrial ADP/ATP translocator protein. CRANT.                            |
| AAA33027.1     | M76669   | <i>Chlorella kessleri</i><br>ATP/ADP translocator. AAT.   |
| AAB72048.1     | AF006490 | <i>Gossypium hirsutum</i><br>adenine nucleotide translocator 2. CANT2.  |
| BAA08104.1     | D45074   | <i>Panicum miliaceum</i><br>2-oxoglutarate/malate translocator. mitochondrial 2-oxoglutarate/malate translocator. |
| BAA08103.1     | D45073   | <i>Panicum miliaceum</i><br>2-oxoglutarate/malate translocator. mitochondrial 2-oxoglutarate/malate translocator. |
| BAA08105.1     | D45075   | <i>Panicum miliaceum</i><br>2-oxoglutarate/malate translocator. mitochondrial 2-oxoglutarate/malate translocator. |
| SEQ ID NO: 447 |          |   |
| BAB16317.1     | AB049589 | <i>Avicennia marina</i><br>secretory peroxidase. PER.   |
| AAC83463.1     | AF039027 | <i>Glycine max</i><br>H2O2 oxidoreductase. cationic peroxidase 2. Prx2. class III plant peroxidase.               |
| AAD37374.1     | AF145348 | <i>Glycine max</i><br>peroxidase. Prx2b.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| CAB71128.2 | AJ271660 | Cicer arietinum         | cationic peroxidase.   |
| AAD33072.1 | AF149251 | Nicotiana tabacum       | secretory peroxidase. PER.   |
| AAA99868.1 | L08199   | Gossypium hirsutum      | peroxidase. putative.  |
| BAA94962.1 | AB042103 | Asparagus officinalis   | peroxidase. AspPOX1.   |
| AAF63027.1 | AF244924 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.   |
| AAF63026.1 | AF244923 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx14 precursor. type III peroxidase.   |
| BAA92500.1 | AP001383 | Oryza sativa            | ESTs D39300(R3292),AU030751(E60187) correspond to a region of the predicted gene. Similar to peroxidase ATP6a. (X98774). |
| AAF63025.1 | AF244922 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx13 precursor. type III peroxidase.   |
| AAG46133.1 | AC082644 | Oryza sativa            | putative peroxidase. OSJNBa0013M12.15.   |
| AAA32973.1 | M73234   | Hordeum vulgare         | peroxidase BP 1. Prx5.   |
| CAA71493.1 | Y10467   | Spinacia oleracea       | peroxidase. prxr6.   |
| AAG46122.1 | AC082644 | Oryza sativa            | putative peroxidase. OSJNBa0013M12.2.  |
| AAG46141.1 | AC082644 | Oryza sativa            | putative peroxidase. OSJNBa0013M12.11.   |
| BAA77387.1 | AB024437 | Scutellaria baicalensis | peroxidase 1.  |
| AAG46142.1 | AC082644 | Oryza sativa            | putative peroxidase. OSJNBa0013M12.9.  |
| AAB02926.1 | U59284   | Linum usitatissimum     | peroxidase. FLXPER3.   |
| CAA50597.1 | X71593   | Lycopersicon esculentum | peroxidase. CEVI-1.  |
| BAA82306.1 | AB027752 | Nicotiana tabacum       | peroxidase.  |
| CAB67121.1 | Y19023   | Lycopersicon esculentum | peroxidase. cevi-1.  |
| AAG46125.1 | AC082644 | Oryza sativa            | putative peroxidase. OSJNBa0013M12.17.   |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| BAA92422.1 | AP001366 | Oryza sativa            | ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene. Similar to A.thaliana mRNA for peroxidase ATP18a. (X98804). |
| BAA92497.1 | AP001383 | Oryza sativa            | ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene. Similar to peroxidase ATP18a. (X98804).                     |
| CAA59487.1 | X85230   | Triticum aestivum       | peroxidase. pox4.  |
| AAA98491.1 | L36981   | Petroselinum crispum    | anionic peroxidase.  |
| CAA71492.1 | Y10466   | Spinacia oleracea       | peroxidase. prxr5.   |
| BAA96643.1 | AP002482 | Oryza sativa            | Similar to Arabidopsis thaliana peroxidase ATP19a (X98805).  |
| BAB12025.1 | AP002820 | Oryza sativa            | putative peroxidase. P0702D12.1.   |
| AAB02554.1 | L37790   | Stylosanthes humilis    | cationic peroxidase.   |
| AAF63024.1 | AF244921 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx12 precursor. type III peroxidase.   |
| AAC84140.1 | AF101427 | Cichorium intybus       | peroxidase.  |
| AAA65637.1 | L13654   | Lycopersicon esculentum | peroxidase. TPX1.  |
| AAC98519.1 | AF007211 | Glycine max             | peroxidase precursor. GMIPER1. pathogen-induced.   |
| BAA14143.1 | D90115   | Armoracia rusticana     | peroxidase isozyme.  |
| CAA62226.1 | X90693   | Medicago sativa         | peroxidase1B. prx1B.   |
| AAD37430.1 | AF149280 | Phaseolus vulgaris      | peroxidase 5 precursor. FBP5. secretory peroxidase.  |
| CAA62615.1 | X91232   | Mercurialis annua       | PRX. peroxidase.   |
| AAG46130.1 | AC082644 | Oryza sativa            | putative peroxidase. OSJNBa0013M12.18.   |
| CAC21392.1 | AJ401275 | Zea mays                | peroxidase. pox2.  |
| BAA07241.1 | D38051   | Populus kitakamiensis   | peroxidase. prxA4a.  |
| BAB39277.1 | AP002971 | Oryza sativa            | putative peroxidase. P0537A05.6. contains ESTs C23550(C52903),C97179(C52903).  |

|                |          |                           |  |
|----------------|----------|---------------------------|--|
| CAB65334.1     | AJ250121 | Picea abies               | peroxidase. SPI2 protein. spi2.  |
| CAA62597.1     | X91172   | Raphanus sativus          | korean-radish isoperoxidase. prxk1.  |
| AAC49820.1     | AF014469 | Oryza sativa              | peroxidase. POX5.1. wound inducible.   |
| CAA71496.1     | Y10470   | Spinacia oleracea         | peroxidase. prxr9.   |
| BAA01950.1     | D11337   | Vigna angularis           | peroxidase.  |
| CAA71491.1     | Y10465   | Spinacia oleracea         | peroxidase. prxr4.   |
| BAA03911.1     | D16442   | Oryza sativa              | peroxidase.  |
| SEQ ID NO: 449 |          |                           |  |
| CAA56325.1     | X80023   | Triticum turgidum         | ATP/ADP carrier protein.   |
| CAC27140.1     | AJ132535 | Picea abies               | ADP, ATP carrier protein precursor.  |
| BAA92520.1     | AP001383 | Oryza sativa              | ESTs AU068633(C30614),AU068634(C30614) correspond to a region of the predicted gene. Similar to Bos taurus mitochondrial solute carrier protein. (AF049236). |
| BAB16462.1     | AP002483 | Oryza sativa              | putative peroxisomal Ca-dependent solute carrier protein. P0019D06.21.   |
| BAB40117.1     | AP003311 | Oryza sativa              | putative peroxisomal Ca-dependent solute carrier protein. P0024G09.9.  |
| AAB71744.1     | U75346   | Chlamydomonas reinhardtii | envelope protein. LIP-36G2. low CO2 inducible carrier protein LIP-36 with a molecular weight of 36 kDa.  |
| AAB71743.1     | U75345   | Chlamydomonas reinhardtii | envelope protein. LIP-36G1. low CO2 inducible carrier protein LIP-36 with a molecular weight of 36 kDa.  |
| AAG48999.1     | AY013246 | Hordeum vulgare           | putative mitochondrial carrier protein. 635P2.1.   |
| CAC12820.1     | AJ299250 | Nicotiana tabacum         | mitochondrial 2-oxoglutarate/malate carrier protein. momc1.  |
| CAA07568.1     | AJ007580 | Ribes nigrum              | Mitochondrial carrier protein. prib7.  |
| AAG45489.1     | AY013245 | Oryza sativa              | 36I5.1. putative mitochondrial carrier protein.  |
| SEQ ID NO: 451 |          |                           |  |
| AAB53099.1     | U68217   | Brassica napus            | iron binding protein. ferritin. LSC30.   |

|            |          |   |
|------------|----------|---|
| AAA33959.1 | M64337   | Glycine max<br>ferritin light chain. ferritin.  |
| AAA34016.1 | M72894   | Glycine max<br>ferritin light chain. SOF-H2.  |
| CAA65771.1 | X97059   | Medicago sativa<br>iron storage. ferritin. FER. abscissic acid regulated.                         |
| AAB18928.1 | U31648   | Glycine max<br>iron storage protein. ferritin.  |
| AAC06026.1 | AF052057 | Vigna unguiculata<br>iron storage and mobilization in plants. ferritin subunit cowpea3 precursor. |
| AAC06027.1 | AF052058 | Vigna unguiculata<br>iron storage and mobilization in plants. ferritin subunit cowpea2 precursor. |
| CAA41213.1 | X58274   | Phaseolus vulgaris<br>ferritin. pfe.  |
| AAD50644.1 | AF133814 | Solanum tuberosum<br>ferritin 1. F1.  |
| CAA51786.1 | X73369   | Pisum sativum<br>ferritin.  |
| CAA45763.1 | X64417   | Pisum sativum<br>ferritin-precursor.  |
| CAA43663.1 | X61391   | Zea mays<br>ferritin.   |
| CAA58146.1 | X83076   | Zea mays<br>ferritin. Fer1.   |
| CAA58147.1 | X83077   | Zea mays<br>ferritin. Fer2.   |
| CAA43664.1 | X61392   | Zea mays<br>ferritin.   |
| AAA33958.1 | M58336   | Glycine max<br>ferritin light chain. SOF-5L.  |
| CAB42587.1 | AJ238628 | Chlorella protothecoides<br>putative ferritin. dee188.  |
| BAB17852.1 | AB042612 | Nicotiana tabacum<br>ferritin 1. tob-fer-1. putative.   |
| AAC15241.1 | AF028072 | Pinus taeda<br>ferritin.  |
| CAA47982.1 | X67754   | Vigna unguiculata<br>ferritin 1. pfe1.  |
| CAA47984.1 | X67756   | Vigna unguiculata<br>ferritin 5. pfe5.  |
| CAA47983.1 | X67755   | Vigna unguiculata<br>ferritin 2. pfe2.  |

|  |          |                              |
|--|----------|------------------------------|
| AAC12282.1   | AF052511 | Glycine max                  |
| iron storage and mobilization. ferritin 2. soybean 2; cowpea 2 homolog.                      |          |                              |
| AAC12281.1   | AF052513 | Glycine max                  |
| iron storage and mobilization. ferritin 1. soybean 1; cowpea 1 homolog.                      |          |                              |
| SEQ ID NO: 452   |          |                              |
| CAB56619.1   | AJ244024 | Nicotiana tabacum            |
| putative role in cell plate formation. phragmoplastin. phragmoplastin. dynamin-like protein. |          |                              |
| SEQ ID NO: 454   |          |                              |
| CAA55039.1   | X78203   | Hyoscyamus muticus           |
| glutathione transferase.   |          |                              |
| AAB65163.1   | AF002692 | Solanum commersonii          |
| glutathione S-transferase, class-phi. GST1. low temperature induced.                         |          |                              |
| BAA01394.1   | D10524   | Nicotiana tabacum            |
| glutathione S-transferase. parB.   |          |                              |
| CAA96431.1   | Z71749   | Nicotiana plumbaginifolia    |
| glutathione S-transferase.   |          |                              |
| AAA33931.1   | M84969   | Silene vulgaris              |
| glutathione-S-transferase.   |          |                              |
| AAA33930.1   | M84968   | Silene vulgaris              |
| glutathione-S-transferase.   |          |                              |
| AAF65767.1   | AF242309 | Euphorbia esula              |
| glutathione S-transferase. putative auxin-binding GST.                                       |          |                              |
| AAF61392.1   | AF133894 | Persea americana             |
| glutathione S-transferase. GTH.  |          |                              |
| CAB38119.1   | AJ010296 | Zea mays                     |
| Glutathione transferase III(b). gst3b.   |          |                              |
| CAB38118.1   | AJ010295 | Zea mays                     |
| Glutathione transferase III(a). gst3a.   |          |                              |
| BAB39935.1   | AP002914 | Oryza sativa                 |
| putative glutathione S-transferase. P0493G01.17.   |          |                              |
| AAG34811.1   | AF243376 | Glycine max                  |
| glutathione S-transferase GST 21.  |          |                              |
| BAB39939.1   | AP002914 | Oryza sativa                 |
| putative glutathione S-transferase. P0493G01.21.   |          |                              |
| AAG34812.1   | AF243377 | Glycine max                  |
| glutathione S-transferase GST 22.  |          |                              |
| AAG34814.1   | AF243379 | Glycine max                  |
| glutathione S-transferase GST 24.  |          |                              |
| CAA09190.1   | AJ010451 | Alopecurus myosuroides       |
| glutathione transferase. GST2a.  |          |                              |
| AAG32476.1   | AF309383 | Oryza sativa subsp. japonica |
| putative glutathione S-transferase OsGSTF4.  |          |                              |



|            |          |                              |   |
|------------|----------|------------------------------|---|
| BAB39929.1 | AP002914 | Oryza sativa                 | putative glutathione transferase. P0493G01.7.   |
| CAA09193.1 | AJ010454 | Alopecurus myosuroides       | glutathione transferase. GST2d.   |
| BAB39941.1 | AP002914 | Oryza sativa                 | putative glutathione S-transferase. P0493G01.23.  |
| CAA09191.1 | AJ010452 | Alopecurus myosuroides       | glutathione transferase. GST2b.   |
| CAA09192.1 | AJ010453 | Alopecurus myosuroides       | glutathione transferase. GST2c.   |
| AAD56395.1 | AF184059 | Triticum aestivum            | glutathione S-transferase. GST1.  |
| CAA39487.1 | X56012   | Triticum aestivum            | glutathione transferase. gstA1.   |
| AAG32477.1 | AF309384 | Oryza sativa subsp. japonica | putative glutathione S-transferase OsGSTF3.   |
| CAA68993.1 | Y07721   | Petunia x hybrida            | conjugates glutathione to anthocyanin to facilitate transport to the vacuole. glutathione S-transferase. an9 locus.           |
| AAA33470.1 | M16901   | Zea mays                     | glutathione S-transferase I.  |
| AAA33469.1 | M16902   | Zea mays                     | glutathione S-transferase I.  |
| AAA20585.1 | U12679   | Zea mays                     | glutathione S-transferase IV. GSTIV.  |
| CAA56047.1 | X79515   | Zea mays                     | glutathione transferase. GST27.   |
| CAB66333.1 | AJ279691 | Betula pendula               | glutathione-S-transferase. gst.   |
| BAB39927.1 | AP002914 | Oryza sativa                 | putative glutathione S-transferase. P0493G01.1. contains ESTs AU031696(R0596), C97559(C60386), C28218(C60386), D28287(R0596). |
| AAC64007.1 | AF062403 | Oryza sativa                 | glutathione S-transferase II.   |
| CAA39480.1 | X56004   | Triticum aestivum            | glutathione transferase. gstA2.   |
| AAG32475.1 | AF309382 | Oryza sativa subsp. japonica | putative glutathione S-transferase OsGSTF5.   |
| AAG34823.1 | AF244680 | Zea mays                     | glutathione S-transferase GST 15.   |
| BAB39940.1 | AP002914 | Oryza sativa                 | putative glutathione S-transferase. P0493G01.22.  |

|                |          |                     |   |
|----------------|----------|---------------------|---|
| AAG34818.1     | AF244675 | Zea mays            | glutathione S-transferase GST 10.   |
| AAG34820.1     | AF244677 | Zea mays            | glutathione S-transferase GST 11.   |
| AAG34817.1     | AF244674 | Zea mays            | glutathione S-transferase GST 9.  |
| AAG34821.1     | AF244678 | Zea mays            | glutathione S-transferase GST 13.   |
| CAA05354.1     | AJ002380 | Oryza sativa        | glutathione S-transferase. Rgst I.  |
| AAG34816.1     | AF244673 | Zea mays            | glutathione S-transferase GST 8.  |
| AAG34822.1     | AF244679 | Zea mays            | glutathione S-transferase GST 14.   |
| CAA05355.1     | AJ002381 | Oryza sativa        | glutathione S-transferase. Rgst II.   |
| SEQ ID NO: 458 |          |                     |   |
| AAD45623.1     | AF084185 | Brassica napus      | dehydration responsive element binding protein. DNA binding protein; DRE binding protein.   |
| BAA90812.1     | AP001168 | Oryza sativa        | Similar to mRNA for DREB1A (AB007787).  |
| AAK01089.1     | AF298231 | Hordeum vulgare     | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.   |
| AAG59618.1     | AF239616 | Hordeum vulgare     | CRT/DRE-binding factor. CBF.  |
| AAK01088.1     | AF298230 | Hordeum vulgare     | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.  |
| AAG59619.1     | AF243384 | Oryza sativa        | CRT/DRE binding factor. CBF. DREB.  |
| AAC29516.1     | U77655   | Solanum tuberosum   | DNA binding protein homolog. STWAAEIRD.   |
| BAA76734.1     | AB024575 | Nicotiana tabacum   | ethylene responsive element binding factor.   |
| CAB93939.1     | AJ238739 | Catharanthus roseus | putative transcription factor. AP2-domain DNA-binding protein. orca1.   |
| BAB03248.1     | AB037183 | Oryza sativa        | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.   |
| BAA97123.1     | AB016265 | Nicotiana glauca    | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3. |
| BAA97124.1     | AB016266 | Nicotiana glauca    | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4. |

|                |          |                                |  |
|----------------|----------|--------------------------------|--|
| AAC50047.1     | U89255   | <i>Lycopersicon esculentum</i> | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti4.   |
| AAD00708.1     | U91857   | <i>Stylosanthes hamata</i>     | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively. |
| AAC14323.1     | AF058827 | <i>Nicotiana tabacum</i>       | TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.   |
| AAG60182.1     | AC084763 | <i>Oryza sativa</i>            | putative ethylene-responsive element binding protein. OSJNBa0027P10.12.  |
| AAK31279.1     | AC079890 | <i>Oryza sativa</i>            | putative ethylene-responsive element binding protein. OSJNBb0089A17.16.  |
| AAC49741.1     | U89257   | <i>Lycopersicon esculentum</i> | DNA-binding protein binds the GCC box pathogenesis-related promoter element. Pti6.   |
| BAA97122.1     | AB016264 | <i>Nicotiana sylvestris</i>    | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2.                |
| CAB96900.1     | AJ251250 | <i>Catharanthus roseus</i>     | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| CAB96899.1     | AJ251249 | <i>Catharanthus roseus</i>     | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| BAB21218.1     | AP002913 | <i>Oryza sativa</i>            | putative DNA binding protein RAV2. P0480E02.24.  |
| CAB93940.1     | AJ238740 | <i>Catharanthus roseus</i>     | putative transcription factor. AP2-domain DNA-binding protein. orca2.  |
| BAA87068.1     | AB035270 | <i>Matricaria chamomilla</i>   | ethylene-responsive element binding protein1 homolog. McEREBP1.  |
| AAC49740.1     | U89256   | <i>Lycopersicon esculentum</i> | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti5.   |
| AAB38748.1     | U81157   | <i>Nicotiana tabacum</i>       | S25-XP1 DNA binding protein.   |
| BAB21211.1     | AP002913 | <i>Oryza sativa</i>            | putative DNA binding protein RAV2. P0480E02.17.  |
| BAA78732.1     | AB023482 | <i>Oryza sativa</i>            | Similar to <i>Arabidopsis thaliana</i> mRNA for DREB1B, complete cds.(AB007788).   |
| SEQ ID NO: 460 |          |                                |  |
| AAB67721.1     | AF015269 | <i>Zea mays</i>                | PL transcription factor. Pl. nonfunctional due to retrotransposon insertion.   |
| SEQ ID NO: 461 |          |                                |  |
| AAD50592.1     | AF093752 | <i>Triticum aestivum</i>       | phytochelatase synthase. PCS1.   |
| AAG22095.1     | AF308658 | <i>Typha latifolia</i>         | phytochelatase synthase.   |

## SEQ ID NO: 463

AAG13629.1 AC078840 *Oryza sativa*  
putative steroid membrane binding protein. OSJNBb0073N24.4.

AAG13623.1 AC078840 *Oryza sativa*  
putative steroid membrane binding protein. OSJNBb0073N24.2.

## SEQ ID NO: 464

CAA56313.1 X79992 *Avena sativa*  
putative pp70 ribosomal protein S6 kinase. Aspk11.

CAB89082.1 AJ277534 *Asparagus officinalis*  
S6 ribosomal protein kinase. pk1. putative.

AAK18843.1 AC082645 *Oryza sativa*  
putative protein kinase. OSJNBb0033N16.3.

AAC05083.1 AF033096 *Avena sativa*  
NPH1-1. NPH1-1. putative serine/threonine protein kinase.

AAC05084.1 AF033097 *Avena sativa*  
NPH1-2. NPH1-2. putative serine/threonine protein kinase.

AAK13156.1 AC078829 *Oryza sativa*  
putative protein kinase. OSJNBa0026O12.14.

CAB82852.1 Z30329 *Mesembryanthemum crystallinum*  
protein kinase MK6.

AAB88817.1 AF033263 *Zea mays*  
signal transduction for phototropism. nonphototropic hypocotyl 1. nph1. NPH1; putative serine/threonine kinase; similar to oat NPH1 proteins.

BAB18105.1 AB042715 *Chlamydomonas reinhardtii*  
cyclic nucleotide dependent protein kinase II. CL-PK2.

BAB18104.1 AB042714 *Chlamydomonas reinhardtii*  
cyclic nucleotide dependent protein kinase. CL-PK1.

BAA83689.1 AB011968 *Oryza sativa*  
OsPK7. OsPK7. protein kinase.

BAA92970.1 AP001551 *Oryza sativa*  
Similar to *Arabidopsis thaliana* chromosome 4 BAC clone F28A21 ; putative protein kinase. (AL035526).

AAD31900.1 AF145482 *Mesembryanthemum crystallinum*  
putative serine/threonine protein kinase.

BAA92972.1 AP001551 *Oryza sativa*  
ESTs AU056183(S20356), AU056881(S20950) correspond to a region of the predicted gene. Similar to *Arabidopsis thaliana* chromosome 4 BAC clone F6I18 ; putative protein kinase. (AL022198).

BAA83688.1 AB011967 *Oryza sativa*  
OsPK4. OsPK4. protein kinase.

CAA73067.1 Y12464 *Sorghum bicolor*  
serine/threonine kinase. SNFL1.

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAF22219.1 | AF141378 | Zea mays                | protein kinase PK4. ZmPK4.  |
| BAA96628.1 | AP002482 | Oryza sativa            | ESTs D41739(S4522),AU055999(S20214), AU057588(S21592 correspond to a region of the predicted gene. Similar to Sorghum bicolor serine/threonine kinase (Y12465).         |
| CAA89202.1 | Z49233   | Chlamydomonas eugametos | calcium-stimulated protein kinase.  |
| CAA73068.1 | Y12465   | Sorghum bicolor         | serine/threonine kinase. SNFL2.   |
| BAA34675.1 | AB011670 | Triticum aestivum       | wpk4 protein kinase. wpk4.  |
| BAA99439.1 | AP002743 | Oryza sativa            | putative protein kinase. P0710E05.26. contains ESTs C22394(C30013),C22393(C30013).  |
| CAA74646.1 | Y14274   | Sorghum bicolor         | putative serine/threonine protein kinase. SNFL3.  |
| BAB12687.1 | AP002746 | Oryza sativa            | putative protein kinase. P0671B11.2. contains ESTs C22394(C30013),C22393(C30013).   |
| AAF06969.1 | AF162661 | Kalanchoe fedtschenkoi  | phosphoenolpyruvate carboxylase kinase. calcium-independent protein kinase.   |
| AAF06970.1 | AF162662 | Kalanchoe fedtschenkoi  | phosphoenolpyruvate carboxylase kinase. calcium-independent protein kinase.   |
| BAA90814.1 | AP001168 | Oryza sativa            | ESTs AU030197(E50746),AU030196(E50746) correspond to a region of the predicted gene.; Similar to calcium-dependent calmodulin-independent protein kinase CDPK (U90262). |
| AAB62693.1 | AF004947 | Oryza sativa            | protein kinase.   |
| AAF21062.1 | AF216527 | Dunaliella tertiolecta  | calcium-dependent protein kinase. CPK1; CDPK.   |
| CAA71142.1 | Y10036   | Cucumis sativus         | SNF1-related protein kinase.  |
| CAA39936.1 | X56599   | Daucus carota           | calcium- dependent protein kinase. DcPK431.   |
| CAA08995.1 | AJ010091 | Brassica napus          | MAP3K alpha 1 protein kinase. MAP3K alpha 1.  |
| BAA05649.1 | D26602   | Nicotiana tabacum       | protein kinase.   |
| AAF19403.1 | AF203481 | Lycopersicon esculentum | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                    |
| AAC25423.1 | AF072908 | Nicotiana tabacum       | calcium-dependent protein kinase. CDPK1.  |

|                |          |                                 |   |
|----------------|----------|---------------------------------|---|
| AAF19402.1     | AF203480 | <i>Lycopersicon esculentum</i>  | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                |
| CAA65244.1     | X95997   | <i>Solanum tuberosum</i>        | SNF1-related protein kinase. PKIN1.   |
| CAA57898.1     | X82548   | <i>Hordeum vulgare</i>          | SNF1-related protein kinase. BKIN2.   |
| AAF19401.1     | AF203479 | <i>Glycine max</i>              | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                |
| AAD23582.1     | AF128443 | <i>Glycine max</i>              | probably involved in plant stress responses possibly regulates gene expression. SNF-1-like serine/threonine protein kinase. expressed in nodules, roots and leaves. |
| AAF34436.1     | AF172282 | <i>Oryza sativa</i>             | similar to mitogen-activated protein kinases. DUPR11.32.  |
| BAA05648.1     | D26601   | <i>Nicotiana tabacum</i>        | protein kinase.   |
| SEQ ID NO: 465 |          |                                 |   |
| CAB85467.1     | AJ250316 | <i>Brassica juncea</i>          | chloroplast omega-3 fatty acid desaturase. fad7.  |
| BAA22441.1     | D63954   | <i>Zea mays</i>                 | fatty acid desaturase. FAD7.  |
| BAA11475.1     | D79979   | <i>Nicotiana tabacum</i>        | omega-3 fatty acid desaturase. NtFAD7.  |
| AAA70334.1     | U25817   | <i>Sesamum indicum</i>          | omega-3 fatty acid desaturase. linoleic acid and hexadecadienoic acids desaturase.  |
| AAB39387.1     | U59477   | <i>Perilla frutescens</i>       | omega-3 fatty acid desaturase.  |
| CAA07638.1     | AJ007739 | <i>Solanum tuberosum</i>        | w-3 desaturase.   |
| AAF27933.1     | AF222989 | <i>Capsicum annuum</i>          | omega-3 fatty acid desaturase. FAD.   |
| AAB72241.1     | U75745   | <i>Petroselinum crispum</i>     | omega-3 fatty acid desaturase. complements the Arabidopsis fad7/8 fatty acid double mutant.   |
| AAA61776.1     | L22965   | Chloroplast <i>Glycine soja</i> | omega-3 fatty acid desaturase. Fadd.  |
| AAF12821.1     | AF200717 | <i>Vernicia fordii</i>          | TnDES2. omega-3 fatty acid desaturase.  |
| AAA86690.1     | U17063   | <i>Limnanthes douglasii</i>     | linoleic acid desaturation. delta-15 lineoyl desaturase. omega-3-fatty acid desaturase.   |
| AAA73511.1     | L25897   | <i>Ricinus communis</i>         | desaturates linoleate to alpha-linolenate. linoleoyl desaturase. fad7. putative.  |

|            |          |                                   |  |
|------------|----------|-----------------------------------|--|
| CAC18722.1 | AJ302017 | <i>Picea abies</i>                | putative plastidial w-3 fatty acid desaturase. fad3.                             |
| AAD13527.1 | AF061027 | <i>Vernicia fordii</i>            | omega-3 fatty acid desaturase precursor.   |
| BAA22442.1 | D84409   | <i>Zea mays</i>                   | fatty acid desaturase. FAD8.   |
| BAA22440.1 | D63953   | <i>Zea mays</i>                   | fatty acid desaturase. FAD8.   |
| BAA07785.2 | D43688   | <i>Triticum aestivum</i>          | plastid omega-3 fatty acid desaturase. TaFAD7.                                   |
| AAA61774.1 | L22963   | Chloroplast <i>Brassica napus</i> | omega-3 fatty acid desaturase. Fadd.   |
| AAC98967.1 | AF047172 | <i>Vernicia fordii</i>            | omega-3 fatty acid desaturase. Fad3.   |
| CAB45155.1 | AJ011004 | <i>Vernicia fordii</i>            | omega-3 desaturase.  |
| AAC16443.1 | AF020204 | <i>Pelargonium x hortorum</i>     | omega-3 desaturase. pxh-15.  |
| AAA61775.1 | L22962   | <i>Brassica napus</i>             | omega-3 fatty acid desaturase. Fad3.   |
| AAA61777.1 | L22964   | Chloroplast <i>Glycine soja</i>   | omega-3 fatty acid desaturase. Fad3.   |
| AAA32994.1 | L01418   | <i>Brassica napus</i>             | desaturation of linoleic acid to linolenic acid. linoleic acid desaturase. fad3. |
| AAD15744.1 | AF047039 | <i>Perilla frutescens</i>         | omega-3 fatty acid desaturase. FAD3.   |
| BAA28358.1 | D84678   | <i>Triticum aestivum</i>          | omega-3 fatty acid desaturase. TaFAD3.   |
| BAA05515.1 | D26509   | <i>Nicotiana tabacum</i>          | desaturation of dienoic fatty acid. microsomal omega-3 acid desaturase. NtFAD3.  |
| BAA11397.1 | D78506   | <i>Oryza sativa</i>               | w-3 fatty acid desaturase.   |
| BAB18135.1 | AB051215 | <i>Glycine max</i>                | microsomal omega-3 fatty acid desaturase.  |
| BAA22439.1 | D63952   | <i>Zea mays</i>                   | fatty acid desaturase. FAD7.   |
| BAA11396.1 | D78505   | <i>Oryza sativa</i>               | w-3 fatty acid desaturase.   |
| CAB71341.1 | AJ250664 | <i>Hordeum vulgare</i>            | omega-3 fatty acid desaturase. bci-8.  |
| AAD48897.1 | AF083613 | <i>Dunaliella salina</i>          | omega-3 fatty acid desaturase. des3-1.   |

|                |          |                                       |  |
|----------------|----------|---------------------------------------|--|
| AAF80560.1     | AF192486 | Sesamum indicum                       | omega-6 fatty acid desaturase. FAD2.   |
| CAB64256.1     | AJ245938 | Calendula officinalis                 | production of calendic acid. (8,11)-linoleoyl desaturase. des8.11.   |
| AAB80696.1     | U86072   | Petroselinum crispum                  | omega-6 fatty acid desaturase. functional expression in yeast results in the formation of dienoic fatty acids (18:2 and 16:2).   |
| SEQ ID NO: 467 |          |                                       |  |
| CAA60016.1     | X86021   | Solanum tuberosum                     | potassium channel. SKT1 gene. putative start codon.  |
| CAA65254.1     | X96390   | Lycopersicon esculentum               | potassium channel. LKT1.   |
| AAF36832.1     | AF207745 | Triticum aestivum                     | AKT1-like potassium channel. TaAKT1.   |
| CAA68912.1     | Y07632   | Zea mays                              | potassium channel. ZMK1.   |
| AAF81249.1     | AF267753 | Mesembryanthemum crystallinum         | putative potassium channel protein Mkt1p.  |
| CAC05489.1     | AJ271447 | Populus tremula x Populus tremuloides | potassium channel. potassium channel 2. ptk2.  |
| AAD16278.1     | AF099095 | Samanea saman                         | pulvinus inward-rectifying channel for potassium SPICK1. similar to Arabidopsis potassium channel AKT3.  |
| CAB54856.1     | AJ132686 | Zea mays                              | potassium channel protein ZMK2. ZMK2.  |
| CAA70870.1     | Y09699   | Solanum tuberosum                     | putative inward rectifying potassium channel. SKT2.  |
| CAA71598.1     | Y10579   | Vicia faba                            | potassium channel.   |
| BAA84085.1     | AB032074 | Nicotiana paniculata                  | potassium channel. NpKT1.  |
| CAA12645.1     | AJ225805 | Egeria densa                          | inward potassium channel alpha subunit. homologous to the sequences of the family of inwardly rectifying potassium channels in plants which is structurally related to the shaker family of outwardly rectifying channels in Drosophila. |
| AAD39492.1     | AF145272 | Samanea saman                         | pulvinus inward-rectifying channel SPICK2. potassium channel; similar to AKT2/3.   |
| CAC10514.1     | AJ299019 | Samanea saman                         | potassium release. outwardly rectifying potassium channel. spork1.   |
| CAC05488.1     | AJ271446 | Populus tremula x Populus tremuloides | potassium channel. outward rectifying potassium channel. ptork.  |
| AAF81251.1     | AF267755 | Mesembryanthemum crystallinum         | potassium channel protein Mkt2p.   |



|                |          |                               |  |
|----------------|----------|-------------------------------|--|
| CAA70900.1     | Y09753   | Secale cereale                | potassium channel. encodes membrane spanning domains H5-S6.                                |
| CAA70899.1     | Y09752   | Secale cereale                | potassium channel. encodes membrane spanning domains H5-S6.                                |
| CAA70947.1     | Y09818   | Solanum tuberosum             | putative inward rectifying potassium channel. SKT3.  |
| CAA70895.1     | Y09748   | Hordeum vulgare               | potassium channel. encodes membrane spanning domains H5-S6.                                |
| CAA70894.1     | Y09747   | Zea mays                      | potassium channel. encodes membrane spanning domains H5-S6.                                |
| CAA70896.1     | Y09749   | Vicia faba                    | potassium channel. encodes membrane spanning domains H5-S6.                                |
| CAA70897.1     | Y09750   | Plantago major                | potassium channel. encodes membrane spanning domains H5-S6.                                |
| AAF33670.1     | AF079872 | Nicotiana tabacum             | cyclic nucleotide-gated calmodulin-binding ion channel. CBP4.                              |
| AAF33669.1     | AF079871 | Nicotiana tabacum             | cyclic nucleotide-gated calmodulin-binding ion channel. CBP7.                              |
| CAA05637.1     | AJ002610 | Hordeum vulgare               | putative calmodulin binding transporter protein. CBT1.                                     |
| CAA70898.1     | Y09751   | Plantago major                | potassium channel. encodes membrane spanning domains H5-S6.                                |
| AAK16188.1     | AC079887 | Oryza sativa                  | putative cyclic nucleotide and calmodulin-regulated ion channel protein. OSJNBa0040E01.13. |
| SEQ ID NO: 468 |          |                               |  |
| AAF76226.1     | AF272572 | Populus x canescens           | 14-3-3 protein. 14-3-3P20-1.   |
| AAC04811.1     | AF037460 | Fritillaria agrestis          | GF14 protein. GRF.   |
| AAB40395.1     | U80070   | Mesembryanthemum crystallinum | G-box binding factor. 14-3-3-like protein. GBF.  |
| AAF05737.1     | AF191746 | Lilium longiflorum            | 14-3-3-like protein.   |
| AAB07457.1     | U65957   | Oryza sativa                  | GF14-c protein. rice 14-3-3 protein homolog; osGF14c.                                      |
| AAB09580.1     | U70533   | Glycine max                   | SGF14A. 14-3-3 related protein.  |
| AAA99431.1     | L29150   | Lycopersicon esculentum       | 14-3-3 protein homologue.  |
| CAA74592.1     | Y14200   | Hordeum vulgare               | 14-3-3 protein.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAB33304.1 | S77133   | Zea mays                | GF14-6. GRF1. 14-3-3 protein homolog; This sequence comes from Fig. 5. |
| CAA66309.1 | X97724   | Solanum tuberosum       | 14-3-3 protein. leaf specific.   |
| AAB07456.1 | U65956   | Oryza sativa            | GF14-b protein. rice 14-3-3 protein homolog; osGF14b.                  |
| AAD27823.2 | AF121194 | Populus x canescens     | 14-3-3 protein. 14-3-3P20-2.   |
| AAC49894.1 | U91726   | Nicotiana tabacum       | 14-3-3 isoform e. T14-3e.  |
| CAA44259.1 | X62388   | Hordeum vulgare         | 14-3-3 protein homologue.  |
| AAD27827.2 | AF121198 | Picea glauca            | 14-3-3 protein. 14-3-3EB9D.  |
| CAA63658.1 | X93170   | Hordeum vulgare         | Hv14-3-3b.   |
| CAB42546.2 | AJ238681 | Pisum sativum           | 14-3-3-like protein. 14-3-3.   |
| AAA33505.1 | M96856   | Zea mays                | regulatory protein. GF14-12.   |
| AAA85817.1 | U15036   | Pisum sativum           | 14-3-3-like protein.   |
| AAK26634.1 | AF342780 | Brassica napus          | GF14 omega. 14-3-3 protein.  |
| CAA72094.1 | Y11211   | Nicotiana tabacum       | 14-3-3-like protein B.   |
| CAA72383.1 | Y11687   | Solanum tuberosum       | 14-3-3 protein. 34G.   |
| AAC49892.1 | U91724   | Nicotiana tabacum       | 14-3-3 isoform c. T14-3c.  |
| CAA53700.1 | X76086   | Cucurbita pepo          | 14-3-3 protein 32kDa endonuclease. A215. single polypeptide.           |
| CAA72382.1 | Y11686   | Solanum tuberosum       | 14-3-3 protein. 30G.   |
| CAB42547.1 | AJ238682 | Pisum sativum           | 14-3-3-like protein. 14-3-3.   |
| CAA65147.1 | X95902   | Lycopersicon esculentum | 14-3-3 protein. tft3 gene.   |
| AAC49891.1 | U91723   | Nicotiana tabacum       | 14-3-3 isoform b. T14-3b.  |
| AAB07458.1 | U65958   | Oryza sativa            | GF14-d protein. rice 14-3-3 protein homolog; osGF14d.                  |

|                |          |                           |  |
|----------------|----------|---------------------------|--|
| CAA72381.1     | Y11685   | Solanum tuberosum         | 14-3-3 protein. 16R.   |
| AAC49895.1     | U91727   | Nicotiana tabacum         | 14-3-3 isoform f. T14-3f.  |
| CAA72095.1     | Y11212   | Nicotiana tabacum         | 14-3-3-like protein A.   |
| BAB11739.1     | AB042193 | Triticum aestivum         | TaWIN1. TaWIN1. TaWIN1 is a member of 14-3-3 protein family.   |
| CAA65146.1     | X95901   | Lycopersicon esculentum   | 14-3-3 protein. tft2 gene.   |
| CAA65148.1     | X95903   | Lycopersicon esculentum   | 14-3-3 protein. tft5 gene.   |
| AAC17447.1     | AF066076 | Helianthus annuus         | 14-3-3-like protein.   |
| CAA60800.1     | X87370   | Solanum tuberosum         | 14-3-3 protein. RA215. root specific.  |
| CAA55964.1     | X79445   | Chlamydomonas reinhardtii | 14-3-3 protein.  |
| CAC03467.1     | Y19105   | Chlamydomonas reinhardtii | 14-3-3 protein.  |
| CAA65149.1     | X95904   | Lycopersicon esculentum   | 14-3-3 protein. tft6 gene.   |
| CAB65693.1     | AJ270959 | Lycopersicon esculentum   | tft3 14-3-3 protein. tft3.   |
| BAB11740.1     | AB042194 | Triticum aestivum         | TaWIN2. TaWIN2. TaWIN2 is a member of 14-3-3 protein family.   |
| AAA99430.1     | L29151   | Lycopersicon esculentum   | 14-3-3 protein homologue.  |
| CAA72384.1     | Y11688   | Solanum tuberosum         | 14-3-3 protein. 35G.   |
| CAA65145.1     | X95900   | Lycopersicon esculentum   | 14-3-3 protein. tft1 gene.   |
| AAC49893.1     | U91725   | Nicotiana tabacum         | 14-3-3 isoform d. T14-3d.  |
| SEQ ID NO: 470 |          |                           |  |
| AAA19701.1     | L24438   | Thlaspi arvense           | cytochrome P450.   |
| AAA32913.1     | M32885   | Persea americana          | cytochrome P-450LXXIA1 (cyp71A1).  |
| AAC39318.1     | AF029858 | Sorghum bicolor           | second multifunctional cytochrome P450 in the biosynthetic pathway of the cyanogenic glucoside dhurrin. Catalyzes the conversion of p-hydroxyphenylacetaldoxime to p-hydroxymandelonitrile. cytochrome P450 CYP71E1. CYP71E1. No EST#s identified. |

|            |          |   |
|------------|----------|---|
| BAB40323.1 | AB037244 | Asparagus officinalis<br>cytochrome P450. ASPI-1.   |
| BAB40324.1 | AB037245 | Asparagus officinalis<br>cytochrome P450. ASPI-2.   |
| AAB94589.1 | AF022460 | Glycine max<br>CYP83D1p. CYP83D1. cytochrome P450 monooxygenase.  |
| CAA70575.1 | Y09423   | Nepeta racemosa<br>cytochrome P450. CYP71A5.  |
| CAA70576.1 | Y09424   | Nepeta racemosa<br>cytochrome P450. CYP71A6.  |
| CAA71513.1 | Y10489   | Glycine max<br>putative cytochrome P450.  |
| AAB61965.1 | U48435   | Solanum chacoense<br>putative cytochrome P450.  |
| AAB94588.1 | AF022459 | Glycine max<br>CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.  |
| CAA50312.1 | X70981   | Solanum melongena<br>P450 hydroxylase. CYPEG2.  |
| CAA71514.1 | Y10490   | Glycine max<br>putative cytochrome P450.  |
| AAB94584.1 | AF022157 | Glycine max<br>capable of catalyzing the metabolism of phenylurea herbicides. CYP71A10. CYP71A10.<br>cytochrome P450 monooxygenase. |
| AAD47832.1 | AF166332 | Nicotiana tabacum<br>cytochrome P450.   |
| AAF27282.1 | AF122821 | Capsicum annuum<br>cytochrome P450. PepCYP.   |
| CAA71517.1 | Y10493   | Glycine max<br>putative cytochrome P450.  |
| CAA83941.1 | Z33875   | Mentha x piperita<br>cytochrome P-450 oxidase.  |
| CAB56503.1 | AJ238612 | Catharanthus roseus<br>cytochrome P450.   |
| AAB61964.1 | U48434   | Solanum chacoense<br>putative cytochrome P450.  |
| AAB69644.1 | AF000403 | Lotus japonicus<br>putative cytochrome P450. LjNP450.   |
| CAA50645.1 | X71654   | Solanum melongena<br>P450 hydroxylase.  |
| BAA03635.1 | D14990   | Solanum melongena<br>Cytochrome P-450EG4.   |
| AAD44151.1 | AF124816 | Mentha x piperita<br>cytochrome p450 isoform PM17.  |

|                |          |  |
|----------------|----------|--|
| AAD44150.1     | AF124815 | Mentha spicata<br>cytochrome p450.   |
| AAD44152.1     | AF124817 | Mentha x piperita<br>cytochrome p450 isoform PM2.  |
| BAB39252.1     | AP002968 | Oryza sativa<br>putative cytochrome P450. P0416G11.1.  |
| AAK38084.1     | AF321860 | Lolium rigidum<br>putative cytochrome P450.  |
| AAK38083.1     | AF321859 | Lolium rigidum<br>putative cytochrome P450.  |
| AAK38087.1     | AF321863 | Lolium rigidum<br>putative cytochrome P450.  |
| AAK38082.1     | AF321858 | Lolium rigidum<br>putative cytochrome P450.  |
| CAC27827.1     | AJ295719 | Catharanthus roseus<br>geraniol hydroxylase. cytochrome P450. cyp71.   |
| AAB94587.1     | AF022458 | Glycine max<br>CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.   |
| AAK38088.1     | AF321864 | Lolium rigidum<br>putative cytochrome P450.  |
| AAD37433.1     | AF150881 | Lycopersicon esculentum x Lycopersicon peruvianum<br>catalyzes the hydroxylation of ferulic acid to 5-hydroxyferulic acid. ferulate-5-hydroxylase.<br>CYP84. cytochrome P450-dependent monooxygenase; F5H; FAH1. |
| AAG14963.1     | AF214009 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H3.   |
| CAA57425.1     | X81831   | Zea mays<br>cytochrome P450. CYP71C4. family CYP71, subfamily CYP71C.  |
| CAA72196.1     | Y11368   | Zea mays<br>cytochrome p450. cyp71c4.  |
| AAG44132.1     | AF218296 | Pisum sativum<br>cytochrome P450. P450 isolog.   |
| AAG14962.1     | AF214008 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H2.   |
| AAG14961.1     | AF214007 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H1.   |
| AAD56282.1     | AF155332 | Petunia x hybrida<br>flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.  |
| SEQ ID NO: 471 |          |  |
| AAD17855.1     | AF099111 | Zea mays<br>sigma factor. sig2. putative sigma subunit of chloroplast Escherichia coli-like RNA<br>polymerase; Sig2.   |
| SEQ ID NO: 472 |          |  |

|            |          |                      |  |
|------------|----------|----------------------|--|
| AAD45623.1 | AF084185 | Brassica napus       | dehydration responsive element binding protein. DNA binding protein; DRE binding protein.  |
| AAG43549.1 | AF211531 | Nicotiana tabacum    | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.   |
| AAG43548.1 | AF211530 | Nicotiana tabacum    | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.   |
| AAK01089.1 | AF298231 | Hordeum vulgare      | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.  |
| AAG59618.1 | AF239616 | Hordeum vulgare      | CRT/DRE-binding factor. CBF.   |
| AAK01088.1 | AF298230 | Hordeum vulgare      | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.   |
| AAG59619.1 | AF243384 | Oryza sativa         | CRT/DRE binding factor. CBF. DREB.   |
| AAK31271.1 | AC079890 | Oryza sativa         | putative transcriptional factor. OSJNBb0089A17.22.   |
| CAC12822.1 | AJ299252 | Nicotiana tabacum    | AP2 domain-containing transcription factor. ap2.   |
| BAA76734.1 | AB024575 | Nicotiana tabacum    | ethylene responsive element binding factor.  |
| AAC29516.1 | U77655   | Solanum tuberosum    | DNA binding protein homolog. STWAAEIRD.  |
| AAF23899.1 | AF193803 | Oryza sativa         | transcription factor EREBP1. EREBP/AP2-like transcription factor.  |
| CAB93939.1 | AJ238739 | Catharanthus roseus  | putative transcription factor. AP2-domain DNA-binding protein. orca1.  |
| BAB16083.1 | AB036883 | Oryza sativa         | transcriptional repressor. osERF3. osERF3. ERF protein family ERF3 associated repression domain.   |
| BAB03248.1 | AB037183 | Oryza sativa         | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.  |
| BAA97124.1 | AB016266 | Nicotiana sylvestris | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4.                |
| BAA97123.1 | AB016265 | Nicotiana sylvestris | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.                |
| AAG43545.1 | AF211527 | Nicotiana tabacum    | Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.   |
| AAD00708.1 | U91857   | Stylosanthes hamata  | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively. |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAC49741.1     | U89257   | Lycopersicon esculentum       | DNA-binding protein binds the GCC box pathogenesis-related promoter element. Pti6.  |
| AAG60182.1     | AC084763 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBa0027P10.12.   |
| AAK31279.1     | AC079890 | Oryza sativa                  | putative ethylene-responsive element binding protein. OSJNBb0089A17.16.   |
| AAC14323.1     | AF058827 | Nicotiana tabacum             | TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.  |
| AAC50047.1     | U89255   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti4.  |
| BAA97122.1     | AB016264 | Nicotiana sylvestris          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2. |
| CAB96899.1     | AJ251249 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| CAB93940.1     | AJ238740 | Catharanthus roseus           | putative transcription factor. AP2-domain DNA-binding protein. orca2.   |
| AAF63205.1     | AF245119 | Mesembryanthemum crystallinum | AP2-related transcription factor. CDBP. stress induced transcription factor.  |
| BAA87068.1     | AB035270 | Matricaria chamomilla         | ethylene-responsive element binding protein1 homolog. McEREBP1.   |
| CAB96900.1     | AJ251250 | Catharanthus roseus           | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| BAB21218.1     | AP002913 | Oryza sativa                  | putative DNA binding protein RAV2. P0480E02.24.   |
| BAA07321.1     | D38123   | Nicotiana tabacum             | ERF1. ethylene-responsive transcription factor.   |
| AAC49740.1     | U89256   | Lycopersicon esculentum       | binds the GCC box pathogenesis-related promoter element DNA-binding protein. Pti5.  |
| AAB38748.1     | U81157   | Nicotiana tabacum             | S25-XP1 DNA binding protein.  |
| BAB21211.1     | AP002913 | Oryza sativa                  | putative DNA binding protein RAV2. P0480E02.17.   |
| SEQ ID NO: 473 |          |                               |   |
| BAA85438.1     | AP000616 | Oryza sativa                  | similar to RING-H2 finger protein RHA1a (AF078683).   |
| AAK00436.1     | AC060755 | Oryza sativa                  | putative zinc finger protein. OSJNBa0003O19.23.   |
| AAG43550.1     | AF211532 | Nicotiana tabacum             | Avr9/Cf-9 rapidly elicited protein 132. ACRE132. similar to RING finger proteins.   |
| SEQ ID NO: 475 |          |                               |   |
| AAG43550.1     | AF211532 | Nicotiana tabacum             | Avr9/Cf-9 rapidly elicited protein 132. ACRE132. similar to RING finger proteins.   |

|                |          |                      |   |
|----------------|----------|----------------------|---|
| BAA78746.1     | AB023482 | Oryza sativa         | Similar to Arabidopsis thaliana RING-H2 finger protein RHX1a mRNA, partial cds.(AF079184).  |
| BAA96875.1     | AB045121 | Oryza sativa         | RING finger 1. RRF1.  |
| CAA74911.1     | Y14573   | Hordeum vulgare      | ring finger protein. putative.  |
| AAK00436.1     | AC060755 | Oryza sativa         | putative zinc finger protein. OSJNBa0003O19.23.   |
| AAG46117.1     | AC073166 | Oryza sativa         | putative ring finger protein. OSJNBb0064P21.7.  |
| BAA90357.1     | AP001080 | Oryza sativa         | EST AU070319(S10788) corresponds to a region of the predicted gene. Similar to RING-H2 finger protein RHA2b (AC006200).   |
| BAA77204.1     | AB026262 | Cicer arietinum      | ring finger protein.  |
| BAA90806.1     | AP001168 | Oryza sativa         | ESTs C26000(C11448),AU082130(C11448) correspond to a region of the predicted gene.; Similar to mRNA for zinc-finger protein (Z36749).   |
| SEQ ID NO: 476 |          |                      |   |
| BAA78764.1     | AB023482 | Oryza sativa         | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAK11674.1     | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| AAF43496.1     | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| AAG16628.1     | AY007545 | Brassica napus       | protein serine/threonine kinase BNK1.   |
| BAA94509.1     | AB041503 | Populus nigra        | protein kinase 1. PnPK1.  |
| BAA94510.1     | AB041504 | Populus nigra        | protein kinase 2. PnPK2.  |
| BAB07999.1     | AP002525 | Oryza sativa         | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB03429.1     | AP002817 | Oryza sativa         | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).             |
| BAA87853.1     | AP000816 | Oryza sativa         | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |



|            |          |                         |   |
|------------|----------|-------------------------|---|
| BAB16871.1 | AP002537 | Oryza sativa            | putative protein kinase APK1A Arabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192), D25110(R3192).          |
| BAB39409.1 | AP002901 | Oryza sativa            | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| AAK00425.1 | AC069324 | Oryza sativa            | Putative protein kinase. OSJNBa0071K19.11.  |
| BAB39873.1 | AP002882 | Oryza sativa            | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808), AU056702(S20808).                                |
| CAB51834.1 | 00069    | Oryza sativa            | 11332.5. contains eukaryotic protein kinase domain PF.  |
| BAB21240.1 | AP002953 | Oryza sativa            | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461), C22360(C11461).                                   |
| AAK21965.1 | AY028699 | Brassica napus          | receptor protein kinase PERK1.  |
| AAG59657.1 | AC084319 | Oryza sativa            | putative protein kinase. OSJNBa0004B24.20.  |
| AAG03090.1 | AC073405 | Oryza sativa            | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).                                      |
| AAG25966.1 | AF302082 | Nicotiana tabacum       | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment. |
| BAB21241.1 | AP002953 | Oryza sativa            | Putative Pto kinase interactor 1. P0426D06.21. contains ESTs AU108280(E0721), D48017(S13927).                         |
| CAB51480.1 | Y14600   | Sorghum bicolor         | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.                                      |
| AAC61805.1 | U28007   | Lycopersicon esculentum | serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.  |
| BAA87852.1 | AP000816 | Oryza sativa            | Similar to putative Ser/Thr protein kinase. (AC004218).   |
| BAA92221.1 | AP001278 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).        |
| AAB09771.1 | U67422   | Zea mays                | CRINKLY4 precursor. cr4. receptor kinase homolog.   |
| BAA90808.1 | AP001168 | Oryza sativa            | Similar to putative receptor-like protein kinase (AL035679).  |
| AAC27894.1 | AF023164 | Zea mays                | leucine-rich repeat transmembrane protein kinase 1. Itk1.   |
| AAB61708.1 | U93048   | Daucus carota           | somatic embryogenesis receptor-like kinase. SERK.   |

|                |          |  |
|----------------|----------|--|
| AAF91336.1     | AF249317 | Glycine max<br>Pti1 kinase-like protein. Pti1a. protein kinase.  |
| AAF91337.1     | AF249318 | Glycine max<br>Pti1 kinase-like protein. Pti1b. protein kinase.  |
| AAG33377.1     | AF290411 | Oryza meyeriana<br>serine/threonine protein kinase. R1.  |
| AAC27895.1     | AF023165 | Zea mays<br>leucine-rich repeat transmembrane protein kinase 2. ltk2.  |
| AAK11566.1     | AF318490 | Lycopersicon hirsutum<br>Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.   |
| CAA97692.1     | Z73295   | Catharanthus roseus<br>receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser.<br>Mechanism: autophosphorylation in cis. |
| AAF34428.1     | AF172282 | Oryza sativa<br>receptor-like protein kinase. DUPR11.18.   |
| AAF76306.1     | AF220602 | Lycopersicon pimpinellifolium<br>Pto kinase.   |
| AAB47423.1     | U59315   | Lycopersicon pimpinellifolium<br>serine/threonine protein kinase Pto. Pto. bacterial speck disease resistance in tomato; disease resistance gene.            |
| AAC48914.1     | U02271   | Lycopersicon pimpinellifolium<br>protein kinase.   |
| AAK11567.1     | AF318491 | Lycopersicon hirsutum<br>Pto-like protein kinase F. LhirPtoF.  |
| AAB47421.1     | U59316   | Lycopersicon esculentum<br>serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.                        |
| AAF76313.1     | AF220603 | Lycopersicon esculentum<br>Pto kinase. LescPth5.   |
| AAF66615.1     | AF142596 | Nicotiana tabacum<br>LRR receptor-like protein kinase.   |
| BAA92954.1     | AP001551 | Oryza sativa<br>Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).  |
| BAA92953.1     | AP001551 | Oryza sativa<br>Similar to Arabidopsis thaliana chromosome 4 BAC clone F10M6 ; S-receptor kinase -like protein. (AL021811).                                  |
| SEQ ID NO: 479 |          |  |
| CAA42622.1     | X60007   | Nicotiana glauca<br>nsGRP-2. putative glycine-rich protein of 19.7kDa.   |
| SEQ ID NO: 480 |          |  |
| BAA83373.1     | AP000391 | Oryza sativa<br>ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484).       |

|            |          |                      |  |
|------------|----------|----------------------|--|
| BAA84787.1 | AP000559 | Oryza sativa         | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| AAF91324.1 | AF244890 | Glycine max          | receptor-like protein kinase 3. RLK3. GmRLK3.  |
| AAF91322.1 | AF244888 | Glycine max          | receptor-like protein kinase 1. RLK1. GmRLK1.  |
| AAF91323.1 | AF244889 | Glycine max          | receptor-like protein kinase 2. RLK2. GmRLK2.  |
| AAB36558.1 | U77888   | Ipomoea nil          | receptor-like protein kinase INRPK1. inrpkl. leucine-rich repeat.  |
| AAC36318.1 | AF053127 | Malus x domestica    | leucine-rich receptor-like protein kinase. LRPKml.   |
| AAF59906.1 | AF197947 | Glycine max          | receptor protein kinase-like protein. CLV1B.   |
| BAB40094.1 | AP003210 | Oryza sativa         | putative receptor protein kinase. OSJNBa0010K01.7.   |
| AAF59905.1 | AF197946 | Glycine max          | receptor protein kinase-like protein. CLV1A.   |
| CAC20842.1 | AJ250467 | Pinus sylvestris     | receptor protein kinase. upk.  |
| BAB03629.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.30.   |
| BAB03621.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.21.   |
| BAB03631.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.32.   |
| AAB82755.1 | U72725   | Oryza longistaminata | receptor kinase-like protein. Xa21 gene family member A1; downstream of microsatellite region; disease resistance gene family member.  |
| AAF34426.1 | AF172282 | Oryza sativa         | leucine rich repeat containing protein kinase. DUPR11.16.  |
| AAK27806.1 | AC022457 | Oryza sativa         | putative protein kinase. OSJNBa0006L06.21.   |
| CAA61510.1 | X89226   | Oryza sativa         | leucine-rich repeat/receptor protein kinase. lrk2.   |
| AAC49123.1 | U37133   | Oryza sativa         | receptor kinase-like protein. Xa21. Xa21 disease resistance gene.  |
| AAC80225.1 | U72723   | Oryza longistaminata | receptor kinase-like protein. Xa21. disease resistance gene.   |
| AAK27817.1 | AC022457 | Oryza sativa         | putative protein kinase. OSJNBa0006L06.16.   |

|                |          |   |
|----------------|----------|---|
| BAA82393.1     | AP000367 | <i>Oryza sativa</i><br>EST C96716(C10608) corresponds to a region of the predicted gene.; Similar to putative<br>receptor protein kinase. (AC002334). |
| BAB03627.1     | AP002522 | <i>Oryza sativa</i><br>putative protein kinase Xa21. P0009G03.27.   |
| AAB82756.1     | U72724   | <i>Oryza sativa</i><br>receptor kinase-like protein. Xa21 gene family member E.   |
| AAG52992.1     | U77888   | <i>Ipomoea nil</i><br>receptor-like protein kinase INRPK1a. inrpk1.   |
| BAA88636.1     | AB029327 | <i>Nicotiana tabacum</i><br>elicitor-inducible LRR receptor-like protein EILP. EILP.  |
| AAB82753.1     | U72726   | <i>Oryza longistaminata</i><br>receptor kinase-like protein. Xa21 gene family member D.   |
| AAG52994.1     | U77888   | <i>Ipomoea nil</i><br>receptor-like protein kinase INRPK1c. inrpk1.   |
| CAB51480.1     | Y14600   | <i>Sorghum bicolor</i><br>putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| AAK21965.1     | AY028699 | <i>Brassica napus</i><br>receptor protein kinase PERK1.   |
| SEQ ID NO: 481 |          |   |
| CAB43505.1     | AJ239051 | <i>Cicer arietinum</i><br>cytochrome P450. cyp81E2.   |
| BAA22422.1     | AB001379 | <i>Glycyrrhiza echinata</i><br>cytochrome P450. CYP81E1.  |
| BAA93634.1     | AB025016 | <i>Lotus japonicus</i><br>cytochrome P450.  |
| BAA74465.1     | AB022732 | <i>Glycyrrhiza echinata</i><br>cytochrome P450. CYP Ge-31.  |
| CAB41490.1     | AJ238439 | <i>Cicer arietinum</i><br>cytochrome P450 monooxygenase. cyp81E3v2.   |
| CAA10067.1     | AJ012581 | <i>Cicer arietinum</i><br>cytochrome P450. cyp81E3.   |
| CAA04117.1     | AJ000478 | <i>Helianthus tuberosus</i><br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-<br>race).                      |
| CAA04116.1     | AJ000477 | <i>Helianthus tuberosus</i><br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.  |
| AAK38080.1     | AF321856 | <i>Lolium rigidum</i><br>putative cytochrome P450.  |
| AAK38079.1     | AF321855 | <i>Lolium rigidum</i><br>putative cytochrome P450.  |
| AAK38081.1     | AF321857 | <i>Lolium rigidum</i><br>putative cytochrome P450.  |

|            |          |  |
|------------|----------|--|
| AAG09208.1 | AF175278 | <i>Pisum sativum</i><br>wound-inducible P450 hydroxylase. CYP82A1.   |
| CAA71876.1 | Y10982   | <i>Glycine max</i><br>putative cytochrome P450.  |
| AAC49188.2 | U29333   | <i>Pisum sativum</i><br>cytochrome P450 monooxygenase. CYP82. new cytochrome P450 family.  |
| CAA65580.1 | X96784   | <i>Nicotiana tabacum</i><br>cytochrome P450. hsr515.   |
| CAA71515.1 | Y10491   | <i>Glycine max</i><br>putative cytochrome P450.  |
| BAA92894.1 | AB006790 | <i>Petunia x hybrida</i><br>cytochrome P450. IMT-2.  |
| AAG44132.1 | AF218296 | <i>Pisum sativum</i><br>cytochrome P450. P450 isolog.  |
| CAA64635.1 | X95342   | <i>Nicotiana tabacum</i><br>cytochrome P450. hsr515. hypersensitivity-related gene.  |
| AAD56282.1 | AF155332 | <i>Petunia x hybrida</i><br>flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.   |
| AAC34853.1 | AF082028 | <i>Hemerocallis</i> hybrid cultivar<br>putative cyt P450-containing fatty acid hydroxylase. senescence-associated protein 3. SA3.<br>mRNA accumulates in senescing petals.   |
| BAA12159.1 | D83968   | <i>Glycine max</i><br>Cytochrome P-450 (CYP93A1).  |
| AAA32913.1 | M32885   | <i>Persea americana</i><br>cytochrome P-450LXXIA1 (cyp71A1).   |
| CAA70575.1 | Y09423   | <i>Nepeta racemosa</i><br>cytochrome P450. CYP71A5.  |
| AAG34695.1 | AF313492 | <i>Matthiola incana</i><br>putative cytochrome P450.   |
| CAA71877.1 | Y10983   | <i>Glycine max</i><br>putative cytochrome P450.  |
| CAA71516.1 | Y10492   | <i>Glycine max</i><br>putative cytochrome P450.  |
| AAB94587.1 | AF022458 | <i>Glycine max</i><br>CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.  |
| AAC39454.1 | AF014802 | <i>Eschscholzia californica</i><br>(S)-N-methylcoclaurine 3'-hydroxylase. CYP82B1. cytochrome P-450-dependent<br>monooxygenase; methyl jasmonate-inducible cytochrome P-450-dependent, homologous to<br>wound-inducible CYP82A1 of <i>Pisum sativum</i> GenBank Accession Number U29333. |
| CAA70576.1 | Y09424   | <i>Nepeta racemosa</i><br>cytochrome P450. CYP71A6.  |
| AAB94590.1 | AF022461 | <i>Glycine max</i><br>CYP82C1p. CYP82C1. cytochrome P450 monooxygenase.  |

BAB40323.1 AB037244 *Asparagus officinalis*  
cytochrome P450. ASPI-1.

BAB40324.1 AB037245 *Asparagus officinalis*  
cytochrome P450. ASPI-2.

AAB61965.1 U48435 *Solanum chacoense*  
putative cytochrome P450.

AAA19701.1 L24438 *Thlaspi arvense*  
cytochrome P450.

BAA84071.1 AB028151 *Antirrhinum majus*  
flavone synthase II. cytochrome P450. AFNS2.

BAA13076.1 D86351 *Glycine max*  
cytochrome P-450 (CYP93A2).

AAD38930.1 AF135485 *Glycine max*  
cytochrome P450 monooxygenase CYP93D1. CYP93E1.

AAC32274.1 AF081575 *Petunia x hybrida*  
flavonoid 3',5'-hydroxylase. Hfl. P450 enzyme.

SEQ ID NO: 487

BAA96181.1 AP002093 *Oryza sativa*  
EST AU056651(S20760) corresponds to a region of the predicted gene. Similar to  
*Arabidopsis thaliana* chromosome II BAC F19I3; putative amino acid transporter  
(AC004238).

BAA96139.1 AP002092 *Oryza sativa*  
EST AU056651(S20760) corresponds to a region of the predicted gene. Similar to  
*Arabidopsis thaliana* chromosome II BAC F19I3; putative amino acid transporter  
(AC004238).

SEQ ID NO: 495

AAB71528.1 U94784 *Helianthus annuus*  
ATPase. unconventional myosin. hamy4. Hamyo4.

AAB71529.1 U94785 *Helianthus annuus*  
ATPase. unconventional myosin. hamy5. Hamyo5.

AAB71527.1 U94782 *Helianthus annuus*  
ATPase. unconventional myosin. hamy2. Hamyo2.

AAD17931.2 AF104924 *Zea mays*  
unconventional myosin heavy chain. MYO1. ATPase; similar to myosin class XI.

AAC27525.1 AF077352 *Chlamydomonas reinhardtii*  
myosin heavy chain. MYO1. class XI.

AAK21311.1 AF338254 *Petroselinum crispum*  
myosin subfamily XI heavy chain. PCM3.

BAA87057.1 AB034154 *Chara corallina*  
unconventional myosin heavy chain. ccm.

BAB03273.1 AB007459 *Chara corallina*  
cytoplasmic streaming. myosin. ccm1. CCM1.

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAF43440.1     | AF233886 | Vallisneria gigantea    | unconventional myosin XI. VMYO1.  |
| AAB53062.1     | U94398   | Acetabularia cliftonii  | myosin. myo2.   |
| AAD34597.1     | AF147739 | Zea mays                | myosin XI. M2. unconventional myosin.                                     |
| AAB53061.1     | U94397   | Acetabularia cliftonii  | myosin. myo1.   |
| AAD31926.1     | AF147738 | Zea mays                | myosin VIII ZMM3. zmm3. unconventional myosin.                            |
| AAB71526.1     | U94781   | Helianthus annuus       | unconventional myosin. hamy1. homologous to ATM1.                         |
| AAG49341.1     | AF319457 | Petroselinum crispum    | myosin subfamily VIII heavy chain.  |
| AAB93521.1     | U94783   | Helianthus annuus       | unconventional myosin. hamy3. Hamyo3; similar to class VIII myosin.       |
| CAA47476.1     | X67102   | Anemia phyllitidis      | myosin heavy chain. myo15.  |
| CAA47477.1     | X67103   | Anemia phyllitidis      | myosin heavy chain. myo22.  |
| AAF43441.1     | AF233887 | Vallisneria gigantea    | unconventional myosin VIII. VMYO2.  |
| AAG13633.1     | AC078840 | Oryza sativa            | putative myosin. OSJNBb0073N24.16.  |
| AAA92120.1     | U48785   | Vigna mungo             | plant myosin MBM1. mbm1. head coding region.                              |
| SEQ ID NO: 496 |          |                         |   |
| AAA34138.1     | M96324   | Lycopersicon esculentum | The calcium ATPase is a calcium ion pump. Ca <sup>2+</sup> -ATPase. LCA1. |
| AAD11618.1     | AF050496 | Lycopersicon esculentum | Ca <sup>2+</sup> -ATPase. LCA1B; alternative transcript.                  |
| AAD11617.1     | AF050495 | Lycopersicon esculentum | Ca <sup>2+</sup> -ATPase. LCA1A; alternative transcript.                  |
| BAA90510.2     | AP001111 | Oryza sativa            | rice EST AU030811, similar to rice Ca <sup>2+</sup> -ATPase (U82966).     |
| CAA63790.1     | X93592   | Dunaliella bioculata    | P-type ATPase. cal. calcium pumping; CA1.                                 |
| AAB58910.1     | U82966   | Oryza sativa            | Ca <sup>2+</sup> -ATPase.   |
| AAF73985.1     | AF096871 | Zea mays                | calcium pump. calcium ATPase. cap1.                                       |

|            |          |                               |  |
|------------|----------|-------------------------------|--|
| AAG28436.1 | AF195029 | Glycine max                   | plasma membrane Ca <sup>2+</sup> -ATPase. SCA2.                                      |
| AAG28435.1 | AF195028 | Glycine max                   | plasma membrane Ca <sup>2+</sup> -ATPase. SCA1.                                      |
| CAA68234.1 | X99972   | Brassica oleracea             | calmodulin-stimulated calcium-ATPase.  |
| AAD31896.1 | AF145478 | Mesembryanthemum crystallinum | calcium ATPase.  |
| CAB85494.1 | AJ132891 | Medicago truncatula           | proton pump. H <sup>+</sup> -ATPase. ha1.  |
| CAB85495.1 | AJ132892 | Medicago truncatula           | proton pump. H <sup>+</sup> -ATPase. ha1.  |
| CAB69824.1 | AJ271439 | Prunus persica                | plasma membrane H <sup>+</sup> ATPase. PPA1.   |
| CAA52107.1 | X73901   | Dunaliella bioculata          | plasma membrane ATPase. pma1.  |
| AAD46188.1 | AF156691 | Nicotiana plumbaginifolia     | plasma membrane proton ATPase. pma9.   |
| BAA01058.1 | D10207   | Oryza sativa                  | H-ATPase. OSA1.  |
| AAA34094.1 | M80489   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> ATPase. pma1.   |
| AAA34098.1 | M80490   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> ATPase. pma3.   |
| AAA34173.1 | M60166   | Lycopersicon esculentum       | H <sup>+</sup> -ATPase. LHA1.  |
| CAA54046.1 | X76536   | Solanum tuberosum             | H(+)-transporting ATPase. PHA1.  |
| AAD55399.1 | AF179442 | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase isoform LHA2. LHA2.                           |
| AAB49042.1 | U54690   | Dunaliella acidophila         | plasma membrane proton ATPase. dha1. DaDHA1; proton pump.                            |
| AAF98344.1 | AF275745 | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase. LHA2. P-type ion pump.                       |
| BAA08134.1 | D45189   | Zostera marina                | plasma membrane H <sup>+</sup> -ATPase. zha1.  |
| AAA34052.1 | M27888   | Nicotiana plumbaginifolia     | H <sup>+</sup> -translocating ATPase.  |
| CAA59800.1 | X85805   | Zea mays                      | H(+)-transporting ATPase. MHA-2.   |
| AAB17186.1 | U72148   | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase. LHA4. plasma membrane proton pumping ATPase. |



|            |          |                               |   |
|------------|----------|-------------------------------|---|
| CAA54045.1 | X76535   | Solanum tuberosum             | H(+)-transporting ATPase. PHA2.   |
| CAB69823.1 | AJ271438 | Prunus persica                | plasma membrane H <sup>+</sup> ATPase. PPA2.  |
| BAA06629.1 | D31843   | Oryza sativa                  | plasma membrane H <sup>+</sup> -ATPase. OSA2.   |
| CAC29435.1 | AJ310523 | Vicia faba                    | P-type H <sup>+</sup> -ATPase. vha4. predominantly expressed in flowers.  |
| CAC29436.1 | AJ310524 | Vicia faba                    | P-type H <sup>+</sup> -ATPase. ha5. predominantly expressed in guard cells and flowers.   |
| CAA59799.1 | X85804   | Phaseolus vulgaris            | H(+)-transporting ATPase. BHA-1.  |
| AAD46186.1 | AF156679 | Nicotiana plumbaginifolia     | plasma membrane proton ATPase. pma6.  |
| AAB84202.2 | AF029256 | Kosteletzkya virginica        | plasma membrane proton ATPase. ATP1.  |
| AAD46187.1 | AF156683 | Nicotiana plumbaginifolia     | plasma membrane proton ATPase. pma8.  |
| AAB60276.1 | U09989   | Zea mays                      | H(+)-transporting ATPase. Mha1.   |
| AAB41898.1 | U84891   | Mesembryanthemum crystallinum | plasma membrane proton pump. H <sup>+</sup> -transporting ATPase. PMA.  |
| AAB35314.2 | S79323   | Vicia faba                    | plasma membrane H(+)-ATPase precursor. plasma membrane H(+)-ATPase. This sequence comes from Fig. 1; conceptual translation presented here differs from translation in publication. |
| AAK31799.1 | AY029190 | Lilium longiflorum            | plasma membrane H <sup>+</sup> ATPase. LILHA1.  |
| CAA47275.1 | X66737   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> -ATPase. pma4.   |
| BAA37150.1 | AB022442 | Vicia faba                    | p-type H <sup>+</sup> -ATPase. VHA2.  |
| AAG01028.1 | AF289025 | Cucumis sativus               | plasma membrane H <sup>+</sup> -ATPase.   |
| AAA81348.1 | U38965   | Vicia faba                    | p-type H <sup>+</sup> -ATPase. VHA2.  |
| AAK32118.1 | AF308816 | Hordeum vulgare               | plasmalemma H <sup>+</sup> -ATPase 1.   |
| AAF97591.1 | AF263917 | Lycopersicon esculentum       | plasma membrane proton ATPase. LHA8.  |
| AAK32119.1 | AF308817 | Hordeum vulgare               | plasmalemma H <sup>+</sup> -ATPase 2.   |

|                |          |                             |   |
|----------------|----------|-----------------------------|---|
| AAA20601.1     | U08985   | Zea mays                    | plasma-membrane H <sup>+</sup> ATPase. Zmpmal.  |
| SEQ ID NO: 497 |          |                             |   |
| AAB41742.1     | U82559   | Lycopersicon esculentum     | aldehyde oxidase 1 homolog. TAO1. has sequence properties of the molybdenum cofactor containing the hydroxylase genes aldehyde oxidase and xanthine dehydrogenase; belongs to a multigene family. |
| AAB41741.1     | U82558   | Lycopersicon esculentum     | aldehyde oxidase 1 homolog. TAO1. has sequence properties of the molybdenum cofactor containing the hydroxylase genes aldehyde oxidase and xanthine dehydrogenase; belongs to a multigene family. |
| SEQ ID NO: 498 |          |                             |   |
| CAA73134.1     | Y12531   | Brassica oleracea           | serine/threonine kinase. BRLK.  |
| AAB93834.1     | U82481   | Zea mays                    | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.  |
| CAA67145.1     | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.   |
| CAA74661.1     | Y14285   | Brassica oleracea           | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.  |
| CAA73133.1     | Y12530   | Brassica oleracea           | serine /threonine kinase. ARLK.   |
| AAA62232.1     | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.  |
| CAB89179.1     | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.  |
| AAA33008.1     | M97667   | Brassica napus              | serine/threonine kinase receptor.   |
| BAA92836.1     | AB032473 | Brassica oleracea           | S18 S-locus receptor kinase. SRK18.   |
| BAA23676.1     | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.   |
| AAC23542.1     | U20948   | Ipomoea trifida             | receptor protein kinase. IRK1.  |
| AAA33000.1     | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.  |
| BAA21132.1     | D88193   | Brassica rapa               | S-receptor kinase. SRK9 (B.c).  |
| BAA06285.1     | D30049   | Brassica rapa               | S-receptor kinase SRK9.   |

|                |          |                    |  |
|----------------|----------|--------------------|--|
| CAB41879.1     | Y18260   | Brassica oleracea  | SRK15 protein. SRK15. receptor-like kinase.  |
| BAA07577.2     | D38564   | Brassica rapa      | receptor protein kinase SRK12.   |
| CAA74662.1     | Y14286   | Brassica oleracea  | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| BAB21001.1     | AB054061 | Brassica rapa      | S locus receptor kinase. SRK22.  |
| BAA92837.1     | AB032474 | Brassica oleracea  | S60 S-locus receptor kinase. SRK60.  |
| CAB41878.1     | Y18259   | Brassica oleracea  | SRK5 protein. SRK5. receptor-like kinase.  |
| CAA79355.1     | Z18921   | Brassica oleracea  | S-receptor kinase-like protein.  |
| BAA07576.1     | D38563   | Brassica rapa      | receptor protein kinase SRK8.  |
| AAF34428.1     | AF172282 | Oryza sativa       | receptor-like protein kinase. DUPR11.18.   |
| AAD21872.1     | AF078082 | Phaseolus vulgaris | receptor-like protein kinase homolog RK20-1.   |
| AAD52097.1     | AF088885 | Nicotiana tabacum  | receptor-like kinase CHRK1. Chrk1.   |
| BAA94516.1     | AP001800 | Oryza sativa       | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB07906.1     | AP002835 | Oryza sativa       | putative S-receptor kinase. P0417G05.14.   |
| BAA94517.1     | AP001800 | Oryza sativa       | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB07905.1     | AP002835 | Oryza sativa       | putative S-receptor kinase. P0417G05.13.   |
| AAA33915.1     | L27821   | Oryza sativa       | receptor type serine/threonine kinase. protein kinase.   |
| BAB18292.1     | AP002860 | Oryza sativa       | putative receptor-like protein kinase. P0409B08.19.  |
| BAA94529.2     | AP001800 | Oryza sativa       | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| SEQ ID NO: 499 |          |                    |  |
| AAC36318.1     | AF053127 | Malus x domestica  | leucine-rich receptor-like protein kinase. LRPKm1.   |
| AAF91324.1     | AF244890 | Glycine max        | receptor-like protein kinase 3. RLK3. GmRLK3.  |

|            |          |  |
|------------|----------|--|
| AAF91323.1 | AF244889 | Glycine max<br>receptor-like protein kinase 2. RLK2. GmRLK2.   |
| CAA61510.1 | X89226   | Oryza sativa<br>leucine-rich repeat/receptor protein kinase. lrk2.   |
| AAF59906.1 | AF197947 | Glycine max<br>receptor protein kinase-like protein. CLV1B.  |
| AAF59905.1 | AF197946 | Glycine max<br>receptor protein kinase-like protein. CLV1A.  |
| AAF91322.1 | AF244888 | Glycine max<br>receptor-like protein kinase 1. RLK1. GmRLK1.   |
| BAB40094.1 | AP003210 | Oryza sativa<br>putative receptor protein kinase. OSJNBa0010K01.7.   |
| CAC20842.1 | AJ250467 | Pinus sylvestris<br>receptor protein kinase. upk.  |
| AAB36558.1 | U77888   | Ipomoea nil<br>receptor-like protein kinase INRPK1. inrpkl. leucine-rich repeat.   |
| BAA83373.1 | AP000391 | Oryza sativa<br>ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| BAA84787.1 | AP000559 | Oryza sativa<br>ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| AAK27806.1 | AC022457 | Oryza sativa<br>putative protein kinase. OSJNBa0006L06.21.   |
| AAF34426.1 | AF172282 | Oryza sativa<br>leucine rich repeat containing protein kinase. DUPR11.16.  |
| BAB03627.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.27.   |
| BAB03631.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.32.   |
| AAK27817.1 | AC022457 | Oryza sativa<br>putative protein kinase. OSJNBa0006L06.16.   |
| BAB03621.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.21.   |
| BAB03629.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.30.   |
| AAC80225.1 | U72723   | Oryza longistaminata<br>receptor kinase-like protein. Xa21. disease resistance gene.   |
| AAC49123.1 | U37133   | Oryza sativa<br>receptor kinase-like protein. Xa21. Xa21 disease resistance gene.  |
| BAA82393.1 | AP000367 | Oryza sativa<br>EST C96716(C10608) corresponds to a region of the predicted gene.; Similar to putative receptor protein kinase. (AC002334).            |

|                |          |                      |   |
|----------------|----------|----------------------|---|
| AAB82755.1     | U72725   | Oryza longistaminata | receptor kinase-like protein. Xa21 gene family member A1; downstream of microsatellite region; disease resistance gene family member.   |
| AAG52992.1     | U77888   | Ipomoea nil          | receptor-like protein kinase INRPK1a. inrpk1.   |
| AAB82756.1     | U72724   | Oryza sativa         | receptor kinase-like protein. Xa21 gene family member E.  |
| AAB82753.1     | U72726   | Oryza longistaminata | receptor kinase-like protein. Xa21 gene family member D.  |
| CAB51480.1     | Y14600   | Sorghum bicolor      | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| BAA88636.1     | AB029327 | Nicotiana tabacum    | elicitor-inducible LRR receptor-like protein EILP. EILP.  |
| AAG52994.1     | U77888   | Ipomoea nil          | receptor-like protein kinase INRPK1c. inrpk1.   |
| BAA94519.1     | AP001800 | Oryza sativa         | ESTs AU032341(R3918),AU071016(R10613) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 4, BAC F9D16; putative receptor kinase (AL035394). |
| BAB07903.1     | AP002835 | Oryza sativa         | putative receptor kinase. P0417G05.10. contains ESTs AU032341(R3918),AU071016(R10613).  |
| BAA87853.1     | AP000816 | Oryza sativa         | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| AAD38286.1     | AC007789 | Oryza sativa         | putative protein kinase. OSJNBa0049B20.13.  |
| BAB40081.1     | AP003074 | Oryza sativa         | putative receptor protein kinase. OSJNBa0004G10.30.   |
| BAB18321.1     | AP002865 | Oryza sativa         | putative receptor protein kinase. P0034C11.11.  |
| SEQ ID NO: 500 |          |                      |   |
| BAB40094.1     | AP003210 | Oryza sativa         | putative receptor protein kinase. OSJNBa0010K01.7.  |
| AAC36318.1     | AF053127 | Malus x domestica    | leucine-rich receptor-like protein kinase. LRPKml.  |
| AAF59906.1     | AF197947 | Glycine max          | receptor protein kinase-like protein. CLV1B.  |
| AAF59905.1     | AF197946 | Glycine max          | receptor protein kinase-like protein. CLV1A.  |
| AAF91323.1     | AF244889 | Glycine max          | receptor-like protein kinase 2. RLK2. GmRLK2.   |

|            |          |  |
|------------|----------|--|
| AAF91322.1 | AF244888 | Glycine max<br>receptor-like protein kinase 1. RLK1. GmRLK1.   |
| AAF91324.1 | AF244890 | Glycine max<br>receptor-like protein kinase 3. RLK3. GmRLK3.   |
| CAA61510.1 | X89226   | Oryza sativa<br>leucine-rich repeat/receptor protein kinase. lrk2.   |
| CAC20842.1 | AJ250467 | Pinus sylvestris<br>receptor protein kinase. upk.  |
| BAA84787.1 | AP000559 | Oryza sativa<br>ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar<br>to receptor protein kinase, ERECTA (AC004484).        |
| BAA83373.1 | AP000391 | Oryza sativa<br>ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar<br>to receptor protein kinase, ERECTA (AC004484).        |
| AAK27806.1 | AC022457 | Oryza sativa<br>putative protein kinase. OSJNBa0006L06.21.   |
| AAB36558.1 | U77888   | Ipomoea nil<br>receptor-like protein kinase INRPK1. inrpkl. leucine-rich repeat.   |
| BAB03627.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.27.   |
| BAB03629.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.30.   |
| BAB03631.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.32.   |
| BAB03621.1 | AP002522 | Oryza sativa<br>putative protein kinase Xa21. P0009G03.21.   |
| AAK27817.1 | AC022457 | Oryza sativa<br>putative protein kinase. OSJNBa0006L06.16.   |
| AAF34426.1 | AF172282 | Oryza sativa<br>leucine rich repeat containing protein kinase. DUPR11.16.  |
| AAB82755.1 | U72725   | Oryza longistaminata<br>receptor kinase-like protein. Xa21 gene family member A1; downstream of microsatellite<br>region; disease resistance gene family member. |
| AAC80225.1 | U72723   | Oryza longistaminata<br>receptor kinase-like protein. Xa21. disease resistance gene.   |
| AAC49123.1 | U37133   | Oryza sativa<br>receptor kinase-like protein. Xa21. Xa21 disease resistance gene.  |
| BAA82393.1 | AP000367 | Oryza sativa<br>EST C96716(C10608) corresponds to a region of the predicted gene.; Similar to putative<br>receptor protein kinase. (AC002334).                   |
| AAB82756.1 | U72724   | Oryza sativa<br>receptor kinase-like protein. Xa21 gene family member E.   |

|                |          |   |
|----------------|----------|---|
| BAA88636.1     | AB029327 | Nicotiana tabacum<br>elicitor-inducible LRR receptor-like protein EILP. EILP.   |
| AAG52992.1     | U77888   | Ipomoea nil<br>receptor-like protein kinase INRPK1a. inrpk1.  |
| AAB82753.1     | U72726   | Oryza longistaminata<br>receptor kinase-like protein. Xa21 gene family member D.  |
| CAB51480.1     | Y14600   | Sorghum bicolor<br>putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.                                 |
| AAB61708.1     | U93048   | Daucus carota<br>somatic embryogenesis receptor-like kinase. SERK.  |
| SEQ ID NO: 501 |          |   |
| CAA73134.1     | Y12531   | Brassica oleracea<br>serine/threonine kinase. BRLK.   |
| AAB93834.1     | U82481   | Zea mays<br>KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.                    |
| CAA67145.1     | X98520   | Brassica oleracea<br>receptor-like kinase. SFR2.  |
| CAA73133.1     | Y12530   | Brassica oleracea<br>serine /threonine kinase. ARLK.  |
| CAA74661.1     | Y14285   | Brassica oleracea<br>SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554. |
| AAA62232.1     | U00443   | Brassica napus<br>S-receptor kinase. protein contains an immunoglobulin-like domain.  |
| BAA92836.1     | AB032473 | Brassica oleracea<br>S18 S-locus receptor kinase. SRK18.  |
| CAB41878.1     | Y18259   | Brassica oleracea<br>SRK5 protein. SRK5. receptor-like kinase.  |
| AAC23542.1     | U20948   | Ipomoea trifida<br>receptor protein kinase. IRK1.   |
| BAA23676.1     | AB000970 | Brassica rapa<br>receptor kinase 1. BcRK1.  |
| CAB41879.1     | Y18260   | Brassica oleracea<br>SRK15 protein. SRK15. receptor-like kinase.  |
| AAA33000.1     | M76647   | Brassica oleracea<br>receptor protein kinase. SKR6.   |
| AAA33008.1     | M97667   | Brassica napus<br>serine/threonine kinase receptor.   |
| CAB89179.1     | AJ245479 | Brassica napus subsp. napus<br>ser /thr kinase. S-locus receptor kinase. srk.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA74662.1     | Y14286   | Brassica oleracea       | SFR3, extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.   |
| BAA06285.1     | D30049   | Brassica rapa           | S-receptor kinase SRK9.  |
| BAA21132.1     | D88193   | Brassica rapa           | S-receptor kinase. SRK9 (B.c).   |
| BAA07577.2     | D38564   | Brassica rapa           | receptor protein kinase SRK12.   |
| CAA79355.1     | Z18921   | Brassica oleracea       | S-receptor kinase-like protein.  |
| BAA07576.1     | D38563   | Brassica rapa           | receptor protein kinase SRK8.  |
| BAB21001.1     | AB054061 | Brassica rapa           | S locus receptor kinase. SRK22.  |
| BAA92837.1     | AB032474 | Brassica oleracea       | S60 S-locus receptor kinase. SRK60.  |
| AAD21872.1     | AF078082 | Phaseolus vulgaris      | receptor-like protein kinase homolog RK20-1.   |
| AAF34428.1     | AF172282 | Oryza sativa            | receptor-like protein kinase. DUPR11.18.   |
| AAD52097.1     | AF088885 | Nicotiana tabacum       | receptor-like kinase CHRK1. Chrkl.   |
| BAB07906.1     | AP002835 | Oryza sativa            | putative S-receptor kinase. P0417G05.14.   |
| BAA94516.1     | AP001800 | Oryza sativa            | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB18292.1     | AP002860 | Oryza sativa            | putative receptor-like protein kinase. P0409B08.19.  |
| BAB07904.1     | AP002835 | Oryza sativa            | putative S-receptor kinase. P0417G05.12.   |
| BAA94518.1     | AP001800 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome 2 section 111 of 255; putative receptor-like protein kinase (AC002392).   |
| SEQ ID NO: 502 |          |                         |  |
| AAC78596.1     | AF053998 | Lycopersicon esculentum | Hcr2-5D. Hcr2-5D. similar to Lycopersicon pimpinellifolium disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |
| AAC78591.1     | AF053993 | Lycopersicon esculentum | disease resistance protein. Cf-5.  |
| AAC78593.1     | AF053995 | Lycopersicon esculentum | Hcr2-0B. Hcr2-0B. similar to Lycopersicon pimpinellifolium disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |



|                |          |   |
|----------------|----------|---|
| AAC78592.1     | AF053994 | <i>Lycopersicon esculentum</i><br>Hcr2-0A. Hcr2-0A. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445.       |
| AAC78594.1     | AF053996 | <i>Lycopersicon pimpinellifolium</i><br>Hcr2-2A. Hcr2-2A. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |
| CAA05274.1     | AJ002236 | <i>Lycopersicon pimpinellifolium</i><br>resistance gene. Cf-9. Cf-9.  |
| AAA65235.1     | U15936   | <i>Lycopersicon pimpinellifolium</i><br>Cf-9 precursor. Cf-9. this is the ninth resistance gene to disease caused by <i>Cladosporium fulvum</i> to be isolated.   |
| CAA05276.1     | AJ002236 | <i>Lycopersicon pimpinellifolium</i><br>resistance gene. Hcr9-9E. Hcr9-9E.  |
| AAC78595.1     | AF053997 | <i>Lycopersicon esculentum</i><br>Hcr2-5B. Hcr2-5B. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445.       |
| CAA05268.1     | AJ002235 | <i>Lycopersicon hirsutum</i><br>Resistance gene. Cf-4. Cf-4.  |
| CAA05279.1     | AJ002237 | <i>Lycopersicon esculentum</i><br>Hcr9-0. Hcr9-0. homologue of <i>Cladosporium fulvum</i> disease resistance gene Cf-9.   |
| BAB08215.1     | AP002539 | <i>Oryza sativa</i><br>Similar to <i>Lycopersicon esculentum</i> disease resistance protein (AF053993).   |
| BAA96776.1     | AP002521 | <i>Oryza sativa</i><br>Similar to <i>Lycopersicon esculentum</i> disease resistance protein (AF053993).   |
| AAG21897.1     | AC026815 | <i>Oryza sativa</i><br>putative disease resistance protein (3' partial). OSJNBa0079L16.21.  |
| AAD50430.1     | AF166121 | <i>Hordeum vulgare</i><br>Cf2/Cf5 disease resistance protein homolog. Big1. leucine rich repeat protein.  |
| AAG21917.1     | AC026815 | <i>Oryza sativa</i><br>putative disease resistance protein. OSJNBa0079L16.5.  |
| CAB55409.1     | AL117265 | <i>Oryza sativa</i><br>zhb0001.1. Incomplete at 5'end, Similar to disease resistance protein; Method: conceptual translation with partial peptide sequencing.   |
| AAG21909.1     | AC026815 | <i>Oryza sativa</i><br>putative disease resistance protein. OSJNBa0079L16.3.  |
| AAC80225.1     | U72723   | <i>Oryza longistaminata</i><br>receptor kinase-like protein. Xa21. disease resistance gene.   |
| AAC49123.1     | U37133   | <i>Oryza sativa</i><br>receptor kinase-like protein. Xa21. Xa21 disease resistance gene.  |
| SEQ ID NO: 503 |          |   |
| CAA05276.1     | AJ002236 | <i>Lycopersicon pimpinellifolium</i><br>resistance gene. Hcr9-9E. Hcr9-9E.  |

|                |          |   |
|----------------|----------|---|
| AAC78592.1     | AF053994 | <i>Lycopersicon esculentum</i><br>Hcr2-0A. Hcr2-0A. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445.       |
| CAA05274.1     | AJ002236 | <i>Lycopersicon pimpinellifolium</i><br>resistance gene. Cf-9. Cf-9.  |
| AAA65235.1     | U15936   | <i>Lycopersicon pimpinellifolium</i><br>Cf-9 precursor. Cf-9. this is the ninth resistance gene to disease caused by <i>Cladosporium fulvum</i> to be isolated.   |
| AAC78594.1     | AF053996 | <i>Lycopersicon pimpinellifolium</i><br>Hcr2-2A. Hcr2-2A. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445. |
| CAA05279.1     | AJ002237 | <i>Lycopersicon esculentum</i><br>Hcr9-0. Hcr9-0. homologue of <i>Cladosporium fulvum</i> disease resistance gene Cf-9.   |
| AAC78595.1     | AF053997 | <i>Lycopersicon esculentum</i><br>Hcr2-5B. Hcr2-5B. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445.       |
| AAC78593.1     | AF053995 | <i>Lycopersicon esculentum</i><br>Hcr2-0B. Hcr2-0B. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445.       |
| CAA05268.1     | AJ002235 | <i>Lycopersicon hirsutum</i><br>Resistance gene. Cf-4. Cf-4.  |
| AAC78596.1     | AF053998 | <i>Lycopersicon esculentum</i><br>Hcr2-5D. Hcr2-5D. similar to <i>Lycopersicon pimpinellifolium</i> disease resistance protein Cf-2.2 encoded by the sequence presented in GenBank Accession Number U42445.       |
| AAC78591.1     | AF053993 | <i>Lycopersicon esculentum</i><br>disease resistance protein. Cf-5.   |
| BAA96776.1     | AP002521 | <i>Oryza sativa</i><br>Similar to <i>Lycopersicon esculentum</i> disease resistance protein (AF053993).   |
| BAB08215.1     | AP002539 | <i>Oryza sativa</i><br>Similar to <i>Lycopersicon esculentum</i> disease resistance protein (AF053993).   |
| CAB55409.1     | AL117265 | <i>Oryza sativa</i><br>zhb0001.1. Incomplete at 5'end, Similar to disease resistance protein; Method: conceptual translation with partial peptide sequencing.   |
| AAD50430.1     | AF166121 | <i>Hordeum vulgare</i><br>Cf2/Cf5 disease resistance protein homolog. Big1. leucine rich repeat protein.  |
| AAG21897.1     | AC026815 | <i>Oryza sativa</i><br>putative disease resistance protein (3' partial). OSJNBa0079L16.21.  |
| AAB82755.1     | U72725   | <i>Oryza longistaminata</i><br>receptor kinase-like protein. Xa21 gene family member A1; downstream of microsatellite region; disease resistance gene family member.  |
| SEQ ID NO: 504 |          |   |
| AAF91324.1     | AF244890 | <i>Glycine max</i><br>receptor-like protein kinase 3. RLK3. GmRLK3.   |

|            |          |                      |  |
|------------|----------|----------------------|--|
| BAA84787.1 | AP000559 | Oryza sativa         | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| BAA83373.1 | AP000391 | Oryza sativa         | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| AAF91322.1 | AF244888 | Glycine max          | receptor-like protein kinase 1. RLK1. GmRLK1.  |
| AAF91323.1 | AF244889 | Glycine max          | receptor-like protein kinase 2. RLK2. GmRLK2.  |
| CAC20842.1 | AJ250467 | Pinus sylvestris     | receptor protein kinase. upk.  |
| AAK27806.1 | AC022457 | Oryza sativa         | putative protein kinase. OSJNBa0006L06.21.   |
| AAB36558.1 | U77888   | Ipomoea nil          | receptor-like protein kinase INRPK1. inrpkl. leucine-rich repeat.  |
| BAA82393.1 | AP000367 | Oryza sativa         | EST C96716(C10608) corresponds to a region of the predicted gene.; Similar to putative receptor protein kinase. (AC002334).            |
| AAF59906.1 | AF197947 | Glycine max          | receptor protein kinase-like protein. CLV1B.   |
| BAB03629.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.30.   |
| BAB03627.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.27.   |
| AAF59905.1 | AF197946 | Glycine max          | receptor protein kinase-like protein. CLV1A.   |
| AAK27817.1 | AC022457 | Oryza sativa         | putative protein kinase. OSJNBa0006L06.16.   |
| BAB03631.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.32.   |
| BAB03621.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.21.   |
| AAF34426.1 | AF172282 | Oryza sativa         | leucine rich repeat containing protein kinase. DUPR11.16.  |
| AAF66615.1 | AF142596 | Nicotiana tabacum    | LRR receptor-like protein kinase.  |
| AAB82755.1 | U72725   | Oryza longistaminata | receptor kinase-like protein. Xa21 gene family member A1; downstream of microsatellite region; disease resistance gene family member.  |
| AAC49123.1 | U37133   | Oryza sativa         | receptor kinase-like protein. Xa21. Xa21 disease resistance gene.  |

|                |          |   |
|----------------|----------|---|
| AAC80225.1     | U72723   | <i>Oryza longistaminata</i><br>receptor kinase-like protein. Xa21. disease resistance gene.   |
| BAB19337.1     | AP003044 | <i>Oryza sativa</i><br>putative protein kinase. P0038C05.10. contains ESTs<br>AU056335(S20481),AU056336(S20481).  |
| BAA88636.1     | AB029327 | <i>Nicotiana tabacum</i><br>elicitor-inducible LRR receptor-like protein EILP. EILP.  |
| CAB51480.1     | Y14600   | <i>Sorghum bicolor</i><br>putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| AAK21965.1     | AY028699 | <i>Brassica napus</i><br>receptor protein kinase PERK1.   |
| BAB39409.1     | AP002901 | <i>Oryza sativa</i><br>putative protein kinase. P0456F08.9. contains EST C23560(R0290).   |
| BAB07903.1     | AP002835 | <i>Oryza sativa</i><br>putative receptor kinase. P0417G05.10. contains ESTs<br>AU032341(R3918),AU071016(R10613).  |
| BAA94519.1     | AP001800 | <i>Oryza sativa</i><br>ESTs AU032341(R3918),AU071016(R10613) correspond to a region of the predicted gene.<br>Similar to <i>Arabidopsis thaliana</i> chromosome 4, BAC F9D16; putative receptor kinase<br>(AL035394). |
| AAK00425.1     | AC069324 | <i>Oryza sativa</i><br>Putative protein kinase. OSJNBa0071K19.11.   |
| AAG59657.1     | AC084319 | <i>Oryza sativa</i><br>putative protein kinase. OSJNBa0004B24.20.   |
| BAA87853.1     | AP000816 | <i>Oryza sativa</i><br>EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative<br>NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB39873.1     | AP002882 | <i>Oryza sativa</i><br>putative protein kinase. P0439B06.8. contains ESTs<br>AU056701(S20808),AU056702(S20808).   |
| AAB82756.1     | U72724   | <i>Oryza sativa</i><br>receptor kinase-like protein. Xa21 gene family member E.   |
| AAB82753.1     | U72726   | <i>Oryza longistaminata</i><br>receptor kinase-like protein. Xa21 gene family member D.   |
| SEQ ID NO: 505 |          |   |
| CAA73134.1     | Y12531   | <i>Brassica oleracea</i><br>serine/threonine kinase. BRLK.  |
| CAA74661.1     | Y14285   | <i>Brassica oleracea</i><br>SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular<br>kinase domain: 1412-2554.   |
| AAC23542.1     | U20948   | <i>Ipomoea trifida</i><br>receptor protein kinase. IRK1.  |

|            |          |                             |  |
|------------|----------|-----------------------------|--|
| BAA23676.1 | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.  |
| AAB93834.1 | U82481   | Zea mays                    | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.             |
| CAA73133.1 | Y12530   | Brassica oleracea           | serine /threonine kinase. ARLK.  |
| CAA67145.1 | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.  |
| CAB89179.1 | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| AAA33008.1 | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| BAA21132.1 | D88193   | Brassica rapa               | S-receptor kinase. SRK9 (B.c).   |
| AAA62232.1 | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| BAA06285.1 | D30049   | Brassica rapa               | S-receptor kinase SRK9.  |
| CAA74662.1 | Y14286   | Brassica oleracea           | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| AAA33000.1 | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| BAA07576.1 | D38563   | Brassica rapa               | receptor protein kinase SRK8.  |
| BAA92836.1 | AB032473 | Brassica oleracea           | S18 S-locus receptor kinase. SRK18.  |
| CAB41878.1 | Y18259   | Brassica oleracea           | SRK5 protein. SRK5. receptor-like kinase.  |
| CAA79355.1 | Z18921   | Brassica oleracea           | S-receptor kinase-like protein.  |
| CAB41879.1 | Y18260   | Brassica oleracea           | SRK15 protein. SRK15. receptor-like kinase.  |
| BAB21001.1 | AB054061 | Brassica rapa               | S locus receptor kinase. SRK22.  |
| BAA92837.1 | AB032474 | Brassica oleracea           | S60 S-locus receptor kinase. SRK60.  |
| BAA07577.2 | D38564   | Brassica rapa               | receptor protein kinase SRK12.   |
| AAD21872.1 | AF078082 | Phaseolus vulgaris          | receptor-like protein kinase homolog RK20-1.   |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| BAA94517.1     | AP001800 | Oryza sativa                  | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| BAB07905.1     | AP002835 | Oryza sativa                  | putative S-receptor kinase. P0417G05.13.  |
| BAB07906.1     | AP002835 | Oryza sativa                  | putative S-receptor kinase. P0417G05.14.  |
| BAA94516.1     | AP001800 | Oryza sativa                  | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| AAD52097.1     | AF088885 | Nicotiana tabacum             | receptor-like kinase CHRK1. Chrkl.  |
| BAB18292.1     | AP002860 | Oryza sativa                  | putative receptor-like protein kinase. P0409B08.19.   |
| SEQ ID NO: 507 |          |                               |   |
| AAA33509.1     | M62985   | Zea mays                      | protein kinase. putative; putative.   |
| CAA62476.1     | X90990   | Solanum tuberosum             | stpk1 protein kinase.   |
| AAK31277.1     | AC079890 | Oryza sativa                  | putative protein kinase. OSJNBb0089A17.15.  |
| CAA66616.1     | X97980   | Solanum berthaultii           | protein kinase.   |
| AAF66637.1     | AF143505 | Lycopersicon esculentum       | viroid symptom modulation protein. PKv. protein kinase; induced by viroid infection.  |
| BAA96593.1     | AP002481 | Oryza sativa                  | Similar to Solanum berthaultii protein kinase. (X97980).  |
| AAB54117.1     | U93559   | Brassica rapa                 | putative serine/threonine protein kinase. Bcpk1.  |
| AAB88817.1     | AF033263 | Zea mays                      | signal transduction for phototropism. nonphototropic hypocotyl 1. nph1. NPH1; putative serine/threonine kinase; similar to oat NPH1 proteins. |
| AAC05084.1     | AF033097 | Avena sativa                  | NPH1-2. NPH1-2. putative serine/threonine protein kinase.   |
| CAA82994.1     | Z30333   | Mesembryanthemum crystallinum | protein kinase.   |
| AAC05083.1     | AF033096 | Avena sativa                  | NPH1-1. NPH1-1. putative serine/threonine protein kinase.   |
| CAA82992.1     | Z30331   | Mesembryanthemum crystallinum | Protein Kinase.   |
| CAA82993.1     | Z30332   | Spinacia oleracea             | protein kinase.   |
| AAA50304.1     | M92989   | Pisum sativum                 | protein kinase. PK5. homologue.   |

|            |          |  |
|------------|----------|--|
| AAB71418.1 | U11553   | <i>Pisum sativum</i><br>PsPK3. putative protein kinase.  |
| BAA93704.1 | AB032564 | <i>Cucumis sativus</i><br>cucumber protein kinase CsPK3. CsPK3. putative.  |
| BAA36192.1 | AB012082 | <i>Adiantum capillus-veneris</i><br>PHY3. PHY3. Ser/Thr protein kinase; chimeric structure of red/far-red light photoreceptive region (phytochrome) and putative blue light photoreceptor in phototropism of hypocotyl (NPH1). |
| AAK18843.1 | AC082645 | <i>Oryza sativa</i><br>putative protein kinase. OSJNBb0033N16.3.   |
| CAB82852.1 | Z30329   | <i>Mesembryanthemum crystallinum</i><br>protein kinase MK6.  |
| CAA50374.1 | X71057   | <i>Nicotiana tabacum</i><br>protein kinase. PKTL7.   |
| AAK13156.1 | AC078829 | <i>Oryza sativa</i><br>putative protein kinase. OSJNBa0026O12.14.  |
| BAB03409.1 | AP002816 | <i>Oryza sativa</i><br>Similar to <i>Spinacia oleracea</i> protein kinase (S42867).  |
| CAA82991.1 | Z30330   | <i>Spinacia oleracea</i><br>protein kinase.  |
| AAD50584.1 | AF089097 | <i>Salvia columbariae</i><br>protein kinase 1. PK1.  |
| AAB93860.1 | U89679   | <i>Lycopersicon esculentum</i><br>protein kinase. LePK2. contains catalytic domain.  |
| AAD50585.1 | AF089099 | <i>Salvia columbariae</i><br>protein kinase 3. PK3.  |
| AAD50586.1 | AF089100 | <i>Salvia columbariae</i><br>protein kinase 4. PK4.  |
| AAB93859.1 | U89678   | <i>Lycopersicon esculentum</i><br>protein kinase. LePK1. contains catalytic domain.  |
| AAA50772.1 | M69030   | <i>Pisum sativum</i><br>protein serine/threonine kinase. PsPK1. putative; putative.  |
| AAB93861.1 | U89680   | <i>Lycopersicon esculentum</i><br>protein kinase. LePK3. contains catalytic domain.  |
| AAD50587.1 | AF089101 | <i>Salvia columbariae</i><br>protein kinase 5. PK5.  |
| AAD50588.1 | AF089102 | <i>Salvia columbariae</i><br>protein kinase 6. PK6.  |
| CAA56313.1 | X79992   | <i>Avena sativa</i><br>putative pp70 ribosomal protein S6 kinase. Aspk11.  |
| AAD50589.1 | AF089103 | <i>Salvia columbariae</i><br>protein kinase 7. PK7.  |

|                |          |   |   |
|----------------|----------|---|---|
| CAB89082.1     | AJ277534 | <i>Asparagus officinalis</i>              | S6 ribosomal protein kinase. pk1. putative.   |
| AAB93862.1     | U89681   | <i>Lycopersicon esculentum</i>            | protein kinase. LePK4. contains catalytic domain.   |
| BAA92972.1     | AP001551 | <i>Oryza sativa</i>                       | ESTs AU056183(S20356),AU056881(S20950) correspond to a region of the predicted gene. Similar to <i>Arabidopsis thaliana</i> chromosome 4 BAC clone F6I18 ; putative protein kinase. (AL022198). |
| BAB12687.1     | AP002746 | <i>Oryza sativa</i>                       | putative protein kinase. P0671B11.2. contains ESTs C22394(C30013),C22393(C30013).   |
| SEQ ID NO: 508 |          |   |   |
| BAA23676.1     | AB000970 | <i>Brassica rapa</i>                      | receptor kinase 1. BcRK1.   |
| CAA74662.1     | Y14286   | <i>Brassica oleracea</i>                  | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.  |
| CAA74661.1     | Y14285   | <i>Brassica oleracea</i>                  | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.  |
| CAA73133.1     | Y12530   | <i>Brassica oleracea</i>                  | serine /threonine kinase. ARLK.   |
| CAA67145.1     | X98520   | <i>Brassica oleracea</i>                  | receptor-like kinase. SFR2.   |
| AAA33008.1     | M97667   | <i>Brassica napus</i>                     | serine/threonine kinase receptor.   |
| CAB89179.1     | AJ245479 | <i>Brassica napus</i> subsp. <i>napus</i> | ser /thr kinase. S-locus receptor kinase. srk.  |
| CAA79355.1     | Z18921   | <i>Brassica oleracea</i>                  | S-receptor kinase-like protein.   |
| CAB41878.1     | Y18259   | <i>Brassica oleracea</i>                  | SRK5 protein. SRK5. receptor-like kinase.   |
| AAA33000.1     | M76647   | <i>Brassica oleracea</i>                  | receptor protein kinase. SKR6.  |
| BAA92837.1     | AB032474 | <i>Brassica oleracea</i>                  | S60 S-locus receptor kinase. SRK60.   |
| BAA21132.1     | D88193   | <i>Brassica rapa</i>                      | S-receptor kinase. SRK9 (B.c).  |
| BAA06285.1     | D30049   | <i>Brassica rapa</i>                      | S-receptor kinase SRK9.   |
| AAA62232.1     | U00443   | <i>Brassica napus</i>                     | S-receptor kinase. protein contains an immunoglobulin-like domain.  |
| BAA92836.1     | AB032473 | <i>Brassica oleracea</i>                  | S18 S-locus receptor kinase. SRK18.   |



|            |          |                    |   |
|------------|----------|--------------------|---|
| BAB21001.1 | AB054061 | Brassica rapa      | S locus receptor kinase. SRK22.   |
| BAA07576.1 | D38563   | Brassica rapa      | receptor protein kinase SRK8.   |
| BAA07577.2 | D38564   | Brassica rapa      | receptor protein kinase SRK12.  |
| CAA73134.1 | Y12531   | Brassica oleracea  | serine/threonine kinase. BRLK.  |
| AAB93834.1 | U82481   | Zea mays           | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.        |
| AAC23542.1 | U20948   | Ipomoea trifida    | receptor protein kinase. IRK1.  |
| CAB41879.1 | Y18260   | Brassica oleracea  | SRK15 protein. SRK15. receptor-like kinase.   |
| AAD52097.1 | AF088885 | Nicotiana tabacum  | receptor-like kinase CHRK1. Chrk1.  |
| AAD21872.1 | AF078082 | Phaseolus vulgaris | receptor-like protein kinase homolog RK20-1.  |
| BAB18292.1 | AP002860 | Oryza sativa       | putative receptor-like protein kinase. P0409B08.19.   |
| AAK02023.1 | AC074283 | Oryza sativa       | Putative protein kinase-like. OSJNBa0087H07.5.  |
| CAA79324.1 | Z18884   | Brassica oleracea  | S-receptor kinase related protein.  |
| BAB21240.1 | AP002953 | Oryza sativa       | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).                          |
| CAB51836.1 | AJ243961 | Oryza sativa       | Putitive Ser/Thr protein kinase. I1332.7.   |
| AAK21965.1 | AY028699 | Brassica napus     | receptor protein kinase PERK1.  |
| BAB39873.1 | AP002882 | Oryza sativa       | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).                       |
| BAA94509.1 | AB041503 | Populus nigra      | protein kinase 1. PnPK1.  |
| AAK00425.1 | AC069324 | Oryza sativa       | Putative protein kinase. OSJNBa0071K19.11.  |
| BAA94510.1 | AB041504 | Populus nigra      | protein kinase 2. PnPK2.  |
| BAA92953.1 | AP001551 | Oryza sativa       | Similar to Arabidopsis thaliana chromosome 4 BAC clone F10M6 ; S-receptor kinase -like protein. (AL021811). |

|  |          |  |
|--|----------|--|
| BAA92954.1   | AP001551 | Oryza sativa                           |
| Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).  |          |  |
| AAG16628.1   | AY007545 | Brassica napus                         |
| protein serine/threonine kinase BNK1.  |          |  |
| BAB16871.1   | AP002537 | Oryza sativa                           |
| putative protein kinase APK1A Arabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192), D25110(R3192). |          |  |
| SEQ ID NO: 510   |          |  |
| AAG30254.1   | AF307333 | Hordeum vulgare                        |
| putative nematode-resistance protein. Hs1. similar to Beta procumbens Hs1pro protein.                        |          |  |
| AAB48305.1   | U79733   | Beta procumbens                        |
| nematode resistance. Hs1pro-1.   |          |  |
| SEQ ID NO: 513   |          |  |
| BAA14144.1   | D90116   | Armoracia rusticana                    |
| peroxidase isozyme.  |          |  |
| BAA14143.1   | D90115   | Armoracia rusticana                    |
| peroxidase isozyme.  |          |  |
| BAA11853.1   | D83225   | Populus nigra                          |
| peroxidase.  |          |  |
| CAA66037.1   | X97351   | Populus balsamifera subsp. trichocarpa |
| signal for ER. peroxidase.   |          |  |
| CAA66034.1   | X97348   | Populus balsamifera subsp. trichocarpa |
| signal for ER. peroxidase.   |          |  |
| CAA66036.1   | X97350   | Populus balsamifera subsp. trichocarpa |
| signal for ER. peroxidase.   |          |  |
| CAA66035.1   | X97349   | Populus balsamifera subsp. trichocarpa |
| signal for ER. peroxidase.   |          |  |
| BAA11852.1   | D83224   | Populus nigra                          |
| peroxidase.  |          |  |
| BAA07241.1   | D38051   | Populus kitakamiensis                  |
| peroxidase. prxA4a.  |          |  |
| BAA06335.1   | D30653   | Populus kitakamiensis                  |
| peroxidase.  |          |  |
| AAB47602.1   | L07554   | Linum usitatissimum                    |
| peroxidase. FLXPER1.   |          |  |
| AAC98519.1   | AF007211 | Glycine max                            |
| peroxidase precursor. GMIPER1. pathogen-induced.   |          |  |
| AAD37427.1   | AF149277 | Phaseolus vulgaris                     |
| peroxidase 1 precursor. FBP1. secretory peroxidase.  |          |  |
| BAA06334.1   | D30652   | Populus kitakamiensis                  |
| peroxidase.  |          |  |
| AAB97734.1   | AF014502 | Glycine max                            |
| seed coat peroxidase precursor. Ep. H2O2 oxidoreductase; class III plant peroxidase.                         |          |  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAB94692.1 | AJ242742 | Ipomoea batatas         | Removal of H <sub>2</sub> O <sub>2</sub> , oxidation of toxic reductants, defence response toward wounding.<br>peroxidase. pod.           |
| CAA62227.1 | X90694   | Medicago sativa         | peroxidase1C. prx1C.  |
| CAA62226.1 | X90693   | Medicago sativa         | peroxidase1B. prx1B.  |
| AAD37430.1 | AF149280 | Phaseolus vulgaris      | peroxidase 5 precursor. FBP5. secretory peroxidase.   |
| CAA62225.1 | X90692   | Medicago sativa         | peroxidase1A. prx1A.  |
| AAB41811.1 | L36157   | Medicago sativa         | peroxidase. pxdC. amino acid feature: conserved domains, aa 123 .. 129, 191 .. 198; amino acid feature: heme-binding domain, aa 68 .. 73. |
| AAB41810.1 | L36156   | Medicago sativa         | peroxidase. pxdA. amino acid feature: conserved motifs, aa 181 .. 188; amino acid feature: heme-binding domain, aa 60 .. 65.              |
| BAA01877.1 | D11102   | Populus kitakamiensis   | peroxidase. prxA1.  |
| CAB67121.1 | Y19023   | Lycopersicon esculentum | peroxidase. cevi-1.   |
| CAA50597.1 | X71593   | Lycopersicon esculentum | peroxidase. CEVI-1.   |
| BAA01992.1 | D11396   | Nicotiana tabacum       | 'peroxidase'.   |
| AAA34108.1 | J02979   | Nicotiana tabacum       | lignin-forming peroxidase precursor (EC 1.11.1.7).  |
| AAA33127.1 | M91373   | Cucumis sativus         | peroxidase. pre-peroxidase. putative.   |
| BAA92500.1 | AP001383 | Oryza sativa            | ESTs D39300(R3292),AU030751(E60187) correspond to a region of the predicted gene. Similar to peroxidase ATP6a. (X98774).                  |
| CAB65334.1 | AJ250121 | Picea abies             | peroxidase. SPI2 protein. spi2.   |
| CAA40796.1 | X57564   | Armoracia rusticana     | peroxidase. peroxidase precursor.   |
| AAB06183.1 | M37636   | Arachis hypogaea        | cationic peroxidase. PNC1.  |
| CAA76680.1 | Y17192   | Cucurbita pepo          | peroxidase. aprx. type III peroxidase.  |
| AAA33129.1 | M91372   | Cucumis sativus         | peroxidase. pre-peroxidase.   |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| CAA71492.1     | Y10466   | Spinacia oleracea       | peroxidase. prxr5.  |
| AAF63027.1     | AF244924 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.  |
| AAD43561.1     | AF155124 | Gossypium hirsutum      | bacterial-induced peroxidase precursor. Perx_Goshiko.   |
| AAF63026.1     | AF244923 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx14 precursor. type III peroxidase.  |
| AAA33121.1     | M32742   | Cucumis sativus         | peroxidase (CuPer2).  |
| BAA92422.1     | AP001366 | Oryza sativa            | ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene. Similar to A.thaliana mRNA for peroxidase ATP18a. (X98804).  |
| BAA92497.1     | AP001383 | Oryza sativa            | ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene. Similar to peroxidase ATP18a. (X98804).  |
| BAA77389.1     | AB024439 | Scutellaria baicalensis | peroxidase 3.   |
| BAA08499.1     | D49551   | Oryza sativa            | peroxidase. poxN.   |
| AAB19129.1     | U41657   | Glycine max             | seed coat peroxidase isozyme. SPOD4.1. H2O2 oxidoreductase.   |
| BAA03373.1     | D14482   | Oryza sativa            | putative peroxidase.  |
| AAB02554.1     | L37790   | Stylosanthes humilis    | cationic peroxidase.  |
| SEQ ID NO: 515 |          |                         |   |
| AAB88134.1     | AF034618 | Spinacia oleracea       | cytosolic heat shock 70 protein. HSC70-1.   |
| AAF34134.1     | AF161180 | Malus x domestica       | high molecular weight heat shock protein. Hsp2.   |
| AAB99745.1     | AF005993 | Triticum aestivum       | HSP70. TaHSP70d. 70 kDa heat shock protein, molecular chaperone.  |
| AAA62325.1     | L32165   | Hordeum vulgare         | Molecular chaperone. HSP70. Heat-shock protein HSP70; The predicted amino acid sequence is highly homologous (more than 80% identity) to other plant heat-shock proteins (HSP70s) in the database; however the C terminus is quite unique.; putative. |
| AAA21808.1     | L23551   | Spinacia oleracea       | molecular chaperone. ER-lumenal protein. HSC70.   |
| AAA34139.1     | L08830   | Lycopersicon esculentum | molecular chaperon (precursor). glucose-regulated protein 78. BiP/grp78. an endoplasmic reticulum residing heat shock protein 70 family member; precursor peptide.  |

|                |          |                       |   |
|----------------|----------|-----------------------|---|
| AAB86942.1     | AF031241 | Glycine max           | endoplasmic reticulum transport protein; molecular chaperone; roles in protein folding, assembly, and transport. endoplasmic reticulum HSC70-cognate binding protein precursor. BIP. BiP; similar to HSC70 and GRP78. |
| AAK21920.1     | AF338252 | Glycine max           | molecular chaperone. BiP-isoform D. BiPD. ER-luminal HSP70; binding protein GRP78.  |
| AAB91473.1     | AF035458 | Spinacia oleracea     | heat shock 70 protein. HSC70-11. mitochondrial protein.   |
| AAB96660.1     | AF039084 | Spinacia oleracea     | heat shock 70 protein. HSC70-11. molecular chaperone.   |
| AAB91472.1     | AF035457 | Spinacia oleracea     | heat shock 70 protein. HSC70-10. mitochondrial protein.   |
| SEQ ID NO: 516 |          |                       |   |
| CAA06927.1     | AJ006233 | Nicotiana tabacum     | putative thaumatin-like protein precursor.  |
| AAF06346.1     | AF195653 | Vitis vinifera        | SCUTL1. thaumatin-like protein.   |
| BAA28872.1     | AB006009 | Pyrus pyrifolia       | thaumatin-like protein precursor. PsTL1.  |
| CAC10270.1     | AJ243427 | Malus x domestica     | thaumatin-like protein. tl. allergen, pathogenesis-related.   |
| AAC36740.1     | AF090143 | Malus x domestica     | thaumatin-like protein precursor Mdtl1. MDTL1. pathogenesis-related.  |
| CAB62167.1     | AJ242828 | Castanea sativa       | antifungal. thaumatin-like protein. tl1.  |
| AAB38064.1     | U32440   | Prunus avium          | thaumatin-like protein precursor.   |
| BAA74546.2     | AB000834 | Nicotiana tabacum     | thaumatin-like protein SE39b.   |
| AAF06347.1     | AF195654 | Vitis vinifera        | SCUTL2. thaumatin-like protein.   |
| CAC09477.1     | AL442113 | Oryza sativa          | thaumatin-like protein. H0806H05.10.  |
| AAB95118.1     | U71244   | Brassica rapa         | pathogenesis-related group 5 protein. BFTP. thaumatin-like protein; PR-5.   |
| CAA10492.1     | AJ131731 | Pseudotsuga menziesii | Thaumatin-like protein. 5A1A.16.  |
| BAA95017.1     | AB031870 | Cestrum elegans       | thaumatin-like protein. CETLP.  |
| BAA95165.1     | AB029918 | Nicotiana tabacum     | pistil transmitting tissue specific thaumatin (SE39b)-like protein. SE39b.  |
| AAB61590.1     | AF003007 | Vitis vinifera        | VVTl1. osmotin-like protein, PR-5 protein; thaumatin-like protein.  |

---

AAD55090.1 AF178653 *Vitis riparia*  
thaumatin. osmotin; pathogenesis-related protein.

---

CAB85637.1 AJ237999 *Vitis vinifera*  
putative thaumatin-like protein. Tl1. alternative name grip 51.

---

AAF82264.1 AF227324 *Vitis vinifera*  
thaumatin-like protein.

---

AAB02259.1 U57787 *Avena sativa*  
permatin precursor. thaumatin-like protein.

---

AAB53368.1 U77657 *Oryza sativa*  
pathogenesis-related thaumatin-like protein.

---

CAA09228.1 AJ010501 *Cicer arietinum*  
thaumatin-like protein PR-5b.

---

CAA33293.1 X15224 *Nicotiana tabacum*  
thaumatin-like protein. E22.

---

CAA33292.1 X15223 *Nicotiana tabacum*  
thaumatin-like protein. E2.

---

## SEQ ID NO: 517

---

CAA71801.1 Y10848 *Brassica juncea*  
gamma-glutamylcysteine synthetase. gsh1.

---

AAB71230.1 AF017983 *Lycopersicon esculentum*  
gamma-glutamylcysteine synthetase. GSH1.

---

AAC82334.1 AF041340 *Medicago truncatula*  
gamma-glutamylcysteine synthetase. putative plastid protein.

---

AAF22137.1 AF128455 *Pisum sativum*  
gamma-glutamylcysteine synthetase precursor. gsh1. putative plastid protein.

---

AAF22136.1 AF128454 *Phaseolus vulgaris*  
gamma-glutamylcysteine synthetase precursor. gsh1. putative plastid protein.

---

CAA06613.1 AJ005587 *Brassica juncea*  
gamma-glutamylcysteine synthetase.

---

AAG13459.1 AF128453 *Glycine max*  
gamma-glutamylcysteine synthetase precursor. gsh1.

---

CAA64808.1 X95563 *Brassica juncea*  
gamma-glutamylcysteine synthetase. gsh1.

---

## SEQ ID NO: 518

---

AAA75414.1 L28005 *Glycine max*  
TGACG-motif-binding protein. STGA1.

---

AAB31250.2 S73827 *Solanum tuberosum*  
mas-binding factor MBF3. transcription factor TGA1a homolog; This sequence comes from Fig. 4.

---

AAB31249.1 S73826 *Solanum tuberosum*  
mas-binding factor MBF2. mas-binding factor MBF2. transcription factor TGA1a homolog; This sequence comes from Fig. 4.

---

|                |          |                   |   |
|----------------|----------|-------------------|---|
| CAA34468.1     | X16449   | Nicotiana sp.     | TGA1a protein (AA 1-359).   |
| AAA34091.1     | M62855   | Nicotiana tabacum | ASF-1/G13. leucine-zipper DNA-binding protein.  |
| AAB31251.2     | S73828   | Solanum tuberosum | mas-binding factor MBF1. transcription factor TGA1a homolog; This sequence comes from Fig. 4. |
| CAA48904.1     | X69152   | Zea mays          | ocs-element binding factor 3.2. OBF3.2.   |
| CAA48905.1     | X69153   | Zea mays          | ocs-element binding factor 3.1. OBF3.1.   |
| AAC24123.1     | AF067187 | Cichorium intybus | cAMP responsive element binding protein. bZIP transcription factor; CREB.                     |
| AAC24122.1     | AF067186 | Cichorium intybus | cAMP responsive element binding protein. CREB1.   |
| AAC49760.1     | AF001454 | Helianthus annuus | Dc3 promoter-binding factor-2. DPBF-2.  |
| SEQ ID NO: 521 |          |                   |   |
| CAA10608.1     | AJ132228 | Ricinus communis  | amino acid carrier. aap3.   |
| CAA07563.1     | AJ007574 | Ricinus communis  | amino acid carrier. aap1.   |
| CAA70778.1     | Y09591   | Vicia faba        | amino acid transporter.   |
| AAD16014.1     | AF080543 | Nepenthes alata   | amino acid transporter. AAP2.   |
| CAA70969.1     | Y09826   | Solanum tuberosum | amino acid transporter. AAP2. transmembrane protein.  |
| AAD16015.1     | AF080544 | Nepenthes alata   | amino acid transporter. AAP3.   |
| CAA70968.1     | Y09825   | Solanum tuberosum | amino acid transporter. AAP1. transmembrane protein.  |
| CAA92992.1     | Z68759   | Ricinus communis  | amino acid carrier.   |
| AAD16013.1     | AF080542 | Nepenthes alata   | amino acid transporter. AAP1.   |
| AAF15945.1     | AF061435 | Vicia faba        | amino acid transporter b. AAPB.   |
| CAA72006.1     | Y11121   | Ricinus communis  | amino acid carrier.   |
| AAF15944.1     | AF061434 | Vicia faba        | amino acid transporter a. AAPA.   |

AAF15946.1 AF061436 *Vicia faba*  
amino acid transporter c. AAPC.

AAB48944.1 U31932 *Nicotiana sylvestris*  
amino acid permease 1. NSAAP1. amino acid transporter; Method: conceptual translation  
supplied by author.

AAB96830.1 U64823 *Nicotiana sylvestris*  
amino acid transporter. amino acid permease. nsaap1.

BAA93437.1 AB022783 *Oryza sativa*  
amino acid permease. OsproT.

AAD25162.1 AF014810 *Lycopersicon esculentum*  
proline transporter 3. LeProT3.

AAD25161.1 AF014809 *Lycopersicon esculentum*  
proline transporter 2. LeProT2.

AAD25160.1 AF014808 *Lycopersicon esculentum*  
proline transporter 1. LeProT1.

AAF76897.1 AF274032 *Atriplex hortensis*  
proline/glycine betaine transporter.

CAB42599.1 AJ238635 *Chlorella protothecoides*  
amino acid carrier. dee4.

## SEQ ID NO: 526

BAA03763.1 D16247 *Nicotiana sylvestris*  
RNA helicase like protein DB10.

AAD46404.1 AF096248 *Lycopersicon esculentum*  
ethylene-responsive RNA helicase. ER68. putative DEAD box/RNA helicase.

AAF75791.1 AF271892 *Pisum sativum*  
DEAD box protein P68. P68. RNA helicase.

AAF40306.1 AF156667 *Vigna radiata*  
RNA helicase. VRH1.

CAA68193.1 X99937 *Spinacia oleracea*  
RNA helicase. prh75. DEAD-box protein; homologous to X99938.

AAG13612.1 AC078840 *Oryza sativa*  
putative RNA helicase. OSJNBb0073N24.12.

AAG34876.1 AF261021 *Nicotiana tabacum*  
putative chloroplast RNA helicase VDL isoform 1. VDL. alternatively spliced.

AAG34873.1 AF261020 *Nicotiana tabacum*  
putative chloroplast RNA helicase VDL isoform 1. VDL. essential for chloroplast  
development; may be involved in post-transcriptional regulation.

AAG34879.1 AF261024 *Nicotiana tabacum*  
putative chloroplast RNA helicase VDL' isoform 1. VDL'. alternatively spliced.

AAD20980.1 AF079782 *Zea mays*  
ATPase and RNA helicase. translation initiation factor 4A2. tif4A2.

AAG34882.1 AF261027 *Nicotiana tabacum*  
putative chloroplast RNA helicase VDL' isoform 4. VDL'. alternatively spliced.



|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAA95705.1     | AB042644 | Oryza sativa            | DEAD box RNA helicase OsPL10b. OsPL10b.  |
| BAA95704.1     | AB042643 | Oryza sativa            | DEAD box RNA helicase OsPL10a. OsPL10a.  |
| AAG34883.1     | AF261028 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 5. VDL'. alternatively spliced. |
| AAG38493.1     | AF261032 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 1. VDL'. alternatively spliced. |
| AAG34884.1     | AF261029 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 6. VDL'. alternatively spliced. |
| AAG38496.1     | AF261032 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 4. VDL'. alternatively spliced. |
| AAG38497.1     | AF261032 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 5. VDL'. alternatively spliced. |
| AAG38498.1     | AF261032 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 6. VDL'. alternatively spliced. |
| AAG38500.1     | AF261032 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 8. VDL'. alternatively spliced. |
| AAG34886.1     | AF261031 | Nicotiana tabacum       | putative chloroplast RNA helicase VDL' isoform 8. VDL'. alternatively spliced. |
| AAG48833.1     | AC084218 | Oryza sativa            | similar to Arabidopsis thaliana DNA helicase (AJ404475).                       |
| SEQ ID NO: 528 |          |                         |  |
| CAA65536.1     | X96761   | Sporobolus stapfianus   | sulphate transporter protein.  |
| AAK27688.1     | AF347614 | Lycopersicon esculentum | sulfate transporter 2. ST2.  |
| AAK27687.1     | AF347613 | Lycopersicon esculentum | sulfate transporter 1. ST1.  |
| CAA57711.1     | X82256   | Stylosanthes hamata     | high affinity sulphate transporter. SHST2.                                     |
| AAG41419.1     | AF309643 | Solanum tuberosum       | high affinity sulfate transporter type 1. ST1.                                 |
| CAA57710.1     | X82255   | Stylosanthes hamata     | high affinity sulphate transporter. SHST1.                                     |
| AAK35215.1     | AF355602 | Zea mays                | sulfate transporter ST1.   |
| CAA65291.1     | X96431   | Hordeum vulgare         | high affinity sulphate transporter. HVST1.                                     |
| AAA97952.1     | U52867   | Hordeum vulgare         | high affinity sulfate transporter HVST1.                                       |

|                |          |                              |   |
|----------------|----------|------------------------------|---|
| CAB42985.1     | AJ238244 | <i>Aegilops tauschii</i>     | putative plasma membrane sulfate transport. putative high affinity sulfate transporter. st1.  |
| CAB42986.1     | AJ238245 | <i>Aegilops tauschii</i>     | putative plasma membrane sulfate transport. putative high affinity sulfate transporter. st2.  |
| CAA57831.1     | X82454   | <i>Stylosanthes hamata</i>   | low affinity sulphate transporter. SHST3.   |
| SEQ ID NO: 531 |          |                              |   |
| AAK00436.1     | AC060755 | <i>Oryza sativa</i>          | putative zinc finger protein. OSJNBa0003O19.23.   |
| BAA85438.1     | AP000616 | <i>Oryza sativa</i>          | similar to RING-H2 finger protein RHA1a (AF078683).   |
| AAG43550.1     | AF211532 | <i>Nicotiana tabacum</i>     | Avr9/Cf-9 rapidly elicited protein 132. ACRE132. similar to RING finger proteins.   |
| BAA90357.1     | AP001080 | <i>Oryza sativa</i>          | EST AU070319(S10788) corresponds to a region of the predicted gene. Similar to RING-H2 finger protein RHA2b (AC006200).   |
| SEQ ID NO: 532 |          |                              |   |
| BAA81751.1     | AB017517 | <i>Marchantia polymorpha</i> | calcium-dependent protein kinase. CDPK1. CDPK-B; alternative splicing.  |
| BAA13232.1     | D87042   | <i>Zea mays</i>              | Calcium-dependent protein kinase.   |
| BAA81749.1     | AB017515 | <i>Marchantia polymorpha</i> | calcium-dependent protein kinase. CDPK1. CDPK-B; alternative splicing.  |
| BAA81750.1     | AB017516 | <i>Marchantia polymorpha</i> | calcium-dependent protein kinase. CDPK1. CDPK-A; alternative splicing.  |
| BAA85396.1     | AP000615 | <i>Oryza sativa</i>          | ESTs C22369(C12239),C22370(C12239), AU057852(S21844),AU057853(S21844) correspond to a region of the predicted gene.; similar to calcium dependent protein kinase. (AF048691). |
| CAA57156.1     | X81393   | <i>Oryza sativa</i>          | calcium-dependent protein kinase. OSCPKII.  |
| AAC05270.1     | AF048691 | <i>Oryza sativa</i>          | calcium dependent protein kinase. CDPK12.   |
| AAB49984.1     | U90262   | <i>Cucurbita pepo</i>        | calcium-dependent calmodulin-independent protein kinase CDPK. cpCPK1. serine/threonine protein kinase that is activated by direct binding of calcium.                         |
| AAB70706.1     | U82087   | <i>Tortula ruralis</i>       | calmodulin-like domain protein kinase. TrCPK1.  |
| AAG46110.1     | AC073166 | <i>Oryza sativa</i>          | calcium-dependent protein kinase. OSJNBb0064P21.2.  |
| BAA02698.1     | D13436   | <i>Oryza sativa</i>          | calcium-dependent protein kinase. spk.  |

|            |          |                               |  |
|------------|----------|-------------------------------|--|
| BAA81748.1 | AB017515 | Marchantia polymorpha         | calcium-dependent protein kinase. CDPK1. CDPK-A; alternative splicing.             |
| CAA07481.1 | AJ007366 | Zea mays                      | calcium-dependent protein kinase.  |
| AAB80692.1 | U69173   | Glycine max                   | calmodulin-like domain protein kinase isoenzyme beta. CDPK beta.                   |
| AAC49405.1 | U08140   | Vigna radiata                 | calcium dependent protein kinase. CDPK.  |
| BAA12338.1 | D84408   | Zea mays                      | calcium dependent protein kinase. ZmCDPK1.   |
| AAA33443.1 | L15390   | Zea mays                      | calcium-dependent protein kinase. CDPK.  |
| CAA57157.1 | X81394   | Oryza sativa                  | calcium-dependent protein kinase. OSCP2.   |
| BAB21081.1 | AP002819 | Oryza sativa                  | putative calcium-dependent protein kinase. P0501G01.10.                            |
| BAA12715.1 | D85039   | Zea mays                      | calcium-dependent protein kinase.  |
| CAA65500.1 | X96723   | Medicago sativa               | protein kinase. CDPK.  |
| AAC25423.1 | AF072908 | Nicotiana tabacum             | calcium-dependent protein kinase. CDPK1.   |
| AAA69507.1 | U28376   | Zea mays                      | calcium-dependent protein kinase. MZECDPK2.  |
| BAA13440.1 | D87707   | Ipomoea batatas               | calcium dependent protein kinase. CDPK.  |
| AAD28192.2 | AF115406 | Solanum tuberosum             | calcium-dependent protein kinase. CDPK; catalytic domain.                          |
| AAB80693.1 | U69174   | Glycine max                   | calmodulin-like domain protein kinase isoenzyme gamma. CDPK gamma.                 |
| AAA61682.1 | L27484   | Zea mays                      | calcium-dependent protein kinase. CDPK.  |
| AAD17800.1 | AF090835 | Mesembryanthemum crystallinum | Ca <sup>2+</sup> -dependent protein kinase. CPK1. serine/threonine protein kinase. |
| CAA39936.1 | X56599   | Daucus carota                 | calcium- dependent protein kinase. DcPK431.  |
| AAK26164.1 | AY027885 | Cucumis sativus               | calcium-dependent calmodulin-independent protein kinase 5. CDPK5. CsCDPK5.         |
| AAB88537.1 | AF035944 | Fragaria x ananassa           | calcium-dependent protein kinase. MAX17.   |
| AAF21062.1 | AF216527 | Dunaliella tertiolecta        | calcium-dependent protein kinase. CPK1; CDPK.                                      |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| CAA89202.1     | Z49233   | Chlamydomonas eugametos | calcium-stimulated protein kinase.  |
| AAC32116.1     | AF051211 | Picea mariana           | probable calcium dependent protein kinase. Sb15. similar to Vigna radiata calcium dependent protein kinase encoded by U08140.   |
| AAF23900.1     | AF194413 | Oryza sativa            | calcium-dependent protein kinase. CDPK1. OsCDPK1.   |
| AAF23901.2     | AF194414 | Oryza sativa            | calcium-dependent protein kinase. CDPK5. OsCDPK5.   |
| CAB46228.1     | Y18055   | Arachis hypogaea        | calcium dependent protein kinase. CDPK.   |
| AAC78558.1     | AF030879 | Solanum tuberosum       | protein kinase CPK1.  |
| CAA58750.1     | X83869   | Daucus carota           | CDPK-related protein kinase. CRK (or PK421).  |
| AAB47181.1     | S82324   | Zea mays                | /gene="calcium/calmodulin-dependent protein kinase. This sequence comes from Fig. 1.  |
| BAA12691.1     | D84507   | Zea mays                | CDPK-related protein kinase. Does not require calcium for its activity (by similarity).   |
| BAA22410.1     | D38452   | Zea mays                | calcium-dependent protein kinase-related kinase.  |
| BAA12692.1     | D84508   | Zea mays                | CDPK-related protein kinase. Does not require calcium for its activity.   |
| AAG01179.1     | AF289237 | Zea mays                | calcium/calmodulin dependent protein kinase MCK2. MCK2.   |
| AAC24961.1     | AF009337 | Tradescantia virginiana | CDPK-related protein kinase. CRK1.  |
| BAA90814.1     | AP001168 | Oryza sativa            | ESTs AU030197(E50746),AU030196(E50746) correspond to a region of the predicted gene.; Similar to calcium-dependent calmodulin-independent protein kinase CDPK (U90262). |
| AAC49008.1     | U24188   | Lilium longiflorum      | calcium/calmodulin-dependent phosphorylation activity. calcium/calmodulin-dependent protein kinase. CCaMK. serine/threonine kinase; binds to calcium and calmodulin.    |
| AAF19402.1     | AF203480 | Lycopersicon esculentum | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                    |
| SEQ ID NO: 535 |          |                         |   |
| AAK19619.1     | AF336286 | Gossypium hirsutum      | GHMYB9. ghmyb9. similar to myb.   |
| CAA64614.1     | X95296   | Lycopersicon esculentum | transcription factor. THM27. myb-related.   |
| CAA50224.1     | X70879   | Hordeum vulgare         | MybHv1. myb1.   |

|            |          |   |
|------------|----------|---|
| CAA50222.1 | X70877   | Hordeum vulgare<br>MybHv1. myb1.  |
| AAA33067.1 | L04497   | Gossypium hirsutum<br>MYB A; putative.  |
| CAA50221.1 | X70876   | Hordeum vulgare<br>MybHv5. myb2.  |
| BAA23337.1 | D88617   | Oryza sativa<br>transfactor. OSMYB1. Osmyb1.  |
| BAA23338.1 | D88618   | Oryza sativa<br>transfactor. OSMYB2. Osmyb2.  |
| AAC04720.1 | AF034134 | Gossypium hirsutum<br>putative MYB-like transcription factor. MYB-like DNA-binding domain protein. Cmy-O.<br>similar to MYB A encoded by GenBank Accession Number L04497.   |
| CAA72218.1 | Y11415   | Oryza sativa<br>myb.  |
| CAA50225.1 | X70880   | Hordeum vulgare<br>MybHv5. myb2.  |
| AAA82943.1 | U39448   | Picea mariana<br>MYB-like transcriptional factor MBF1. putative DNA binding region highly similar to the<br>maize C1.   |
| CAA78386.1 | Z13996   | Petunia x hybrida<br>DNA binding protein; transcriptional activator. protein 1. myb.Ph3. Product related to animal<br>myb proto-oncoproteins. Sequence from nucleotide 992 is not included in clone cPF1 and has<br>been obtained by PCR amplification of cDNA. |
| BAA81732.1 | AB029161 | Glycine max<br>GmMYB29A2.   |
| AAK19616.1 | AF336283 | Gossypium hirsutum<br>GHMYB25. ghmyb25. similar to myb; contains an unspliced intron.   |
| AAK19611.1 | AF336278 | Gossypium hirsutum<br>BNLGH1233. bnlghi6233. similar to myb.  |
| AAK19617.1 | AF336284 | Gossypium hirsutum<br>GHMYB36. ghmyb36. similar to myb.   |
| BAA93038.1 | AP001552 | Oryza sativa<br>EST AU082058(C12976) corresponds to a region of the predicted gene. Similar to<br>Arabidopsis thaliana putative transcription factor (AF062916).  |
| AAK19615.1 | AF336282 | Gossypium hirsutum<br>GHMYB10. ghmyb10. similar to myb.   |
| BAB39987.1 | AP003020 | Oryza sativa<br>putative transcription factor (myb). P0498A12.16. contains ESTs<br>AU097474(S5087),D40175(S1959).   |
| BAB39972.1 | AP003018 | Oryza sativa<br>putative transcription factor (myb). OSJNBa0004B13.27. contains ESTs<br>AU097474(S5087),D40175(S1959).  |

|            |          |                                   |  |
|------------|----------|-----------------------------------|--|
| CAB43399.1 | AJ006292 | <i>Antirrhinum majus</i>          | Myb-related transcription factor mixta-like 1. mybml1.   |
| CAA72185.1 | Y11350   | <i>Oryza sativa</i>               | myb factor. myb.   |
| AAF22256.1 | AF161711 | <i>Pimpinella brachycarpa</i>     | myb-related transcription factor.  |
| AAG13574.1 | AC037425 | <i>Oryza sativa</i>               | myb factor. OSJNBa0055P24.4.   |
| BAA81731.1 | AB029160 | <i>Glycine max</i>                | GmMYB29A1.   |
| BAA81730.1 | AB029159 | <i>Glycine max</i>                | GmMYB29A1.   |
| CAA72186.1 | Y11351   | <i>Oryza sativa</i>               | myb factor. myb.   |
| CAA67600.1 | X99210   | <i>Lycopersicon esculentum</i>    | myb-related transcription factor. THM16.   |
| CAA75509.1 | Y15219   | <i>Oryza sativa</i> subsp. indica | transcriptional activator. C1.   |
| AAC04718.1 | AF034132 | <i>Gossypium hirsutum</i>         | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. Cmy-J. similar to MYB A encoded by GenBank Accession Number L04497. |
| CAA78387.1 | Z13997   | <i>Petunia x hybrida</i>          | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.                                    |
| BAA81736.1 | AB029165 | <i>Glycine max</i>                | GmMYB29B2.   |
| AAK19618.1 | AF336285 | <i>Gossypium hirsutum</i>         | GHMYB38. ghmyb38. similar to myb.  |
| CAA72217.1 | Y11414   | <i>Oryza sativa</i>               | myb.   |
| AAC49394.1 | U57002   | <i>Zea mays</i>                   | P protein. P. allele P-wr; Myb-like transcriptional regulator with a putative zinc-finger at the C-terminal.                                     |
| AAA33500.1 | M73028   | <i>Zea mays</i>                   | myb-like transcription factor. P.  |
| BAB20661.1 | AP002871 | <i>Oryza sativa</i>               | putative myb-related protein P. P0475H04.31.   |
| AAG36774.1 | AF210616 | <i>Zea mays</i>                   | P2 protein. P2. myb-like transcriptional factor; similar to <i>Zea mays</i> P gene.  |
| BAA88222.1 | AB028650 | <i>Nicotiana tabacum</i>          | myb-related transcription factor LBM2. lbm2.   |
| BAA81733.2 | AB029162 | <i>Glycine max</i>                | GmMYB29A2.   |

BAA23339.1 D88619 *Oryza sativa*  
transfactor. OSMYB3. Osmyb3.

BAA88224.1 AB028652 *Nicotiana tabacum*  
myb-related transcription factor LBM4. lbm4.

BAA88221.1 AB028649 *Nicotiana tabacum*  
myb-related transcription factor LBM1. lbm1.

AAB41101.1 U72762 *Nicotiana tabacum*  
transcription factor Myb1. myb1. TMV-inducible Myb homolog; contains helix-turn-helix motif; contains redox-sensitive cysteine.

## SEQ ID NO: 536

CAA70968.1 Y09825 *Solanum tuberosum*  
amino acid transporter. AAP1. transmembrane protein.

AAF15946.1 AF061436 *Vicia faba*  
amino acid transporter c. AAPC.

CAA70969.1 Y09826 *Solanum tuberosum*  
amino acid transporter. AAP2. transmembrane protein.

AAB96830.1 U64823 *Nicotiana sylvestris*  
amino acid transporter. amino acid permease. nsaap1.

CAA07563.1 AJ007574 *Ricinus communis*  
amino acid carrier. aap1.

AAB48944.1 U31932 *Nicotiana sylvestris*  
amino acid permease 1. NSAAP1. amino acid transporter; Method: conceptual translation supplied by author.

AAD16015.1 AF080544 *Nepenthes alata*  
amino acid transporter. AAP3.

CAA70778.1 Y09591 *Vicia faba*  
amino acid transporter.

AAF15944.1 AF061434 *Vicia faba*  
amino acid transporter a. AAPA.

AAF15945.1 AF061435 *Vicia faba*  
amino acid transporter b. AAPB.

AAF76897.1 AF274032 *Atriplex hortensis*  
proline/glycine betaine transporter.

AAD16014.1 AF080543 *Nepenthes alata*  
amino acid transporter. AAP2.

AAD25161.1 AF014809 *Lycopersicon esculentum*  
proline transporter 2. LeProT2.

CAA10608.1 AJ132228 *Ricinus communis*  
amino acid carrier. aap3.

## SEQ ID NO: 537

BAA89009.1 AB027455 *Petunia x hybrida*  
anthocyanin 5-O-glucosyltransferase. PH1.

|            |          |                            |   |
|------------|----------|----------------------------|---|
| BAA36423.1 | AB013598 | Verbena x hybrida          | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| BAA36421.1 | AB013596 | Perilla frutescens         | UDP-glucose:anthocynin 5-O-glucosyltransferase. PF3R4.  |
| BAB07962.1 | AP002524 | Oryza sativa               | putative anthocyanin 5-O-glucosyltransferase. P0406H10.16. contains ESTs AU067881(C10481),AU067882(C10481).   |
| BAA36422.1 | AB013597 | Perilla frutescens         | UDP-glucose:anthocyanin 5-O-glucosyltransferase homologue. PF3R6.   |
| BAA93039.1 | AB033758 | Citrus unshiu              | limonoid UDP-glucosyltransferase. LGTase.   |
| AAF61647.1 | AF190634 | Nicotiana tabacum          | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| AAF98390.1 | AF287143 | Brassica napus             | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT. |
| AAK16175.1 | AC079887 | Oryza sativa               | putative glucosyltransferase. OSJNBa0040E01.15.   |
| AAF17077.1 | AF199453 | Sorghum bicolor            | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |
| AAK16172.1 | AC079887 | Oryza sativa               | putative glucosyltransferase. OSJNBa0040E01.14.   |
| BAA83484.1 | AB031274 | Scutellaria baicalensis    | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.   |
| AAK16181.1 | AC079887 | Oryza sativa               | putative glucosyltransferase. OSJNBa0040E01.16.   |
| AAG25643.1 | AF303396 | Phaseolus vulgaris         | UDP-glucosyltransferase HRA25. putative; defense associated.  |
| AAK16178.1 | AC079887 | Oryza sativa               | putative glucosyltransferase. OSJNBa0040E01.5.  |
| AAD21086.1 | AF127218 | Forsythia x intermedia     | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.   |
| BAA12737.1 | D85186   | Gentiana triflora          | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| AAK28303.1 | AF346431 | Nicotiana tabacum          | phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.  |
| CAB56231.1 | Y18871   | Dortheanthus bellidiformis | betanidin-5-O-glucosyltransferase.  |
| CAA54612.1 | X77462   | Manihot esculenta          | UTP-glucose glucosyltransferase. CGT5.  |



|                |          |                                 |  |
|----------------|----------|---------------------------------|--|
| BAB41019.1     | AB047092 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT1.  |
| BAB41020.1     | AB047093 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT2.  |
| AAB36653.1     | U32644   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS5a.  |
| BAB41025.1     | AB047098 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.  |
| BAB41023.1     | AB047096 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.  |
| BAB41022.1     | AB047095 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT2.  |
| BAB41021.1     | AB047094 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT1.  |
| BAA19659.1     | AB002818 | Perilla frutescens              | flavonoid 3-O-glucosyltransferase. UDP glucose.  |
| BAB41026.1     | AB047099 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT2.  |
| BAB41024.1     | AB047097 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT2.  |
| AAK16180.1     | AC079887 | Oryza sativa                    | putative glucosyltransferase. OSJNBa0040E01.21.  |
| AAK28304.1     | AF346432 | Nicotiana tabacum               | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.   |
| AAB36652.1     | U32643   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS10a.   |
| BAB41017.1     | AB047090 | Vitis labrusca x Vitis vinifera | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT1. The gene was derived from one of the parents V. labruscana cv. Ishiharawase. |
| BAA89008.1     | AB027454 | Petunia x hybrida               | anthocyanidin 3-O-glucosyltransferase. PGT8.   |
| CAA59450.1     | X85138   | Lycopersicon esculentum         | twil. homologous to glucosyltransferases.  |
| CAA54614.1     | X77464   | Manihot esculenta               | UTP-glucose glucosyltransferase. CGT7.   |
| CAA54611.1     | X77461   | Manihot esculenta               | UTP-glucose glucosyltransferase. CGT2.   |
| CAA54613.1     | X77463   | Manihot esculenta               | UTP-glucose glucosyltransferase. CGT6.   |
| SEQ ID NO: 540 |          |                                 |  |
| AAK27688.1     | AF347614 | Lycopersicon esculentum         | sulfate transporter 2. ST2.  |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAG41419.1     | AF309643 | Solanum tuberosum       | high affinity sulfate transporter type 1. ST1.   |
| AAK27687.1     | AF347613 | Lycopersicon esculentum | sulfate transporter 1. ST1.  |
| AAA97952.1     | U52867   | Hordeum vulgare         | high affinity sulfate transporter HVST1.   |
| CAA57711.1     | X82256   | Stylosanthes hamata     | high affinity sulphate transporter. SHST2.   |
| CAA65291.1     | X96431   | Hordeum vulgare         | high affinity sulphate transporter. HVST1.   |
| CAA57710.1     | X82255   | Stylosanthes hamata     | high affinity sulphate transporter. SHST1.   |
| CAB42985.1     | AJ238244 | Aegilops tauschii       | putative plasma membrane sulfate transport. putative high affinity sulfate transporter. st1. |
| AAK35215.1     | AF355602 | Zea mays                | sulfate transporter ST1.   |
| CAB42986.1     | AJ238245 | Aegilops tauschii       | putative plasma membrane sulfate transport. putative high affinity sulfate transporter. st2. |
| CAA65536.1     | X96761   | Sporobolus stapfianus   | sulphate transporter protein.  |
| CAA57831.1     | X82454   | Stylosanthes hamata     | low affinity sulphate transporter. SHST3.  |
| CAA11413.1     | AJ223495 | Brassica juncea         | sulfate permease. sp1.   |
| AAB94543.1     | AF016306 | Zea mays                | sulfate permease.  |
| SEQ ID NO: 541 |          |                         |  |
| AAF36491.1     | AF129479 | Hordeum vulgare         | HAK2. HAK2. similar to Hordeum vulgare K+ transporter HAK1.                                  |
| BAB32443.1     | AB055630 | Phragmites australis    | high-affinity potassium transporter. PcnHAK1.  |
| BAB32444.1     | AB055631 | Phragmites australis    | high-affinity potassium transporter. PceHAK1A.   |
| BAB32445.1     | AB055632 | Phragmites australis    | high-affinity potassium transporter. PceHAK1B.   |
| BAB32442.1     | AB055629 | Phragmites australis    | high-affinity potassium transporter. PcuHAK1.  |
| AAC39315.1     | AF025292 | Hordeum vulgare         | putative high-affinity potassium transporter. HvHAK1.  |
| AAF36497.1     | AF129485 | Oryza sativa            | HAK4. HAK4. OsHAK4; similar to Hordeum vulgare K+ transporter HAK1.                          |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAF36496.1     | AF129484 | Hordeum vulgare         | HAK4. HAK4. HvHAK4; similar to Hordeum vulgare K <sup>+</sup> transporter HAK1.                         |
| CAC14883.1     | AJ297888 | Hordeum vulgare         | putative potassium transporter. hak1.   |
| CAC14787.1     | AJ297886 | Hordeum vulgare         | putative potassium transporter. hak1.   |
| AAF36492.1     | AF129480 | Hordeum vulgare         | HAK1B. HAK1B. HvHAK1B; similar to Hordeum vulgare K <sup>+</sup> transporter HAK1.                      |
| CAC15061.1     | AJ300161 | Hordeum vulgare         | potassium transporter. hak4.  |
| SEQ ID NO: 542 |          |                         |   |
| AAA91063.1     | M88254   | Hevea brasiliensis      | ethylene-inducible protein. ER1.  |
| SEQ ID NO: 546 |          |                         |   |
| CAA75386.1     | Y15113   | Morinda citrifolia      | 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase. DS3. 2-dehydro-3-deoxyphosphoheptonate aldolase.  |
| CAA79855.1     | Z21792   | Lycopersicon esculentum | phospho-2-dehydro-3-deoxyheptonate aldolase.  |
| CAA79856.1     | Z21793   | Lycopersicon esculentum | phospho-2-dehydro-3-deoxyheptonate aldolase.  |
| SEQ ID NO: 548 |          |                         |   |
| BAA96751.1     | AP002521 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome4, BAC clone T16H5; lectin like protein (AL024486).           |
| SEQ ID NO: 551 |          |                         |   |
| BAB19096.1     | AP002839 | Oryza sativa            | putative DNA-binding protein homolog. P0688A04.2.   |
| BAB19075.1     | AP002744 | Oryza sativa            | putative DNA-binding protein homolog. P0006C01.17.  |
| AAK16170.1     | AC079887 | Oryza sativa            | putative DNA binding protein. OSJNBa0040E01.4.  |
| AAD32677.1     | AF140554 | Avena sativa            | DNA-binding protein WRKY1. wrky1. putative transcription factor.  |
| AAD16139.1     | AF096299 | Nicotiana tabacum       | DNA-binding protein 2. WRKY2. transcription factor.   |
| CAB97004.1     | AJ278507 | Solanum tuberosum       | putative transcription factor. WRKY DNA binding protein. WRKY1.   |
| CAA88326.1     | Z48429   | Avena fatua             | binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein.                      |
| AAC49527.1     | U48831   | Petroselinum crispum    | WRKY1. contains two WRKY domains; WRKY-type DNA-binding protein; sequence-specific DNA-binding protein. |

|                |          |                       |  |
|----------------|----------|-----------------------|--|
| AAC49529.1     | U58540   | Petroselinum crispum  | WRKY2. Contains two WRKY domains; WRKY-type DNA-binding protein.   |
| AAK16171.1     | AC079887 | Oryza sativa          | putative DNA-binding protein. OSJNBa0040E01.10.  |
| AAC37515.1     | L44134   | Cucumis sativus       | SPF1-like DNA-binding protein.   |
| AAD16138.1     | AF096298 | Nicotiana tabacum     | DNA-binding protein 1. WRKY1. transcription factor.  |
| AAD38283.1     | AC007789 | Oryza sativa          | putative WRKY DNA binding protein. OSJNBa0049B20.9.  |
| BAB18313.1     | AP002865 | Oryza sativa          | putative WRKY DNA binding protein. P0034C11.1. contains EST C26525(C12525).  |
| BAB40073.1     | AP003074 | Oryza sativa          | putative WRKY DNA binding protein. OSJNBa0004G10.20. contains EST C26525(C12525).                                    |
| AAC49528.1     | U56834   | Petroselinum crispum  | DNA-binding. WRKY3. WRKY-type DNA-binding protein.   |
| BAA77358.1     | AB020023 | Nicotiana tabacum     | WRKY domain Zn-finger type DNA-binding protein. DNA-binding protein NtWRKY3.   |
| AAG46150.1     | AC018727 | Oryza sativa          | putative DNA-binding protein. OSJNBa0056G17.18.  |
| CAA88331.1     | Z48431   | Avena fatua           | binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein.                                   |
| AAD32676.1     | AF140553 | Avena sativa          | DNA-binding protein WRKY3. wrky3. putative transcription factor.   |
| CAB66338.1     | AJ279697 | Betula pendula        | wrky-type DNA binding protein. wrky.   |
| AAF61863.1     | AF193770 | Nicotiana tabacum     | DNA-binding protein 3. WRKY3. transcription factor.  |
| AAF61864.1     | AF193771 | Nicotiana tabacum     | DNA-binding protein 4. WRKY4. transcription factor.  |
| AAD27591.1     | AF121354 | Petroselinum crispum  | binds sequence specifically to W Boxes (TTGACC). transcription factor. WRKY3. sequence specific DNA-binding protein. |
| BAA87069.1     | AB035271 | Matricaria chamomilla | elicitor-induced DNA-binding protein homolog. McWRKY1.   |
| SEQ ID NO: 553 |          |                       |  |
| AAF34428.1     | AF172282 | Oryza sativa          | receptor-like protein kinase. DUPR11.18.   |
| BAA94518.1     | AP001800 | Oryza sativa          | Similar to Arabidopsis thaliana chromosome 2 section 111 of 255; putative receptor-like protein kinase (AC002392).   |
| BAB07904.1     | AP002835 | Oryza sativa          | putative S-receptor kinase. P0417G05.12.   |

|            |          |                             |  |
|------------|----------|-----------------------------|--|
| BAA94516.1 | AP001800 | Oryza sativa                | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB07906.1 | AP002835 | Oryza sativa                | putative S-receptor kinase. P0417G05.14.   |
| BAA94517.1 | AP001800 | Oryza sativa                | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB07905.1 | AP002835 | Oryza sativa                | putative S-receptor kinase. P0417G05.13.   |
| BAA94528.1 | AP001800 | Oryza sativa                | Similar to Arabidopsis thaliana chromosome 2 BAC T20K24; putative receptor-like protein kinase (AC002392).       |
| BAA94529.2 | AP001800 | Oryza sativa                | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| CAA67145.1 | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.  |
| CAA73133.1 | Y12530   | Brassica oleracea           | serine /threonine kinase. ARLK.  |
| CAB89179.1 | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| AAA33008.1 | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| AAC23542.1 | U20948   | Ipomoea trifida             | receptor protein kinase. IRK1.   |
| BAA23676.1 | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.  |
| BAA92836.1 | AB032473 | Brassica oleracea           | S18 S-locus receptor kinase. SRK18.  |
| CAB41878.1 | Y18259   | Brassica oleracea           | SRK5 protein. SRK5. receptor-like kinase.  |
| CAB41879.1 | Y18260   | Brassica oleracea           | SRK15 protein. SRK15. receptor-like kinase.  |
| BAA21132.1 | D88193   | Brassica rapa               | S-receptor kinase. SRK9 (B.c).   |
| BAA06285.1 | D30049   | Brassica rapa               | S-receptor kinase SRK9.  |
| CAA74662.1 | Y14286   | Brassica oleracea           | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| CAA79355.1 | Z18921   | Brassica oleracea           | S-receptor kinase-like protein.  |
| BAB21001.1 | AB054061 | Brassica rapa               | S locus receptor kinase. SRK22.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAA33915.1     | L27821   | Oryza sativa            | receptor type serine/threonine kinase. protein kinase.          |
| BAA92954.1     | AP001551 | Oryza sativa            | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821). |
| SEQ ID NO: 554 |          |                         |   |
| CAC05658.1     | AJ250919 | Brassica napus          | endopolygalacturonase. pegaz.                                   |
| CAC05657.1     | AJ250918 | Brassica napus          | endopolygalacturonase. pgaz.                                    |
| CAA65072.1     | X95800   | Brassica napus          | polygalacturonase.  |
| CAA90272.1     | Z49971   | Brassica napus          | Hydrolytic enzyme. Polygalacturonase. pga.                      |
| CAA54448.1     | X77231   | Prunus persica          | polygalacturonase. PG.  |
| AAC14453.1     | L12019   | Actinidia deliciosa     | polygalacturonase.  |
| AAF71160.1     | AF152758 | Actinidia chinensis     | polygalacturonase A. PGA.                                       |
| AAA34178.1     | M37304   | Lycopersicon esculentum | polygalacturonase.  |
| CAA32235.1     | X14074   | Lycopersicon esculentum | polygalacturonase.  |
| CAA29148.1     | X05656   | Lycopersicon esculentum | polygalacturonase (AA 1-457).                                   |
| AAA32914.1     | L06094   | Persea americana        | cell wall degradation. polygalacturonase.                       |
| CAA47055.1     | X66426   | Persea americana        | polygalacturonase.  |
| AAC26512.1     | AF062467 | Cucumis melo            | polygalacturonase precursor. MPG3.                              |
| CAA11846.1     | AJ224147 | Rubus idaeus            | polygalacturonase. RAS3.  |
| AAF61444.1     | AF138858 | Lycopersicon esculentum | hydrolyses polygalacturonic acid. polygalacturonase. XOPG1.     |
| BAA88472.1     | AB035890 | Cucumis sativus         | polygalacturonase. CUPG1.                                       |
| AAD46483.1     | AF128266 | Glycine max             | polygalacturonase PG1.  |
| AAD46484.1     | AF128267 | Glycine max             | polygalacturonase PG2.  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAC28905.1 | AF001002 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 4. TAPG4. expressed in abscission.    |
| AAC28906.1 | AF001003 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 5. TAPG5. expressed in abscission.    |
| AAC28903.1 | AF001000 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 1. TAPG1. expressed in abscission.    |
| AAA80489.1 | U23053   | Lycopersicon esculentum | polygalacturonase precursor.  |
| AAC64184.1 | AF095577 | Prunus persica          | endopolygalacturonase.  |
| AAC28902.2 | AF000999 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 3. TAPG3. expressed in abscission.    |
| AAB09575.1 | U70480   | Lycopersicon esculentum | abscission polygalacturonase. TAPG2.  |
| AAC28904.1 | AF001001 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 2. TAPG2. expressed in abscission.    |
| CAA54150.1 | X76735   | Prunus persica          | endopolygalacturonase.  |
| AAC70951.1 | AF072732 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella. polygalacturonase 7. specifically and abundantly expressed in pistils. |
| AAC26511.1 | AF062466 | Cucumis melo            | polygalacturonase precursor. MPG2.  |
| AAB09576.1 | U70481   | Lycopersicon esculentum | abscission polygalacturonase. TAPG4.  |
| AAC28947.1 | AF029230 | Lycopersicon esculentum | polygalacturonase. TPG6.  |
| AAA82167.1 | U09717   | Gossypium hirsutum      | polygalacturonase.  |
| AAA58322.1 | U09805   | Gossypium barbadense    | polygalacturonase.  |
| CAA47052.1 | X66422   | Zea mays                | polygalacturonase. PG.  |
| AAG14416.1 | AF248538 | Nicotiana tabacum       | NTS1 protein. similar to polygalacturonase.   |
| CAB42886.1 | AJ238848 | Phleum pratense         | polygalacturonase. pg.  |

|                |          |                    |   |
|----------------|----------|--------------------|---|
| CAA40850.1     | X57627   | Zea mays           | polygalacturonase.                                      |
| CAA46679.1     | X65844   | Zea mays           | polygalacturonase. PGg6.                                |
| CAA40910.1     | X57743   | Zea mays           | polygalacturonase.                                      |
| CAA44249.1     | X62385   | Zea mays           | polygalacturonase.                                      |
| CAA45751.1     | X64408   | Zea mays           | polygalacturonase. PG. pollen-preferentially expressed. |
| CAA40851.1     | X57628   | Zea mays           | polygalacturonase.                                      |
| CAA46680.1     | X65845   | Zea mays           | polygalacturonase. PGg14.                               |
| AAC26510.1     | AF062465 | Cucumis melo       | polygalacturonase precursor. MPG1.                      |
| SEQ ID NO: 556 |          |                    |   |
| CAA51734.1     | X73301   | Vigna mungo        | alpha-amylase. amyVm1.                                  |
| CAA37217.1     | X53049   | Vigna mungo        | alpha-amylase (AA 1-421).                               |
| BAA33879.1     | AB015131 | Phaseolus vulgaris | alpha-amylase.  |
| AAA16513.1     | U06754   | Cuscuta reflexa    | starch hydrolysis. alpha amylase precursor. CUS AMY2.   |
| AAA98615.1     | J04202   | Hordeum vulgare    | alpha-amylase. Amy46. precursor.                        |
| AAA98790.1     | K02637   | Hordeum vulgare    | alpha-amylase type B. Amy6-4. precursor.                |
| CAA33298.1     | X15226   | Hordeum vulgare    | alpha-amylase.  |
| AAA33885.1     | M24286   | Oryza sativa       | alpha-amylase (EC 3.2.1.1).                             |
| CAA34516.1     | X16509   | Oryza sativa       | alpha-amylase.  |
| CAA72144.1     | Y11277   | Hordeum vulgare    | alpha-amylase. amy.                                     |
| CAA33299.1     | X15227   | Hordeum vulgare    | alpha amylase.  |
| AAA32925.1     | M17126   | Hordeum vulgare    | alpha-amylase 1.  |



|            |          |  |
|------------|----------|--|
| AAA32927.1 | M17128   | Hordeum vulgare<br>alpha-amylase 2.  |
| AAA32929.1 | J01236   | Hordeum vulgare<br>alpha-amylase type A, EC 3.2.1.1.                             |
| CAA39777.1 | X56337   | Oryza sativa<br>alpha-amylase. RAm3B.  |
| CAA09323.1 | AJ010728 | Avena fatua<br>alpha amylase. alpha-Amy2A.                                       |
| AAA33897.1 | M24941   | Oryza sativa<br>alpha-amylase precursor (EC 3.2.1.1).                            |
| CAA09324.1 | AJ010729 | Avena fatua<br>alpha-amylase. alpha-Amy2D.                                       |
| CAA28803.1 | X05166   | Hordeum vulgare<br>alpha-amylase type A. Amy32b.                                 |
| AAA32926.1 | M17125   | Hordeum vulgare<br>alpha-amylase 1.  |
| CAA39778.1 | X56338   | Oryza sativa<br>alpha-amylase. RAm3C.  |
| AAA32935.1 | M15208   | Hordeum vulgare<br>alpha-amylase.  |
| AAA33895.1 | M59351   | Oryza sativa<br>alpha-amylase. RAm3D.  |
| AAA33896.1 | M59352   | Oryza sativa<br>alpha-amylase. RAm3E.  |
| AAA50161.1 | L25805   | Zea mays<br>alpha-amylase.   |
| AAA33894.1 | M74177   | Oryza sativa<br>alpha-amylase. amy2A.  |
| CAA72143.1 | Y11276   | Hordeum vulgare<br>alpha-amylase. amy.   |
| CAA39776.1 | X56336   | Oryza sativa<br>alpha-amylase. RAm3A.  |
| CAA45903.1 | X64619   | Oryza sativa<br>alpha-amylase. Amyc2. starts hydrolysis during seed germination. |
| AAA33886.1 | M24287   | Oryza sativa<br>alpha-amylase (EC 3.2.1.1).                                      |
| AAA32928.1 | M17127   | Hordeum vulgare<br>alpha-amylase 2.  |
| AAA32933.1 | K02638   | Hordeum vulgare<br>pre-alpha-amylase type B, EC 3.2.1.1.                         |
| AAA91883.1 | M81682   | Solanum tuberosum<br>alpha-amylase. Amy23.                                       |

|                |          |   |
|----------------|----------|---|
| AAA34259.1     | M16991   | Triticum aestivum<br>alpha-amylase.   |
| CAA29252.1     | X05809   | Triticum aestivum<br>alpha-amylase.   |
| AAA91884.1     | M79328   | Solanum tuberosum<br>alpha-amylase.   |
| AAF63239.1     | AF153828 | Malus x domestica<br>degrades starch. alpha-amylase. alpha-amylase by similarity. |
| AAA32930.1     | M10056   | Hordeum vulgare<br>alpha-amylase type B, EC 3.2.1.1.                              |
| AAA32931.1     | K02635   | Hordeum vulgare<br>alpha-amylase type B, EC 3.2.1.1.                              |
| CAA36485.1     | X52240   | Oryza sativa<br>alpha-amylase. OSamy-c.   |
| AAA32932.1     | K02636   | Hordeum vulgare<br>alpha-amylase type B, EC 3.2.1.1.                              |
| SEQ ID NO: 557 |          |   |
| CAA90272.1     | Z49971   | Brassica napus<br>Hydrolytic enzyme. Polygalacturonase. pga.                      |
| CAA65072.1     | X95800   | Brassica napus<br>polygalacturonase.  |
| CAA67020.1     | X98373   | Brassica napus<br>endo-polygalacturonidase.                                       |
| CAC05658.1     | AJ250919 | Brassica napus<br>endopolygalacturonase. pegaz.                                   |
| CAC05657.1     | AJ250918 | Brassica napus<br>endopolygalacturonase. pgaz.                                    |
| AAC14453.1     | L12019   | Actinidia deliciosa<br>polygalacturonase.   |
| AAF71160.1     | AF152758 | Actinidia chinensis<br>polygalacturonase A. PGA.                                  |
| CAA54448.1     | X77231   | Prunus persica<br>polygalacturonase. PG.  |
| AAA34178.1     | M37304   | Lycopersicon esculentum<br>polygalacturonase.                                     |
| CAA29148.1     | X05656   | Lycopersicon esculentum<br>polygalacturonase (AA 1-457).                          |
| CAA32235.1     | X14074   | Lycopersicon esculentum<br>polygalacturonase.                                     |
| AAA32914.1     | L06094   | Persea americana<br>cell wall degradation. polygalacturonase.                     |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAA47055.1 | X66426   | Persea americana        | polygalacturonase.  |
| AAC26512.1 | AF062467 | Cucumis melo            | polygalacturonase precursor. MPG3.  |
| CAA11846.1 | AJ224147 | Rubus idaeus            | polygalacturonase. RAS3.  |
| BAA88472.1 | AB035890 | Cucumis sativus         | polygalacturonase. CUPG1.   |
| AAF61444.1 | AF138858 | Lycopersicon esculentum | hydrolyses polygalacturonic acid. polygalacturonase. XOPG1.   |
| AAA80489.1 | U23053   | Lycopersicon esculentum | polygalacturonase precursor.  |
| AAC28903.1 | AF001000 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 1. TAPG1. expressed in abscission.    |
| AAB09575.1 | U70480   | Lycopersicon esculentum | abscission polygalacturonase. TAPG2.  |
| AAC28904.1 | AF001001 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 2. TAPG2. expressed in abscission.    |
| AAC28905.1 | AF001002 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 4. TAPG4. expressed in abscission.    |
| AAC64184.1 | AF095577 | Prunus persica          | endopolygalacturonase.  |
| AAD46483.1 | AF128266 | Glycine max             | polygalacturonase PG1.  |
| AAC28906.1 | AF001003 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 5. TAPG5. expressed in abscission.    |
| AAD46484.1 | AF128267 | Glycine max             | polygalacturonase PG2.  |
| CAA54150.1 | X76735   | Prunus persica          | endopolygalacturonase.  |
| AAC26511.1 | AF062466 | Cucumis melo            | polygalacturonase precursor. MPG2.  |
| AAC28902.2 | AF000999 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella of plant cells. polygalacturonase 3. TAPG3. expressed in abscission.    |
| AAC70951.1 | AF072732 | Lycopersicon esculentum | hydrolyzes pectin in the cell wall and middle lamella. polygalacturonase 7. specifically and abundantly expressed in pistils. |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA47052.1     | X66422   | Zea mays                | polygalacturonase. PG.   |
| AAC28947.1     | AF029230 | Lycopersicon esculentum | polygalacturonase. TPG6.   |
| AAA82167.1     | U09717   | Gossypium hirsutum      | polygalacturonase.   |
| AAB09576.1     | U70481   | Lycopersicon esculentum | abscission polygalacturonase. TAPG4.                                     |
| CAA40910.1     | X57743   | Zea mays                | polygalacturonase.   |
| CAA44249.1     | X62385   | Zea mays                | polygalacturonase.   |
| CAA40850.1     | X57627   | Zea mays                | polygalacturonase.   |
| AAG14416.1     | AF248538 | Nicotiana tabacum       | NTS1 protein. similar to polygalacturonase.                              |
| CAA46679.1     | X65844   | Zea mays                | polygalacturonase. PGg6.   |
| AAC26510.1     | AF062465 | Cucumis melo            | polygalacturonase precursor. MPG1.                                       |
| AAA58322.1     | U09805   | Gossypium barbadense    | polygalacturonase.   |
| CAA46680.1     | X65845   | Zea mays                | polygalacturonase. PGg14.  |
| CAA40851.1     | X57628   | Zea mays                | polygalacturonase.   |
| CAA45751.1     | X64408   | Zea mays                | polygalacturonase. PG. pollen-preferentially expressed.                  |
| CAB42886.1     | AJ238848 | Phleum pratense         | polygalacturonase. pg.   |
| SEQ ID NO: 559 |          |                         |  |
| CAB43937.1     | AJ006348 | Fragaria x ananassa     | cell wall hydrolysis. endo-beta-1,4-glucanase. egl.                      |
| AAC95009.1     | AF074923 | Fragaria x ananassa     | endo-1,4-beta-glucanase precursor. Cell. 1,4-beta-glucanohydrolase.      |
| AAC12684.1     | U76725   | Pinus radiata           | endo-beta-1,4-glucanase. PrCell. cellulase; PRCEL1.                      |
| AAA69909.1     | U13055   | Lycopersicon esculentum | cell wall hydrolase. endo-1,4-beta-glucanase precursor. Cel2. cellulase. |
| CAA65600.1     | X96856   | Prunus persica          | endo-beta-1,4-glucanase. ppEG1.  |

|            |          |   |
|------------|----------|---|
| CAA65597.1 | X96853   | <i>Prunus persica</i><br>endo-beta-1,4-glucanase. pcell.  |
| CAA65827.1 | X97189   | <i>Capsicum annuum</i><br>endo-beta-1,4-glucanase. ccel3. cellulase.  |
| AAC62241.1 | AF077339 | <i>Lycopersicon esculentum</i><br>endo-1,4-beta-glucanase. cel5. cellulase; also expressed in fruit and pistils.              |
| BAA85150.1 | AB032830 | <i>Pisum sativum</i><br>endo-1,4-beta-glucanase. EGL2.  |
| AAA80495.1 | U20590   | <i>Lycopersicon esculentum</i><br>endo-1,4-beta-glucanase precursor. cellulase.   |
| BAB32662.1 | AB055886 | <i>Atriplex lentiformis</i><br>beta-1,4-glucanase. Al-cell1. cellulase.   |
| CAA65828.1 | X97190   | <i>Capsicum annuum</i><br>endo-beta-1,4-glucanase. ccel2.   |
| AAC12685.1 | U76756   | <i>Pinus radiata</i><br>endo-beta-1,4-glucanase. PrCel2. cellulase; PRCEL2.   |
| BAA77239.1 | AB025796 | <i>Populus alba</i><br>endo-1,4-beta glucanase. POPCEL2. cellulase.   |
| CAB59900.1 | AJ010950 | <i>Capsicum annuum</i><br>cell wall degradation. endo-beta-1,4-glucanase. eg2.  |
| BAB39483.1 | AB049200 | <i>Populus alba</i><br>endo-1,4-beta-glucanase. PopCel2.  |
| BAB39482.1 | AB049199 | <i>Populus alba</i><br>endo-1,4-beta glucanase. PopCel1.  |
| CAA72133.1 | Y11268   | <i>Lycopersicon esculentum</i><br>endo-1,4-beta-D-glucanase. cel7.  |
| AAA96135.1 | L41046   | <i>Pisum sativum</i><br>endo-1,4-beta-glucanase. EGL1.  |
| AAC78504.1 | U34754   | <i>Phaseolus vulgaris</i><br>endo-1,4-beta-D-glucanase. cellulase.  |
| AAA02563.1 | M57400   | <i>Phaseolus vulgaris</i><br>cellulase precursor. endo-1,4-beta-D-glucanase.  |
| CAA60737.1 | X87323   | <i>Capsicum annuum</i><br>catalyzes hydrolysis of cell wall polysaccharides. cellulase. cell1. Beta-1,4-endoglycanohydrolase. |
| CAA65826.1 | X97188   | <i>Capsicum annuum</i><br>endo-beta-1,4-glucanase. ccell1. cellulase.   |
| AAA69908.1 | U13054   | <i>Lycopersicon esculentum</i><br>cell wall hydrolase. endo-1,4-beta-glucanase precursor. Cell1. cellulase.                   |
| CAB43938.1 | AJ006349 | <i>Fragaria x ananassa</i><br>cell wall hydrolysis. endo-beta-1,4-glucanase. eg3.   |

---

BAA96209.1 AP002094 *Oryza sativa*  
EST C72268(E1328) corresponds to a region of the predicted gene. Similar to *Fragaria x ananassa* endo-beta-1,4-glucanase (AJ006349).

---

BAA96207.1 AP002094 *Oryza sativa*  
EST C72268(E1328) corresponds to a region of the predicted gene. Similar to *Fragaria x ananassa* endo-beta-1,4-glucanase (AJ006349).

---

AAD08699.1 AF098292 *Lycopersicon esculentum*  
endo-beta-1,4-D-glucanase. Cel8.

---

BAA94257.1 AB040769 *Hordeum vulgare*  
endo-1,4-beta-glucanase Cel1. Cel1.

---

AAC49704.1 U78526 *Lycopersicon esculentum*  
endo-1,4-beta-glucanase. Cel3.

---

CAB51903.1 AJ242807 *Brassica napus*  
endo-1,4-beta-D-glucanase. Cell6. cellulase.

---

AAA20082.1 U00730 *Glycine max*  
CMCase; cellulase; endo-1,4-beta-D-glucanase.

---

CAA11301.1 AJ223386 *Fragaria x ananassa*  
endo-beta-1,4-glucanase. faEG1. partial cDNA obtained by RT-PCR amplification.

---

CAA11302.1 AJ223387 *Fragaria x ananassa*  
endo-beta-1,4-glucanase. partial cDNA obtained by RT-PCR amplification.

---

BAA21111.1 D88417 *Gossypium hirsutum*  
endo-1,4-beta-glucanase.

---

AAA20083.1 U00731 *Glycine max*  
CMCase; cellulase; endo-1,4-beta-D-glucanase.

---

SEQ ID NO: 560

---

AAB97617.1 U83687 *Apium graveolens*  
NADPH-dependent mannose 6-phosphate reductase. m6pr. aldo-keto reductase; similar to aldose 6-phosphate reductase also known as NADP-sorbitol-6-phosphate dehydrogenase encoded by GenBank Accession Number D11080.

---

AAC97607.1 AF057134 *Malus x domestica*  
synthesizes sorbitol, a major photosynthetic product in many members of the Rosaceae family. NADP-dependent sorbitol 6-phosphate dehydrogenase. S6PDH.

---

BAA01853.1 D11080 *Malus x domestica*  
NADP-dependent D-sorbitol-6-phosphate dehydrogenase. S6PDH.

---

SEQ ID NO: 561

---

CAB43938.1 AJ006349 *Fragaria x ananassa*  
cell wall hydrolysis. endo-beta-1,4-glucanase. eg3.

---

BAA96209.1 AP002094 *Oryza sativa*  
EST C72268(E1328) corresponds to a region of the predicted gene. Similar to *Fragaria x ananassa* endo-beta-1,4-glucanase (AJ006349).

---

BAA96207.1 AP002094 *Oryza sativa*  
EST C72268(E1328) corresponds to a region of the predicted gene. Similar to *Fragaria x ananassa* endo-beta-1,4-glucanase (AJ006349).

---

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAD08699.1 | AF098292 | Lycopersicon esculentum | endo-beta-1,4-D-glucanase. Cel8.   |
| BAA21111.1 | D88417   | Gossypium hirsutum      | endo-1,4-beta-glucanase.   |
| CAA65828.1 | X97190   | Capsicum annuum         | endo-beta-1,4-glucanase. ccel2.  |
| CAB59900.1 | AJ010950 | Capsicum annuum         | cell wall degradation. endo-beta-1,4-glucanase. eg2.                           |
| AAC95009.1 | AF074923 | Fragaria x ananassa     | endo-1,4-beta-glucanase precursor. Cell1. 1,4-beta-glucanohydrolase.           |
| CAB43937.1 | AJ006348 | Fragaria x ananassa     | cell wall hydrolysis. endo-beta-1,4-glucanase. eg1.                            |
| AAA80495.1 | U20590   | Lycopersicon esculentum | endo-1,4-beta-glucanase precursor. cellulase.                                  |
| BAA85150.1 | AB032830 | Pisum sativum           | endo-1,4-beta-glucanase. EGL2.   |
| AAC12684.1 | U76725   | Pinus radiata           | endo-beta-1,4-glucanase. PrCell1. cellulase; PRCEL1.                           |
| BAB32662.1 | AB055886 | Atriplex lentiformis    | beta-1,4-glucanase. Al-cell1. cellulase.                                       |
| AAC62241.1 | AF077339 | Lycopersicon esculentum | endo-1,4-beta-glucanase. cel5. cellulase; also expressed in fruit and pistils. |
| CAA65826.1 | X97188   | Capsicum annuum         | endo-beta-1,4-glucanase. ccell1. cellulase.                                    |
| AAA69909.1 | U13055   | Lycopersicon esculentum | cell wall hydrolase. endo-1,4-beta-glucanase precursor. Cel2. cellulase.       |
| CAA65597.1 | X96853   | Prunus persica          | endo-beta-1,4-glucanase. pcell1.   |
| CAA65600.1 | X96856   | Prunus persica          | endo-beta-1,4-glucanase. ppEG1.  |
| AAC12685.1 | U76756   | Pinus radiata           | endo-beta-1,4-glucanase. PrCel2. cellulase; PRCEL2.                            |
| BAB39482.1 | AB049199 | Populus alba            | endo-1,4-beta glucanase. PopCell1.   |
| CAA65827.1 | X97189   | Capsicum annuum         | endo-beta-1,4-glucanase. ccel3. cellulase.                                     |
| BAB39483.1 | AB049200 | Populus alba            | endo-1,4-beta-glucanase. PopCel2.  |
| BAA77239.1 | AB025796 | Populus alba            | endo-1,4-beta glucanase. POPCEL2. cellulase.                                   |
| AAA69908.1 | U13054   | Lycopersicon esculentum | cell wall hydrolase. endo-1,4-beta-glucanase precursor. Cell1. cellulase.      |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAC78504.1     | U34754   | Phaseolus vulgaris      | endo-1,4-beta-D-glucanase. cellulase.  |
| AAA02563.1     | M57400   | Phaseolus vulgaris      | cellulase precursor. endo-1,4-beta-D-glucanase.                                |
| CAA72133.1     | Y11268   | Lycopersicon esculentum | endo-1,4-beta-D-glucanase. cel7.   |
| AAA96135.1     | L41046   | Pisum sativum           | endo-1,4-beta-glucanase. EGL1.   |
| CAB51903.1     | AJ242807 | Brassica napus          | endo-1,4-beta-D-glucanase. Cell6. cellulase.                                   |
| AAC49704.1     | U78526   | Lycopersicon esculentum | endo-1,4-beta-glucanase. Cel3.   |
| BAA94257.1     | AB040769 | Hordeum vulgare         | endo-1,4-beta-glucanase Cell. Cell.  |
| CAA11302.1     | AJ223387 | Fragaria x ananassa     | endo-beta-1,4-glucanase. partial cDNA obtained by RT-PCR amplification.        |
| AAA20082.1     | U00730   | Glycine max             | CMCase; cellulase; endo-1,4-beta-D-glucanase.                                  |
| CAA11301.1     | AJ223386 | Fragaria x ananassa     | endo-beta-1,4-glucanase. faEG1. partial cDNA obtained by RT-PCR amplification. |
| AAC64045.1     | AF077340 | Lycopersicon esculentum | endo-1,4-beta-glucanase. cel5. cellulase.                                      |
| SEQ ID NO: 562 |          |                         |  |
| AAD53011.1     | AF089848 | Brassica napus          | senescence-specific cysteine protease. SAG12-1. BnSAG12-1.                     |
| AAD53012.1     | AF089849 | Brassica napus          | senescence-specific cysteine protease. SAG12-2. BnSAG12-2.                     |
| AAK27968.1     | AF242372 | Ipomoea batatas         | cysteine protease. SPCP1.  |
| AAA50755.1     | U13940   | Alnus glutinosa         | cysteine proteinase. putative preproprotein.                                   |
| BAB13759.1     | AB040454 | Astragalus sinicus      | cysteine proteinase. AsNODf32. preproprotein putative.                         |
| AAC62396.1     | AF050756 | Ricinus communis        | cysteine endopeptidase precursor. CysEP.                                       |
| CAB09698.1     | Z97022   | Hordeum vulgare         | cysteine proteinase. putative.   |
| CAA52425.1     | X74406   | Hemerocallis sp.        | thiol-protease. SEN102.  |
| BAA83472.1     | AB004648 | Oryza sativa            | cysteine endopeptidase. RepA.  |



|            |          |                              |   |
|------------|----------|------------------------------|---|
| CAA56844.1 | X80876   | Oryza sativa                 | cysteine protease.  |
| BAA88898.1 | AB020961 | Zea mays                     | cysteine protease component of protease-inhibitor complex. CPPIC.   |
| AAC35211.1 | U12637   | Hemerocallis hybrid cultivar | cysteine proteinase. SEN11.   |
| CAB09697.1 | Z97021   | Hordeum vulgare              | cysteine endopeptidase EP-A. precursor.   |
| AAB88263.1 | AF019147 | Zea mays                     | cysteine proteinase Mir3. mir3.   |
| AAD28477.1 | AF133839 | Sandersonia aurantiaca       | papain-like cysteine protease. PRT5. senescence-related.  |
| CAB16317.1 | Z99173   | Nicotiana tabacum            | storage protein hydrolysis. cysteine proteinase precursor.  |
| CAB09699.1 | Z97023   | Hordeum vulgare              | cysteine endopeptidase EP-A.  |
| AAD10337.1 | U94591   | Hordeum vulgare              | cysteine proteinase precursor. EPA.   |
| CAA06243.1 | AJ004958 | Pisum sativum                | thiol-protease. pre-pro-TPE4A protein. tpE4A.   |
| CAB53515.1 | AJ245924 | Solanum tuberosum            | proteolysis. cysteine protease. cyp.  |
| AAD48496.1 | AF172856 | Lycopersicon esculentum      | cysteine protease TDI-65. tdi-65. induced by drought; localized in the nuclei and chloroplast (Tabaeizadeh, Z. et al., 1995. Protoplasma, 186:208-219). |
| CAA05894.1 | AJ003137 | Lycopersicon esculentum      | cysteine protease. CYP1. C14.   |
| AAA79915.1 | U17135   | Dianthus caryophyllus        | cysteine proteinase. DCCP1. expressed in senescing flower petals.   |
| AAB37233.1 | U34747   | Phalaenopsis sp. SM9108      | cysteine proteinase.  |
| CAB17076.1 | Z99954   | Phaseolus vulgaris           | protein hydrolysis. cysteine proteinase precursor.  |
| CAA84378.1 | Z34895   | Vicia sativa                 | storage protein degradation. cysteine proteinase.   |
| CAA53377.1 | X75749   | Vicia sativa                 | storage protein hydrolysis. cysteine protease.  |
| CAB17074.1 | Z99952   | Phaseolus vulgaris           | degradation of storage proteins. cysteine proteinase precursor.   |
| AAB68374.1 | U52970   | Phaseolus vulgaris           | suggested to initiate phaseolin degradation during germination. cysteine endopeptidase 1. PvCEP-1.  |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA12118.1     | AJ224766 | Phaseolus vulgaris      | phaseolin degradation. cysteine protease.  |
| AAC49455.1     | U41902   | Pseudotsuga menziesii   | cysteine protease. Pseudotsain. PM33cysP.  |
| CAA46863.1     | X66061   | Pisum sativum           | thiolprotease. tpp. start codon ttg.   |
| AAB41816.1     | U44947   | Pisum sativum           | NTH1. PsCyp1. cysteine protease homolog.   |
| BAA83473.1     | AB004819 | Oryza sativa            | cysteine endopeptidase. Rep1.  |
| BAA22544.1     | D38532   | Ananas comosus          | precursor of cysteine proteinase. FBSB precursor. stem bromelain precursor in fruit.                               |
| BAA11170.1     | D76415   | Oryza sativa            | cysteine proteinase.   |
| AAD20453.1     | AF099203 | Oryza sativa            | cysteine endopeptidase precursor. EP3A.  |
| CAA08860.1     | AJ009829 | Ananas comosus          | cysteine proteinase precursor, AN8. an8.   |
| AAA85036.1     | U19384   | Hordeum vulgare         | cysteine proteinase EPB2 precursor.  |
| AAA85035.1     | U19359   | Hordeum vulgare         | cysteine proteinase EPB1 precursor.  |
| SEQ ID NO: 565 |          |                         |  |
| AAA97411.1     | U51918   | Pisum sativum           | pyruvate dehydrogenase E1 alpha subunit.   |
| CAA81558.1     | Z26949   | Solanum tuberosum       | subunit of the mitochondrial pyruvate dehydrogenase complex. E1 alpha subunit of pyruvate dehydrogenase precursor. |
| AAC72195.1     | AF069911 | Zea mays                | pyruvate dehydrogenase E1 alpha subunit.   |
| AAG43499.1     | AF209924 | Lycopersicon esculentum | pyruvate dehydrogenase.  |
| CAB08111.1     | Z94180   | Lycopersicon esculentum | branched chain alpha-keto acid dehydrogenase E1-alpha subunit.   |
| SEQ ID NO: 566 |          |                         |  |
| AAD55090.1     | AF178653 | Vitis riparia           | thaumatin. osmotin; pathogenesis-related protein.  |
| CAA51432.1     | X72928   | Solanum commersonii     | osmotin-like protein.  |
| CAA47601.1     | X67121   | Solanum commersonii     | osmotin-like protein.  |
| CAC34055.1     | AJ297410 | Capsicum annuum         | osmotin-like protein. pr5 p23.   |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| CAA47047.1 | X66416   | Lycopersicon esculentum | tpm 1. Induced during viroid infection. Osmotin-like, antifungal protein homologue.  |
| AAB23375.1 | S44889   | Nicotiana tabacum       | osmotin. osmotin. pathogenesis-related protein homolog; This sequence comes from Fig. 3.   |
| AAB22459.2 | S40046   | Nicotiana tabacum       | osmotin. abscisic acid-activated. basic PR-like protein; This sequence comes from Fig. 1A; conceptual translation presented here differs from translation in publication; mismatch(181[R->G]). |
| AAG16625.1 | AY007309 | Solanum dulcamara       | cryoprotective osmotin-like protein.   |
| CAA46623.1 | X65701   | Nicotiana tabacum       | osmotin. AP24.   |
| CAA46622.1 | X65700   | Nicotiana tabacum       | osmotin. AP24.   |
| CAA51431.1 | X72927   | Solanum commersonii     | osmotin-like protein.  |
| CAA64620.1 | X95308   | Nicotiana tabacum       | PR protein. osmotin.   |
| CAA51430.1 | X72926   | Solanum commersonii     | osmotin-like protein.  |
| AAC64171.1 | AF093743 | Lycopersicon esculentum | pathogenesis-related protein osmotin precursor. NP24.  |
| AAB61590.1 | AF003007 | Vitis vinifera          | VVTL1. osmotin-like protein, PR-5 protein; thaumatin-like protein.   |
| BAA11180.1 | D76437   | Nicotiana sylvestris    | antifungal activity. neutral PR-5 (osmotin-like protein, PR-5d). neutral isoform of pathogenesis-related protein group 5.  |
| AAA34087.1 | M64081   | Nicotiana tabacum       | osmotin-like protein. OLP1.  |
| CAA47669.1 | X67244   | Solanum commersonii     | osmotin-like protein.  |
| CAA71883.1 | Y10992   | Vitis vinifera          | osmotin-like protein. OSM1.  |
| AAF13707.1 | AF199508 | Fragaria x ananassa     | osmotin-like protein. olp.   |
| AAA34089.1 | M29279   | Nicotiana tabacum       | osmotin.   |
| CAA43854.1 | X61679   | Nicotiana tabacum       | osmotin.   |
| CAA04642.1 | AJ001268 | Hordeum vulgare         | antifungal. basic pathogenesis-related protein PR5. osmotin/permatin-like.   |
| CAC22330.1 | AJ298304 | Fagus sylvatica         | stress protein. osmotin-like protein. olp2.  |

CAC22329.1 AJ298303 *Fagus sylvatica*  
stress protein. osmotin-like protein. olp.

AAB67852.1 L76377 *Oryza sativa*  
osmotin. 14b.

CAB36911.1 AJ000692 *Quercus suber*  
stress protein. osmotin-like protein. olp.

SEQ ID NO: 569

BAA95814.1 AP002069 *Oryza sativa*  
ESTs AU082579(S2069),D40238(S2069) correspond to a region of the predicted gene.  
Similar to *Arabidopsis thaliana* vesicle-associated membrane protein 7C; synaptobrevin 7C.  
(AF025332).

SEQ ID NO: 572

AAA92677.1 U13736 *Pisum sativum*  
binds calcium. calmodulin-like protein.

AAA34015.1 L01433 *Glycine max*  
calcium-binding regulatory protein. calmodulin. SCaM-4. putative.

AAA33948.1 L19359 *Glycine max*  
calcium-binding regulatory protein. calmodulin. SCaM-5. putative.

CAA66159.1 X97558 *Capsicum annuum*  
calmodulin-1.

CAA09302.1 AJ010645 *Capsicum annuum*  
calcium binding protein. calmodulin 3 protein. calmodulin 3.

AAA34144.1 M67472 *Lycopersicon esculentum*  
calmodulin. CALM1LE.

CAA62150.1 X90560 *Physcomitrella patens*  
Calmodulin. CaM.

AAB46588.1 U83402 *Capsicum annuum*  
calmodulin.

BAA87825.1 AP000815 *Oryza sativa*  
ESTs AU030013(E50493),AU081341(E50493) correspond to a region of the predicted gene.  
Similar to *O.sativa* gene encoding calmodulin. (Z12828).

AAF65511.1 AF108889 *Capsicum annuum*  
calmodulin.

AAA85157.1 U20297 *Solanum tuberosum*  
calcium-binding protein. calmodulin.

AAA85156.1 U20296 *Solanum tuberosum*  
calcium-binding protein. calmodulin.

AAA62351.1 U20295 *Solanum tuberosum*  
calcium-binding protein. calmodulin.

AAA85155.1 U20294 *Solanum tuberosum*  
calcium-binding protein. calmodulin.

AAA33900.1 L18914 *Oryza sativa*  
calcium binding protein, signal transduction. calmodulin.

|            |          |                           |  |
|------------|----------|---------------------------|--|
| CAA78288.1 | Z12828   | Oryza sativa              | calcium binding protein, signal transduction. calmodulin.  |
| AAC49583.1 | U48692   | Triticum aestivum         | calmodulin TaCaM2-3. calcium-binding protein.  |
| AAC49582.1 | U48691   | Triticum aestivum         | calmodulin TaCaM2-2. calcium-binding protein.  |
| CAA61980.1 | X89890   | Bidens pilosa             | Calmodulin.  |
| CAA67054.1 | X98404   | Capsicum annuum           | calmodulin-2.  |
| AAA33083.1 | M20729   | Chlamydomonas reinhardtii | calmodulin.  |
| AAG11418.1 | AF292108 | Prunus avium              | calmodulin.  |
| AAF33852.1 | AF231026 | Oryza sativa              | calmodulin-like protein.   |
| AAA92681.1 | U13882   | Pisum sativum             | calcium-binding protein. calmodulin.   |
| AAA33706.1 | M80836   | Petunia x hybrida         | calmodulin. CAM81.   |
| AAA33705.1 | M80831   | Petunia x hybrida         | calmodulin-related protein. CAM53.   |
| AAA98933.1 | U37936   | Oryza sativa              | novel calmodulin-like protein. C-terminal of this sequence contains a CAAX box which is common in signal transport protein, but not in calmodulin. |
| CAA43143.1 | X60738   | Malus x domestica         | Calmodulin. CaM.   |
| CAA78301.1 | Z12839   | Lilium longiflorum        | calcium binding protein, signal transduction. calmodulin.  |
| AAA33397.1 | L18912   | Lilium longiflorum        | calcium binding protein, signal transduction. calmodulin. putative.  |
| CAA42423.1 | X59751   | Daucus carota             | calmodulin. Ccam-1.  |
| AAF73157.1 | AF150059 | Brassica napus            | calmodulin. CaM1. involved in seed germination.  |
| AAA19571.1 | U10150   | Brassica napus            | calcium binding. calmodulin. bcml.   |
| AAA87347.1 | M88307   | Brassica juncea           | calmodulin.  |
| AAG27432.1 | AF295637 | Elaeis guineensis         | calmodulin.  |
| BAA94697.1 | AB041712 | Chara corallina           | calmodulin. cccam2.  |

|                |          |   |
|----------------|----------|---|
| BAA94696.1     | AB041711 | Chara corallina<br>calmodulin. cccam1.  |
| BAA96536.1     | AB044286 | Chara corallina<br>calmodulin. ccam.  |
| AAC18355.1     | AF064456 | Oryza sativa subsp. indica<br>calmodulin-like protein. CAM-like.                        |
| AAA34237.1     | L20691   | Vigna radiata<br>calmodulin.  |
| CAA52602.1     | X74490   | Zea mays<br>Calmodulin. ZMCALM1.  |
| CAA54583.1     | X77397   | Zea mays<br>calmodulin. CaM2.   |
| AAC49585.1     | U49103   | Triticum aestivum<br>calmodulin TaCaM3-2. calcium-binding protein.                      |
| AAC49586.1     | U49104   | Triticum aestivum<br>calmodulin TaCaM3-3. calcium-binding protein.                      |
| AAC49587.1     | U49105   | Triticum aestivum<br>calmodulin TaCaM4-1. calcium-binding protein.                      |
| AAC49580.1     | U48689   | Triticum aestivum<br>calmodulin TaCaM1-3. calcium-binding protein.                      |
| AAC49584.1     | U48693   | Triticum aestivum<br>calmodulin TaCaM3-1. calcium-binding protein.                      |
| SEQ ID NO: 575 |          |   |
| AAK28303.1     | AF346431 | Nicotiana tabacum<br>phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase. |
| AAB36653.1     | U32644   | Nicotiana tabacum<br>immediate-early salicylate-induced glucosyltransferase. IS5a.      |
| AAB36652.1     | U32643   | Nicotiana tabacum<br>immediate-early salicylate-induced glucosyltransferase. IS10a.     |
| AAK28304.1     | AF346432 | Nicotiana tabacum<br>phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase. |
| CAA59450.1     | X85138   | Lycopersicon esculentum<br>twil. homologous to glucosyltransferases.                    |
| CAB56231.1     | Y18871   | Dorotheanthus bellidiformis<br>betanidin-5-O-glucosyltransferase.                       |
| BAA83484.1     | AB031274 | Scutellaria baicalensis<br>UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.        |
| AAB48444.1     | U82367   | Solanum tuberosum<br>UDP-glucose glucosyltransferase.                                   |
| CAA54610.1     | X77460   | Manihot esculenta<br>UTP-glucose glucosyltransferase. CGT4.                             |

|            |          |   |   |
|------------|----------|---|---|
| AAD04166.1 | AF101972 | <i>Phaseolus lunatus</i>                      | catalyzes formation of O-glucosylzeatin from zeatin and UDPG; catalyzes formation of O-xylosylzeatin from zeatin and UDPX. zeatin O-glucosyltransferase. ZOG1. cytokinin O-glucosyltransferase. |
| BAB17182.1 | AP002843 | <i>Oryza sativa</i>                           | putative UTP-glucose glucosyltransferase. P0407B12.19.  |
| BAA93039.1 | AB033758 | <i>Citrus unshiu</i>                          | limonoid UDP-glucosyltransferase. LGTase.   |
| AAF61647.1 | AF190634 | <i>Nicotiana tabacum</i>                      | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAB17176.1 | AP002843 | <i>Oryza sativa</i>                           | putative UTP-glucose glucosyltransferase. P0407B12.13.  |
| CAC09351.1 | AL442007 | <i>Oryza sativa</i>                           | putative glucosyltransferase. H0212B02.7.   |
| CAA54611.1 | X77461   | <i>Manihot esculenta</i>                      | UTP-glucose glucosyltransferase. CGT2.  |
| AAF98390.1 | AF287143 | <i>Brassica napus</i>                         | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT.         |
| AAK16181.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.16.   |
| AAK16178.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.5.  |
| CAA54612.1 | X77462   | <i>Manihot esculenta</i>                      | UTP-glucose glucosyltransferase. CGT5.  |
| AAK16180.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.21.   |
| AAF17077.1 | AF199453 | <i>Sorghum bicolor</i>                        | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |
| CAA54609.1 | X77459   | <i>Manihot esculenta</i>                      | UTP-glucose glucosyltransferase. CGT1.  |
| BAB41025.1 | AB047098 | <i>Vitis vinifera</i>                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.   |
| AAK16172.1 | AC079887 | <i>Oryza sativa</i>                           | putative glucosyltransferase. OSJNBa0040E01.14.   |
| BAB41019.1 | AB047092 | <i>Vitis vinifera</i>                         | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT1.   |
| BAB41018.1 | AB047091 | <i>Vitis labrusca</i> x <i>Vitis vinifera</i> | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT2. The gene originated in one of the parents <i>V. vinifera</i> cv. Centennial.  |

|                |          |  |   |
|----------------|----------|--|---|
| BAB41023.1     | AB047096 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.   |
| BAB41021.1     | AB047094 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT1.   |
| BAA89009.1     | AB027455 | <i>Petunia x hybrida</i>               | anthocyanin 5-O-glucosyltransferase. PH1.   |
| AAB81682.1     | AF000371 | <i>Vitis vinifera</i>                  | UDP glucose:flavonoid 3-o-glucosyltransferase.  |
| BAB41017.1     | AB047090 | <i>Vitis labrusca x Vitis vinifera</i> | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT1. The gene was derived from one of the parents <i>V. labruscana</i> cv. Ishiharawase. |
| BAB41022.1     | AB047095 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT2.   |
| BAB41020.1     | AB047093 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT2.   |
| BAB41024.1     | AB047097 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT2.   |
| AAB81683.1     | AF000372 | <i>Vitis vinifera</i>                  | UDP glucose:flavonoid 3-o-glucosyltransferase.  |
| BAB41026.1     | AB047099 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT2.   |
| CAA54613.1     | X77463   | <i>Manihot esculenta</i>               | UTP-glucose glucosyltransferase. CGT6.  |
| BAA89008.1     | AB027454 | <i>Petunia x hybrida</i>               | anthocyanidin 3-O-glucosyltransferase. PGT8.  |
| BAA12737.1     | D85186   | <i>Gentiana triflora</i>               | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| SEQ ID NO: 576 |          |  |   |
| CAB60277.1     | AJ002586 | <i>Solanum tuberosum</i>               | UCP.  |
| CAA72107.1     | Y11220   | <i>Solanum tuberosum</i>               | mitochondrial uncoupling protein.   |
| BAA92172.1     | AB024733 | <i>Symplocarpus renifolius</i>         | SfUCPa. SfUCPa.   |
| BAB40658.1     | AB049998 | <i>Oryza sativa</i>                    | uncoupling protein. OsUCP2.   |
| BAA92173.1     | AB024734 | <i>Symplocarpus renifolius</i>         | SfUCPb. SfUCPb.   |
| BAB16385.1     | AB042429 | <i>Triticum aestivum</i>               | uncoupling protein. WhUCP1b.  |
| BAB16384.1     | AB042428 | <i>Triticum aestivum</i>               | uncoupling protein. WhUCP1a.  |



|                |          |                    |   |
|----------------|----------|--------------------|---|
| BAB40657.1     | AB049997 | Oryza sativa       | uncoupling protein. OsUCP1.   |
| SEQ ID NO: 577 |          |                    |   |
| AAA19571.1     | U10150   | Brassica napus     | calcium binding. calmodulin. bcm1.                                  |
| AAA87347.1     | M88307   | Brassica juncea    | calmodulin.   |
| AAA92681.1     | U13882   | Pisum sativum      | calcium-binding protein. calmodulin.                                |
| AAA33706.1     | M80836   | Petunia x hybrida  | calmodulin. CAM81.  |
| CAA43143.1     | X60738   | Malus x domestica  | Calmodulin. CaM.  |
| CAA78301.1     | Z12839   | Lilium longiflorum | calcium binding protein, signal transduction. calmodulin.           |
| AAA33397.1     | L18912   | Lilium longiflorum | calcium binding protein, signal transduction. calmodulin. putative. |
| CAA42423.1     | X59751   | Daucus carota      | calmodulin. Ccam-1.   |
| AAG27432.1     | AF295637 | Elaeis guineensis  | calmodulin.   |
| AAG11418.1     | AF292108 | Prunus avium       | calmodulin.   |
| AAA34237.1     | L20691   | Vigna radiata      | calmodulin.   |
| AAC49587.1     | U49105   | Triticum aestivum  | calmodulin TaCaM4-1. calcium-binding protein.                       |
| AAC49586.1     | U49104   | Triticum aestivum  | calmodulin TaCaM3-3. calcium-binding protein.                       |
| AAC49585.1     | U49103   | Triticum aestivum  | calmodulin TaCaM3-2. calcium-binding protein.                       |
| AAC49584.1     | U48693   | Triticum aestivum  | calmodulin TaCaM3-1. calcium-binding protein.                       |
| AAC49580.1     | U48689   | Triticum aestivum  | calmodulin TaCaM1-3. calcium-binding protein.                       |
| AAC49579.1     | U48688   | Triticum aestivum  | calmodulin TaCaM1-2. calcium binding protein.                       |
| AAC49578.1     | U48242   | Triticum aestivum  | calmodulin TaCaM1-1. calcium-binding.                               |
| AAA03580.1     | L01431   | Glycine max        | calcium-binding regulatory protein. calmodulin. SCaM-2. putative.   |

|            |          |                    |   |
|------------|----------|--------------------|---|
| AAB36130.1 | S81594   | Vigna radiata      | auxin-regulated calmodulin. auxin-regulated calmodulin, arCaM. This sequence comes from Fig. 1; arCaM.                |
| AAA33901.1 | L18913   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin. putative.   |
| CAA78287.1 | Z12827   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin.   |
| AAB46588.1 | U83402   | Capsicum annuum    | calmodulin.   |
| AAA32938.1 | M27303   | Hordeum vulgare    | calmodulin.   |
| BAA88540.1 | AP000969 | Oryza sativa       | ESTs AU081349(E61253), D41425(S3918) correspond to a region of the predicted gene. Similar to calmodulin. (AF042840). |
| AAF65511.1 | AF108889 | Capsicum annuum    | calmodulin.   |
| AAC36059.1 | AF042840 | Oryza sativa       | calmodulin. CaM1.   |
| AAA34238.1 | L20507   | Vigna radiata      | calmodulin.   |
| AAA34014.1 | L01432   | Glycine max        | calcium-binding regulatory protein. calmodulin. SCaM-3. putative.   |
| AAA34013.1 | L01430   | Glycine max        | calcium-binding regulatory protein. calmodulin. SCaM-1. putative.   |
| CAA36644.1 | X52398   | Medicago sativa    | calmodulin (AA 1-149).  |
| AAB68399.1 | U79736   | Helianthus annuus  | calmodulin. HaCaM.  |
| AAD10244.1 | AF030032 | Phaseolus vulgaris | calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal transduction pathways.       |
| AAA33705.1 | M80831   | Petunia x hybrida  | calmodulin-related protein. CAM53.  |
| AAA16320.1 | L14071   | Bryonia dioica     | calmodulin. Bc329. calcium-binding sites (amino acid #): 1. (21..33); 2. (57..68); 3. (94..106); 4. (130..141).       |
| CAA61980.1 | X89890   | Bidens pilosa      | Calmodulin.   |
| AAA33900.1 | L18914   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin.   |
| CAA78288.1 | Z12828   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin.   |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| CAA74307.1     | Y13974   | Zea mays                | calmodulin.   |
| AAC49583.1     | U48692   | Triticum aestivum       | calmodulin TaCaM2-3. calcium-binding protein.   |
| AAC49582.1     | U48691   | Triticum aestivum       | calmodulin TaCaM2-2. calcium-binding protein.   |
| AAA85157.1     | U20297   | Solanum tuberosum       | calcium-binding protein. calmodulin.  |
| AAA85156.1     | U20296   | Solanum tuberosum       | calcium-binding protein. calmodulin.  |
| AAA62351.1     | U20295   | Solanum tuberosum       | calcium-binding protein. calmodulin.  |
| AAA85155.1     | U20294   | Solanum tuberosum       | calcium-binding protein. calmodulin.  |
| CAA46150.1     | X65016   | Oryza sativa            | calmodulin. cam.  |
| AAD10246.1     | AF030034 | Phaseolus vulgaris      | calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal transduction pathways. |
| AAF73157.1     | AF150059 | Brassica napus          | calmodulin. CaM1. involved in seed germination.   |
| CAA54583.1     | X77397   | Zea mays                | calmodulin. CaM2.   |
| SEQ ID NO: 578 |          |                         |   |
| AAF01764.2     | AF184277 | Glycine max             | homeodomain-leucine zipper protein 56. Hdl56. transcription factor.   |
| BAA21017.1     | D26578   | Daucus carota           | transcriptional regulator. DNA-binding protein. homeodomain at nt 761-940; leucine zipper at nt 941-1048.       |
| AAD37697.1     | AF145728 | Oryza sativa            | homeodomain leucine zipper protein. Oshox4. transcription factor.   |
| CAB67118.1     | Y17306   | Lycopersicon esculentum | homeodomain protein. h52.   |
| AAF01765.1     | AF184278 | Glycine max             | homeodomain-leucine zipper protein 57. Hdl57. transcription factor.   |
| BAA05624.1     | D26575   | Daucus carota           | transcriptional regulator. DNA-binding protein. homeodomain at nt 520-699; leucine zipper at nt 700-805.        |
| BAA93466.1     | AB028078 | Physcomitrella patens   | homeobox protein PpHB7. PpHB7. homeodomain-leucine zipper gene.   |
| BAA93465.1     | AB028077 | Physcomitrella patens   | homeobox protein PpHB6. PpHB6. homeodomain-leucine zipper gene.   |

|                |          |                            |  |
|----------------|----------|----------------------------|--|
| BAA93464.1     | AB028076 | Physcomitrella patens      | homeobox protein PpHB5. PpHB5. homeodomain-leucine zipper gene.  |
| BAA05622.1     | D26573   | Daucus carota              | transcriptional regulator. DNA-binding protein. homeodomain at nt 585-764; leucine zipper at nt 765-851. |
| BAA05625.1     | D26576   | Daucus carota              | transcriptional regulator. DNA-binding protein. homeodomain at nt 300-479; leucine zipper at nt 480-587. |
| BAA05623.1     | D26574   | Daucus carota              | transcriptional regulator. DNA-binding protein. homeodomain at nt 498-677; leucine zipper at nt 678-785. |
| AAD37698.1     | AF145729 | Oryza sativa               | homeodomain leucine zipper protein. Oshox5. transcription factor.  |
| BAA93461.1     | AB028073 | Physcomitrella patens      | homeobox protein PpHB2. PpHB2. homeodomain-leucine zipper gene.  |
| BAA93467.1     | AB028079 | Physcomitrella patens      | homeobox protein PpHB8. PpHB8. homeodomain-leucine zipper gene.  |
| BAA93468.1     | AB028080 | Physcomitrella patens      | homeobox protein PpHB9. PpHB9. homeodomain-leucine zipper gene.  |
| BAA93460.1     | AB028072 | Physcomitrella patens      | homeobox protein PpHB1. PpHB1. homeodomain-leucine zipper gene.  |
| AAD37699.1     | AF145730 | Oryza sativa               | homeodomain leucine zipper protein. Oshox6. transcription factor.  |
| CAA06717.1     | AJ005820 | Craterostigma plantagineum | transcription factor. homeodomain leucine zipper protein. hb-1.  |
| BAA93463.1     | AB028075 | Physcomitrella patens      | homeobox protein PpHB4. PpHB4. homeodomain-leucine zipper gene.  |
| CAA65456.2     | X96681   | Oryza sativa               | transcription factor. DNA-binding protein. Oshox1. homeodomain leucine zipper gene.                      |
| AAF19980.1     | AF211193 | Oryza sativa               | homeodomain-leucine zipper transcription factor. Hox1. hox1.   |
| AAK31270.1     | AC079890 | Oryza sativa               | homeodomain leucine zipper protein hox1. OSJNBb0089A17.12.   |
| CAA06728.1     | AJ005833 | Craterostigma plantagineum | transcription factor. homeodomain leucine zipper protein. hb-2.  |
| AAD37696.1     | AF145727 | Oryza sativa               | homeodomain leucine zipper protein. Oshox3. transcription factor.  |
| SEQ ID NO: 580 |          |                            |  |
| AAD32141.1     | AF123503 | Nicotiana tabacum          | Nt-gh3 deduced protein.  |
| CAA42636.1     | X60033   | Glycine max                | auxin-responsive GH3 product. GH3.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAA96221.1     | AP002094 | Oryza sativa            | ESTs C19814(E10971),AU090481(E10971) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, sequence from clones T3F17, F11C10, F13A10; auxin-responsive GH3-like protein (AC006526).   |
| SEQ ID NO: 581 |          |                         |  |
| AAD13632.1     | AF059488 | Lycopersicon esculentum | expansin precursor. Exp4.  |
| AAF32410.1     | AF230277 | Triphysaria versicolor  | alpha-expansin 2.  |
| CAA04385.1     | AJ000885 | Brassica napus          | Cell wall extension in plants. Expansin.   |
| CAB46492.1     | AJ243340 | Lycopersicon esculentum | expansin9. exp9.   |
| AAC63088.1     | U82123   | Lycopersicon esculentum | expansin. LeEXP1. fruit ripening regulated expansin.   |
| CAA06271.2     | AJ004997 | Lycopersicon esculentum | expansin18. exp18.   |
| AAF62182.1     | AF247164 | Oryza sativa            | alpha-expansin OsEXP7. cell wall loosening factor; expressed in internodes and leaves.   |
| AAF35900.1     | AF230331 | Zinnia elegans          | expansin. Exp1.  |
| AAG13982.1     | AF297521 | Prunus avium            | expansin 1. Exp1. PruavExp1.   |
| AAC33529.1     | U93167   | Prunus armeniaca        | expansin. PA-Exp1.   |
| AAF21101.1     | AF159563 | Fragaria x ananassa     | expansin. Exp2. ripening regulated.  |
| AAF35901.1     | AF230332 | Zinnia elegans          | expansin 2.  |
| AAG13983.1     | AF297522 | Prunus avium            | expansin 2. Exp2. PruavExp2.   |
| AAC33530.1     | AF038815 | Prunus armeniaca        | expansin. Exp2.  |
| BAB19676.1     | AB029083 | Prunus persica          | expansin. PchExp1.   |
| AAB40637.1     | U64893   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAB37746.1     | U30382   | Cucumis sativus         | expansin S1 precursor. Cs-EXP1. similar to pollen allergen Lol pI, Lolium perenne, Swiss-Prot Accession Number P14946; former gene name CuExS1; expansin-29 (Ex29) protein.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| CAC19184.1 | AJ291817 | Cicer arietinum         | expansin.  |
| AAD47901.1 | AF085330 | Pinus taeda             | expansin.  |
| AAB40634.1 | U64890   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAF32409.1 | AF230276 | Triphysaria versicolor  | alpha-expansin 3.  |
| AAC39512.1 | AF043284 | Gossypium hirsutum      | expansin. GhEX1. contains N-terminal signal peptide.   |
| AAB40635.1 | U64891   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAC96081.1 | AF049354 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP5. cell wall protein.  |
| AAB38074.1 | U30477   | Oryza sativa            | induces extension (creep) in plant cell walls. expansin Os-EXP2. Os-EXP2. former gene name RiExB.  |
| AAG32921.1 | AF184233 | Lycopersicon esculentum | expansin. Exp10.   |
| AAB81662.1 | U85246   | Oryza sativa            | expansin. Os-EXP4.   |
| AAD49956.1 | AF167360 | Rumex palustris         | expansin. EXP1.  |
| AAB40636.1 | U64892   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAF32411.1 | AF230278 | Triphysaria versicolor  | alpha-expansin 1.  |
| AAC96080.1 | AF049353 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP4. cell wall protein.  |
| AAF35902.1 | AF230333 | Zinnia elegans          | expansin 3.  |
| CAB43197.1 | AJ239068 | Lycopersicon esculentum | cell wall loosening enzyme. expansin2. exp2.   |
| AAC64201.1 | AF096776 | Lycopersicon esculentum | expansin. LeEXP2.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAF17570.1     | AF202119 | Marsilea quadrifolia    | alpha-expansin. EXP1. Mq-EXP1.  |
| AAD13633.1     | AF059489 | Lycopersicon esculentum | expansin precursor. Exp5.   |
| AAF62181.1     | AF247163 | Oryza sativa            | alpha-expansin OsEXP6. cell wall loosening factor; expressed in internodes and leaves.  |
| AAF17571.1     | AF202120 | Regnellidium diphyllum  | alpha-expansin. EXP1. Rd-EXP1.  |
| AAF62180.1     | AF247162 | Oryza sativa            | alpha-expansin OsEXP5. cell wall loosening factor; expressed in internodes, leaves, coleoptiles, and roots.   |
| CAC19183.1     | AJ291816 | Cicer arietinum         | expansin.   |
| CAC06433.1     | AJ276007 | Festuca pratensis       | expansin. exp2.   |
| BAB32732.1     | AB049406 | Eustoma grandiflorum    | expansin. Eg Expansin.  |
| AAG01875.1     | AF291659 | Striga asiatica         | alpha-expansin 3. Exp3.   |
| AAB37749.1     | U30460   | Cucumis sativus         | expansin S2 precursor. Cs-EXP2. similar to pollen allergen Lol pI, Lolium perenne, Swiss-Prot Accession Number P14946; former gene name CuExS2; expansin-30 (Ex30) protein. |
| AAC96077.1     | AF049350 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP1. cell wall protein.   |
| AAC96079.1     | AF049352 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP3. cell wall protein.   |
| AAG32920.1     | AF184232 | Lycopersicon esculentum | expansin. Exp8.   |
| CAA69105.1     | Y07782   | Oryza sativa            | expansin. RiExA.  |
| AAC96078.1     | AF049351 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP2. cell wall protein.   |
| SEQ ID NO: 583 |          |                         |   |
| BAA85412.1     | AP000615 | Oryza sativa            | ESTs AU065232(E60855), C23624(S1554), AU078241(E60855) correspond to a region of the predicted gene.; similar to putative adenylate kinase. (AC005896).                     |
| BAA01181.1     | D10335   | Oryza sativa            | adenylate kinase-b. Adk-b.  |
| BAA01180.1     | D10334   | Oryza sativa            | adenylate kinase-a. Adk-a.  |
| BAA94761.1     | AB041773 | Oryza sativa            | adenylate kinase. Adk-a.  |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| AAB68604.1     | U82330   | Prunus armeniaca          | adenylate kinase homolog.   |
| AAF23372.1     | AF187063 | Oryza sativa              | catalyzes the transfer of phosphate from ATP to UMP or CMP to form ADP and UDP or CDP. UMP/CMP kinase b. ura6.  |
| AAF23371.1     | AF187062 | Oryza sativa              | catalyzes the transfer of phosphate from ATP to UMP or CMP to form ADP and UDP or CDP. UMP/CMP kinase a. ura6.  |
| AAD41679.1     | AF086603 | Ceratopteris richardii    | adenylate kinase. ADK1.   |
| BAA85443.1     | AP000616 | Oryza sativa              | similar to UMP/CMP kinase (AF000147).   |
| SEQ ID NO: 584 |          |                           |   |
| BAA87052.2     | D88273   | Hordeum vulgare           | nicotianamine aminotransferase A. naat-A.   |
| BAA87055.1     | AB024006 | Hordeum vulgare           | nicotianamine aminotransferase. naat-A.   |
| BAA87053.1     | AB005788 | Hordeum vulgare           | nicotianamine aminotransferase B. naat-b. NAAT-B.   |
| BAA87054.1     | AB024006 | Hordeum vulgare           | nicotianamine aminotransferase. naat-B.   |
| BAA77261.1     | AB007405 | Oryza sativa              | alanine aminotransferase. AlaAT.  |
| BAA77260.1     | AB007404 | Oryza sativa              | alanine aminotransferase. AlaAT.  |
| CAA49199.1     | X69421   | Panicum miliaceum         | alanine aminotransferase. pAlaAT-2.   |
| AAB01685.1     | U31975   | Chlamydomonas reinhardtii | catalyzes the transfer of -NH <sub>2</sub> from ala to 2-oxoglutarate. alanine aminotransferase. The translation start site has not been experimentally tested, but a 55 kDa product can be detected in Western blot. |
| AAC62456.1     | AF055898 | Zea mays                  | alanine aminotransferase. alt. AlaAT.   |
| CAA81231.1     | Z26322   | Hordeum vulgare           | alanine aminotransferase.   |
| SEQ ID NO: 586 |          |                           |   |
| AAB94589.1     | AF022460 | Glycine max               | CYP83D1p. CYP83D1. cytochrome P450 monooxygenase.   |
| AAA32913.1     | M32885   | Persea americana          | cytochrome P-450LXXIA1 (cyp71A1).   |
| AAA19701.1     | L24438   | Thlaspi arvense           | cytochrome P450.  |



|            |          |   |  |
|------------|----------|---|--|
| BAB40324.1 | AB037245 | Asparagus officinalis                             | cytochrome P450. ASPI-2.   |
| AAC39318.1 | AF029858 | Sorghum bicolor                                   | second multifunctional cytochrome P450 in the biosynthetic pathway of the cyanogenic glucoside dhurrin. Catalyzes the conversion of p-hydroxyphenylacetaldoxime to p-hydroxymandelonitrile. cytochrome P450 CYP71E1. CYP71E1. No EST#s identified. |
| BAB40323.1 | AB037244 | Asparagus officinalis                             | cytochrome P450. ASPI-1.   |
| CAA71517.1 | Y10493   | Glycine max                                       | putative cytochrome P450.  |
| AAB61965.1 | U48435   | Solanum chacoense                                 | putative cytochrome P450.  |
| CAA70575.1 | Y09423   | Nepeta racemosa                                   | cytochrome P450. CYP71A5.  |
| CAA71513.1 | Y10489   | Glycine max                                       | putative cytochrome P450.  |
| AAB61964.1 | U48434   | Solanum chacoense                                 | putative cytochrome P450.  |
| AAD47832.1 | AF166332 | Nicotiana tabacum                                 | cytochrome P450.   |
| AAB94588.1 | AF022459 | Glycine max                                       | CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.  |
| AAF27282.1 | AF122821 | Capsicum annuum                                   | cytochrome P450. PepCYP.   |
| CAA50312.1 | X70981   | Solanum melongena                                 | P450 hydroxylase. CYPEG2.  |
| CAA83941.1 | Z33875   | Mentha x piperita                                 | cytochrome P-450 oxidase.  |
| AAG44132.1 | AF218296 | Pisum sativum                                     | cytochrome P450. P450 isolog.  |
| CAA70576.1 | Y09424   | Nepeta racemosa                                   | cytochrome P450. CYP71A6.  |
| AAD44151.1 | AF124816 | Mentha x piperita                                 | cytochrome p450 isoform PM17.  |
| BAA03635.1 | D14990   | Solanum melongena                                 | Cytochrome P-450EG4.   |
| CAA50645.1 | X71654   | Solanum melongena                                 | P450 hydroxylase.  |
| AAB69644.1 | AF000403 | Lotus japonicus                                   | putative cytochrome P450. LjNP450.   |
| AAD37433.1 | AF150881 | Lycopersicon esculentum x Lycopersicon peruvianum | catalyzes the hydroxylation of ferulic acid to 5-hydroxyferulic acid. ferulate-5-hydroxylase. CYP84. cytochrome P450-dependent monooxygenase; F5H; FAH1.   |

|                |          |                     |   |
|----------------|----------|---------------------|---|
| CAA65580.1     | X96784   | Nicotiana tabacum   | cytochrome P450. hsr515.  |
| AAD44152.1     | AF124817 | Mentha x piperita   | cytochrome p450 isoform PM2.  |
| CAA64635.1     | X95342   | Nicotiana tabacum   | cytochrome P450. hsr515. hypersensitivity-related gene.   |
| AAG14962.1     | AF214008 | Brassica napus      | cytochrome p450-dependent monooxygenase. BNF5H2.  |
| AAG14961.1     | AF214007 | Brassica napus      | cytochrome p450-dependent monooxygenase. BNF5H1.  |
| CAB56503.1     | AJ238612 | Catharanthus roseus | cytochrome P450.  |
| AAG14963.1     | AF214009 | Brassica napus      | cytochrome p450-dependent monooxygenase. BNF5H3.  |
| AAK38088.1     | AF321864 | Lolium rigidum      | putative cytochrome P450.   |
| AAD44150.1     | AF124815 | Mentha spicata      | cytochrome p450.  |
| AAK38084.1     | AF321860 | Lolium rigidum      | putative cytochrome P450.   |
| AAD56282.1     | AF155332 | Petunia x hybrida   | flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.  |
| AAK38087.1     | AF321863 | Lolium rigidum      | putative cytochrome P450.   |
| AAK38083.1     | AF321859 | Lolium rigidum      | putative cytochrome P450.   |
| AAB94584.1     | AF022157 | Glycine max         | capable of catalyzing the metabolism of phenylurea herbicides. CYP71A10. CYP71A10. cytochrome P450 monooxygenase. |
| CAA71514.1     | Y10490   | Glycine max         | putative cytochrome P450.   |
| BAA12159.1     | D83968   | Glycine max         | Cytochrome P-450 (CYP93A1).   |
| BAB39252.1     | AP002968 | Oryza sativa        | putative cytochrome P450. P0416G11.1.   |
| CAA50155.1     | X70824   | Solanum melongena   | flavonoid hydroxylase (P450). CYP75.  |
| SEQ ID NO: 587 |          |                     |   |
| CAA60120.1     | X86222   | Pisum sativum       | heat shock protein. hsp22.  |
| AAF37726.1     | AF237957 | Euphorbia esula     | LMW heat shock protein. putative 22 kDa mitochondrial heat shock protein.   |

|            |          |                                     |   |
|------------|----------|-------------------------------------|---|
| BAA32547.1 | AB017134 | Lycopersicon esculentum             | mitochondrial small heat shock protein. LEMTSHP.  |
| CAA33388.1 | X15333   | Chenopodium rubrum                  | heat shock protein (AA 1-204).  |
| AAB03096.1 | U21722   | Glycine max                         | Hsp23.9. Gmhsp23.9. low molecular weight heat shock protein.  |
| AAC12279.1 | AF035460 | Zea mays                            | low molecular weight heat shock protein precursor. hsp22.   |
| AAD03604.1 | AF104107 | Triticum aestivum                   | small heat shock protein Hsp23.5. mRNA not present, or only at very low levels, prior to heat stress; mRNA abundant after two hours at 40C.                                 |
| AAB01557.1 | L47741   | Picea glauca                        | mitochondria-localized low molecular weight heat shock protein 23.5. EMB22, SMW HSP23.5.  |
| AAD03605.1 | AF104108 | Triticum aestivum                   | small heat shock protein Hsp23.6.   |
| CAA38037.1 | X54103   | Plastid Petunia x hybrida           | heat shock protein. hsp21.  |
| AAF19022.1 | AF197942 | Funaria hygrometrica                | chloroplast-localized small heat shock protein 22. CPsHSP22.  |
| AAB49626.1 | U59917   | Lycopersicon esculentum             | chromoplast-associated hsp20. pTOM111. small molecular weight heat shock protein.   |
| AAB07023.1 | U66300   | Chloroplast Lycopersicon esculentum | heat shock protein. HSP21.  |
| BAA29064.1 | D88584   | Nicotiana tabacum                   | heat shock protein 26 (Type I). Nthsp26a. plastid-localized small heat shock protein.   |
| AAF19021.1 | AF197941 | Funaria hygrometrica                | chloroplast-localized small heat shock protein. CPsHSP21. CPsHS 21.   |
| BAA78385.1 | AB020973 | Oryza sativa                        | heat shock protein 26. Oshsp26. chloroplast-localized small heat shock protein.   |
| CAA41219.1 | X58280   | Triticum aestivum                   | heat shock protein 26.6. Tahsp26.6.   |
| AAC96315.1 | AF097657 | Triticum aestivum                   | heat shock protein HSP26. hsp26.6. similar to the Triticum aestivum hsp26.6G2 protein encoded by the sequence presented in GenBank Accession Number L41503; 26 kDa protein. |
| AAC96316.1 | AF097658 | Triticum aestivum                   | heat shock protein HSP26. hsp26.6. 26 kDa protein.  |
| AAC96314.1 | AF097656 | Triticum aestivum                   | heat shock protein HSP26. hsp26.6. 26 kDa protein.  |
| CAA47745.1 | X67328   | Triticum aestivum                   | heat shock protein 26.6B. hsp 26.6B.  |
| AAC96317.1 | AF097659 | Triticum aestivum                   | heat shock protein HSP26. hsp26.6. 26 kDa protein.  |

|            |          |   |  |
|------------|----------|---|--|
| AAA33477.1 | L28712   | <i>Zea mays</i>                                   | heat shock protein 26. HSP26. plastid-localized low molecular weight hsp.                |
| BAA29066.1 | AB006043 | <i>Nicotiana sylvestris</i>                       | heat shock protein 26. hsp26. plastid-localized small heat shock protein.                |
| BAA29067.1 | AB006044 | <i>Nicotiana tomentosiformis</i>                  | heat shock protein 26. hsp26. plastid-localized small heat shock protein.                |
| BAA29065.1 | AB006041 | <i>Nicotiana tabacum</i>                          | heat shock protein 26 (Type I). Nthsp26a. plastid-localized small heat shock protein 26. |
| AAC01570.1 | AF019144 | <i>Agrostis stolonifera</i> var. <i>palustris</i> | low molecular weight heat shock protein.   |
| AAD30452.1 | AF123255 | <i>Lycopersicon esculentum</i>                    | 17.7 kD class I small heat shock protein. HSP17.7.                                       |
| CAA39603.1 | X56138   | <i>Lycopersicon esculentum</i>                    | small heat shock protein (class I).  |
| AAD30453.1 | AF123256 | <i>Lycopersicon esculentum</i>                    | 17.8 kD class I small heat shock protein. HSP17.8.                                       |
| AAD30454.1 | AF123257 | <i>Lycopersicon esculentum</i>                    | 17.6 kD class I small heat shock protein. HSP17.6.                                       |
| AAD49336.1 | AF166277 | <i>Nicotiana tabacum</i>                          | low molecular weight heat-shock protein. LHS-1. TLHS-1.                                  |
| AAF34133.1 | AF161179 | <i>Malus x domestica</i>                          | low molecular weight heat shock protein. Hsp1.   |
| AAC39360.1 | U63631   | <i>Fragaria x ananassa</i>                        | LMW heat shock protein.  |
| AAA61632.1 | U08601   | <i>Papaver somniferum</i>                         | low molecular weight heat-shock protein.   |
| CAA08908.1 | AJ009880 | <i>Castanea sativa</i>                            | molecular chaperone. cytosolic class I small heat-shock protein HSP17.5. hsp17.5.        |
| AAA33975.1 | M11395   | <i>Glycine max</i>                                | small heat shock protein.  |
| CAA63570.1 | X92983   | <i>Pseudotsuga menziesii</i>                      | low molecular weight heat-shock protein.   |
| AAB03893.1 | M11318   | <i>Glycine max</i>                                | 17.5 kd heat shock protein Gmhsp17.6L.   |
| CAA25578.1 | X01104   | <i>Glycine max</i>                                | heat shock protein 6871 (aa 1-153).  |
| AAB72109.1 | AF022217 | <i>Brassica rapa</i>                              | low molecular weight heat-shock protein. BcHSP17.6. 17.6 kDa; cytosolic class I.         |
| CAA46641.1 | X65725   | <i>Zea mays</i>                                   | heat shock protein 17.2. Zmhsp 17.2. Class I low molecular weight heat shock protein.    |
| CAB08441.1 | Z95153   | <i>Helianthus annuus</i>                          | 17.6 kD class I small heat-shock protein HSP17.6. Ha hsp17.6.                            |

|                |          |  |
|----------------|----------|--|
| AAA33974.1     | M11317   | Glycine max<br>17.6 kd heat shock protein Gmhsp17.6L.  |
| SEQ ID NO: 588 |          |  |
| AAA32913.1     | M32885   | Persea americana<br>cytochrome P-450LXXIA1 (cyp71A1).  |
| AAA19701.1     | L24438   | Thlaspi arvense<br>cytochrome P450.  |
| AAC39318.1     | AF029858 | Sorghum bicolor<br>second multifunctional cytochrome P450 in the biosynthetic pathway of the cyanogenic<br>glucoside dhurrin. Catalyzes the conversion of p-hydroxyphenylacetaldoxime to p-<br>hydroxymandelonitrile. cytochrome P450 CYP71E1. CYP71E1. No EST#s identified. |
| BAB40323.1     | AB037244 | Asparagus officinalis<br>cytochrome P450. ASPI-1.  |
| BAB40324.1     | AB037245 | Asparagus officinalis<br>cytochrome P450. ASPI-2.  |
| AAB94589.1     | AF022460 | Glycine max<br>CYP83D1p. CYP83D1. cytochrome P450 monooxygenase.   |
| CAA71513.1     | Y10489   | Glycine max<br>putative cytochrome P450.   |
| CAA70575.1     | Y09423   | Nepeta racemosa<br>cytochrome P450. CYP71A5.   |
| AAB94584.1     | AF022157 | Glycine max<br>capable of catalyzing the metabolism of phenylurea herbicides. CYP71A10. CYP71A10.<br>cytochrome P450 monooxygenase.  |
| AAB61965.1     | U48435   | Solanum chacoense<br>putative cytochrome P450.   |
| AAF27282.1     | AF122821 | Capsicum annuum<br>cytochrome P450. PepCYP.  |
| CAB56503.1     | AJ238612 | Catharanthus roseus<br>cytochrome P450.  |
| AAD47832.1     | AF166332 | Nicotiana tabacum<br>cytochrome P450.  |
| AAB94588.1     | AF022459 | Glycine max<br>CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.   |
| AAB61964.1     | U48434   | Solanum chacoense<br>putative cytochrome P450.   |
| CAA71517.1     | Y10493   | Glycine max<br>putative cytochrome P450.   |
| CAA50645.1     | X71654   | Solanum melongena<br>P450 hydroxylase.   |
| BAA03635.1     | D14990   | Solanum melongena<br>Cytochrome P-450EG4.  |

|            |          |                     |   |
|------------|----------|---------------------|---|
| CAA50312.1 | X70981   | Solanum melongena   | P450 hydroxylase. CYPEG2.                                 |
| CAA71514.1 | Y10490   | Glycine max         | putative cytochrome P450.                                 |
| AAB69644.1 | AF000403 | Lotus japonicus     | putative cytochrome P450. LjNP450.                        |
| CAA70576.1 | Y09424   | Nepeta racemosa     | cytochrome P450. CYP71A6.                                 |
| AAD44151.1 | AF124816 | Mentha x piperita   | cytochrome p450 isoform PM17.                             |
| AAK38084.1 | AF321860 | Lolium rigidum      | putative cytochrome P450.                                 |
| CAA83941.1 | Z33875   | Mentha x piperita   | cytochrome P-450 oxidase.                                 |
| AAK38083.1 | AF321859 | Lolium rigidum      | putative cytochrome P450.                                 |
| AAD44152.1 | AF124817 | Mentha x piperita   | cytochrome p450 isoform PM2.                              |
| BAB39252.1 | AP002968 | Oryza sativa        | putative cytochrome P450. P0416G11.1.                     |
| AAD44150.1 | AF124815 | Mentha spicata      | cytochrome p450.  |
| AAK38087.1 | AF321863 | Lolium rigidum      | putative cytochrome P450.                                 |
| AAK38082.1 | AF321858 | Lolium rigidum      | putative cytochrome P450.                                 |
| CAA57423.1 | X81829   | Zea mays            | cytochrome P450. CYP71C2. family CYP71, subfamily CYP71C. |
| CAA72208.1 | Y11404   | Zea mays            | cytochrome p450. cyp71c2.                                 |
| CAC27827.1 | AJ295719 | Catharanthus roseus | geraniol hydroxylase. cytochrome P450. cyp71.             |
| AAG44132.1 | AF218296 | Pisum sativum       | cytochrome P450. P450 isolog.                             |
| AAK38088.1 | AF321864 | Lolium rigidum      | putative cytochrome P450.                                 |
| AAG14963.1 | AF214009 | Brassica napus      | cytochrome p450-dependent monooxygenase. BNF5H3.          |
| CAA57424.2 | X81830   | Zea mays            | cytochrome P450. CYP71C3. family CYP71, subfamily CYP71C. |
| CAA72207.1 | Y11403   | Zea mays            | cytochrome p450. cyp71c3.                                 |

AAG14962.1 AF214008 *Brassica napus*  
cytochrome p450-dependent monooxygenase. BNF5H2.

AAG14961.1 AF214007 *Brassica napus*  
cytochrome p450-dependent monooxygenase. BNF5H1.

BAA12159.1 D83968 *Glycine max*  
Cytochrome P-450 (CYP93A1).

BAB40322.1 AB036772 *Triticum aestivum*  
cytochrome P450. N-1.

SEQ ID NO: 593

BAA09645.1 D63331 *Nicotiana tabacum*  
Indole-3-acetonitrile hydrolysis. nitrilase. TNIT4. Indole-3-acetonitrile hydrolysis, indole-3-acetic acid biosynthesis.

BAA11770.1 D83078 *Nicotiana tabacum*  
indole-3-acetonitrile hydrolysis. nitrilase. TNIT4B. indole-3-acetonitrile hydrolysis, indole-3-acetic acid biosynthesis.

BAA77679.1 AB027054 *Oryza sativa*  
nitrilase-like protein. ONIT4.

SEQ ID NO: 595

AAG01147.1 AF283816 *Pinus taeda*  
calreticulin. calcium-binding protein.

CAA05161.1 AJ002057 *Beta vulgaris*  
calreticulin.

AAB71419.1 U74630 *Ricinus communis*  
calreticulin.

AAB71420.1 U74631 *Ricinus communis*  
endoplasmic reticulum calcium binding protein. calreticulin.

CAA95999.1 Z71395 *Nicotiana glauca*  
ER Ca<sup>2+</sup> binding protein. calreticulin. cal1.

AAD17490.1 AF052040 *Berberis stolonifera*  
possible molecular chaperon. calreticulin. calcium binding protein of the endoplasmic reticulum.

AAD32207.1 AF134733 *Prunus armeniaca*  
calcium-binding protein calreticulin.

CAA61939.1 X89813 *Zea mays*  
Calreticulin precursor.

AAA32948.1 L27348 *Hordeum vulgare*  
calcium binding protein. calreticulin. CRH1.

AAA32949.1 L27349 *Hordeum vulgare*  
calcium binding protein. calreticulin. CRH2.

AAF01470.1 AF190454 *Zea mays*  
calreticulin. CRT. calcium binding protein.

CAB54526.1 AJ000765 *Chlamydomonas reinhardtii*  
calreticulin.

AAB70919.1 AF019376 *Brassica napus*  
calreticulin. Crt1. calcium binding protein; similar to *A. thaliana* calreticulin encoded by  
GenBank Accession Number U66343.

BAA85118.1 AB018243 *Solanum melongena*  
calreticulin-like protein. EEF22.

CAA54975.1 X78057 *Zea mays*  
calreticulin. CRH.

CAA57914.1 X82578 *Parthenium argentatum*  
calreticulin. 111R.

AAK15502.1 AF325720 *Pennisetum ciliare*  
calreticulin-like protein. Pcp4.

BAA77025.1 AB026251 *Lithospermum erythrorhizon*  
calreticulin.

SEQ ID NO: 596

BAA12206.1 D84061 *Spinacia oleracea*  
phosphoserine aminotransferase.

SEQ ID NO: 597

AAF73124.1 AF148534 *Lycopersicon esculentum*  
whitefly-induced gp91-phox. Wf1. mammalian gp91-phox homolog; respiratory burst  
oxidase homolog; putative NADPH oxidase; similar to neutrophil NADPH oxidase gp91-  
phox subunit; similar to *Arabidopsis thaliana* Rboh and *Oryza sativa* Rboh; similar to  
*Caenorhabditis elegans* PhoX-like proteins.

AAF73104.1 AF147783 *Lycopersicon esculentum*  
whitefly-induced gp91-phox. Wf1. mammalian gp91-phox homolog; respiratory burst  
oxidase homolog; putative NADPH oxidase; similar to neutrophil NADPH oxidase gp1-phox  
subunit; similar to *Arabidopsis thaliana* Rboh and *Oryza sativa* Rboh.

AAD25300.1 AF088276 *Lycopersicon esculentum*  
NADPH oxidase. RBOH1. gp91; phox homolog.

AAB87790.1 AF015302 *Oryza sativa*  
RbohA<sub>OSP</sub>. putative NADPH oxidase; intrinsic plasma membrane protein; similar to the  
neutrophil NADPH oxidase gp91phox subunit; contains RanGAP1 domain; similar to *O.*  
*sativa* NAD(P)H oxidase sequence with GenBank Accession Number X93301.

CAA63704.1 X93301 *Oryza sativa*  
NAD(P)H oxidase. rbohA.

AAD24966.1 AF109150 *Lycopersicon esculentum*  
NADPH oxidase. gp91; phox homolog.

AAD25225.1 AF088279 *Potamogeton crispus*  
NADPH oxidase. RBOH1. gp91; phox homolog.

SEQ ID NO: 599

CAC21424.1 AJ278332 *Lycopersicon esculentum*  
involved in octadecanoid biosynthesis. 12-oxophytodienoate reductase 3. opr3.

BAB40340.1 AB044940 *Pisum sativum*  
12-oxophytodienoic acid 10, 11-reductase. OPDAR.



|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| CAB43506.1     | AJ242551 | Lycopersicon esculentum       | involved in jasmonic acid biosynthesis. 12-oxophytodienoate reductase. opr.                 |
| CAC21423.1     | AJ278331 | Lycopersicon esculentum       | involved in octadecanoid biosynthesis. putative 12-oxophytodienoate reductase 2. opr2.      |
| CAB97122.1     | AJ297421 | Daucus carota                 | oxidoreductase. putative 12-oxophytodienoate reductase. opr2.                               |
| SEQ ID NO: 602 |          |                               |   |
| AAB37246.1     | U58971   | Nicotiana tabacum             | calmodulin-binding protein. TCB60.  |
| SEQ ID NO: 603 |          |                               |   |
| AAK15006.1     | AF233433 | Brassica napus                | arginase. similar to Arabidopsis thaliana and Glycine max arginase.                         |
| AAK07744.1     | AF130440 | Pinus taeda                   | converts L-arginine to L-ornithine and urea. arginase. ARS20. L-arginine amidinohydrolase.  |
| AAC04613.1     | AF035671 | Glycine max                   | hydrolysis of L-Arg to L-Orn and urea. arginase. pAG1. L-Arg amidinohydrolase.              |
| SEQ ID NO: 605 |          |                               |   |
| CAA81210.1     | Z26251   | Helianthus tuberosus          | Catalyzes the reduction of cytochrome P450 in microsomes. NADPH-ferrihemoprotein reductase. |
| AAB02721.1     | U58629   | Helianthus tuberosus          | NADPH-ferrihemoprotein oxidoreductase. HTR2. NADPH-cytochrome P450 reductase isoform.       |
| CAC27143.1     | AJ132538 | Picea abies                   | NADPH-cytochrome P450 reductase.  |
| AAA79131.1     | U10545   | Chlamydomonas reinhardtii     | ferredoxin-NADP+ reductase. fnr.  |
| CAA55406.1     | X78851   | Chlamydomonas reinhardtii     | ferredoxin NADP reductase.  |
| AAB40978.1     | U22328   | Volvox carteri                | ferredoxin-NADP+ reductase. fnr. FNR.   |
| AAA33029.1     | M25528   | Mesembryanthemum crystallinum | ferredoxin-NADP+ reductase precursor (fnrA; EC 1.6.7.1).                                    |
| CAA30978.1     | X12446   | Pisum sativum                 | ferredoxin-NADH+ reductase preprotein (AA -52 to 308).                                      |
| BAA13417.1     | D87547   | Oryza sativa                  | precursor ferredoxin-NADP+ oxidoreductase.  |
| AAA21758.1     | U14956   | Vicia faba                    | photosynthetic electron transport. ferredoxin NADP+ reductase precursor. fnr.               |
| AAA34029.1     | M86349   | Spinacia oleracea             | ferredoxin-NADP oxidoreductase. precursor protein.  |
| CAB71293.1     | AJ250378 | Capsicum annuum               | chloroplast ferredoxin-NADP+ oxidoreductase precursor. fnr.                                 |

|                |          |                        |  |
|----------------|----------|------------------------|--|
| CAA74359.1     | Y14032   | Nicotiana tabacum      | ferredoxin--NADP(+) reductase. fnr.  |
| BAA88236.1     | AB035644 | Zea mays               | NADP+ oxidoreductase. ferredoxin. L-FNRI.  |
| BAA88237.1     | AB035645 | Zea mays               | NADP+ oxidoreductase. ferredoxin. L-FNRII.   |
| BAA20365.1     | AB004307 | Nicotiana tabacum      | ferredoxin-NADP oxidoreductase.  |
| BAA07479.1     | D38445   | Oryza sativa           | root ferredoxin-NADP+ reductase. root FNR. nitrate assimilation pathway.   |
| BAA02248.1     | D12815   | Oryza sativa           | ferredoxin-NADP+ reductase enzyme.   |
| BAA04232.1     | D17410   | Oryza sativa           | ferredoxin-NADP+ reductase.  |
| BAA90642.1     | AP001129 | Oryza sativa           | ESTs AU078647(E1557),C72400(E1557) correspond to a region of the predicted gene.; Similar to Oryza sativa, ferredoxin-NADP+ reductase. (D17790). |
| BAA85425.1     | AP000616 | Oryza sativa           | ESTs AU078647(E1557),C72400(E1557) correspond to a region of the predicted gene.; similar to ferredoxin-NADP+ reductase (D17790).                |
| BAA04616.1     | D17790   | Oryza sativa           | ferredoxin-NADP+ reductase.  |
| AAB40034.1     | U10418   | Zea mays               | ferredoxin-NADP reductase precursor.   |
| CAA67796.1     | X99419   | Pisum sativum          | ferredoxin NADP oxidoreductase.  |
| AAK09367.1     | AF321525 | Pisum sativum          | ferredoxin-NADP+ reductase. FNR.   |
| AAK09370.1     | AF321528 | Pisum sativum          | ferredoxin-NADP+ reductase. FNR.   |
| AAK09369.1     | AF321527 | Pisum sativum          | ferredoxin-NADP+ reductase. FNR.   |
| AAK09368.1     | AF321526 | Pisum sativum          | ferredoxin-NADP+ reductase. FNR.   |
| SEQ ID NO: 606 |          |                        |  |
| BAA82107.1     | AB022693 | Nicotiana tabacum      | transcription factor. NtWRKY1.   |
| AAC31956.1     | AF080595 | Pimpinella brachycarpa | zinc finger protein. ZFP1. WRKY1.  |
| AAC49527.1     | U48831   | Petroselinum crispum   | WRKY1. contains two WRKY domains; WRKY-type DNA-binding protein; sequence-specific DNA-binding protein.  |

|            |          |                      |  |
|------------|----------|----------------------|--|
| AAD55974.1 | AF121353 | Petroselinum crispum | zinc-finger type transcription factor WRKY1. WRKY1.                                |
| BAA77383.1 | AB020590 | Nicotiana tabacum    | transcription factor NtWRKY2.  |
| AAD32677.1 | AF140554 | Avena sativa         | DNA-binding protein WRKY1. wrky1. putative transcription factor.                   |
| CAA88326.1 | Z48429   | Avena fatua          | binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein. |
| BAA86031.1 | AB026890 | Nicotiana tabacum    | transcription factor NtWRKY4.  |
| AAD16139.1 | AF096299 | Nicotiana tabacum    | DNA-binding protein 2. WRKY2. transcription factor.                                |
| AAC37515.1 | L44134   | Cucumis sativus      | SPF1-like DNA-binding protein.   |
| AAF23898.1 | AF193802 | Oryza sativa         | zinc finger transcription factor WRKY1.  |
| AAK16171.1 | AC079887 | Oryza sativa         | putative DNA-binding protein. OSJNBa0040E01.10.                                    |
| AAC49529.1 | U58540   | Petroselinum crispum | WRKY2. Contains two WRKY domains; WRKY-type DNA-binding protein.                   |
| AAD16138.1 | AF096298 | Nicotiana tabacum    | DNA-binding protein 1. WRKY1. transcription factor.                                |
| BAB19096.1 | AP002839 | Oryza sativa         | putative DNA-binding protein homolog. P0688A04.2.                                  |
| BAB19075.1 | AP002744 | Oryza sativa         | putative DNA-binding protein homolog. P0006C01.17.                                 |
| AAK16170.1 | AC079887 | Oryza sativa         | putative DNA binding protein. OSJNBa0040E01.4.                                     |
| CAB97004.1 | AJ278507 | Solanum tuberosum    | putative transcription factor. WRKY DNA binding protein. WRKY1.                    |
| BAB18313.1 | AP002865 | Oryza sativa         | putative WRKY DNA binding protein. P0034C11.1. contains EST C26525(C12525).        |
| AAD38283.1 | AC007789 | Oryza sativa         | putative WRKY DNA binding protein. OSJNBa0049B20.9.                                |
| BAB40073.1 | AP003074 | Oryza sativa         | putative WRKY DNA binding protein. OSJNBa0004G10.20. contains EST C26525(C12525).  |
| AAC49528.1 | U56834   | Petroselinum crispum | DNA-binding. WRKY3. WRKY-type DNA-binding protein.                                 |
| BAB16432.1 | AB041520 | Nicotiana tabacum    | WRKY transcription factor Nt-SubD48. Nt-SubD48.                                    |
| BAA77358.1 | AB020023 | Nicotiana tabacum    | WRKY domain Zn-finger type DNA-binding protein. DNA-binding protein NtWRKY3.       |

|                |          |                             |  |
|----------------|----------|-----------------------------|--|
| AAD32676.1     | AF140553 | <i>Avena sativa</i>         | DNA-binding protein WRKY3. wrky3. putative transcription factor.   |
| CAA88331.1     | Z48431   | <i>Avena fatua</i>          | binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein.   |
| AAG35658.1     | AF204925 | <i>Petroselinum crispum</i> | transcription factor WRKY4. WRKY4. binds to W box (TTGACC) elements.   |
| AAG46150.1     | AC018727 | <i>Oryza sativa</i>         | putative DNA-binding protein. OSJNBa0056G17.18.  |
| AAD27591.1     | AF121354 | <i>Petroselinum crispum</i> | binds sequence specifically to W Boxes (TTGACC). transcription factor. WRKY3. sequence specific DNA-binding protein.                 |
| AAG35659.1     | AF204926 | <i>Petroselinum crispum</i> | transcription factor WRKY5. WRKY5. binds to W box (TTGACC) elements.   |
| CAB66338.1     | AJ279697 | <i>Betula pendula</i>       | wrky-type DNA binding protein. wrky.   |
| AAF61864.1     | AF193771 | <i>Nicotiana tabacum</i>    | DNA-binding protein 4. WRKY4. transcription factor.  |
| AAF61863.1     | AF193770 | <i>Nicotiana tabacum</i>    | DNA-binding protein 3. WRKY3. transcription factor.  |
| SEQ ID NO: 608 |          |                             |  |
| BAA87853.1     | AP000816 | <i>Oryza sativa</i>         | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308). |
| AAK21965.1     | AY028699 | <i>Brassica napus</i>       | receptor protein kinase PERK1.   |
| BAB21240.1     | AP002953 | <i>Oryza sativa</i>         | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).   |
| BAB19337.1     | AP003044 | <i>Oryza sativa</i>         | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).   |
| BAA92836.1     | AB032473 | <i>Brassica oleracea</i>    | S18 S-locus receptor kinase. SRK18.  |
| AAB61708.1     | U93048   | <i>Daucus carota</i>        | somatic embryogenesis receptor-like kinase. SERK.  |
| BAA23676.1     | AB000970 | <i>Brassica rapa</i>        | receptor kinase 1. BcRK1.  |
| AAK00425.1     | AC069324 | <i>Oryza sativa</i>         | Putative protein kinase. OSJNBa0071K19.11.   |
| BAB39873.1     | AP002882 | <i>Oryza sativa</i>         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| AAD21872.1     | AF078082 | <i>Phaseolus vulgaris</i>   | receptor-like protein kinase homolog RK20-1.   |

|            |          |                               |  |
|------------|----------|-------------------------------|--|
| CAB41878.1 | Y18259   | Brassica oleracea             | SRK5 protein. SRK5. receptor-like kinase.  |
| AAA62232.1 | U00443   | Brassica napus                | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| AAA33000.1 | M76647   | Brassica oleracea             | receptor protein kinase. SKR6.   |
| AAG59657.1 | AC084319 | Oryza sativa                  | putative protein kinase. OSJNBa0004B24.20.   |
| CAB41879.1 | Y18260   | Brassica oleracea             | SRK15 protein. SRK15. receptor-like kinase.  |
| CAA73134.1 | Y12531   | Brassica oleracea             | serine/threonine kinase. BRLK.   |
| BAB16871.1 | AP002537 | Oryza sativa                  | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).       |
| BAB18292.1 | AP002860 | Oryza sativa                  | putative receptor-like protein kinase. P0409B08.19.  |
| BAA06285.1 | D30049   | Brassica rapa                 | S-receptor kinase SRK9.  |
| BAA21132.1 | D88193   | Brassica rapa                 | S-receptor kinase. SRK9 (B.c).   |
| CAB89179.1 | AJ245479 | Brassica napus subsp. napus   | ser /thr kinase. S-locus receptor kinase. srk.   |
| AAA33008.1 | M97667   | Brassica napus                | serine/threonine kinase receptor.  |
| CAA67145.1 | X98520   | Brassica oleracea             | receptor-like kinase. SFR2.  |
| CAA73133.1 | Y12530   | Brassica oleracea             | serine /threonine kinase. ARLK.  |
| AAF66615.1 | AF142596 | Nicotiana tabacum             | LRR receptor-like protein kinase.  |
| AAF76314.1 | AF220603 | Lycopersicon esculentum       | Fen kinase. Lescfen.   |
| AAK11568.1 | AF318492 | Lycopersicon hirsutum         | Pto-like protein kinase B. LhirPtoB.   |
| CAA74662.1 | Y14286   | Brassica oleracea             | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| AAB47424.1 | U59317   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Fen. Fen. fenthion sensitivity gene from tomato.                                 |
| AAB47422.1 | U59318   | Lycopersicon esculentum       | serine/threonine protein kinase Fen. fen. allele of Fen, fenthion sensitivity gene from tomato.                  |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAC48932.1     | U13923   | Lycopersicon pimpinellifolium | Fen. putative serine/threonine protein kinase; similar to product encoded by Lycopersicon pimpinellifolium Pto gene, GenBank Accession Number U02271; Fen is a member of the Pto gene family. |
| AAF76307.1     | AF220602 | Lycopersicon pimpinellifolium | Fen kinase.   |
| BAA07576.1     | D38563   | Brassica rapa                 | receptor protein kinase SRK8.   |
| AAG16628.1     | AY007545 | Brassica napus                | protein serine/threonine kinase BNK1.   |
| CAA97692.1     | Z73295   | Catharanthus roseus           | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.  |
| CAA79355.1     | Z18921   | Brassica oleracea             | S-receptor kinase-like protein.   |
| AAB93834.1     | U82481   | Zea mays                      | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.  |
| BAA92837.1     | AB032474 | Brassica oleracea             | S60 S-locus receptor kinase. SRK60.   |
| AAA33915.1     | L27821   | Oryza sativa                  | receptor type serine/threonine kinase. protein kinase.  |
| SEQ ID NO: 610 |          |                               |   |
| BAB17348.1     | AP002747 | Oryza sativa                  | putative receptor kinase. P0698G03.32.  |
| BAB17345.1     | AP002747 | Oryza sativa                  | putative receptor kinase. P0698G03.29.  |
| BAB17126.1     | AP002867 | Oryza sativa                  | putative receptor kinase. P0463F06.16.  |
| BAB39451.1     | AP003338 | Oryza sativa                  | putative receptor kinase. OJ1212_B09.24.  |
| BAB17342.1     | AP002747 | Oryza sativa                  | putative receptor kinase. P0698G03.26.  |
| AAC27489.1     | AF077130 | Oryza sativa                  | receptor-like protein kinase.   |
| AAF78018.1     | AF238474 | Oryza sativa                  | receptor-like kinase. RLG16. protein kinase.  |
| AAC02535.1     | AF044260 | Oryza sativa                  | receptor serine/threonine kinase. protein kinase.   |
| BAB39435.1     | AP003338 | Oryza sativa                  | putative receptor kinase. OJ1212_B09.2.   |
| AAD46917.1     | AF164021 | Oryza sativa                  | receptor kinase.  |

|            |          |                   |   |
|------------|----------|-------------------|---|
| BAB39438.1 | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.7.   |
| BAB17339.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.23.  |
| AAC49629.1 | U51330   | Triticum aestivum | rust resistance kinase Lr10. LRK10.   |
| BAB17321.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.1.   |
| BAB17129.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.20.  |
| AAF68398.1 | AF237568 | Oryza sativa      | receptor-like protein kinase. RLG2.   |
| BAB17331.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.12.  |
| BAB17139.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.31.  |
| AAC01746.1 | AF044489 | Oryza sativa      | receptor-like protein kinase. drpk1.  |
| AAF78016.1 | AF238472 | Oryza sativa      | receptor-like kinase. RLG15. protein kinase.  |
| BAB39441.1 | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.11.  |
| BAB17116.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.3.   |
| AAF78044.1 | AF248493 | Oryza sativa      | receptor-like kinase. RLG18. protein kinase.  |
| AAD44029.1 | AF085164 | Hordeum vulgare   | receptor-like kinase LRK10.   |
| BAB17332.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.13. contains ESTs D47575(S13157),AU032665(S13157). |
| BAB17347.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.31. contains ESTs D47575(S13157),AU032665(S13157). |
| BAB17344.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.28. contains ESTs D47575(S13157),AU032665(S13157). |
| BAB17337.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.21. contains ESTs D47575(S13157),AU032665(S13157). |
| BAB17335.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.17. contains ESTs D47575(S13157),AU032665(S13157). |
| BAB17115.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.2.   |
| BAB39440.1 | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.10.  |

|                |          |                      |   |
|----------------|----------|----------------------|---|
| BAB39437.1     | AP003338 | Oryza sativa         | receptor-like kinase. OJ1212_B09.6.   |
| AAF78019.1     | AF238475 | Oryza sativa         | receptor-like kinase. RLG17. protein kinase.  |
| BAB17127.1     | AP002867 | Oryza sativa         | putative receptor kinase. P0463F06.17.  |
| AAD46916.1     | AF164020 | Oryza sativa         | receptor kinase.  |
| AAF78021.1     | AF238477 | Oryza sativa         | receptor-like kinase. RLG5. protein kinase.   |
| AAD46417.1     | AF100767 | Oryza sativa         | receptor-like kinase. 8ARK3. similar to wheat ARK1AS.   |
| AAD46415.1     | AF100765 | Oryza sativa         | receptor-like kinase. 8ARK1. similar to wheat ARK1AS.   |
| BAB17128.1     | AP002867 | Oryza sativa         | putative receptor kinase. P0463F06.18.  |
| AAF68400.1     | AF237570 | Oryza sativa         | receptor-like protein kinase. RLG3.   |
| BAB17323.1     | AP002747 | Oryza sativa         | putative receptor kinase. P0698G03.3.   |
| BAB17131.1     | AP002867 | Oryza sativa         | putative receptor kinase. P0463F06.22.  |
| AAF68397.1     | AF237567 | Oryza sativa         | receptor-like protein kinase. RLG1.   |
| AAF78015.1     | AF238471 | Oryza sativa         | receptor-like kinase. RLG10. protein kinase.  |
| AAD43962.1     | U78762   | Triticum aestivum    | receptor-like kinase ARK1AS. ARK1AS.  |
| SEQ ID NO: 611 |          |                      |   |
| BAA78764.1     | AB023482 | Oryza sativa         | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAK11674.1     | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| AAF43496.1     | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| AAG16628.1     | AY007545 | Brassica napus       | protein serine/threonine kinase BNK1.   |
| BAA94509.1     | AB041503 | Populus nigra        | protein kinase 1. PnPK1.  |
| BAA94510.1     | AB041504 | Populus nigra        | protein kinase 2. PnPK2.  |



|            |          |                               |   |
|------------|----------|-------------------------------|---|
| BAB07999.1 | AP002525 | Oryza sativa                  | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB03429.1 | AP002817 | Oryza sativa                  | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).                         |
| BAA87853.1 | AP000816 | Oryza sativa                  | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB16871.1 | AP002537 | Oryza sativa                  | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).  |
| BAB39409.1 | AP002901 | Oryza sativa                  | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| CAB51834.1 | 00069    | Oryza sativa                  | 11332.5. contains eukaryotic protein kinase domain PF.  |
| BAB21240.1 | AP002953 | Oryza sativa                  | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| AAK00425.1 | AC069324 | Oryza sativa                  | Putative protein kinase. OSJNBa0071K19.11.  |
| BAB39873.1 | AP002882 | Oryza sativa                  | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| AAK11567.1 | AF318491 | Lycopersicon hirsutum         | Pto-like protein kinase F. LhirPtoF.  |
| BAA90808.1 | AP001168 | Oryza sativa                  | Similar to putative receptor-like protein kinase (AL035679).  |
| AAK21965.1 | AY028699 | Brassica napus                | receptor protein kinase PERK1.  |
| AAC48932.1 | U13923   | Lycopersicon pimpinellifolium | Fen. putative serine/threonine protein kinase; similar to product encoded by Lycopersicon pimpinellifolium Pto gene, GenBank Accession Number U02271; Fen is a member of the Pto gene family. |
| AAB47424.1 | U59317   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Fen. Fen. fenthion sensitivity gene from tomato.  |
| AAF76307.1 | AF220602 | Lycopersicon pimpinellifolium | Fen kinase.   |
| AAK11566.1 | AF318490 | Lycopersicon hirsutum         | Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.   |
| AAG03090.1 | AC073405 | Oryza sativa                  | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| AAF76314.1 | AF220603 | Lycopersicon esculentum       | Fen kinase. Lescfen.  |

|            |          |  |
|------------|----------|--|
| AAB47421.1 | U59316   | <i>Lycopersicon esculentum</i><br>serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.                     |
| AAF76313.1 | AF220603 | <i>Lycopersicon esculentum</i><br>Pto kinase. LescPth5.  |
| AAB47422.1 | U59318   | <i>Lycopersicon esculentum</i><br>serine/threonine protein kinase Fen. fen. allele of Fen, fenthion sensitivity gene from tomato.                                |
| AAG59657.1 | AC084319 | <i>Oryza sativa</i><br>putative protein kinase. OSJNBa0004B24.20.  |
| AAF76306.1 | AF220602 | <i>Lycopersicon pimpinellifolium</i><br>Pto kinase.  |
| AAB47423.1 | U59315   | <i>Lycopersicon pimpinellifolium</i><br>serine/threonine protein kinase Pto. Pto. bacterial speck disease resistance in tomato; disease resistance gene.         |
| AAC48914.1 | U02271   | <i>Lycopersicon pimpinellifolium</i><br>protein kinase.  |
| AAK11568.1 | AF318492 | <i>Lycopersicon hirsutum</i><br>Pto-like protein kinase B. LhirPtoB.   |
| BAB21241.1 | AP002953 | <i>Oryza sativa</i><br>Putative Pto kinase interactor 1. P0426D06.21. contains ESTs AU108280(E0721),D48017(S13927).  |
| AAC27894.1 | AF023164 | <i>Zea mays</i><br>leucine-rich repeat transmembrane protein kinase 1. ltk1.   |
| AAF91337.1 | AF249318 | <i>Glycine max</i><br>Pti1 kinase-like protein. Pti1b. protein kinase.   |
| AAC61805.1 | U28007   | <i>Lycopersicon esculentum</i><br>serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.   |
| CAA97692.1 | Z73295   | <i>Catharanthus roseus</i><br>receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis. |
| AAF91336.1 | AF249317 | <i>Glycine max</i><br>Pti1 kinase-like protein. Pti1a. protein kinase.   |
| AAF66615.1 | AF142596 | <i>Nicotiana tabacum</i><br>LRR receptor-like protein kinase.  |
| AAB61708.1 | U93048   | <i>Daucus carota</i><br>somatic embryogenesis receptor-like kinase. SERK.  |
| AAK11569.1 | AF318493 | <i>Lycopersicon hirsutum</i><br>Pto-like protein kinase D. LhirPtoD.   |
| BAA92221.1 | AP001278 | <i>Oryza sativa</i><br>Similar to <i>Arabidopsis thaliana</i> chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).                     |
| BAA87852.1 | AP000816 | <i>Oryza sativa</i><br>Similar to putative Ser/Thr protein kinase. (AC004218).   |

|                |          |                             |   |
|----------------|----------|-----------------------------|---|
| AAC27895.1     | AF023165 | Zea mays                    | leucine-rich repeat transmembrane protein kinase 2. ltk2.   |
| SEQ ID NO: 612 |          |                             |   |
| CAA54613.1     | X77463   | Manihot esculenta           | UTP-glucose glucosyltransferase. CGT6.  |
| CAA54609.1     | X77459   | Manihot esculenta           | UTP-glucose glucosyltransferase. CGT1.  |
| CAA54611.1     | X77461   | Manihot esculenta           | UTP-glucose glucosyltransferase. CGT2.  |
| CAA54612.1     | X77462   | Manihot esculenta           | UTP-glucose glucosyltransferase. CGT5.  |
| BAB17182.1     | AP002843 | Oryza sativa                | putative UTP-glucose glucosyltransferase. P0407B12.19.  |
| BAB17176.1     | AP002843 | Oryza sativa                | putative UTP-glucose glucosyltransferase. P0407B12.13.  |
| CAB56231.1     | Y18871   | Dorotheanthus bellidiformis | betanidin-5-O-glucosyltransferase.  |
| BAA83484.1     | AB031274 | Scutellaria baicalensis     | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.   |
| AAB36653.1     | U32644   | Nicotiana tabacum           | immediate-early salicylate-induced glucosyltransferase. IS5a.                                       |
| AAK28303.1     | AF346431 | Nicotiana tabacum           | phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.                                  |
| CAA59450.1     | X85138   | Lycopersicon esculentum     | twil. homologous to glucosyltransferases.   |
| AAB36652.1     | U32643   | Nicotiana tabacum           | immediate-early salicylate-induced glucosyltransferase. IS10a.                                      |
| AAK28304.1     | AF346432 | Nicotiana tabacum           | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.                                  |
| BAA89009.1     | AB027455 | Petunia x hybrida           | anthocyanin 5-O-glucosyltransferase. PH1.   |
| AAG25643.1     | AF303396 | Phaseolus vulgaris          | UDP-glucosyltransferase HRA25. putative; defense associated.  |
| AAF61647.1     | AF190634 | Nicotiana tabacum           | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAA93039.1     | AB033758 | Citrus unshiu               | limonoid UDP-glucosyltransferase. LGTase.   |
| AAB48444.1     | U82367   | Solanum tuberosum           | UDP-glucose glucosyltransferase.  |
| AAD21086.1     | AF127218 | Forsythia x intermedia      | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT. |

|                |          |                    |   |
|----------------|----------|--------------------|---|
| AAK16175.1     | AC079887 | Oryza sativa       | putative glucosyltransferase. OSJNBa0040E01.15.   |
| CAC09351.1     | AL442007 | Oryza sativa       | putative glucosyltransferase. H0212B02.7.   |
| AAD04166.1     | AF101972 | Phaseolus lunatus  | catalyzes formation of O-glucosylzeatin from zeatin and UDPG; catalyzes formation of O-xylosylzeatin from zeatin and UDPX. zeatin O-glucosyltransferase. ZOG1. cytokinin O-glucosyltransferase. |
| AAF17077.1     | AF199453 | Sorghum bicolor    | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |
| BAA12737.1     | D85186   | Gentiana triflora  | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| BAB07962.1     | AP002524 | Oryza sativa       | putative anthocyanin 5-O-glucosyltransferase. P0406H10.16. contains ESTs AU067881(C10481),AU067882(C10481).   |
| BAA36421.1     | AB013596 | Perilla frutescens | UDP-glucose:anthocynin 5-O-glucosyltransferase. PF3R4.  |
| CAA54610.1     | X77460   | Manihot esculenta  | UTP-glucose glucosyltransferase. CGT4.  |
| AAF98390.1     | AF287143 | Brassica napus     | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT.         |
| BAA36423.1     | AB013598 | Verbena x hybrida  | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| CAA30760.1     | X07937   | Zea mays           | UDPglucose flavonoid glycosyl transferase. Bz-W22.  |
| CAA30761.1     | X07940   | Zea mays           | UDPglucose flavonoid glycosyl-transferase. Bz-McC.  |
| CAA31855.1     | X13500   | Zea mays           | UDPglucose:flavonol 3-O-glucosyltransferase.  |
| BAA19659.1     | AB002818 | Perilla frutescens | flavonoid 3-O-glucosyltransferase. UDP glucose.   |
| BAA89008.1     | AB027454 | Petunia x hybrida  | anthocyanidin 3-O-glucosyltransferase. PGT8.  |
| AAK16172.1     | AC079887 | Oryza sativa       | putative glucosyltransferase. OSJNBa0040E01.14.   |
| SEQ ID NO: 613 |          |                    |   |
| CAA54609.1     | X77459   | Manihot esculenta  | UTP-glucose glucosyltransferase. CGT1.  |
| CAA54613.1     | X77463   | Manihot esculenta  | UTP-glucose glucosyltransferase. CGT6.  |

|            |          |                             |   |
|------------|----------|-----------------------------|---|
| CAA54611.1 | X77461   | Manihot esculenta           | UTP-glucose glucosyltransferase. CGT2.  |
| CAA54612.1 | X77462   | Manihot esculenta           | UTP-glucose glucosyltransferase. CGT5.  |
| BAB17182.1 | AP002843 | Oryza sativa                | putative UTP-glucose glucosyltransferase. P0407B12.19.  |
| BAB17176.1 | AP002843 | Oryza sativa                | putative UTP-glucose glucosyltransferase. P0407B12.13.  |
| CAB56231.1 | Y18871   | Dorotheanthus bellidiformis | betanidin-5-O-glucosyltransferase.  |
| AAB36653.1 | U32644   | Nicotiana tabacum           | immediate-early salicylate-induced glucosyltransferase. IS5a.   |
| AAK28303.1 | AF346431 | Nicotiana tabacum           | phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.  |
| AAB36652.1 | U32643   | Nicotiana tabacum           | immediate-early salicylate-induced glucosyltransferase. IS10a.  |
| AAK28304.1 | AF346432 | Nicotiana tabacum           | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.  |
| CAA59450.1 | X85138   | Lycopersicon esculentum     | twil. homologous to glucosyltransferases.   |
| BAA83484.1 | AB031274 | Scutellaria baicalensis     | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.   |
| AAF61647.1 | AF190634 | Nicotiana tabacum           | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAA93039.1 | AB033758 | Citrus unshiu               | limonoid UDP-glucosyltransferase. LGTase.   |
| AAB48444.1 | U82367   | Solanum tuberosum           | UDP-glucose glucosyltransferase.  |
| BAA89009.1 | AB027455 | Petunia x hybrida           | anthocyanin 5-O-glucosyltransferase. PH1.   |
| BAA36421.1 | AB013596 | Perilla frutescens          | UDP-glucose:anthocynsin 5-O-glucosyltransferase. PF3R4.   |
| CAC09351.1 | AL442007 | Oryza sativa                | putative glucosyltransferase. H0212B02.7.   |
| BAA12737.1 | D85186   | Gentiana triflora           | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| AAG25643.1 | AF303396 | Phaseolus vulgaris          | UDP-glucosyltransferase HRA25. putative; defense associated.  |
| AAD04166.1 | AF101972 | Phaseolus lunatus           | catalyzes formation of O-glucosylzeatin from zeatin and UDPG; catalyzes formation of O-xylosylzeatin from zeatin and UDPX. zeatin O-glucosyltransferase. ZOG1. cytokinin O-glucosyltransferase. |

|                |          |                             |   |
|----------------|----------|-----------------------------|---|
| CAA54610.1     | X77460   | Manihot esculenta           | UTP-glucose glucosyltransferase. CGT4.  |
| AAF98390.1     | AF287143 | Brassica napus              | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT. |
| AAK16172.1     | AC079887 | Oryza sativa                | putative glucosyltransferase. OSJNBa0040E01.14.   |
| BAA36423.1     | AB013598 | Verbena x hybrida           | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| AAD21086.1     | AF127218 | Forsythia x intermedia      | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.   |
| AAF17077.1     | AF199453 | Sorghum bicolor             | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |
| AAK16175.1     | AC079887 | Oryza sativa                | putative glucosyltransferase. OSJNBa0040E01.15.   |
| BAB07962.1     | AP002524 | Oryza sativa                | putative anthocyanin 5-O-glucosyltransferase. P0406H10.16. contains ESTs AU067881(C10481),AU067882(C10481).   |
| CAA31855.1     | X13500   | Zea mays                    | UDPglucose:flavonol 3-O-glucosyltransferase.  |
| BAA36422.1     | AB013597 | Perilla frutescens          | UDP-glucose:anthocyanin 5-O-glucosyltransferase homologue. PF3R6.   |
| BAA89008.1     | AB027454 | Petunia x hybrida           | anthocyanidin 3-O-glucosyltransferase. PGT8.  |
| SEQ ID NO: 614 |          |                             |   |
| AAD21872.1     | AF078082 | Phaseolus vulgaris          | receptor-like protein kinase homolog RK20-1.  |
| CAA73134.1     | Y12531   | Brassica oleracea           | serine/threonine kinase. BRLK.  |
| AAB93834.1     | U82481   | Zea mays                    | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.  |
| AAA33008.1     | M97667   | Brassica napus              | serine/threonine kinase receptor.   |
| CAB89179.1     | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.  |
| AAA33000.1     | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.  |
| BAA23676.1     | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.   |

|            |          |                   |  |
|------------|----------|-------------------|--|
| BAA06285.1 | D30049   | Brassica rapa     | S-receptor kinase SRK9.  |
| BAA21132.1 | D88193   | Brassica rapa     | S-receptor kinase. SRK9 (B.c).   |
| CAA74661.1 | Y14285   | Brassica oleracea | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.   |
| BAA92837.1 | AB032474 | Brassica oleracea | S60 S-locus receptor kinase. SRK60.  |
| BAA07577.2 | D38564   | Brassica rapa     | receptor protein kinase SRK12.   |
| BAA07576.1 | D38563   | Brassica rapa     | receptor protein kinase SRK8.  |
| CAA73133.1 | Y12530   | Brassica oleracea | serine /threonine kinase. ARLK.  |
| AAC23542.1 | U20948   | Ipomoea trifida   | receptor protein kinase. IRK1.   |
| CAA79355.1 | Z18921   | Brassica oleracea | S-receptor kinase-like protein.  |
| CAA67145.1 | X98520   | Brassica oleracea | receptor-like kinase. SFR2.  |
| CAA74662.1 | Y14286   | Brassica oleracea | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| BAB21001.1 | AB054061 | Brassica rapa     | S locus receptor kinase. SRK22.  |
| AAA62232.1 | U00443   | Brassica napus    | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| CAB41879.1 | Y18260   | Brassica oleracea | SRK15 protein. SRK15. receptor-like kinase.  |
| CAB41878.1 | Y18259   | Brassica oleracea | SRK5 protein. SRK5. receptor-like kinase.  |
| BAA92836.1 | AB032473 | Brassica oleracea | S18 S-locus receptor kinase. SRK18.  |
| BAB18292.1 | AP002860 | Oryza sativa      | putative receptor-like protein kinase. P0409B08.19.  |
| AAD52097.1 | AF088885 | Nicotiana tabacum | receptor-like kinase CHRK1. Chrk1.   |
| AAA33915.1 | L27821   | Oryza sativa      | receptor type serine/threonine kinase. protein kinase.   |
| BAB39873.1 | AP002882 | Oryza sativa      | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).                            |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| BAB16871.1     | AP002537 | Oryza sativa                  | putative protein kinase APK1A Arabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192), D25110(R3192).  |
| AAK21965.1     | AY028699 | Brassica napus                | receptor protein kinase PERK1.  |
| AAK00425.1     | AC069324 | Oryza sativa                  | Putative protein kinase. OSJNBa0071K19.11.  |
| BAA94529.2     | AP001800 | Oryza sativa                  | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| BAA94509.1     | AB041503 | Populus nigra                 | protein kinase 1. PnPK1.  |
| BAA94510.1     | AB041504 | Populus nigra                 | protein kinase 2. PnPK2.  |
| BAA87853.1     | AP000816 | Oryza sativa                  | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB39409.1     | AP002901 | Oryza sativa                  | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| AAG03090.1     | AC073405 | Oryza sativa                  | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| BAA92954.1     | AP001551 | Oryza sativa                  | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).   |
| BAA94518.1     | AP001800 | Oryza sativa                  | Similar to Arabidopsis thaliana chromosome 2 section 111 of 255; putative receptor-like protein kinase (AC002392).  |
| BAB07904.1     | AP002835 | Oryza sativa                  | putative S-receptor kinase. P0417G05.12.  |
| SEQ ID NO: 619 |          |                               |   |
| CAA72092.1     | Y11209   | Nicotiana tabacum             | protein disulfide-isomerase precursor. PDI.   |
| AAG13988.1     | AF298829 | Prunus avium                  | putative protein disulfide-isomerase. PDI.  |
| AAD02069.1     | AF036939 | Chlamydomonas reinhardtii     | redox-regulator of 5'UTR psbA mRNA binding complex and translation. protein disulfide isomerase. localized to ER and chloroplast.   |
| AAC49896.1     | AF027727 | Chlamydomonas reinhardtii     | involved in the redox-regulated binding of chloroplast poly(A)-binding protein to the 5'-UTR of psbA mRNA; regulates chloroplast translational activation. protein disulfide isomerase RB60. PDI. |
| AAD55566.1     | AF110784 | Volvox carteri f. nagariensis | protein disulfide isomerase precursor. pdi.   |
| AAB08519.1     | L39014   | Zea mays                      | protein disulfide isomerase. pdi. putative.   |



|                |          |                                |  |
|----------------|----------|--------------------------------|--|
| AAA70344.1     | L33250   | Hordeum vulgare                | catalyze the formation of disulfide bonds. disulfide isomerase. PDI. putative. |
| AAA70345.1     | L33251   | Hordeum vulgare                | catalyze the formation of disulfide bonds. disulfide isomerase. PDI. putative. |
| CAC21230.1     | AJ277379 | Triticum turgidum subsp. durum | catalyzes formation of disulfide bonds. protein disulfide isomerase. Pdi.      |
| AAA19660.1     | U11496   | Triticum aestivum              | protein disulfide isomerase. PDI.  |
| CAC21231.1     | AJ277380 | Triticum turgidum subsp. durum | catalyzes formation of disulfide bonds. protein disulfide isomerase. Pdi.      |
| CAC21229.1     | AJ277378 | Triticum turgidum subsp. durum | catalyzes formation of disulfide bonds. protein disulfide isomerase. Pdi.      |
| CAC21228.1     | AJ277377 | Triticum turgidum subsp. durum | catalyzes the formation of disulfide bonds. protein disulfide isomerase. Pdi.  |
| AAB05641.1     | U41385   | Ricinus communis               | protein disulphide isomerase PDI. molecular chaperone.                         |
| CAA77575.1     | Z11499   | Medicago sativa                | protein disulfide isomerase.   |
| AAD28260.1     | AF131223 | Datisca glomerata              | protein disulfide isomerase homolog. PDI.                                      |
| AAA70346.1     | L33252   | Hordeum vulgare                | catalyze the formation of disulfide bonds. disulfide isomerase. PDI. putative. |
| BAA92322.1     | AB039278 | Oryza sativa                   | protein disulfide isomerase. Pdi.  |
| BAA77026.1     | AB026252 | Lithospermum erythrorhizon     | disulfide-isomerase precursor.   |
| AAC79709.1     | AF093614 | Acetabularia acetabulum        | putative protein disulfide isomerase.  |
| SEQ ID NO: 620 |          |                                |  |
| CAA64413.1     | X94943   | Lycopersicon esculentum        | peroxidase. cevil6.  |
| AAA32676.1     | M37637   | Arachis hypogaea               | cationic peroxidase. PNC2.   |
| BAA82307.1     | AB027753 | Nicotiana tabacum              | peroxidase.  |
| AAB67737.1     | L77080   | Stylosanthes humilis           | cationic peroxidase.   |
| AAD37429.2     | AF149279 | Phaseolus vulgaris             | peroxidase 4 precursor. FBP4. secretory peroxidase.                            |
| CAA71494.1     | Y10468   | Spinacia oleracea              | peroxidase. prxr7.   |

|            |          |  |   |
|------------|----------|--|---|
| AAD37375.1 | AF145349 | Glycine max                            | peroxidase. Prx3.   |
| AAF63024.1 | AF244921 | Spinacia oleracea                      | hydrogen peroxide catabolism. peroxidase prx12 precursor. type III peroxidase.  |
| CAA66037.1 | X97351   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.  |
| BAB39274.1 | AP002971 | Oryza sativa                           | putative peroxidase. P0537A05.2.  |
| AAA65637.1 | L13654   | Lycopersicon esculentum                | peroxidase. TPX1.   |
| CAA40796.1 | X57564   | Armoracia rusticana                    | peroxidase. peroxidase precursor.   |
| AAD11482.1 | U51192   | Glycine max                            | peroxidase precursor. sEPa2.  |
| CAA80502.1 | Z22920   | Spirodela polyrrhiza                   | peroxidase.   |
| BAA77387.1 | AB024437 | Scutellaria baicalensis                | peroxidase 1.   |
| CAA59485.1 | X85228   | Triticum aestivum                      | peroxidase. POX2.   |
| BAA07663.1 | D42064   | Nicotiana tabacum                      | cationic peroxidase isozyme 38K precursor.  |
| BAA11853.1 | D83225   | Populus nigra                          | peroxidase.   |
| BAA07664.1 | D42065   | Nicotiana tabacum                      | cationic peroxidase isozyme 40K precursor.  |
| AAD37430.1 | AF149280 | Phaseolus vulgaris                     | peroxidase 5 precursor. FBP5. secretory peroxidase.   |
| AAD11481.1 | U51191   | Glycine max                            | peroxidase precursor. sEPa1.  |
| CAB94692.1 | AJ242742 | Ipomoea batatas                        | Removal of H <sub>2</sub> O <sub>2</sub> , oxidation of toxic reductants, defence response toward wounding.<br>peroxidase. pod. |
| BAA03644.1 | D14997   | Oryza sativa                           | peroxidase.   |
| AAD43561.1 | AF155124 | Gossypium hirsutum                     | bacterial-induced peroxidase precursor. Perx_Goshiko.   |
| BAA06334.1 | D30652   | Populus kitakamiensis                  | peroxidase.   |
| BAA92500.1 | AP001383 | Oryza sativa                           | ESTs D39300(R3292),AU030751(E60187) correspond to a region of the predicted gene.<br>Similar to peroxidase ATP6a. (X98774).     |

|            |          |  |   |
|------------|----------|--|---|
| BAA90365.1 | AP001081 | Oryza sativa                           | ESTs D24550(R2151),D24265(R1609),AU031848(R2151) correspond to a region of the predicted gene. Similar to cationic peroxidase isozyme 40K precursor (D42065). |
| BAA89584.1 | AP001073 | Oryza sativa                           | ESTs D24550(R2151),D24265(R1609),AU031848(R2151) correspond to a region of the predicted gene. Similar to cationic peroxidase isozyme 40K precursor (D42065). |
| AAF34416.1 | AF172282 | Oryza sativa                           | putative peroxidase. DUPR11.5.  |
| AAC49820.1 | AF014469 | Oryza sativa                           | peroxidase. POX5.1. wound inducible.  |
| CAA66034.1 | X97348   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.  |
| BAB39281.1 | AP002971 | Oryza sativa                           | putative peroxidase. P0537A05.10. contains ESTs D24657(R2329),AU082066(R2329).  |
| BAA14144.1 | D90116   | Armoracia rusticana                    | peroxidase isozyme.   |
| AAB97734.1 | AF014502 | Glycine max                            | seed coat peroxidase precursor. Ep. H <sub>2</sub> O <sub>2</sub> oxidoreductase; class III plant peroxidase.   |
| CAA37713.1 | X53675   | Triticum aestivum                      | peroxidase.   |
| AAC05277.1 | AF049881 | Linum usitatissimum                    | peroxidase FLXPER4. PER4.   |
| CAA66035.1 | X97349   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.  |
| BAA06335.1 | D30653   | Populus kitakamiensis                  | peroxidase.   |
| CAA39486.1 | X56011   | Triticum aestivum                      | peroxidase.   |
| BAA03911.1 | D16442   | Oryza sativa                           | peroxidase.   |
| BAA94962.1 | AB042103 | Asparagus officinalis                  | peroxidase. AspPOX1.  |
| AAC49821.1 | AF014470 | Oryza sativa                           | peroxidase. POXgX9. expressed in roots.   |
| AAD37427.1 | AF149277 | Phaseolus vulgaris                     | peroxidase 1 precursor. FBP1. secretory peroxidase.   |
| CAA76374.2 | Y16776   | Spinacia oleracea                      | peroxidase. prx10.  |
| BAA08499.1 | D49551   | Oryza sativa                           | peroxidase. poxN.   |
| CAA66036.1 | X97350   | Populus balsamifera subsp. trichocarpa | signal for ER. peroxidase.  |

|                |          |                           |  |
|----------------|----------|---------------------------|--|
| AAA34108.1     | J02979   | Nicotiana tabacum         | lignin-forming peroxidase precursor (EC 1.11.1.7).   |
| CAA62226.1     | X90693   | Medicago sativa           | peroxidase1B. prx1B.   |
| CAA59487.1     | X85230   | Triticum aestivum         | peroxidase. pox4.  |
| SEQ ID NO: 626 |          |                           |  |
| CAA98160.1     | Z73932   | Lotus japonicus           | GTP-binding protein. RAB1C. rab1C.   |
| BAA76422.1     | AB024994 | Cicer arietinum           | rab-type small GTP-binding protein.  |
| BAA02116.1     | D12548   | Pisum sativum             | GTP-binding protein.   |
| CAA69701.1     | Y08425   | Nicotiana plumbaginifolia | small GTP-binding protein. Rab1 subfamily.   |
| AAA80678.1     | U38464   | Lycopersicon esculentum   | small GTP-binding protein. LeRab1A. ; YPT1/Rab1A homolog LeRab1A.                                  |
| BAA02118.1     | D12550   | Pisum sativum             | GTP-binding protein.   |
| AAB97115.1     | U58854   | Glycine max               | small GTP-binding protein. sra2.   |
| CAA51011.1     | X72212   | Nicotiana tabacum         | ras-related GTP-binding protein. ypt2 homologue.   |
| AAF65510.1     | AF108883 | Capsicum annuum           | small GTP-binding protein.   |
| AAA80680.1     | U38466   | Lycopersicon esculentum   | small GTP-binding protein. LeRab1C. ; YPT1/Rab1A homolog LeRab1C.                                  |
| CAA98161.1     | Z73933   | Lotus japonicus           | GTP-binding protein. RAB1D. rab1D.   |
| BAA02117.1     | D12549   | Pisum sativum             | GTP-binding protein.   |
| CAA98162.1     | Z73934   | Lotus japonicus           | GTP-binding protein. RAB1E. rab1E.   |
| AAA50159.1     | L27417   | Glycine max               | GTP binding protein.   |
| AAB28535.1     | S66160   | Oryza sativa              | ric1. ras-related GTP binding protein possessing GTPase activity; This sequence comes from Fig. 1. |
| CAA98159.1     | Z73931   | Lotus japonicus           | GTP-binding protein. RAB1B. rab1B.   |
| BAA02115.1     | D12547   | Pisum sativum             | GTP-binding protein.   |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAA66447.1 | X97853   | Lotus japonicus         | GTP-binding protein. RAB1A. rab1A.  |
| AAD10389.1 | U35026   | Petunia x hybrida       | Rab1-like small GTP-binding protein.  |
| AAA80679.1 | U38465   | Lycopersicon esculentum | small GTP-binding protein. LeRab1B. ; Ypt1/Rab1A homolog LeRab1B.                       |
| BAB07961.1 | AP002524 | Oryza sativa            | putative GTP-binding protein. P0406H10.17. contains ESTs D23874(R0480),AU031678(R0480). |
| CAA98176.1 | Z73948   | Lotus japonicus         | GTP-binding protein. RAB8E. rab8E.  |
| CAA89021.1 | Z49152   | Beta vulgaris           | GTP-binding. small G protein.   |
| CAA98172.1 | Z73944   | Lotus japonicus         | GTP-binding protein. RAB8A. rab8A.  |
| CAA04701.1 | AJ001367 | Daucus carota           | small GTP-binding protein. Dc-Rab8.   |
| CAA90080.1 | Z49900   | Pisum sativum           | small GTP-binding protein.  |
| AAD46405.1 | AF096249 | Lycopersicon esculentum | ethylene-responsive small GTP-binding protein. ER43.                                    |
| CAA98174.1 | Z73946   | Lotus japonicus         | GTP-binding protein. RAB8C. rab8C.  |
| CAA90082.1 | Z49902   | Pisum sativum           | small GTP-binding protein.  |
| CAA49600.1 | X69980   | Lycopersicon esculentum | GTP-binding protein. ypt2.  |
| CAA98175.1 | Z73947   | Lotus japonicus         | GTP-binding protein. RAB8D. rab8D.  |
| CAA90081.1 | Z49901   | Pisum sativum           | small GTP-binding protein.  |
| AAB17726.1 | U38471   | Brassica rapa           | small GTP-binding protein rab. BRAB-1. small GTP-binding protein rab family.            |
| AAA34251.1 | L08128   | Volvox carteri          | GTP-binding protein. yptV2.   |
| CAA98173.1 | Z73945   | Lotus japonicus         | GTP-binding protein. RAB8B. rab8B.  |
| CAA89049.1 | Z49190   | Beta vulgaris           | GTP-binding. small G protein.   |
| CAA98179.1 | Z73951   | Lotus japonicus         | GTP-binding protein. RAB11C. rab11C.  |
| AAA34253.1 | L08130   | Volvox carteri          | GTP-binding protein. yptV4.   |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| CAA98165.1     | Z73937   | Lotus japonicus               | GTP-binding protein. RAB2A. rab2A.  |
| AAA63902.1     | U22433   | Zea mays                      | GTP binding protein. rab2.  |
| AAA90955.1     | U32185   | Glycine max                   | vesicular transport. guanine nucleotide regulatory protein. rab2. GTP-binding protein; soyrab.  |
| SEQ ID NO: 628 |          |                               |   |
| AAA87456.1     | U22147   | Hevea brasiliensis            | beta-1,3-glucanase. HGN1. hydrolytic enzyme.  |
| CAB38443.1     | AJ133470 | Hevea brasiliensis            | beta-1,3-glucanase. hgn1.   |
| AAG24921.1     | AF311749 | Hevea brasiliensis            | beta-1,3-glucanase.   |
| AAF44667.1     | AF239617 | Vitis vinifera                | hydrolysis of 1,3-beta-D-glucosidic linkages in 1,3-beta-D-glucans. beta-1,3-glucanase. fungal pathogen defense-related protein.  |
| AAA33648.1     | L02212   | Pisum sativum                 | beta-1,3-glucan hydrolysis. beta-1,3-glucanase. putative.   |
| AAB41551.1     | U27179   | Medicago sativa subsp. sativa | acidic glucanase.   |
| AAB24398.1     | S51479   | Pisum sativum                 | beta-1,3-glucanase. beta-1,3-glucanase. This sequence comes from Fig. 1B.   |
| CAA37289.1     | X53129   | Phaseolus vulgaris            | 1,3,-beta-D-glucanase.  |
| AAA34078.1     | M63634   | Nicotiana plumbaginifolia     | regulator of beta(1,3)-glucanase. beta(1,3)-glucanase regulator.  |
| AAA51643.1     | M23120   | Nicotiana plumbaginifolia     | beta-glucanase precursor.   |
| CAA30261.1     | X07280   | Nicotiana plumbaginifolia     | beta-glucanase.   |
| AAA03618.1     | M80608   | Lycopersicon esculentum       | beta-1,3-glucanase.   |
| AAC19114.1     | AF067863 | Solanum tuberosum             | 1,3-beta-glucan glucanohydrolase. glucanase.  |
| AAA18928.1     | U01901   | Solanum tuberosum             | catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). glub2. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| AAA63539.1     | M60402   | Nicotiana tabacum             | glucan beta-1,3-glucanase. glucanase GLA.   |
| AAA63540.1     | M60403   | Nicotiana tabacum             | glucan-1,3-beta-glucosidase. glucanase GLB.   |

|            |          |   |
|------------|----------|---|
| AAA88794.1 | U01900   | <i>Solanum tuberosum</i><br>catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). gluB1. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| AAA63541.1 | M59442   | <i>Nicotiana tabacum</i><br>basic beta-1,3-glucanase. glucanase.  |
| AAB82772.2 | AF001523 | <i>Musa acuminata</i><br>beta-1, 3-glucanase. similar to beta-1, 3-glucanase.   |
| AAF08679.1 | AF004838 | <i>Musa acuminata</i><br>beta-1,3-glucanase.  |
| AAA19111.1 | U01902   | <i>Solanum tuberosum</i><br>catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). gluB3. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| AAC04710.1 | AF034106 | <i>Glycine max</i><br>beta-1,3-glucanase 1. SGlu1.  |
| AAC04714.1 | AF034113 | <i>Glycine max</i><br>beta-1,3-glucanase 8. SGlu8.  |
| CAB91554.1 | AJ277900 | <i>Vitis vinifera</i><br>beta 1-3 glucanase. gl.  |
| AAA34082.1 | M20620   | <i>Nicotiana tabacum</i><br>prepro-beta-1,3-glucanase precursor.  |
| CAA03908.1 | AJ000081 | <i>Citrus sinensis</i><br>glucan hydrolase. beta-1,3-glucanase. gns1.   |
| AAB03501.1 | U41323   | <i>Glycine max</i><br>beta-1,3-glucanase. SGN1.   |
| AAA92013.1 | U49454   | <i>Prunus persica</i><br>beta-1,3-glucanase. Gns1.  |
| AAA33946.1 | M37753   | <i>Glycine max</i><br>beta-1,3-endoglucanase (EC 3.2.1.39).   |
| AAA63542.1 | M59443   | <i>Nicotiana tabacum</i><br>acidic beta-1,3-glucanase. glucanase.   |
| AAF34761.1 | AF227953 | <i>Capsicum annuum</i><br>basic beta-1,3-glucanase. BGLU.   |
| AAD33881.1 | AF141654 | <i>Nicotiana tabacum</i><br>beta-1,3-glucanase. GGL4.   |
| AAG34080.1 | AF294849 | <i>Capsicum annuum</i><br>beta-1,3-glucanase-like protein.  |
| AAF33405.1 | AF230109 | <i>Populus x canescens</i><br>beta-1,3 glucanase. BGLUC.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAD33880.1     | AF141653 | Nicotiana tabacum       | beta-1,3-glucanase. GGL1.   |
| CAA57255.1     | X81560   | Nicotiana tabacum       | (1-)-beta-glucanase. Sp41a.   |
| AAA34053.1     | M60464   | Nicotiana tabacum       | beta-1,3-glucanase.   |
| SEQ ID NO: 630 |          |                         |   |
| AAD37698.1     | AF145729 | Oryza sativa            | homeodomain leucine zipper protein. Oshox5. transcription factor.   |
| BAA05624.1     | D26575   | Daucus carota           | transcriptional regulator. DNA-binding protein. homeodomain at nt 520-699; leucine zipper at nt 700-805.  |
| AAF01765.1     | AF184278 | Glycine max             | homeodomain-leucine zipper protein 57. Hdl57. transcription factor.                                       |
| CAA64417.1     | X94947   | Lycopersicon esculentum | homeobox. VAHOX1.   |
| BAA93465.1     | AB028077 | Physcomitrella patens   | homeobox protein PpHB6. PpHB6. homeodomain-leucine zipper gene.   |
| BAB18171.1     | AB042769 | Zinnia elegans          | homeobox-leucine zipper protein. ZeHB3. full length.  |
| BAA93460.1     | AB028072 | Physcomitrella patens   | homeobox protein PpHB1. PpHB1. homeodomain-leucine zipper gene.   |
| BAA93466.1     | AB028078 | Physcomitrella patens   | homeobox protein PpHB7. PpHB7. homeodomain-leucine zipper gene.   |
| BAA93461.1     | AB028073 | Physcomitrella patens   | homeobox protein PpHB2. PpHB2. homeodomain-leucine zipper gene.   |
| BAA05625.1     | D26576   | Daucus carota           | transcriptional regulator. DNA-binding protein. homeodomain at nt 300-479; leucine zipper at nt 480-587.  |
| BAA05622.1     | D26573   | Daucus carota           | transcriptional regulator. DNA-binding protein. homeodomain at nt 585-764; leucine zipper at nt 765-851.  |
| BAA93467.1     | AB028079 | Physcomitrella patens   | homeobox protein PpHB8. PpHB8. homeodomain-leucine zipper gene.   |
| BAA93464.1     | AB028076 | Physcomitrella patens   | homeobox protein PpHB5. PpHB5. homeodomain-leucine zipper gene.   |
| AAD37697.1     | AF145728 | Oryza sativa            | homeodomain leucine zipper protein. Oshox4. transcription factor.   |
| AAF01764.2     | AF184277 | Glycine max             | homeodomain-leucine zipper protein 56. Hdl56. transcription factor.                                       |
| BAA21017.1     | D26578   | Daucus carota           | transcriptional regulator. DNA-binding protein. homeodomain at nt 761-940; leucine zipper at nt 941-1048. |



|                |          |                                   |  |
|----------------|----------|-----------------------------------|--|
| BAB18168.1     | AB042766 | <i>Zinnia elegans</i>             | homeobox-leucine zipper protein. ZeHB7. 3'RACE product.  |
| BAA93468.1     | AB028080 | <i>Physcomitrella patens</i>      | homeobox protein PpHB9. PpHB9. homeodomain-leucine zipper gene.  |
| BAA05623.1     | D26574   | <i>Daucus carota</i>              | transcriptional regulator. DNA-binding protein. homeodomain at nt 498-677; leucine zipper at nt 678-785. |
| AAD37699.1     | AF145730 | <i>Oryza sativa</i>               | homeodomain leucine zipper protein. Oshox6. transcription factor.  |
| AAD38144.1     | AF139497 | <i>Prunus armeniaca</i>           | DNA-binding protein. homeobox leucine zipper protein. HBLZP.   |
| AAA63768.2     | AF339748 | <i>Helianthus annuus</i>          | homeobox-leucine zipper protein HAHB-4. Hahb-4.  |
| BAA93463.1     | AB028075 | <i>Physcomitrella patens</i>      | homeobox protein PpHB4. PpHB4. homeodomain-leucine zipper gene.  |
| CAA64491.1     | X95193   | <i>Pimpinella brachycarpa</i>     | transcription activator. homeobox-leucine zipper protein.  |
| CAA64221.1     | X94449   | <i>Pimpinella brachycarpa</i>     | transcription activator. homeobox-leucine zipper protein. PHZ4.  |
| CAA64152.1     | X94375   | <i>Pimpinella brachycarpa</i>     | transcription activator. homeobox-leucine zipper protein.  |
| AAD37700.1     | AF145731 | <i>Oryza sativa</i>               | homeodomain leucine zipper protein. Oshox7. transcription factor.  |
| AAD37695.1     | AF145726 | <i>Oryza sativa</i>               | homeodomain leucine zipper protein. Oshox2. transcription factor.  |
| CAA06728.1     | AJ005833 | <i>Craterostigma plantagineum</i> | transcription factor. homeodomain leucine zipper protein. hb-2.  |
| CAA62608.1     | X91212   | <i>Lycopersicon esculentum</i>    | HD-ZIP protein. THOM1.   |
| CAA63222.1     | X92489   | <i>Glycine max</i>                | transcription activator. homeobox-leucine zipper protein.  |
| CAA65456.2     | X96681   | <i>Oryza sativa</i>               | transcription factor. DNA-binding protein. Oshox1. homeodomain leucine zipper gene.                      |
| AAF19980.1     | AF211193 | <i>Oryza sativa</i>               | homeodomain-leucine zipper transcription factor. Hox1. hox1.   |
| SEQ ID NO: 631 |          |                                   |  |
| CAC19183.1     | AJ291816 | <i>Cicer arietinum</i>            | expansin.  |
| AAD13633.1     | AF059489 | <i>Lycopersicon esculentum</i>    | expansin precursor. Exp5.  |
| AAG13983.1     | AF297522 | <i>Prunus avium</i>               | expansin 2. Exp2. PruavExp2.   |

|            |          |  |
|------------|----------|--|
| AAF35902.1 | AF230333 | <i>Zinnia elegans</i><br>expansin 3.   |
| AAF32409.1 | AF230276 | <i>Triphysaria versicolor</i><br>alpha-expansin 3.   |
| AAC96080.1 | AF049353 | <i>Nicotiana tabacum</i><br>involved in acid-growth response. alpha-expansin precursor. Nt-EXP4. cell wall protein.  |
| AAG13982.1 | AF297521 | <i>Prunus avium</i><br>expansin 1. Exp1. PruavExp1.  |
| AAC33529.1 | U93167   | <i>Prunus armeniaca</i><br>expansin. PA-Exp1.  |
| AAF32411.1 | AF230278 | <i>Triphysaria versicolor</i><br>alpha-expansin 1.   |
| AAF35901.1 | AF230332 | <i>Zinnia elegans</i><br>expansin 2.   |
| AAF21101.1 | AF159563 | <i>Fragaria x ananassa</i><br>expansin. Exp2. ripening regulated.  |
| BAB19676.1 | AB029083 | <i>Prunus persica</i><br>expansin. PchExp1.  |
| AAD47901.1 | AF085330 | <i>Pinus taeda</i><br>expansin.  |
| AAC33530.1 | AF038815 | <i>Prunus armeniaca</i><br>expansin. Exp2.   |
| CAC19184.1 | AJ291817 | <i>Cicer arietinum</i><br>expansin.  |
| AAB40635.1 | U64891   | <i>Pinus taeda</i><br>expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAG32921.1 | AF184233 | <i>Lycopersicon esculentum</i><br>expansin. Exp10.   |
| AAD49956.1 | AF167360 | <i>Rumex palustris</i><br>expansin. EXP1.  |
| AAB40637.1 | U64893   | <i>Pinus taeda</i><br>expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAB40634.1 | U64890   | <i>Pinus taeda</i><br>expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAB37746.1 | U30382   | Cucumis sativus         | expansin S1 precursor. Cs-EXP1. similar to pollen allergen Lol pI, Lolium perenne, Swiss-Prot Accession Number P14946; former gene name CuExS1; expansin-29 (Ex29) protein.  |
| AAB40636.1 | U64892   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAC39512.1 | AF043284 | Gossypium hirsutum      | expansin. GhEX1. contains N-terminal signal peptide.   |
| CAB43197.1 | AJ239068 | Lycopersicon esculentum | cell wall loosening enzyme. expansin2. exp2.   |
| AAC64201.1 | AF096776 | Lycopersicon esculentum | expansin. LeEXP2.  |
| AAC96081.1 | AF049354 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP5. cell wall protein.  |
| CAB46492.1 | AJ243340 | Lycopersicon esculentum | expansin9. exp9.   |
| AAF17570.1 | AF202119 | Marsilea quadrifolia    | alpha-expansin. EXP1. Mq-EXP1.   |
| AAB81662.1 | U85246   | Oryza sativa            | expansin. Os-EXP4.   |
| AAF62180.1 | AF247162 | Oryza sativa            | alpha-expansin OsEXP5. cell wall loosening factor; expressed in internodes, leaves, coleoptiles, and roots.  |
| AAB38074.1 | U30477   | Oryza sativa            | induces extension (creep) in plant cell walls. expansin Os-EXP2. Os-EXP2. former gene name RiExB.  |
| AAD13632.1 | AF059488 | Lycopersicon esculentum | expansin precursor. Exp4.  |
| AAF32410.1 | AF230277 | Triphysaria versicolor  | alpha-expansin 2.  |
| AAG01875.1 | AF291659 | Striga asiatica         | alpha-expansin 3. Exp3.  |
| CAA04385.1 | AJ000885 | Brassica napus          | Cell wall extension in plants. Expansin.   |
| CAA06271.2 | AJ004997 | Lycopersicon esculentum | expansin18. exp18.   |
| BAB32732.1 | AB049406 | Eustoma grandiflorum    | expansin. Eg Expansin.   |
| AAC63088.1 | U82123   | Lycopersicon esculentum | expansin. LeEXP1. fruit ripening regulated expansin.   |

|            |          |                                |   |
|------------|----------|--------------------------------|---|
| AAF62182.1 | AF247164 | <i>Oryza sativa</i>            | alpha-expansin OsEXP7. cell wall loosening factor; expressed in internodes and leaves.  |
| AAB37749.1 | U30460   | <i>Cucumis sativus</i>         | expansin S2 precursor. Cs-EXP2. similar to pollen allergen Lol pI, <i>Lolium perenne</i> , Swiss-Prot Accession Number P14946; former gene name CuExS2; expansin-30 (Ex30) protein. |
| AAG32920.1 | AF184232 | <i>Lycopersicon esculentum</i> | expansin. Exp8.   |
| CAC06433.1 | AJ276007 | <i>Festuca pratensis</i>       | expansin. exp2.   |
| AAF62181.1 | AF247163 | <i>Oryza sativa</i>            | alpha-expansin OsEXP6. cell wall loosening factor; expressed in internodes and leaves.  |
| AAC96079.1 | AF049352 | <i>Nicotiana tabacum</i>       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP3. cell wall protein.   |
| AAC96077.1 | AF049350 | <i>Nicotiana tabacum</i>       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP1. cell wall protein.   |
| BAA88200.1 | AP000837 | <i>Oryza sativa</i>            | EST AU078708(E60526) corresponds to a region of the predicted gene. Similar to expansin (U85246).   |
| AAF17571.1 | AF202120 | <i>Regnellidium diphyllum</i>  | alpha-expansin. EXP1. Rd-EXP1.  |
| AAC96078.1 | AF049351 | <i>Nicotiana tabacum</i>       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP2. cell wall protein.   |
| AAG01874.1 | AF291658 | <i>Striga asiatica</i>         | alpha-expansin 2. Exp2.   |

## SEQ ID NO: 634

|            |          |                                      |   |
|------------|----------|--------------------------------------|---|
| AAF63205.1 | AF245119 | <i>Mesembryanthemum crystallinum</i> | AP2-related transcription factor. CDBP. stress induced transcription factor.  |
| BAA97122.1 | AB016264 | <i>Nicotiana sylvestris</i>          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2. |
| BAA07321.1 | D38123   | <i>Nicotiana tabacum</i>             | ERF1. ethylene-responsive transcription factor.   |
| AAG43545.1 | AF211527 | <i>Nicotiana tabacum</i>             | Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.  |
| BAA87068.1 | AB035270 | <i>Matricaria chamomilla</i>         | ethylene-responsive element binding protein1 homolog. McEREBP1.   |
| BAA97124.1 | AB016266 | <i>Nicotiana sylvestris</i>          | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4. |
| AAC62619.1 | AF057373 | <i>Nicotiana tabacum</i>             | transcription factor. ethylene response element binding protein 1. EREBP1.  |
| CAB93940.1 | AJ238740 | <i>Catharanthus roseus</i>           | putative transcription factor. AP2-domain DNA-binding protein. orca2.   |

|                |          |                      |   |
|----------------|----------|----------------------|---|
| AAK31279.1     | AC079890 | Oryza sativa         | putative ethylene-responsive element binding protein. OSJNBb0089A17.16.   |
| AAG60182.1     | AC084763 | Oryza sativa         | putative ethylene-responsive element binding protein. OSJNBa0027P10.12.   |
| CAB96900.1     | AJ251250 | Catharanthus roseus  | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| CAB96899.1     | AJ251249 | Catharanthus roseus  | transcription factor. AP2-domain DNA-binding protein. orca3.  |
| CAC12822.1     | AJ299252 | Nicotiana tabacum    | AP2 domain-containing transcription factor. ap2.  |
| AAF23899.1     | AF193803 | Oryza sativa         | transcription factor EREBP1. EREBP/AP2-like transcription factor.   |
| AAF05606.1     | AF190770 | Oryza sativa         | EREBP-like protein. tsh1. TSH1; induced by ethylene.  |
| BAA97123.1     | AB016265 | Nicotiana sylvestris | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.   |
| BAA76734.1     | AB024575 | Nicotiana tabacum    | ethylene responsive element binding factor.   |
| BAB03248.1     | AB037183 | Oryza sativa         | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.   |
| CAB93939.1     | AJ238739 | Catharanthus roseus  | putative transcription factor. AP2-domain DNA-binding protein. orca1.   |
| AAD00708.1     | U91857   | Stylosanthes hamata  | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively.  |
| AAG43548.1     | AF211530 | Nicotiana tabacum    | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.  |
| AAG43549.1     | AF211531 | Nicotiana tabacum    | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.  |
| AAK31271.1     | AC079890 | Oryza sativa         | putative transcriptional factor. OSJNBb0089A17.22.  |
| AAK01088.1     | AF298230 | Hordeum vulgare      | CBF1-like protein BCBF1. BCBF1. AP2 domain protein; putative DRE binding factor.  |
| AAC49567.1     | U41466   | Zea mays             | Glossy15. Glossy15. AP2 DNA-binding domain protein; similar to the Arabidopsis floral homeotic gene APETALA2; maize AP2-domain regulator of leaf epidermal traits; homeotic regulator of leaf epidermal cell identity; allele: W64A; putative transcription factor. |
| SEQ ID NO: 635 |          |                      |   |
| CAC19789.1     | AJ251686 | Catharanthus roseus  | putative transcription factor. MYB-like DNA-binding protein. bpf-1.   |

|                |          |                              |  |
|----------------|----------|------------------------------|--|
| CAA55693.1     | X79086   | Zea mays                     | initiator-binding protein. IBP2.   |
| CAA55691.1     | X79085   | Zea mays                     | initiator binding protein. IBP1.   |
| AAF97508.1     | AF242298 | Oryza sativa                 | telomere binding protein-1. TBP1.  |
| SEQ ID NO: 636 |          |                              |  |
| CAB43505.1     | AJ239051 | Cicer arietinum              | cytochrome P450. cyp81E2.  |
| BAA22422.1     | AB001379 | Glycyrrhiza echinata         | cytochrome P450. CYP81E1.  |
| BAA74465.1     | AB022732 | Glycyrrhiza echinata         | cytochrome P450. CYP Ge-31.  |
| CAA10067.1     | AJ012581 | Cicer arietinum              | cytochrome P450. cyp81E3.  |
| CAB41490.1     | AJ238439 | Cicer arietinum              | cytochrome P450 monooxygenase. cyp81E3v2.  |
| BAA93634.1     | AB025016 | Lotus japonicus              | cytochrome P450.   |
| CAA04117.1     | AJ000478 | Helianthus tuberosus         | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-race).                                    |
| CAA04116.1     | AJ000477 | Helianthus tuberosus         | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.  |
| AAK38080.1     | AF321856 | Lolium rigidum               | putative cytochrome P450.  |
| AAK38079.1     | AF321855 | Lolium rigidum               | putative cytochrome P450.  |
| AAK38081.1     | AF321857 | Lolium rigidum               | putative cytochrome P450.  |
| AAC34853.1     | AF082028 | Hemerocallis hybrid cultivar | putative cyt P450-containing fatty acid hydroxylase. senescence-associated protein 3. SA3. mRNA accumulates in senescing petals. |
| CAA65580.1     | X96784   | Nicotiana tabacum            | cytochrome P450. hsr515.   |
| CAB56742.1     | AJ249800 | Cicer arietinum              | cytochrome P450 monooxygenase. cyp81E5.  |
| AAA32913.1     | M32885   | Persea americana             | cytochrome P-450LXXIA1 (cyp71A1).  |
| BAA12159.1     | D83968   | Glycine max                  | Cytochrome P-450 (CYP93A1).  |
| AAD56282.1     | AF155332 | Petunia x hybrida            | flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.   |

|            |          |  |
|------------|----------|--|
| CAA71515.1 | Y10491   | Glycine max<br>putative cytochrome P450.   |
| AAB94590.1 | AF022461 | Glycine max<br>CYP82C1p. CYP82C1. cytochrome P450 monooxygenase.   |
| CAA71516.1 | Y10492   | Glycine max<br>putative cytochrome P450.   |
| CAA64635.1 | X95342   | Nicotiana tabacum<br>cytochrome P450. hsr515. hypersensitivity-related gene.   |
| BAA13076.1 | D86351   | Glycine max<br>cytochrome P-450 (CYP93A2).   |
| AAG44132.1 | AF218296 | Pisum sativum<br>cytochrome P450. P450 isolog.   |
| AAG09208.1 | AF175278 | Pisum sativum<br>wound-inducible P450 hydroxylase. CYP82A1.  |
| AAC49188.2 | U29333   | Pisum sativum<br>cytochrome P450 monooxygenase. CYP82. new cytochrome P450 family.   |
| AAD38930.1 | AF135485 | Glycine max<br>cytochrome P450 monooxygenase CYP93D1. CYP93E1.   |
| CAA71876.1 | Y10982   | Glycine max<br>putative cytochrome P450.   |
| CAA70575.1 | Y09423   | Nepeta racemosa<br>cytochrome P450. CYP71A5.   |
| CAA70576.1 | Y09424   | Nepeta racemosa<br>cytochrome P450. CYP71A6.   |
| CAA71513.1 | Y10489   | Glycine max<br>putative cytochrome P450.   |
| CAA71877.1 | Y10983   | Glycine max<br>putative cytochrome P450.   |
| AAB94587.1 | AF022458 | Glycine max<br>CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.   |
| AAC39454.1 | AF014802 | Eschscholzia californica<br>(S)-N-methylcoclaurine 3'-hydroxylase. CYP82B1. cytochrome P-450-dependent monooxygenase; methyl jasmonate-inducible cytochrome P-450-dependent, homologous to wound-inducible CYP82A1 of Pisum sativum GenBank Accession Number U29333. |
| BAA92894.1 | AB006790 | Petunia x hybrida<br>cytochrome P450. IMT-2.   |
| BAA84072.1 | AB028152 | Torenia hybrida<br>flavone synthase II. cytochrome P450. TFNS5.  |
| BAA35080.1 | AB015762 | Nicotiana tabacum<br>putative cytochrome P450. CYP82E1.  |
| CAB56743.1 | AJ249801 | Cicer arietinum<br>cytochrome P450 monooxygenase. cyp81E4.   |

|                |          |  |
|----------------|----------|--|
| AAB17562.1     | U72654   | <i>Eustoma grandiflorum</i><br>flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.   |
| AAG34695.1     | AF313492 | <i>Matthiola incana</i><br>putative cytochrome P450.   |
| BAA74466.1     | AB022733 | <i>Glycyrrhiza echinata</i><br>cytochrome P450. CYP Ge-51.   |
| BAA22423.1     | AB001380 | <i>Glycyrrhiza echinata</i><br>cytochrome P450. CYP93B1.   |
| SEQ ID NO: 639 |          |  |
| AAC06319.1     | AF053084 | <i>Malus x domestica</i><br>putative cinnamyl alcohol dehydrogenase. CAD.  |
| CAA61275.1     | X88797   | <i>Eucalyptus gunnii</i><br>cinnamyl alcohol dehydrogenase. CAD1.  |
| SEQ ID NO: 640 |          |  |
| BAA92916.1     | AP001539 | <i>Oryza sativa</i><br>EST C26826(C50159) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome II BAC F13A10; putative ARF1 family auxin responsive transcription factor. (AC006418). |
| AAK21342.1     | AC024594 | <i>Oryza sativa</i><br>putative transcription factor. OSJNBa0093B11.2.   |
| AAG43286.1     | AF140228 | <i>Oryza sativa</i><br>auxin response factor 1.  |
| SEQ ID NO: 641 |          |  |
| CAC24691.1     | AJ132363 | <i>Brassica juncea</i><br>efflux carrier of polar auxin transport. pina.   |
| AAG17172.1     | AF190881 | <i>Populus tremula x Populus tremuloides</i><br>PIN1-like auxin transport protein. ppl1.   |
| AAC39514.1     | AF056027 | <i>Oryza sativa</i><br>auxin transport protein REH1. REH1. potential membrane protein.   |
| SEQ ID NO: 642 |          |  |
| AAG22044.1     | AF305783 | <i>Pisum sativum</i><br>apyrase 2. apy2. phosphatase.  |
| AAF00610.1     | AF156781 | <i>Dolichos biflorus</i><br>apyrase. apyrase-2.  |
| AAG32959.1     | AF207687 | <i>Glycine soja</i><br>apyrase GS50.   |
| AAG32960.1     | AF207688 | <i>Glycine soja</i><br>apyrase GS52.   |
| AAF00609.1     | AF156780 | <i>Lotus japonicus</i><br>apyrase. nod factor binding lectin-nucleotide phosphohydrolase. LNP.   |
| AAD31285.1     | AF139807 | <i>Dolichos biflorus</i><br>apyrase. nod factor binding lectin-nucleotide phosphohydrolase. LNP.   |



|                |          |                     |   |
|----------------|----------|---------------------|---|
| AAF00611.1     | AF156782 | Medicago sativa     | apyrase. nod factor binding lectin-nucleotide phosphohydrolase. LNP.  |
| AAK15160.1     | AF288132 | Medicago truncatula | putative apyrase. apy1. nucleotide phosphohydrolase; Mtapy1.  |
| BAB18896.1     | AB038669 | Pisum sativum       | apyrase.  |
| BAB18895.1     | AB038668 | Pisum sativum       | apyrase.  |
| BAB18894.1     | AB038555 | Pisum sativum       | apyrase H-type.   |
| BAB18893.1     | AB038554 | Pisum sativum       | apyrase S-type.   |
| BAB18900.1     | AB027614 | Pisum sativum       | apyrase.  |
| BAB40230.1     | AB027613 | Pisum sativum       | S-type apyras. ATP diphosphohydrolase (apyrase) S-type.   |
| BAB18890.1     | AB023621 | Pisum sativum       | apyrase S-type. ATP diphosphohydrolase (apyrase) S-type.  |
| BAA75506.1     | AB022319 | Pisum sativum       | apyrase. cytoskeleton associated.   |
| BAA89275.1     | AB027616 | Pisum sativum       | apyrase.  |
| BAB40231.1     | AB027615 | Pisum sativum       | S-type apyrase. ATP diphosphohydrolase (apyrase) S-type.  |
| AAB02720.1     | U58597   | Solanum tuberosum   | catalyzes the hydrolysis of phosphoanhydride bonds of nucleoside tri- and di- phosphates in the presence of divalent cations. ATP-diphosphohydrolase. RROP1. apyrase, Ecto-ATPase, E-type ATPase; NTP-diphosphohydrolase. |
| AAK15161.1     | AF288133 | Medicago truncatula | putative apyrase. apy4. nucleotide phosphohydrolase; Mtapy4.  |
| BAB18891.1     | AB030444 | Pisum sativum       | apyrase. ATP diphosphohydrolase, (EC 3.6.1.5) this sequence is reported in Acc#:AB022319.   |
| BAB18892.1     | AB030445 | Pisum sativum       | apyrase. ATP diphosphohydrolase, EC 3.6.1.5 this sequence is reported in Acc#:AB022319, Acc#:AB027613.  |
| SEQ ID NO: 645 |          |                     |   |
| AAG22044.1     | AF305783 | Pisum sativum       | apyrase 2. apy2. phosphatase.   |
| AAF00610.1     | AF156781 | Dolichos biflorus   | apyrase. apyrase-2.   |
| AAG32959.1     | AF207687 | Glycine soja        | apyrase GS50.   |

|                |          |                     |   |
|----------------|----------|---------------------|---|
| AAG32960.1     | AF207688 | Glycine soja        | apyrase GS52.   |
| AAD31285.1     | AF139807 | Dolichos biflorus   | apyrase. nod factor binding lectin-nucleotide phosphohydrolase. LNP.  |
| AAK15160.1     | AF288132 | Medicago truncatula | putative apyrase. apy1. nucleotide phosphohydrolase; Mtapy1.  |
| AAF00609.1     | AF156780 | Lotus japonicus     | apyrase. nod factor binding lectin-nucleotide phosphohydrolase. LNP.  |
| BAB18896.1     | AB038669 | Pisum sativum       | apyrase.  |
| BAB18895.1     | AB038668 | Pisum sativum       | apyrase.  |
| BAB18894.1     | AB038555 | Pisum sativum       | apyrase H-type.   |
| BAB18893.1     | AB038554 | Pisum sativum       | apyrase S-type.   |
| BAB18900.1     | AB027614 | Pisum sativum       | apyrase.  |
| BAB40230.1     | AB027613 | Pisum sativum       | S-type apyras. ATP diphosphohydrolase (apyrase) S-type.   |
| BAB18890.1     | AB023621 | Pisum sativum       | apyrase S-type. ATP diphosphohydrolase (apyrase) S-type.  |
| BAA75506.1     | AB022319 | Pisum sativum       | apyrase. cytoskeleton associated.   |
| AAF00611.1     | AF156782 | Medicago sativa     | apyrase. nod factor binding lectin-nucleotide phosphohydrolase. LNP.  |
| BAA89275.1     | AB027616 | Pisum sativum       | apyrase.  |
| BAB40231.1     | AB027615 | Pisum sativum       | S-type apyrase. ATP diphosphohydrolase (apyrase) S-type.  |
| AAB02720.1     | U58597   | Solanum tuberosum   | catalyzes the hydrolysis of phosphoanhydride bonds of nucleoside tri- and di- phosphates in the presence of divalent cations. ATP-diphosphohydrolase. RROP1. apyrase, Ecto-ATPase, E-type ATPase; NTP-diphosphohydrolase. |
| AAK15161.1     | AF288133 | Medicago truncatula | putative apyrase. apy4. nucleotide phosphohydrolase; Mtapy4.  |
| BAB18891.1     | AB030444 | Pisum sativum       | apyrase. ATP diphosphohydrolase, (EC 3.6.1.5) this sequence is reported in Acc#:AB022319.   |
| BAB18892.1     | AB030445 | Pisum sativum       | apyrase. ATP diphosphohydrolase, EC 3.6.1.5 this sequence is reported in Acc#:AB022319, Acc#:AB027613.  |
| SEQ ID NO: 646 |          |                     |   |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| AAB80947.1     | AF022915 | Triticum aestivum         | ornithine/acetylornithine aminotransferase.   |
| CAA69936.1     | Y08680   | Alnus glutinosa           | acetylornithine aminotransferase. ag118.  |
| AAK11219.1     | AF324485 | Oryza sativa              | aminotransferase-like protein.  |
| AAG09278.1     | AF177590 | Vitis vinifera            | ornithine aminotransferase.   |
| AAC78480.1     | AF085149 | Capsicum chinense         | putative aminotransferase. pyridoxal phosphate dependent.   |
| AAA02916.1     | L08400   | Vigna aconitifolia        | production of pyrroline-5-carboxylate by deamination of ornithine. ornithine aminotransferase.  |
| AAB59330.1     | M31545   | Hordeum vulgare           | glutamate 1-semialdehyde aminotransferase. GSA. precursor.  |
| AAA18861.1     | U03632   | Chlamydomonas reinhardtii | glutamate-1-semialdehyde aminotransferase. gsa.   |
| AAA33968.1     | L12453   | Glycine max               | catalyzes 5-aminolevulinic acid formation from GSA. glutamate 1-semialdehyde aminotransferase. Gsa. putative.                             |
| AAC48996.1     | U20260   | Glycine max               | converts GSA to 5-aminolevulinic acid. glutamate 1-semialdehyde aminotransferase. Gsa1.   |
| SEQ ID NO: 650 |          |                           |   |
| AAF66982.1     | AF247646 | Zea mays                  | transposase. similar to Mutator family transposases.  |
| SEQ ID NO: 652 |          |                           |   |
| AAB41812.1     | L36158   | Medicago sativa           | peroxidase. pxdD. amino acid feature: conserved domains, aa 120 .. 126, 188 .. 195; amino acid feature: heme-binding domain, aa 63 .. 68. |
| CAA71495.1     | Y10469   | Spinacia oleracea         | peroxidase. prxr8.  |
| CAA09881.1     | AJ011939 | Trifolium repens          | peroxidase. prx2.   |
| CAA62228.1     | X90695   | Medicago sativa           | peroxidase2. prx2.  |
| AAA98491.1     | L36981   | Petroselinum crispum      | anionic peroxidase.   |
| BAB39281.1     | AP002971 | Oryza sativa              | putative peroxidase. P0537A05.10. contains ESTs D24657(R2329),AU082066(R2329).  |
| AAB02926.1     | U59284   | Linum usitatissimum       | peroxidase. FLXPER3.  |
| BAA77387.1     | AB024437 | Scutellaria baicalensis   | peroxidase 1.   |

|            |          |   |
|------------|----------|---|
| CAA66037.1 | X97351   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.  |
| CAA71488.1 | Y10462   | Spinacia oleracea<br>peroxidase. prxr1.   |
| BAA01950.1 | D11337   | Vigna angularis<br>peroxidase.  |
| BAA14143.1 | D90115   | Armoracia rusticana<br>peroxidase isozyme.  |
| CAA71490.1 | Y10464   | Spinacia oleracea<br>peroxidase. prxr3.   |
| CAB94692.1 | AJ242742 | Ipomoea batatas<br>Removal of H <sub>2</sub> O <sub>2</sub> , oxidation of toxic reductants, defence response toward wounding.<br>peroxidase. pod.                |
| BAA92497.1 | AP001383 | Oryza sativa<br>ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene.<br>Similar to peroxidase ATP18a. (X98804).                     |
| AAC36707.1 | AF078691 | Manihot esculenta<br>peroxidase.  |
| BAA92422.1 | AP001366 | Oryza sativa<br>ESTs AU081576(R0541),AU032412(R4029) correspond to a region of the predicted gene.<br>Similar to A.thaliana mRNA for peroxidase ATP18a. (X98804). |
| CAA62226.1 | X90693   | Medicago sativa<br>peroxidase1B. prx1B.   |
| AAF63024.1 | AF244921 | Spinacia oleracea<br>hydrogen peroxide catabolism. peroxidase prx12 precursor. type III peroxidase.   |
| AAA32676.1 | M37637   | Arachis hypogaea<br>cationic peroxidase. PNC2.  |
| BAA11853.1 | D83225   | Populus nigra<br>peroxidase.  |
| CAC21393.1 | AJ401276 | Zea mays<br>peroxidase. pox3.   |
| AAA65636.1 | L13653   | Lycopersicon esculentum<br>peroxidase. TPX2.  |
| CAB67121.1 | Y19023   | Lycopersicon esculentum<br>peroxidase. cevi-1.  |
| CAA62227.1 | X90694   | Medicago sativa<br>peroxidase1C. prx1C.   |
| CAA50597.1 | X71593   | Lycopersicon esculentum<br>peroxidase. CEVI-1.  |
| AAD37376.1 | AF145350 | Glycine max<br>peroxidase. Prx4.  |
| AAB67737.1 | L77080   | Stylosanthes humilis<br>cationic peroxidase.  |

|                |          |  |
|----------------|----------|--|
| CAA71489.1     | Y10463   | Spinacia oleracea<br>peroxidase. prxr2.  |
| CAA71496.1     | Y10470   | Spinacia oleracea<br>peroxidase. prxr9.  |
| CAA71494.1     | Y10468   | Spinacia oleracea<br>peroxidase. prxr7.  |
| BAA06334.1     | D30652   | Populus kitakamiensis<br>peroxidase.   |
| CAA66034.1     | X97348   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA94962.1     | AB042103 | Asparagus officinalis<br>peroxidase. AspPOX1.  |
| CAA80502.1     | Z22920   | Spirodela polyrrhiza<br>peroxidase.  |
| CAA66035.1     | X97349   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA11852.1     | D83224   | Populus nigra<br>peroxidase.   |
| CAA66036.1     | X97350   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA07241.1     | D38051   | Populus kitakamiensis<br>peroxidase. prxA4a.   |
| AAD11481.1     | U51191   | Glycine max<br>peroxidase precursor. sEPa1.  |
| AAC98519.1     | AF007211 | Glycine max<br>peroxidase precursor. GMIPER1. pathogen-induced.  |
| AAA32973.1     | M73234   | Hordeum vulgare<br>peroxidase BP 1. Prx5.  |
| AAB47602.1     | L07554   | Linum usitatissimum<br>peroxidase. FLXPER1.  |
| AAB41810.1     | L36156   | Medicago sativa<br>peroxidase. pxdA. amino acid feature: conserved motifs, aa 181 .. 188; amino acid feature:<br>heme-binding domain, aa 60 .. 65. |
| AAF63027.1     | AF244924 | Spinacia oleracea<br>hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.  |
| AAA34108.1     | J02979   | Nicotiana tabacum<br>lignin-forming peroxidase precursor (EC 1.11.1.7).  |
| AAD43561.1     | AF155124 | Gossypium hirsutum<br>bacterial-induced peroxidase precursor. Perx_Goshiko.  |
| SEQ ID NO: 653 |          |  |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| AAB97617.1     | U83687   | <i>Apium graveolens</i>   | NADPH-dependent mannose 6-phosphate reductase. m6pr. aldo-keto reductase; similar to aldose 6-phosphate reductase also known as NADP-sorbitol-6-phosphate dehydrogenase encoded by GenBank Accession Number D11080. |
| BAA01853.1     | D11080   | <i>Malus x domestica</i>  | NADP-dependent D-sorbitol-6-phosphate dehydrogenase. S6PDH.   |
| AAC97607.1     | AF057134 | <i>Malus x domestica</i>  | synthesizes sorbitol, a major photosynthetic product in many members of the Rosaceae family. NADP-dependent sorbitol 6-phosphate dehydrogenase. S6PDH.  |
| SEQ ID NO: 654 |          |                           |   |
| BAA82556.1     | AB030083 | <i>Populus nigra</i>      | lectin-like protein kinase. PnLPK.  |
| AAB61708.1     | U93048   | <i>Daucus carota</i>      | somatic embryogenesis receptor-like kinase. SERK.   |
| BAB19337.1     | AP003044 | <i>Oryza sativa</i>       | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).  |
| AAK21965.1     | AY028699 | <i>Brassica napus</i>     | receptor protein kinase PERK1.  |
| BAB39873.1     | AP002882 | <i>Oryza sativa</i>       | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| AAB93834.1     | U82481   | <i>Zea mays</i>           | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.  |
| BAB18292.1     | AP002860 | <i>Oryza sativa</i>       | putative receptor-like protein kinase. P0409B08.19.   |
| AAK00425.1     | AC069324 | <i>Oryza sativa</i>       | Putative protein kinase. OSJNBa0071K19.11.  |
| AAD21872.1     | AF078082 | <i>Phaseolus vulgaris</i> | receptor-like protein kinase homolog RK20-1.  |
| AAC23542.1     | U20948   | <i>Ipomoea trifida</i>    | receptor protein kinase. IRK1.  |
| CAA73134.1     | Y12531   | <i>Brassica oleracea</i>  | serine/threonine kinase. BRLK.  |
| CAB51480.1     | Y14600   | <i>Sorghum bicolor</i>    | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| AAG59657.1     | AC084319 | <i>Oryza sativa</i>       | putative protein kinase. OSJNBa0004B24.20.  |
| BAB16871.1     | AP002537 | <i>Oryza sativa</i>       | putative protein kinase APK1A <i>Arabidopsis thaliana</i> . P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).   |

|            |          |                   |   |
|------------|----------|-------------------|---|
| CAB51834.1 | 00069    | Oryza sativa      | 11332.5. contains eukaryotic protein kinase domain PF.  |
| BAA94509.1 | AB041503 | Populus nigra     | protein kinase 1. PnPK1.  |
| AAG16628.1 | AY007545 | Brassica napus    | protein serine/threonine kinase BNK1.   |
| BAA87853.1 | AP000816 | Oryza sativa      | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB03429.1 | AP002817 | Oryza sativa      | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).             |
| BAB07999.1 | AP002525 | Oryza sativa      | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB21240.1 | AP002953 | Oryza sativa      | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| BAA92954.1 | AP001551 | Oryza sativa      | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).   |
| BAA94510.1 | AB041504 | Populus nigra     | protein kinase 2. PnPK2.  |
| AAG03090.1 | AC073405 | Oryza sativa      | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| AAF91322.1 | AF244888 | Glycine max       | receptor-like protein kinase 1. RLK1. GmRLK1.   |
| BAA94517.1 | AP001800 | Oryza sativa      | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| BAB07905.1 | AP002835 | Oryza sativa      | putative S-receptor kinase. P0417G05.13.  |
| CAB51836.1 | AJ243961 | Oryza sativa      | Putitive Ser/Thr protein kinase. 11332.7.   |
| BAA78764.1 | AB023482 | Oryza sativa      | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAA33915.1 | L27821   | Oryza sativa      | receptor type serine/threonine kinase. protein kinase.  |
| AAF91323.1 | AF244889 | Glycine max       | receptor-like protein kinase 2. RLK2. GmRLK2.   |
| CAA67145.1 | X98520   | Brassica oleracea | receptor-like kinase. SFR2.   |
| CAA73133.1 | Y12530   | Brassica oleracea | serine /threonine kinase. ARLK.   |
| AAF91324.1 | AF244890 | Glycine max       | receptor-like protein kinase 3. RLK3. GmRLK3.   |

|                |          |                      |  |
|----------------|----------|----------------------|--|
| CAA74661.1     | Y14285   | Brassica oleracea    | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554. |
| BAA92953.1     | AP001551 | Oryza sativa         | Similar to Arabidopsis thaliana chromosome 4 BAC clone F10M6 ; S-receptor kinase -like protein. (AL021811).    |
| BAB40094.1     | AP003210 | Oryza sativa         | putative receptor protein kinase. OSJNBa0010K01.7.   |
| AAF43496.1     | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.    |
| AAK11674.1     | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.   |
| BAB07906.1     | AP002835 | Oryza sativa         | putative S-receptor kinase. P0417G05.14.   |
| BAA94516.1     | AP001800 | Oryza sativa         | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| CAB41878.1     | Y18259   | Brassica oleracea    | SRK5 protein. SRK5. receptor-like kinase.  |
| BAA23676.1     | AB000970 | Brassica rapa        | receptor kinase 1. BcRK1.  |
| CAB41879.1     | Y18260   | Brassica oleracea    | SRK15 protein. SRK15. receptor-like kinase.  |
| SEQ ID NO: 655 |          |                      |  |
| AAD21872.1     | AF078082 | Phaseolus vulgaris   | receptor-like protein kinase homolog RK20-1.   |
| CAA73134.1     | Y12531   | Brassica oleracea    | serine/threonine kinase. BRLK.   |
| CAA67145.1     | X98520   | Brassica oleracea    | receptor-like kinase. SFR2.  |
| CAA73133.1     | Y12530   | Brassica oleracea    | serine /threonine kinase. ARLK.  |
| AAB93834.1     | U82481   | Zea mays             | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.           |
| BAA23676.1     | AB000970 | Brassica rapa        | receptor kinase 1. BcRK1.  |
| AAC23542.1     | U20948   | Ipomoea trifida      | receptor protein kinase. IRK1.   |
| CAB41879.1     | Y18260   | Brassica oleracea    | SRK15 protein. SRK15. receptor-like kinase.  |



|            |          |                             |  |
|------------|----------|-----------------------------|--|
| CAA74662.1 | Y14286   | Brassica oleracea           | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| CAB89179.1 | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| AAA33008.1 | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| AAA62232.1 | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| CAA74661.1 | Y14285   | Brassica oleracea           | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.   |
| AAA33000.1 | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| CAA79355.1 | Z18921   | Brassica oleracea           | S-receptor kinase-like protein.  |
| CAB41878.1 | Y18259   | Brassica oleracea           | SRK5 protein. SRK5. receptor-like kinase.  |
| BAA07576.1 | D38563   | Brassica rapa               | receptor protein kinase SRK8.  |
| BAA06285.1 | D30049   | Brassica rapa               | S-receptor kinase SRK9.  |
| BAA21132.1 | D88193   | Brassica rapa               | S-receptor kinase. SRK9 (B.c).   |
| BAA07577.2 | D38564   | Brassica rapa               | receptor protein kinase SRK12.   |
| BAA92836.1 | AB032473 | Brassica oleracea           | S18 S-locus receptor kinase. SRK18.  |
| BAB21001.1 | AB054061 | Brassica rapa               | S locus receptor kinase. SRK22.  |
| BAA92837.1 | AB032474 | Brassica oleracea           | S60 S-locus receptor kinase. SRK60.  |
| BAB18292.1 | AP002860 | Oryza sativa                | putative receptor-like protein kinase. P0409B08.19.  |
| AAD52097.1 | AF088885 | Nicotiana tabacum           | receptor-like kinase CHRK1. Chrk1.   |
| BAB39873.1 | AP002882 | Oryza sativa                | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).                            |
| AAK21965.1 | AY028699 | Brassica napus              | receptor protein kinase PERK1.   |
| AAA33915.1 | L27821   | Oryza sativa                | receptor type serine/threonine kinase. protein kinase.   |

|                |          |                                   |  |
|----------------|----------|-----------------------------------|--|
| AAK00425.1     | AC069324 | Oryza sativa                      | Putative protein kinase. OSJNBa0071K19.11.   |
| BAA94529.2     | AP001800 | Oryza sativa                      | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB16871.1     | AP002537 | Oryza sativa                      | putative protein kinase APK1A Arabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192), D25110(R3192).                         |
| AAF34428.1     | AF172282 | Oryza sativa                      | receptor-like protein kinase. DUPR11.18.   |
| AAG03090.1     | AC073405 | Oryza sativa                      | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).   |
| BAB21240.1     | AP002953 | Oryza sativa                      | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461), C22360(C11461).  |
| BAA87853.1     | AP000816 | Oryza sativa                      | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308). |
| BAA94516.1     | AP001800 | Oryza sativa                      | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).  |
| BAB07905.1     | AP002835 | Oryza sativa                      | putative S-receptor kinase. P0417G05.13.   |
| BAB07906.1     | AP002835 | Oryza sativa                      | putative S-receptor kinase. P0417G05.14.   |
| SEQ ID NO: 657 |          |                                   |  |
| AAF43869.1     | AF166114 | Chloroplast Mesostigma viride     | probable transport protein. cysA.  |
| BAB17113.1     | AP002866 | Oryza sativa                      | putative white protein; ATP-binding cassette transporter. P0410E01.34.   |
| BAA57907.1     | AB001684 | Chlorella vulgaris                | sulfate transport system permease protein. cysA.   |
| AAD54843.1     | AF137379 | Chloroplast Nephroselmis olivacea | probable transport protein. cysA.  |
| BAA90508.1     | AP001111 | Oryza sativa                      | similar to ABC transporter of Arabidopsis thaliana (AC004697).   |
| BAA90507.1     | AP001111 | Oryza sativa                      | similar to ABC transporter of Arabidopsis thaliana (AC004697).   |
| BAA83352.1     | AP000391 | Oryza sativa                      | ESTs AU067992(C11433), AU077424(C11433) correspond to a region of the predicted gene.; Similar to ABC transporter-7 (U43892).        |
| AAG49003.1     | AY013246 | Hordeum vulgare                   | putative ABC transporter. 635P2.4b; GC splice donor confirmed by cDNA alignment and comparative sequence.                            |
| AAD10836.1     | U52079   | Solanum tuberosum                 | P-glycoprotein. pmdr1. binds ATP; ATPase; transporter; transmembrane protein.  |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| BAA96612.1     | AP002482 | Oryza sativa              | Similar to Arabidopsis thaliana chromosome 2, BAC F14M4 ; putative ABC transporter (AC004411).          |
| BAB40032.1     | AP003046 | Oryza sativa              | putative ABC transporter. P0445D12.3.   |
| AAG45492.1     | AY013245 | Oryza sativa              | 36I5.4. putative ABC transporter; GC splice donor confirmed by cDNA alignment and comparative sequence. |
| AAG49002.1     | AY013246 | Hordeum vulgare           | putative ABC transporter. 635P2.4a; GC splice donor confirmed by cDNA and comparative sequencing.       |
| BAB21275.1     | AP002844 | Oryza sativa              | putative ABC transporter protein. P0410E03.6.   |
| BAB21273.1     | AP002844 | Oryza sativa              | putative ABC transporter protein. P0410E03.4.   |
| CAA94437.1     | Z70524   | Spirodela polyrrhiza      | multidrug resistance protein. PDR5-like ABC transporter.  |
| BAB21276.1     | AP002844 | Oryza sativa              | putative ABC transporter protein. P0410E03.7. contains EST D22472(C1173).                               |
| BAB21279.1     | AP002844 | Oryza sativa              | putative ABC transporter protein. P0410E03.10. contains ESTs AU065360(R3463),AU101680(R3463).           |
| SEQ ID NO: 658 |          |                           |   |
| AAD10386.1     | U72255   | Oryza sativa              | beta-1,3-glucanase precursor. Gns9.   |
| BAA89481.1     | AB029462 | Salix gilgiana            | beta-1,3-glucanase. SgGN1.  |
| CAB85903.1     | AJ251646 | Pisum sativum             | hydrolysis of beta-1,3 conjugated glucans. beta-1,3 glucanase. gns2.                                    |
| CAA49513.1     | X69887   | Brassica napus            | beta-1,3-glucanase homologue.   |
| AAA90953.1     | U30323   | Triticum aestivum         | beta 1,3-glucanase. Glc1.   |
| BAB19363.1     | AP002542 | Oryza sativa              | putative beta-1,3-glucanase. P0679C08.2.  |
| CAA82271.1     | Z28697   | Nicotiana tabacum         | beta-1,3-glucanase.   |
| CAA30261.1     | X07280   | Nicotiana plumbaginifolia | beta-glucanase.   |
| AAA51643.1     | M23120   | Nicotiana plumbaginifolia | beta-glucanase precursor.   |
| AAA34078.1     | M63634   | Nicotiana plumbaginifolia | regulator of beta(1,3)-glucanase. beta(1,3)-glucanase regulator.  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAB82772.2 | AF001523 | Musa acuminata          | beta-1, 3-glucanase. similar to beta-1, 3-glucanase.  |
| AAF08679.1 | AF004838 | Musa acuminata          | beta-1,3-glucanase.   |
| CAB71021.1 | AJ271598 | Hieracium piloselloides | putative role in callose degradation. putative beta-1,3-glucanase. gluc.  |
| AAD10383.1 | U72252   | Oryza sativa            | beta-1,3-glucanase precursor. Gns6.   |
| AAD28732.1 | AF112965 | Triticum aestivum       | beta-1,3-glucanase precursor. Glb3.   |
| BAB40807.1 | AB052291 | Pyrus pyrifolia         | catalyzing the hydrolysis of 1,3-beta-glucosyl linkages. endo-1,3-beta-glucanase-like protein. bgn-1. Amino acid alignment of the protein(BGN-1) encoded by pear pollen bgn-1 with barley endo-1,3-beta-glucanase(GII. accession number: pdb/1GHS-B/2.3/2/306/N/)(40% identity) and their hydrophobic cluster analysis(HCA)(a overall HCA homology score of 87.1%) showed that it was most likely that the bgn-1 encoded a endo-1,3-beta-glucanase. A higher identity(59.3%) was found between BGN-1 and a putative pea endo-1,3-beta-glucanase (accession number: dad/AJ251646-1). |
| AAA63539.1 | M60402   | Nicotiana tabacum       | glucan beta-1,3-glucanase. glucanase GLA.   |
| AAA63541.1 | M59442   | Nicotiana tabacum       | basic beta-1,3-glucanase. glucanase.  |
| AAA63540.1 | M60403   | Nicotiana tabacum       | glucan-1,3-beta-glucosidase. glucanase GLB.   |
| AAA32939.1 | M62907   | Hordeum vulgare         | hydrolysis of beta-(1-3)-glucan. (1-3)-beta-glucanase. cBGL32.  |
| AAC14399.1 | AF030771 | Hordeum vulgare         | beta-1,3-glucanase 2. BGL32.  |
| AAA87456.1 | U22147   | Hevea brasiliensis      | beta-1,3-glucanase. HGN1. hydrolytic enzyme.  |
| BAA77784.1 | AB027429 | Oryza sativa            | beta-1,3-glucanase.   |
| BAA77785.1 | AB027430 | Oryza sativa            | beta-1,3-glucanase.   |
| CAB91554.1 | AJ277900 | Vitis vinifera          | beta 1-3 glucanase. gl.   |
| AAD10381.1 | U72250   | Oryza sativa            | beta-1,3-glucanase precursor. Gns4.   |
| AAA33946.1 | M37753   | Glycine max             | beta-1,3-endoglucanase (EC 3.2.1.39).   |
| AAD33881.1 | AF141654 | Nicotiana tabacum       | beta-1,3-glucanase. GGL4.   |

|                |          |                    |   |
|----------------|----------|--------------------|---|
| AAB86541.1     | AF030166 | Oryza sativa       | glucanase. glu1.  |
| AAD10384.1     | U72253   | Oryza sativa       | beta-1,3-glucanase precursor. Gns7.   |
| CAB38443.1     | AJ133470 | Hevea brasiliensis | beta-1,3-glucanase. hgn1.   |
| AAB03501.1     | U41323   | Glycine max        | beta-1,3-glucanase. SGN1.   |
| AAA18928.1     | U01901   | Solanum tuberosum  | catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). glub2. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| AAA88794.1     | U01900   | Solanum tuberosum  | catalyzes the hydrolysis of 1,3-beta-D-glucoside linkages in 1,3,-beta-D-glucans (such as laminarin). 1,3-beta-D-glucan glucanohydrolase; endo-1,3-beta-D-glucanase; 1,3-beta-glucanase (basic, class I). gluB1. plant defense gene; induced expression in response to infection, elicitor, ethylene, wounding. |
| AAC19114.1     | AF067863 | Solanum tuberosum  | 1,3-beta-glucan glucanohydrolase. glucanase.  |
| AAG24921.1     | AF311749 | Hevea brasiliensis | beta-1,3-glucanase.   |
| CAA03908.1     | AJ000081 | Citrus sinensis    | glucan hydrolase. beta-1,3-glucanase. gns1.   |
| CAA37289.1     | X53129   | Phaseolus vulgaris | 1,3,-beta-D-glucanase.  |
| CAA57255.1     | X81560   | Nicotiana tabacum  | (1-)-beta-glucanase. Sp41a.   |
| SEQ ID NO: 659 |          |                    |   |
| AAB65776.1     | U97521   | Vitis vinifera     | class IV endochitinase. VvChi4A.  |
| AAB65777.1     | U97522   | Vitis vinifera     | class IV endochitinase. VvChi4B.  |
| BAA03751.1     | D16223   | Oryza sativa       | endochitinase. Cht-3.   |
| CAA30142.1     | X07130   | Solanum tuberosum  | endochitinase.  |
| BAA03749.1     | D16221   | Oryza sativa       | endochitinase. Cht-1.   |
| SEQ ID NO: 660 |          |                    |   |
| AAF07221.1     | AF072519 | Nicotiana tabacum  | centrin. CEN1. caltractin; EF-hand domain calcium-binding protein.  |

|                |          |                           |  |
|----------------|----------|---------------------------|--|
| AAF07222.1     | AF072520 | Nicotiana tabacum         | centrin. CEN2. caltractin; EF-hand domain calcium-binding protein.   |
| CAA49153.1     | X69220   | Scherffelia dubia         | caltractin.  |
| AAC04626.1     | U92973   | Marsilea vestita          | calcium-binding protein. centrin. MvCen1. caltractin.  |
| AAB67855.1     | U53812   | Dunaliella salina         | caltractin-like protein.   |
| CAA41039.1     | X57973   | Chlamydomonas reinhardtii | caltractin.  |
| CAA31163.1     | X12634   | Chlamydomonas reinhardtii | caltractin (AA 1 - 169).   |
| SEQ ID NO: 664 |          |                           |  |
| BAB16432.1     | AB041520 | Nicotiana tabacum         | WRKY transcription factor Nt-SubD48. Nt-SubD48.  |
| AAC49528.1     | U56834   | Petroselinum crispum      | DNA-binding. WRKY3. WRKY-type DNA-binding protein.   |
| AAD27591.1     | AF121354 | Petroselinum crispum      | binds sequence specifically to W Boxes (TTGACC). transcription factor. WRKY3. sequence specific DNA-binding protein. |
| AAD32676.1     | AF140553 | Avena sativa              | DNA-binding protein WRKY3. wrky3. putative transcription factor.   |
| BAA77358.1     | AB020023 | Nicotiana tabacum         | WRKY domain Zn-finger type DNA-binding protein. DNA-binding protein NtWRKY3.   |
| AAG46150.1     | AC018727 | Oryza sativa              | putative DNA-binding protein. OSJNBa0056G17.18.  |
| BAB40073.1     | AP003074 | Oryza sativa              | putative WRKY DNA binding protein. OSJNBa0004G10.20. contains EST C26525(C12525).                                    |
| BAB18313.1     | AP002865 | Oryza sativa              | putative WRKY DNA binding protein. P0034C11.1. contains EST C26525(C12525).  |
| AAD38283.1     | AC007789 | Oryza sativa              | putative WRKY DNA binding protein. OSJNBa0049B20.9.  |
| BAA77383.1     | AB020590 | Nicotiana tabacum         | transcription factor NtWRKY2.  |
| BAA82107.1     | AB022693 | Nicotiana tabacum         | transcription factor. NtWRKY1.   |
| CAA88326.1     | Z48429   | Avena fatua               | binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein.                                   |
| AAD32677.1     | AF140554 | Avena sativa              | DNA-binding protein WRKY1. wrky1. putative transcription factor.   |
| AAD55974.1     | AF121353 | Petroselinum crispum      | zinc-finger type transcription factor WRKY1. WRKY1.  |

|                |          |  |
|----------------|----------|--|
| AAC49529.1     | U58540   | <i>Petroselinum crispum</i><br>WRKY2. Contains two WRKY domains; WRKY-type DNA-binding protein.  |
| AAC49527.1     | U48831   | <i>Petroselinum crispum</i><br>WRKY1. contains two WRKY domains; WRKY-type DNA-binding protein; sequence-specific DNA-binding protein. |
| AAC31956.1     | AF080595 | <i>Pimpinella brachycarpa</i><br>zinc finger protein. ZFP1. WRKY1.   |
| BAA86031.1     | AB026890 | <i>Nicotiana tabacum</i><br>transcription factor NtWRKY4.  |
| AAD16139.1     | AF096299 | <i>Nicotiana tabacum</i><br>DNA-binding protein 2. WRKY2. transcription factor.  |
| AAF23898.1     | AF193802 | <i>Oryza sativa</i><br>zinc finger transcription factor WRKY1.   |
| AAD16138.1     | AF096298 | <i>Nicotiana tabacum</i><br>DNA-binding protein 1. WRKY1. transcription factor.  |
| AAC37515.1     | L44134   | <i>Cucumis sativus</i><br>SPF1-like DNA-binding protein.   |
| AAG35658.1     | AF204925 | <i>Petroselinum crispum</i><br>transcription factor WRKY4. WRKY4. binds to W box (TTGACC) elements.                                    |
| CAA88331.1     | Z48431   | <i>Avena fatua</i><br>binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein.                               |
| BAB19075.1     | AP002744 | <i>Oryza sativa</i><br>putative DNA-binding protein homolog. P0006C01.17.  |
| BAB19096.1     | AP002839 | <i>Oryza sativa</i><br>putative DNA-binding protein homolog. P0688A04.2.   |
| AAK16170.1     | AC079887 | <i>Oryza sativa</i><br>putative DNA binding protein. OSJNBa0040E01.4.  |
| AAK16171.1     | AC079887 | <i>Oryza sativa</i><br>putative DNA-binding protein. OSJNBa0040E01.10.   |
| AAG35659.1     | AF204926 | <i>Petroselinum crispum</i><br>transcription factor WRKY5. WRKY5. binds to W box (TTGACC) elements.                                    |
| CAB97004.1     | AJ278507 | <i>Solanum tuberosum</i><br>putative transcription factor. WRKY DNA binding protein. WRKY1.  |
| AAF61864.1     | AF193771 | <i>Nicotiana tabacum</i><br>DNA-binding protein 4. WRKY4. transcription factor.  |
| BAA87069.1     | AB035271 | <i>Matricaria chamomilla</i><br>elicitor-induced DNA-binding protein homolog. McWRKY1.   |
| AAF61863.1     | AF193770 | <i>Nicotiana tabacum</i><br>DNA-binding protein 3. WRKY3. transcription factor.  |
| SEQ ID NO: 665 |          |  |
| AAD02558.1     | AF049933 | <i>Petunia x hybrida</i><br>PGPS/NH17. PGPS/NH17. protein sec61 gamma subunit homolog; protein transport to ER.                        |
| SEQ ID NO: 667 |          |  |

---

AAD51623.1 AF169020 Glycine max  
seed maturation protein PM35. PM35. similar to Phaseolus vulgaris putative osmoprotector  
PvLEA-18.

---

AAC49859.1 U72764 Phaseolus vulgaris  
putative osmoprotector. PvLEA-18. Pvlea-18. atypical late embryogenesis abundant protein.

---

AAF81194.1 AF240774 Phaseolus vulgaris  
LEA-18.

---

SEQ ID NO: 672

---

CAB40743.1 AJ011885 Solanum tuberosum  
starch branching enzyme II. sbe II.

---

CAB40746.1 AJ011888 Solanum tuberosum  
starch branching enzyme II. SBE II.

---

CAB40748.1 AJ011890 Solanum tuberosum  
starch branching enzyme II. SBE II.

---

AAD30186.1 AF076679 Triticum aestivum  
starch branching enzyme-I. SBE-I.

---

AAD30187.1 AF076680 Aegilops tauschii  
starch branching enzyme-I. SBE-I.

---

BAA82348.1 AB029548 Phaseolus vulgaris  
branching enzyme 1. kbe1.

---

CAA56319.1 X80009 Pisum sativum  
starch branching enzyme I. SBEI.

---

CAB40747.1 AJ011889 Solanum tuberosum  
starch branching enzyme II. SBE II.

---

CAA03846.1 AJ000004 Solanum tuberosum  
branches 1,4-alpha glucans. starch branching enzyme II, SBE-II. Sbe-II.

---

BAA03738.1 D16201 Oryza sativa  
branching enzyme-3 precursor.

---

AAG27623.1 AF286319 Triticum aestivum  
starch branching enzyme 2. Sbe2. glucosyltransferase.

---

CAA72154.1 Y11282 Triticum aestivum  
1,4-alpha-glucan branching enzyme II. sbe2.

---

AAK26821.1 AF338431 Aegilops tauschii  
starch branching enzyme IIa. SBEIIa.

---

AAK26822.1 AF338432 Triticum aestivum  
starch branching enzyme IIa variant. SBEIIa variant.

---

AAC33764.1 AF072725 Zea mays  
starch branching enzyme IIb. ae. SBEIIb.

---

AAA18571.1 L08065 Zea mays  
starch branching enzyme II.

---

BAA82828.1 AB023498 Oryza sativa  
starch branching enzyme rbe4. RBE4.

---



|            |          |                              |   |
|------------|----------|------------------------------|---|
| AAC69753.1 | AF064560 | Hordeum vulgare              | starch branching enzyme IIa. sbellA.  |
| CAA56320.1 | X80010   | Pisum sativum                | starch branching enzyme II. SBEII.  |
| AAC69754.1 | AF064561 | Hordeum vulgare              | starch branching enzyme IIb. sbellB.  |
| AAC36471.1 | AF072724 | Zea mays                     | starch branching enzyme I. sbel. confirmed by partial peptide sequencing.   |
| AAA82735.1 | U17897   | Zea mays                     | starch branching enzyme I. sbel.  |
| AAD50279.2 | AF169833 | Sorghum bicolor              | seed starch branching enzyme. SBE.  |
| BAA01854.1 | D11081   | Zea mays                     | branching enzyme-I precursor.   |
| CAA49463.1 | X69805   | Solanum tuberosum            | 1,4-alpha-glucan branching enzyme. SBE.   |
| CAA70038.1 | Y08786   | Solanum tuberosum            | 1,4-alpha-glucan branching enzyme. sbel.  |
| AAB17086.1 | U66376   | Triticum aestivum            | 1,4-alpha-D-glucan 6-alpha-D-(1,4-alpha-D-glucanotransferase. branching enzyme.   |
| AAB67316.1 | U65948   | Zea mays                     | formation of alpha-1-6 glucosidic linkage in starch biosynthesis. starch branching enzyme IIa. Sbe2a. starch branching enzyme isozyme SBEIIa. |
| BAB40334.1 | AB042937 | Ipomoea batatas              | starch branching enzyme. IBE.   |
| BAA01584.1 | D10752   | Oryza sativa                 | branching enzyme.   |
| AAD28284.1 | AF136268 | Oryza sativa subsp. japonica | starch-branching enzyme I. Rbel.  |
| BAA01616.1 | D10838   | Oryza sativa                 | 1,4-alpha-glucan branching enzyme. sbel.  |
| BAA01855.1 | D11082   | Oryza sativa                 | branching enzyme-I precursor.   |
| CAB40981.1 | AJ237897 | Triticum aestivum            | starch branching enzyme I. sbel. alternative.   |
| CAB40979.1 | AJ237897 | Triticum aestivum            | starch branching enzyme I. sbel.  |
| CAB40980.1 | AJ237897 | Triticum aestivum            | starch branching enzyme I. sbel. alternative.   |
| AAG27622.1 | AF286318 | Triticum aestivum            | starch branching enzyme I. SbelA. glucosyltransferase.  |
| CAA54308.1 | X77012   | Manihot esculenta            | 1,4-alpha-glucan branching enzyme. SBE.   |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| CAA72987.1     | Y12320   | Triticum aestivum       | starch branching enzyme I. Sbe1.  |
| AAG27621.1     | AF286317 | Triticum aestivum       | starch branching enzyme 1. Sbe1D. glucosyltransferase.  |
| BAA82349.1     | AB029549 | Phaseolus vulgaris      | branching enzyme 3. kbe3.   |
| AAB61925.1     | AF002820 | Triticum aestivum       | starch branching enzyme I. wSBE I-D2.   |
| CAB40749.1     | AJ011891 | Solanum tuberosum       | starch branching enzyme II. SBE II.   |
| CAB40745.1     | AJ011887 | Solanum tuberosum       | starch branching enzyme II. SBE II.   |
| CAB40744.1     | AJ011886 | Solanum tuberosum       | starch branching enzyme II. SBE II.   |
| BAA85762.1     | AB028067 | Nicotiana tabacum       | starch branching enzyme. SBE.   |
| CAA49371.1     | X69713   | Manihot esculenta       | branching enzyme. r-2.  |
| BAB40335.1     | AB042940 | Ipomoea batatas         | starch branching enzyme. IBE.   |
| CAA49370.1     | X69712   | Manihot esculenta       | branching enzyme. r-1.  |
| AAC72336.1     | AF064563 | Hordeum vulgare         | starch branching enzyme IIb. sbeIIb.  |
| SEQ ID NO: 673 |          |                         |   |
| CAA09881.1     | AJ011939 | Trifolium repens        | peroxidase. prx2.   |
| CAA62228.1     | X90695   | Medicago sativa         | peroxidase2. prx2.  |
| CAA71495.1     | Y10469   | Spinacia oleracea       | peroxidase. prxr8.  |
| AAB41812.1     | L36158   | Medicago sativa         | peroxidase. pxdD. amino acid feature: conserved domains, aa 120 .. 126, 188 .. 195; amino acid feature: heme-binding domain, aa 63 .. 68. |
| BAA77387.1     | AB024437 | Scutellaria baicalensis | peroxidase 1.   |
| AAF63024.1     | AF244921 | Spinacia oleracea       | hydrogen peroxide catabolism. peroxidase prx12 precursor. type III peroxidase.  |
| AAD11483.1     | U51193   | Glycine max             | peroxidase. sEPb1.  |
| AAB67737.1     | L77080   | Stylosanthes humilis    | cationic peroxidase.  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| BAA07663.1 | D42064   | Nicotiana tabacum       | cationic peroxidase isozyme 38K precursor.  |
| BAA07664.1 | D42065   | Nicotiana tabacum       | cationic peroxidase isozyme 40K precursor.  |
| CAB94692.1 | AJ242742 | Ipomoea batatas         | Removal of H <sub>2</sub> O <sub>2</sub> , oxidation of toxic reductants, defence response toward wounding.<br>peroxidase. pod. |
| CAB67121.1 | Y19023   | Lycopersicon esculentum | peroxidase. cevi-1.   |
| CAA62226.1 | X90693   | Medicago sativa         | peroxidase1B. prx1B.  |
| CAA50597.1 | X71593   | Lycopersicon esculentum | peroxidase. CEVI-1.   |
| AAD11481.1 | U51191   | Glycine max             | peroxidase precursor. sEPa1.  |
| AAD11484.1 | U51194   | Glycine max             | peroxidase. sEPb2.  |
| BAA82306.1 | AB027752 | Nicotiana tabacum       | peroxidase.   |
| AAA65637.1 | L13654   | Lycopersicon esculentum | peroxidase. TPX1.   |
| AAD11482.1 | U51192   | Glycine max             | peroxidase precursor. sEPa2.  |
| CAA62225.1 | X90692   | Medicago sativa         | peroxidase1A. prx1A.  |
| AAC98519.1 | AF007211 | Glycine max             | peroxidase precursor. GMIPER1. pathogen-induced.  |
| AAD37427.1 | AF149277 | Phaseolus vulgaris      | peroxidase 1 precursor. FBP1. secretory peroxidase.   |
| BAA14144.1 | D90116   | Armoracia rusticana     | peroxidase isozyme.   |
| AAA98491.1 | L36981   | Petroselinum crispum    | anionic peroxidase.   |
| CAA71488.1 | Y10462   | Spinacia oleracea       | peroxidase. prxr1.  |
| AAD43561.1 | AF155124 | Gossypium hirsutum      | bacterial-induced peroxidase precursor. Perx_Goshiko.   |
| BAA14143.1 | D90115   | Armoracia rusticana     | peroxidase isozyme.   |
| CAA71490.1 | Y10464   | Spinacia oleracea       | peroxidase. prxr3.  |
| AAB02554.1 | L37790   | Stylosanthes humilis    | cationic peroxidase.  |

|                |          |  |
|----------------|----------|--|
| CAA66037.1     | X97351   | Populus balsamifera subsp. trichocarpa<br>signal for ER. peroxidase.   |
| BAA01877.1     | D11102   | Populus kitakamiensis<br>peroxidase. prxA1.  |
| CAA62227.1     | X90694   | Medicago sativa<br>peroxidase1C. prx1C.  |
| BAA07241.1     | D38051   | Populus kitakamiensis<br>peroxidase. prxA4a.   |
| BAA01950.1     | D11337   | Vigna angularis<br>peroxidase.   |
| AAF65464.2     | AF247700 | Oryza sativa<br>peroxidase POC1.   |
| AAF63027.1     | AF244924 | Spinacia oleracea<br>hydrogen peroxide catabolism. peroxidase prx15 precursor. type III peroxidase.  |
| CAC21393.1     | AJ401276 | Zea mays<br>peroxidase. pox3.  |
| CAA59487.1     | X85230   | Triticum aestivum<br>peroxidase. pox4.   |
| AAD37430.1     | AF149280 | Phaseolus vulgaris<br>peroxidase 5 precursor. FBP5. secretory peroxidase.  |
| CAA71491.1     | Y10465   | Spinacia oleracea<br>peroxidase. prxr4.  |
| AAB41811.1     | L36157   | Medicago sativa<br>peroxidase. pxdC. amino acid feature: conserved domains, aa 123 .. 129, 191 .. 198; amino<br>acid feature: heme-binding domain, aa 68 .. 73.    |
| BAA03644.1     | D14997   | Oryza sativa<br>peroxidase.  |
| AAA32676.1     | M37637   | Arachis hypogaea<br>cationic peroxidase. PNC2.   |
| CAA71494.1     | Y10468   | Spinacia oleracea<br>peroxidase. prxr7.  |
| AAA34050.1     | M74103   | Nicotiana sylvestris<br>anionic peroxidase.  |
| CAA40796.1     | X57564   | Armoracia rusticana<br>peroxidase. peroxidase precursor.   |
| SEQ ID NO: 674 |          |  |
| AAG31438.1     | AF241793 | Perilla frutescens<br>limonene synthase.   |
| BAA08367.1     | D49368   | Perilla frutescens<br>limonene cyclase.  |
| BAA21629.1     | AB005744 | Perilla frutescens<br>catalyzing the cyclization of geranyl pyrophosphate to 1-limonene. 1-limonene synthase.<br>gPFLC. similar to perilla PFLC-1:DDBJ Acc#D49368. |

|                |          |                                    |  |
|----------------|----------|------------------------------------|--|
| AAG31437.1     | AF241792 | Perilla frutescens                 | limonene synthase.   |
| AAK06663.1     | AF317695 | Perilla frutescens var. frutescens | limonene synthase.   |
| AAG31435.1     | AF241790 | Perilla citriodora                 | limonene synthase.   |
| AAF65545.1     | AF233894 | Perilla citriodora                 | limonene synthase.   |
| AAD50304.1     | AF175323 | Mentha longifolia                  | limonene synthase, monoterpene synthase.   |
| AAC37366.1     | L13459   | Mentha spicata                     | 4S-limonene synthase.  |
| AAG01140.1     | AF282875 | Schizonepeta tenuifolia            | (+)-4R-limonene synthase.  |
| AAC61260.1     | AF061285 | Capsicum annuum                    | sesquiterpene cyclase. UV induced.   |
| AAG09949.1     | AF171216 | Lycopersicon esculentum            | vetispiradiene synthase. LEVS2. sesquiterpene cyclase.   |
| AAF74977.1     | AF270425 | Gossypium hirsutum                 | (E,E)-farnesyl diphosphate cyclase. (+)-delta-cadinene synthase. cdn2. sesquiterpene cyclase.                            |
| BAA82141.1     | AB023816 | Solanum tuberosum                  | vetispiradiene synthase. PVS4. potato sesquiterpene cyclase.   |
| BAA82092.1     | AB022598 | Solanum tuberosum                  | vetispiradiene synthase. PVS1. potato sesquiterpene cyclase.   |
| AAF21053.1     | AF212433 | Capsicum annuum                    | UV-induced sesquiterpene cyclase. SC2.   |
| BAA82109.1     | AB022720 | Solanum tuberosum                  | vetispiradiene synthase. PVS3. potato sesquiterpene cyclase; PVS3.   |
| BAA82108.1     | AB022719 | Solanum tuberosum                  | vetispiradiene synthase. PVS2. potato sesquiterpene cyclase; PVS2.   |
| AAC12784.1     | U88318   | Gossypium hirsutum                 | (E,E)-farnesyl diphosphate cyclizing. (+)-delta-cadinene synthase. cdn1. sesquiterpene cyclase; delta-cadinene synthase. |
| AAG24640.2     | AF304444 | Artemisia annua                    | sesquiterpene cyclase.   |
| AAK15641.1     | AF326117 | Capsicum annuum                    | sesquiterpene cyclase. PSC2.   |
| CAC12731.1     | AJ271792 | Artemisia annua                    | putative sesquiterpene cyclase. cASC125.   |
| SEQ ID NO: 675 |          |                                    |  |
| CAA50609.1     | X71609   | Nicotiana tabacum                  | ras-related GTP-binding protein.   |

|  |          |                                       |
|--|----------|---------------------------------------|
| CAA98166.1   | Z73938   | Lotus japonicus                       |
| GTP-binding protein. RAB5A. rab5A.                           |          |                                       |
| CAC24477.1   | AJ296336 | Cichorium intybus x Cichorium endivia |
| GTP binding protein. chi3154.                                |          |                                       |
| CAC24476.1   | AJ296335 | Cichorium intybus x Cichorium endivia |
| GTP binding protein. chi3152.                                |          |                                       |
| CAB57220.1   | AJ249866 | Cichorium intybus x Cichorium endivia |
| GTP binding protein. gtp2.                                   |          |                                       |
| CAB57219.1   | AJ249865 | Cichorium intybus x Cichorium endivia |
| GTP binding protein. gtp1.                                   |          |                                       |
| CAC24475.1   | AJ296334 | Cichorium intybus x Cichorium endivia |
| GTP binding protein. chi3153.                                |          |                                       |
| CAA46112.1   | X64941   | Nicotiana plumbaginifolia             |
| small GTP binding protein.                                   |          |                                       |
| CAC24474.1   | AJ296333 | Cichorium intybus x Cichorium endivia |
| GTP binding protein. chi3151.                                |          |                                       |
| CAC19792.1   | AJ292320 | Oryza sativa                          |
| small GTP-binding protein, RAB family. RAB5A protein. rab5A. |          |                                       |
| AAD28731.1   | AF112964 | Triticum aestivum                     |
| small GTP-binding protein. Sgp.                              |          |                                       |
| CAA98167.1   | Z73939   | Lotus japonicus                       |
| GTP-binding protein. RAB5B. rab5B.                           |          |                                       |
| CAA06922.1   | AJ006225 | Mesembryanthemum crystallinum         |
| small GTP-binding protein. rab5B.                            |          |                                       |
| AAG42497.1   | AF323991 | Oryza sativa                          |
| small GTP-binding protein RAB5B. rab5B.                      |          |                                       |
| BAA84717.1   | AB032761 | Oryza sativa                          |
| GTP-binding protein. rab5B.                                  |          |                                       |
| AAG24438.1   | AF304518 | Oryza sativa                          |
| small GTP-binding protein RAB5B. rab5B.                      |          |                                       |
| CAB57221.1   | AJ249867 | Cichorium intybus x Cichorium endivia |
| GTP binding protein. bl.5.                                   |          |                                       |
| CAA98180.1   | Z73952   | Lotus japonicus                       |
| GTP-binding protein. RAB11D. rab11D.                         |          |                                       |
| CAA98181.1   | Z73953   | Lotus japonicus                       |
| GTP-binding protein. RAB11E. rab11E.                         |          |                                       |
| BAA02114.1   | D12546   | Pisum sativum                         |
| GTP-binding protein.   |          |                                       |
| BAA02113.1   | D12545   | Pisum sativum                         |
| GTP-binding protein.   |          |                                       |
| BAA02116.1   | D12548   | Pisum sativum                         |
| GTP-binding protein.   |          |                                       |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAA02112.1     | D12544   | Pisum sativum           | GTP-binding protein.   |
| BAA02118.1     | D12550   | Pisum sativum           | GTP-binding protein.   |
| CAA98159.1     | Z73931   | Lotus japonicus         | GTP-binding protein. RAB1B. rab1B.   |
| CAB65172.1     | AJ245570 | Lycopersicon esculentum | putative role in secretion of cell wall modifying enzymes. Rab11 GTPase. Rab11a. |
| CAA89049.1     | Z49190   | Beta vulgaris           | GTP-binding. small G protein.  |
| BAA02110.1     | D12542   | Pisum sativum           | GTP-binding protein.   |
| AAB97114.1     | U58853   | Glycine max             | small GTP-binding protein. sral.   |
| BAA02904.1     | D13758   | Oryza sativa            | ras-related GTP binding protein. ss230.  |
| CAA98178.1     | Z73950   | Lotus japonicus         | GTP-binding protein. RAB11B. rab11B.   |
| CAA98177.1     | Z73949   | Lotus japonicus         | GTP-binding protein. RAB11A. rab11A.   |
| AAK15703.1     | AF327517 | Oryza sativa            | GTP-binding protein.   |
| AAF65510.1     | AF108883 | Capsicum annuum         | small GTP-binding protein.   |
| BAA76422.1     | AB024994 | Cicer arietinum         | rab-type small GTP-binding protein.  |
| CAA98160.1     | Z73932   | Lotus japonicus         | GTP-binding protein. RAB1C. rab1C.   |
| AAA80680.1     | U38466   | Lycopersicon esculentum | small GTP-binding protein. LeRab1C. ; YPT1/Rab1A homolog LeRab1C.                |
| AAA80678.1     | U38464   | Lycopersicon esculentum | small GTP-binding protein. LeRab1A. ; YPT1/Rab1A homolog LeRab1A.                |
| SEQ ID NO: 676 |          |                         |  |
| AAA80499.1     | U20594   | Lycopersicon esculentum | leucine aminopeptidase.  |
| CAA54314.1     | X77015   | Solanum tuberosum       | leucine aminopeptidase. LAP.   |
| CAA48038.1     | X67845   | Solanum tuberosum       | leucine aminopeptidase. LAP.   |
| AAC49457.1     | U50152   | Lycopersicon esculentum | peptidase. leucine aminopeptidase. lap2.   |

|                |          |   |
|----------------|----------|---|
| AAC49456.1     | U50151   | Lycopersicon esculentum<br>exoprotease in the defense response. leucine aminopeptidase. lap.  |
| AAA80498.1     | U20593   | Lycopersicon esculentum<br>leucine aminopeptidase.  |
| CAA68143.1     | X99825   | Petroselinum crispum<br>cytosol aminopeptidase. leucine aminopeptidase.   |
| BAA90521.1     | AB037678 | Phaseolus vulgaris<br>leucine aminopeptidase.   |
| SEQ ID NO: 678 |          |   |
| CAC09580.1     | AJ298992 | Fagus sylvatica<br>Absciscic acid (ABA) and calcium induced protein kinase. protein kinase (PK). pk1.   |
| AAA34002.1     | M67449   | Glycine max<br>protein kinase. PK6.   |
| AAK11734.1     | AY027437 | Arachis hypogaea<br>serine/threonine/tyrosine kinase.   |
| BAB16918.1     | AP002863 | Oryza sativa<br>putative protein kinase. P0005A05.22.   |
| CAA06334.1     | AJ005077 | Lycopersicon esculentum<br>protein kinase. TCTR2 protein. TCTR2.  |
| AAG31141.1     | AF305911 | Oryza sativa<br>EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.   |
| CAA73722.1     | Y13273   | Lycopersicon esculentum<br>putative protein kinase.   |
| AAD46406.1     | AF096250 | Lycopersicon esculentum<br>ethylene-responsive protein kinase TCTR1. ER50. serine/threonine kinase; similar to Arabidopsis thaliana negative regulator of the ethylene response pathway encoded by GenBank Accession Number L08789. |
| AAG31142.1     | AF305912 | Hordeum vulgare<br>EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.  |
| AAD10056.1     | AF110518 | Lycopersicon esculentum<br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1.  |
| AAD10057.1     | AF110519 | Lycopersicon esculentum<br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1v.   |
| AAK30005.1     | AY029067 | Rosa hybrid cultivar<br>CTR2 protein kinase.  |
| BAB39409.1     | AP002901 | Oryza sativa<br>putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| BAA87853.1     | AP000816 | Oryza sativa<br>EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |



|            |          |                   |  |
|------------|----------|-------------------|--|
| BAA06538.1 | D31737   | Nicotiana tabacum | protein-serine/threonine kinase.   |
| BAB21240.1 | AP002953 | Oryza sativa      | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).                         |
| BAA94528.1 | AP001800 | Oryza sativa      | Similar to Arabidopsis thaliana chromosome 2 BAC T20K24; putative receptor-like protein kinase (AC002392). |
| BAB17126.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.16.   |
| BAB39451.1 | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.24.   |
| BAB17348.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.32.   |
| BAB17321.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.1.  |
| BAB17129.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.20.   |
| AAF68398.1 | AF237568 | Oryza sativa      | receptor-like protein kinase. RLG2.  |
| BAB07905.1 | AP002835 | Oryza sativa      | putative S-receptor kinase. P0417G05.13.   |
| BAA94517.1 | AP001800 | Oryza sativa      | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).                                      |
| BAB17335.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.17. contains ESTs D47575(S13157),AU032665(S13157).                      |
| BAB39441.1 | AP003338 | Oryza sativa      | putative receptor kinase. OJ1212_B09.11.   |
| BAB17116.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.3.  |
| AAF78044.1 | AF248493 | Oryza sativa      | receptor-like kinase. RLG18. protein kinase.   |
| CAB51834.1 | 00069    | Oryza sativa      | l1332.5. contains eukaryotic protein kinase domain PF.   |
| BAB17127.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.17.   |
| AAF78021.1 | AF238477 | Oryza sativa      | receptor-like kinase. RLG5. protein kinase.  |
| AAD46420.1 | AF100771 | Hordeum vulgare   | receptor-like kinase. Hv3ARK. similar to wheat ARK1AS.   |
| AAD46917.1 | AF164021 | Oryza sativa      | receptor kinase.   |
| BAB17345.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.29.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAB39438.1     | AP003338 | Oryza sativa            | putative receptor kinase. OJ1212_B09.7.  |
| AAF78018.1     | AF238474 | Oryza sativa            | receptor-like kinase. RLG16. protein kinase.   |
| AAC01746.1     | AF044489 | Oryza sativa            | receptor-like protein kinase. drpk1.   |
| BAB17337.1     | AP002747 | Oryza sativa            | putative receptor kinase. P0698G03.21. contains ESTs D47575(S13157),AU032665(S13157).  |
| BAB17347.1     | AP002747 | Oryza sativa            | putative receptor kinase. P0698G03.31. contains ESTs D47575(S13157),AU032665(S13157).  |
| BAB17344.1     | AP002747 | Oryza sativa            | putative receptor kinase. P0698G03.28. contains ESTs D47575(S13157),AU032665(S13157).  |
| BAB39437.1     | AP003338 | Oryza sativa            | receptor-like kinase. OJ1212_B09.6.  |
| AAF78019.1     | AF238475 | Oryza sativa            | receptor-like kinase. RLG17. protein kinase.   |
| BAA05648.1     | D26601   | Nicotiana tabacum       | protein kinase.  |
| BAB17332.1     | AP002747 | Oryza sativa            | putative receptor kinase. P0698G03.13. contains ESTs D47575(S13157),AU032665(S13157).  |
| BAB39435.1     | AP003338 | Oryza sativa            | putative receptor kinase. OJ1212_B09.2.  |
| SEQ ID NO: 679 |          |                         |  |
| CAA78386.1     | Z13996   | Petunia x hybrida       | DNA binding protein; transcriptional activator. protein 1. myb.Ph3. Product related to animal myb proto-oncoproteins. Sequence from nucleotide 992 is not included in clone cPF1 and has been obtained by PCR amplification of cDNA. |
| CAB43399.1     | AJ006292 | Antirrhinum majus       | Myb-related transcription factor mixta-like 1. mybml1.   |
| CAA67600.1     | X99210   | Lycopersicon esculentum | myb-related transcription factor. THM16.   |
| BAA23337.1     | D88617   | Oryza sativa            | transfactor. OSMYB1. Osmyb1.   |
| AAA82943.1     | U39448   | Picea mariana           | MYB-like transcriptional factor MBF1. putative DNA binding region highly similar to the maize C1.  |
| CAA64614.1     | X95296   | Lycopersicon esculentum | transcription factor. THM27. myb-related.  |
| BAA93038.1     | AP001552 | Oryza sativa            | EST AU082058(C12976) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana putative transcription factor (AF062916).  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAC04720.1 | AF034134 | Gossypium hirsutum      | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-O. similar to MYB A encoded by GenBank Accession Number L04497. |
| CAA72186.1 | Y11351   | Oryza sativa            | myb factor. myb.   |
| AAF22256.1 | AF161711 | Pimpinella brachycarpa  | myb-related transcription factor.  |
| BAB39987.1 | AP003020 | Oryza sativa            | putative transcription factor (myb). P0498A12.16. contains ESTs AU097474(S5087),D40175(S1959).   |
| BAB39972.1 | AP003018 | Oryza sativa            | putative transcription factor (myb). OSJNBa0004B13.27. contains ESTs AU097474(S5087),D40175(S1959).  |
| BAA23338.1 | D88618   | Oryza sativa            | transfactor. OSMYB2. Osmyb2.   |
| BAA88222.1 | AB028650 | Nicotiana tabacum       | myb-related transcription factor LBM2. lbm2.   |
| CAA67575.1 | X99134   | Lycopersicon esculentum | transcription factor. THM6. myb-related.   |
| CAA78387.1 | Z13997   | Petunia x hybrida       | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.                                    |
| CAA66952.1 | X98308   | Lycopersicon esculentum | THM18. myb-related transcription factor.   |
| BAA23339.1 | D88619   | Oryza sativa            | transfactor. OSMYB3. Osmyb3.   |
| AAC49394.1 | U57002   | Zea mays                | P protein. P. allele P-wr; Myb-like transcriptional regulator with a putative zinc-finger at the C-terminal.                                     |
| AAB41101.1 | U72762   | Nicotiana tabacum       | transcription factor Myb1. myb1. TMV-inducible Myb homolog; contains helix-turn-helix motif; contains redox-sensitive cysteine.                  |
| AAA33500.1 | M73028   | Zea mays                | myb-like transcription factor. P.  |
| AAG36774.1 | AF210616 | Zea mays                | P2 protein. P2. myb-like transcriptional factor; similar to Zea mays P gene.   |
| BAA88223.1 | AB028651 | Nicotiana tabacum       | myb-related transcription factor LBM3. lbm3.   |
| BAA88224.1 | AB028652 | Nicotiana tabacum       | myb-related transcription factor LBM4. lbm4.   |
| CAA72187.1 | Y11352   | Oryza sativa            | myb factor. myb.   |

|                |          |                    |  |
|----------------|----------|--------------------|--|
| BAA88221.1     | AB028649 | Nicotiana tabacum  | myb-related transcription factor LBM1. lbm1.   |
| CAA72185.1     | Y11350   | Oryza sativa       | myb factor. myb.   |
| AAG13574.1     | AC037425 | Oryza sativa       | myb factor. OSJNBa0055P24.4.   |
| SEQ ID NO: 680 |          |                    |  |
| AAA34238.1     | L20507   | Vigna radiata      | calmodulin.  |
| AAA34014.1     | L01432   | Glycine max        | calcium-binding regulatory protein. calmodulin. SCaM-3. putative.  |
| AAA34013.1     | L01430   | Glycine max        | calcium-binding regulatory protein. calmodulin. SCaM-1. putative.  |
| CAA36644.1     | X52398   | Medicago sativa    | calmodulin (AA 1-149).   |
| AAD10245.1     | AF030033 | Phaseolus vulgaris | calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal transduction pathways.                              |
| AAD10244.1     | AF030032 | Phaseolus vulgaris | calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal transduction pathways.                              |
| CAA74307.1     | Y13974   | Zea mays           | calmodulin.  |
| CAA46150.1     | X65016   | Oryza sativa       | calmodulin. cam.   |
| AAC36058.1     | AF042839 | Oryza sativa       | calmodulin. CaM2.  |
| AAD10246.1     | AF030034 | Phaseolus vulgaris | calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal transduction pathways.                              |
| CAA54583.1     | X77397   | Zea mays           | calmodulin. CaM2.  |
| BAA87825.1     | AP000815 | Oryza sativa       | ESTs AU030013(E50493),AU081341(E50493) correspond to a region of the predicted gene. Similar to O.sativa gene encoding calmodulin. (Z12828). |
| AAA92681.1     | U13882   | Pisum sativum      | calcium-binding protein. calmodulin.   |
| AAA33706.1     | M80836   | Petunia x hybrida  | calmodulin. CAM81.   |
| AAA33705.1     | M80831   | Petunia x hybrida  | calmodulin-related protein. CAM53.   |
| CAA43143.1     | X60738   | Malus x domestica  | Calmodulin. CaM.   |

|            |          |  |
|------------|----------|--|
| CAA78301.1 | Z12839   | <i>Lilium longiflorum</i><br>calcium binding protein, signal transduction. calmodulin.   |
| AAA33397.1 | L18912   | <i>Lilium longiflorum</i><br>calcium binding protein, signal transduction. calmodulin. putative.                               |
| CAA42423.1 | X59751   | <i>Daucus carota</i><br>calmodulin. Ccam-1.  |
| CAA67054.1 | X98404   | <i>Capsicum annuum</i><br>calmodulin-2.  |
| AAG27432.1 | AF295637 | <i>Elaeis guineensis</i><br>calmodulin.  |
| AAG11418.1 | AF292108 | <i>Prunus avium</i><br>calmodulin.   |
| AAA34237.1 | L20691   | <i>Vigna radiata</i><br>calmodulin.  |
| AAC49587.1 | U49105   | <i>Triticum aestivum</i><br>calmodulin TaCaM4-1. calcium-binding protein.  |
| AAC49586.1 | U49104   | <i>Triticum aestivum</i><br>calmodulin TaCaM3-3. calcium-binding protein.  |
| AAC49585.1 | U49103   | <i>Triticum aestivum</i><br>calmodulin TaCaM3-2. calcium-binding protein.  |
| AAC49584.1 | U48693   | <i>Triticum aestivum</i><br>calmodulin TaCaM3-1. calcium-binding protein.  |
| AAC49583.1 | U48692   | <i>Triticum aestivum</i><br>calmodulin TaCaM2-3. calcium-binding protein.  |
| AAC49582.1 | U48691   | <i>Triticum aestivum</i><br>calmodulin TaCaM2-2. calcium-binding protein.  |
| AAC49580.1 | U48689   | <i>Triticum aestivum</i><br>calmodulin TaCaM1-3. calcium-binding protein.  |
| AAC49579.1 | U48688   | <i>Triticum aestivum</i><br>calmodulin TaCaM1-2. calcium binding protein.  |
| AAC49578.1 | U48242   | <i>Triticum aestivum</i><br>calmodulin TaCaM1-1. calcium-binding.  |
| AAA03580.1 | L01431   | <i>Glycine max</i><br>calcium-binding regulatory protein. calmodulin. SCaM-2. putative.  |
| AAB36130.1 | S81594   | <i>Vigna radiata</i><br>auxin-regulated calmodulin. auxin-regulated calmodulin, arCaM. This sequence comes from Fig. 1; arCaM. |
| AAA33901.1 | L18913   | <i>Oryza sativa</i><br>calcium binding protein, signal transduction. calmodulin. putative.                                     |
| CAA78287.1 | Z12827   | <i>Oryza sativa</i><br>calcium binding protein, signal transduction. calmodulin.   |
| AAB46588.1 | U83402   | <i>Capsicum annuum</i><br>calmodulin.  |

|                |          |                   |   |
|----------------|----------|-------------------|---|
| CAA61980.1     | X89890   | Bidens pilosa     | Calmodulin.   |
| AAA32938.1     | M27303   | Hordeum vulgare   | calmodulin.   |
| BAA88540.1     | AP000969 | Oryza sativa      | ESTs AU081349(E61253), D41425(S3918) correspond to a region of the predicted gene. Similar to calmodulin. (AF042840). |
| AAF65511.1     | AF108889 | Capsicum annuum   | calmodulin.   |
| AAC36059.1     | AF042840 | Oryza sativa      | calmodulin. CaM1.   |
| AAA33900.1     | L18914   | Oryza sativa      | calcium binding protein, signal transduction. calmodulin.   |
| CAA78288.1     | Z12828   | Oryza sativa      | calcium binding protein, signal transduction. calmodulin.   |
| AAA34015.1     | L01433   | Glycine max       | calcium-binding regulatory protein. calmodulin. SCaM-4. putative.   |
| AAA16320.1     | L14071   | Bryonia dioica    | calmodulin. Bc329. calcium-binding sites (amino acid #): 1. (21..33); 2. (57..68); 3. (94..106); 4. (130..141).       |
| AAA19571.1     | U10150   | Brassica napus    | calcium binding. calmodulin. bcm1.  |
| AAA87347.1     | M88307   | Brassica juncea   | calmodulin.   |
| CAA52602.1     | X74490   | Zea mays          | Calmodulin. ZMCALM1.  |
| SEQ ID NO: 683 |          |                   |   |
| CAA71800.1     | Y10847   | Brassica juncea   | O-acetylserine(thiol) lyase.  |
| CAA71798.1     | Y10845   | Brassica juncea   | O-acetylserine(thiol) lyase.  |
| AAC25635.1     | AF044172 | Solanum tuberosum | cysteine synthase. CS-A; O-acetylserine (thiol) lyase; cytosolic isoform.   |
| BAA01279.1     | D10476   | Spinacia oleracea | O-acetylserine(thiol) lyase.  |
| BAA02438.1     | D13153   | Triticum aestivum | O-acetylserine (thiol) lyase. cys1.   |
| CAA59798.1     | X85803   | Zea mays          | O-acetylserine (thiol) lyase. Mcysp. cysteine synthase.   |
| AAD23907.1     | AF073695 | Oryza sativa      | cysteine synthase. rcs1. O-acetylserine(thiol)-lyase.   |
| AAD23909.1     | AF073697 | Oryza sativa      | cysteine synthase. rcs3. O-acetylserine(thiol)-lyase.   |

|                |          |                               |  |
|----------------|----------|-------------------------------|--|
| AAC25636.1     | AF044173 | Solanum tuberosum             | cysteine synthase. CS-B; O-acetylserine (thiol) lyase; plastidic isoform.  |
| AAC27794.1     | AF078693 | Chlamydomonas reinhardtii     | cysteine biosynthesis. putative O-acetylserine(thiol)lyase precursor. Crcys-1A.  |
| CAA06819.1     | AJ006024 | Cicer arietinum               | cysteine synthase, O-acetyl-L-serine (thiol)-lyase.  |
| CAA46086.1     | X64874   | Capsicum annuum               | O-acetylserine (thiol)-lyase.  |
| CAA71799.1     | Y10846   | Brassica juncea               | O-acetylserine(thiol) lyase.   |
| AAA16973.1     | L05184   | Chloroplast Spinacia oleracea | O-acetylserine-(thiol)-lyase.  |
| AAD23908.1     | AF073696 | Oryza sativa                  | cysteine synthase. rcs2. O-acetylserine(thiol)-lyase.  |
| AAD23910.1     | AF073698 | Oryza sativa                  | cysteine synthase. rcs4. O-acetylserine(thiol)-lyase.  |
| SEQ ID NO: 684 |          |                               |  |
| AAA68983.1     | L12395   | Brassica napus                | signal transduction, membrane vehicle traffic. small GTP-binding protein. bra. putative.   |
| SEQ ID NO: 685 |          |                               |  |
| AAB61961.1     | L81152   | Oryza sativa                  | integral membrane protein. OsNramp2.   |
| AAB36424.1     | S81897   | Oryza sativa                  | OsNramp1. OsNramp1. Nramp1 homolog/Bcg product homolog; This sequence comes from Fig. 2.   |
| AAB62273.1     | L41217   | Oryza sativa                  | integral membrane protein. OsNramp1. putative.   |
| AAC49720.1     | U60767   | Oryza sativa                  | integral membrane protein OsNramp3. OsNramp3.  |
| SEQ ID NO: 686 |          |                               |  |
| AAC04719.1     | AF034133 | Gossypium hirsutum            | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. Cmy-N. similar to MYB A encoded by GenBank Accession Number L04497. |
| AAB58314.1     | U33917   | Craterostigma plantagineum    | Cpm7. cpm7. putative DNA-binding protein; myb-like gene; myb-related transcription factor.   |
| AAB58313.1     | U33916   | Craterostigma plantagineum    | Cpm5. cpm5. putative DNA-binding protein; myb-related gene; myb-related transcription factor.  |
| AAC13876.1     | U33915   | Craterostigma plantagineum    | myb-related transcription factor Cpm10. cpm10. putative DNA-binding protein.   |

|            |          |                          |  |
|------------|----------|--------------------------|--|
| AAK08983.1 | AY026332 | Oryza sativa             | Myb transcription factor JAMyb. related to host cell death and defense responses; induced by jasmonic acid, wounding, or infection of rice blast fungus, but not by salicylic acid or abscisic acid. |
| BAB18296.1 | AP002860 | Oryza sativa             | putative myb-related transcription factor. P0409B08.23.  |
| CAA71992.1 | Y11105   | Pisum sativum            | Myb26.   |
| BAB40790.1 | AB058642 | Lilium hybrid division I | LhMyb.   |
| AAK19618.1 | AF336285 | Gossypium hirsutum       | GHMYB38. ghmyb38. similar to myb.  |
| CAA78387.1 | Z13997   | Petunia x hybrida        | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.  |
| BAA88221.1 | AB028649 | Nicotiana tabacum        | myb-related transcription factor LBM1. lbm1.   |
| BAA88224.1 | AB028652 | Nicotiana tabacum        | myb-related transcription factor LBM4. lbm4.   |
| AAK19617.1 | AF336284 | Gossypium hirsutum       | GHMYB36. ghmyb36. similar to myb.  |
| AAA33067.1 | L04497   | Gossypium hirsutum       | MYB A; putative.   |
| AAK19615.1 | AF336282 | Gossypium hirsutum       | GHMYB10. ghmyb10. similar to myb.  |
| BAA81733.2 | AB029162 | Glycine max              | GmMYB29A2.   |
| BAA81732.1 | AB029161 | Glycine max              | GmMYB29A2.   |
| BAA81730.1 | AB029159 | Glycine max              | GmMYB29A1.   |
| BAA88223.1 | AB028651 | Nicotiana tabacum        | myb-related transcription factor LBM3. lbm3.   |
| AAB41101.1 | U72762   | Nicotiana tabacum        | transcription factor Myb1. myb1. TMV-inducible Myb homolog; contains helix-turn-helix motif; contains redox-sensitive cysteine.  |
| BAA81736.1 | AB029165 | Glycine max              | GmMYB29B2.   |
| BAA81731.1 | AB029160 | Glycine max              | GmMYB29A1.   |
| CAA72218.1 | Y11415   | Oryza sativa             | myb.   |



|            |          |   |
|------------|----------|---|
| CAA67000.1 | X98355   | <i>Oryza sativa</i><br>activator of alpha-amylase gene promoter. transcription factor GAMyb. Gam1. Myb-like; expression is regulated by gibberellin.                          |
| AAK19611.1 | AF336278 | <i>Gossypium hirsutum</i><br>BNLGH1233. bnlghi6233. similar to myb.   |
| CAA67575.1 | X99134   | <i>Lycopersicon esculentum</i><br>transcription factor. THM6. myb-related.  |
| CAA64614.1 | X95296   | <i>Lycopersicon esculentum</i><br>transcription factor. THM27. myb-related.   |
| AAD31395.1 | AF114162 | <i>Lolium temulentum</i><br>gibberellin MYB transcription factor. GAMyb. R2/R3-MYB.   |
| BAA93038.1 | AP001552 | <i>Oryza sativa</i><br>EST AU082058(C12976) corresponds to a region of the predicted gene. Similar to <i>Arabidopsis thaliana</i> putative transcription factor (AF062916).   |
| AAG22863.1 | AY008692 | <i>Hordeum vulgare</i><br>transcription factor GAMyb. Gamyb.  |
| BAA96421.1 | AB044084 | <i>Triticum aestivum</i><br>transcription activator for gibberellin response. GAMyb protein. Ta-GAMyb.  |
| CAA61021.1 | X87690   | <i>Hordeum vulgare</i><br>transcriptional activator of alpha-amylase gene promoter. GAMyb protein. Gam1.  |
| CAA66952.1 | X98308   | <i>Lycopersicon esculentum</i><br>THM18. myb-related transcription factor.  |
| AAC49394.1 | U57002   | <i>Zea mays</i><br>P protein. P. allele P-wr; Myb-like transcriptional regulator with a putative zinc-finger at the C-terminal.   |
| BAA88222.1 | AB028650 | <i>Nicotiana tabacum</i><br>myb-related transcription factor LBM2. lbm2.  |
| AAG36774.1 | AF210616 | <i>Zea mays</i><br>P2 protein. P2. myb-like transcriptional factor; similar to <i>Zea mays</i> P gene.  |
| AAA33500.1 | M73028   | <i>Zea mays</i><br>myb-like transcription factor. P.  |
| CAC19439.1 | AJ237661 | <i>Oryza sativa</i><br>Myb factor protein. myb.   |
| CAB40189.1 | AJ133638 | <i>Avena sativa</i><br>transcriptional activator. myb protein. gamyb.   |
| AAK19619.1 | AF336286 | <i>Gossypium hirsutum</i><br>GHMYB9. ghmyb9. similar to myb.  |
| AAC04720.1 | AF034134 | <i>Gossypium hirsutum</i><br>putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-O. similar to MYB A encoded by GenBank Accession Number L04497. |
| AAC04716.1 | AF034130 | <i>Gossypium hirsutum</i><br>putative MYB-like transcription factor. MYB-like DNA-binding domain protein. CmY-D. similar to MYB A encoded by GenBank Accession Number L04497. |

|                |          |                               |  |
|----------------|----------|-------------------------------|--|
| AAG28526.1     | AF198499 | Nicotiana tabacum             | anther-specific myb-related protein 1. myBAS1. NtMYBAS1; contains N-terminal R2, R3 myb domain repeats similar to c-myb.   |
| CAA78388.1     | Z13998   | Petunia x hybrida             | DNA-binding protein; transcriptional activator. protein 3. myb.Ph3. related to animal myb proto-oncoproteins.  |
| AAA33482.1     | M37153   | Zea mays                      | c1 locus myb homologue; putative.  |
| AAK09327.1     | AF320614 | Zea mays                      | activates anthocyanin transcription. anthocyanin regulatory C1. c1. transcription factor.  |
| AAK09326.1     | AF320613 | Zea mays                      | activates anthocyanin transcription. anthocyanin regulatory C1. c1. transcription factor.  |
| SEQ ID NO: 687 |          |                               |  |
| BAA92972.1     | AP001551 | Oryza sativa                  | ESTs AU056183(S20356),AU056881(S20950) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 4 BAC clone F6I18 ; putative protein kinase. (AL022198). |
| BAA83689.1     | AB011968 | Oryza sativa                  | OsPK7. OsPK7. protein kinase.  |
| BAA83688.1     | AB011967 | Oryza sativa                  | OsPK4. OsPK4. protein kinase.  |
| AAF22219.1     | AF141378 | Zea mays                      | protein kinase PK4. ZmPK4.   |
| BAA34675.1     | AB011670 | Triticum aestivum             | wpk4 protein kinase. wpk4.   |
| CAA73068.1     | Y12465   | Sorghum bicolor               | serine/threonine kinase. SNFL2.  |
| CAA73067.1     | Y12464   | Sorghum bicolor               | serine/threonine kinase. SNFL1.  |
| BAA92970.1     | AP001551 | Oryza sativa                  | Similar to Arabidopsis thaliana chromosome 4 BAC clone F28A21 ; putative protein kinase. (AL035526).   |
| CAA74646.1     | Y14274   | Sorghum bicolor               | putative serine/threonine protein kinase. SNFL3.   |
| AAD31900.1     | AF145482 | Mesembryanthemum crystallinum | putative serine/threonine protein kinase.  |
| AAB62693.1     | AF004947 | Oryza sativa                  | protein kinase.  |
| BAA96628.1     | AP002482 | Oryza sativa                  | ESTs D41739(S4522),AU055999(S20214), AU057588(S21592) correspond to a region of the predicted gene. Similar to Sorghum bicolor serine/threonine kinase (Y12465).                         |

|            |          |                   |   |
|------------|----------|-------------------|---|
| AAD23582.1 | AF128443 | Glycine max       | probably involved in plant stress responses possibly regulates gene expression. SNF-1-like serine/threonine protein kinase. expressed in nodules, roots and leaves. |
| BAA05649.1 | D26602   | Nicotiana tabacum | protein kinase.   |
| CAA71142.1 | Y10036   | Cucumis sativus   | SNF1-related protein kinase.  |
| AAC99329.1 | AF062479 | Oryza sativa      | protein kinase SNF1. Snf1. similar to yeast SNF1.   |
| CAA65244.1 | X95997   | Solanum tuberosum | SNF1-related protein kinase. PKIN1.   |
| CAA57898.1 | X82548   | Hordeum vulgare   | SNF1-related protein kinase. BKIN2.   |
| CAA07813.1 | AJ007990 | Hordeum vulgare   | SnRK1-type protein kinase. kin12a.  |
| CAA46556.1 | X65606   | Hordeum vulgare   | protein kinase. BKIN12.   |
| AAB05457.1 | U55768   | Oryza sativa      | SNF1-related protein kinase. RSk1. Ser/Thr protein kinase homolog.  |
| CAA46554.1 | X65604   | Hordeum vulgare   | protein kinase. BKIN12.   |
| AAD00239.1 | U73938   | Nicotiana tabacum | protein kinase. PK11-C1. PK11-C1. induced at the transcriptional level by the abscisic acid plant hormone; similar to serine/threonine protein kinase.              |
| AAC69450.1 | AF032465 | Nicotiana tabacum | putative serine/threonine protein kinase. WAPK.   |
| AAD00240.1 | U73939   | Nicotiana tabacum | protein kinase. PK11-C5. PK11-C5. induced at the transcriptional level by the abscisic acid plant hormone; similar to serine/threonine protein kinase.              |
| BAA13608.1 | D88399   | Oryza sativa      | serine-threonine kinase. endosperm kinase. REK.   |
| AAA34017.1 | L19360   | Glycine max       | protein kinase 2. SPK-2. putative.  |
| AAG60195.1 | AC084763 | Oryza sativa      | protein kinase REK. OSJNBa0027P10.6.  |
| AAA33004.1 | L12394   | Brassica napus    | serine/threonine protein kination. serine/threonine protein kinase. BSK2. putative.   |
| BAA19573.1 | AB002109 | Oryza sativa      | protein kinase. a novel protein kinase.   |
| AAA33979.1 | L01453   | Glycine max       | protein phosphorylation, regulatory protein. protein kinase. SPK-1. putative.   |
| AAA33003.1 | L12393   | Brassica napus    | protein kination. serine/threonine protein kinase. BSK1. putative.  |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAB68962.1 | L38855   | Glycine max                   | protein kinase. SPK-4.  |
| AAB68961.1 | L19361   | Glycine max                   | protein kinase 3. SPK-3. putative.  |
| AAB58348.1 | U29095   | Triticum aestivum             | serine-threonine protein kinase. TaPK3.   |
| AAF27340.1 | AF186020 | Vicia faba                    | abscisic acid-activated protein kinase. AAPK.                                   |
| AAA96325.1 | M94726   | Triticum aestivum             | protein kinase. abscisic acid inducible.  |
| CAA81443.1 | Z26846   | Mesembryanthemum crystallinum | protein kinase. MCPK9.  |
| AAG31326.1 | AF178575 | Vitis vinifera                | putative serine/threonine kinase GDBrPK. GDBrPK. similar to SNF protein kinase. |
| CAA06503.1 | AJ005373 | Craterostigma plantagineum    | protein kinase. cppk1.  |
| AAF21062.1 | AF216527 | Dunaliella tertiolecta        | calcium-dependent protein kinase. CPK1; CDPK.                                   |
| CAA89202.1 | Z49233   | Chlamydomonas eugametos       | calcium-stimulated protein kinase.  |

SEQ ID NO: 689

|            |          |                       |  |
|------------|----------|-----------------------|--|
| AAC49826.1 | U71604   | Catharanthus roseus   | involved in the second to last step in vindoline biosynthesis. desacetoxyvindoline 4-hydroxylase. CRSD4H. 2-oxoglutarate dependent dioxygenase.                  |
| AAB97311.1 | AF008597 | Catharanthus roseus   | desacetoxyvindoline-4-hydroxylase. CRSD4H. 2-oxoglutarate dependent dioxygenase; involved in the second to last step in vindoline biosynthesis.                  |
| AAC49827.1 | U71605   | Catharanthus roseus   | involved in the second to last step in vindoline biosynthesis. desacetoxyvindoline 4-hydroxylase. CRSD4H. 2-oxoglutarate dependent dioxygenase.                  |
| BAA37127.1 | AB012203 | Lactuca sativa        | 2-oxoglutarate-dependent dioxygenase. gibberelin 20-oxidase. Ls20ox1.  |
| BAA95828.1 | AP002069 | Oryza sativa          | ESTs D47168(S12332),D46350(S10967) correspond to a region of the predicted gene. Similar to Prunus armeniaca ethylene-forming-enzyme-like dioxygenase. (U97530). |
| AAD56580.1 | AF184273 | Daucus carota         | leucoanthocyanidin dioxygenase 1. LDOX1. 2-oxoglutarate dependent dioxygenase.   |
| AAB39995.1 | U82432   | Dianthus caryophyllus | anthocyanidin synthase. allele: S; 2-oxoglutarat-dependent dioxygenase.  |
| AAD56581.1 | AF184274 | Daucus carota         | leucoanthocyanidin dioxygenase 2. LDOX. 2-oxoglutarate dependent dioxygenase.  |

SEQ ID NO: 691

|            |          |   |
|------------|----------|---|
| AAA34002.1 | M67449   | Glycine max<br>protein kinase. PK6.   |
| CAC09580.1 | AJ298992 | Fagus sylvatica<br>Absciscic acid (ABA) and calcium induced protein kinase. protein kinase (PK). pk1.   |
| AAK11734.1 | AY027437 | Arachis hypogaea<br>serine/threonine/tyrosine kinase.   |
| BAB16918.1 | AP002863 | Oryza sativa<br>putative protein kinase. P0005A05.22.   |
| CAA06334.1 | AJ005077 | Lycopersicon esculentum<br>protein kinase. TCTR2 protein. TCTR2.  |
| AAD10057.1 | AF110519 | Lycopersicon esculentum<br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit<br>ripening inducible CTR1-like protein kinase; TCTR1v.  |
| AAD10056.1 | AF110518 | Lycopersicon esculentum<br>ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit<br>ripening inducible CTR1-like protein kinase; TCTR1.   |
| CAA73722.1 | Y13273   | Lycopersicon esculentum<br>putative protein kinase.   |
| AAD46406.1 | AF096250 | Lycopersicon esculentum<br>ethylene-responsive protein kinase TCTR1. ER50. serine/threonine kinase; similar to<br>Arabidopsis thaliana negative regulator of the ethylene response pathway encoded by<br>GenBank Accession Number L08789. |
| AAG31141.1 | AF305911 | Oryza sativa<br>EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.   |
| AAG31142.1 | AF305912 | Hordeum vulgare<br>EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.  |
| AAK30005.1 | AY029067 | Rosa hybrid cultivar<br>CTR2 protein kinase.  |
| AAK21965.1 | AY028699 | Brassica napus<br>receptor protein kinase PERK1.  |
| BAB16871.1 | AP002537 | Oryza sativa<br>putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs<br>C22608(R3192),D25110(R3192).   |
| AAF91323.1 | AF244889 | Glycine max<br>receptor-like protein kinase 2. RLK2. GmRLK2.  |
| AAF91324.1 | AF244890 | Glycine max<br>receptor-like protein kinase 3. RLK3. GmRLK3.  |
| AAK00425.1 | AC069324 | Oryza sativa<br>Putative protein kinase. OSJNBa0071K19.11.  |
| BAB39409.1 | AP002901 | Oryza sativa<br>putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| AAF91322.1 | AF244888 | Glycine max<br>receptor-like protein kinase 1. RLK1. GmRLK1.  |

|            |          |                      |  |
|------------|----------|----------------------|--|
| BAB18292.1 | AP002860 | Oryza sativa         | putative receptor-like protein kinase. P0409B08.19.  |
| AAK16409.1 | AF320086 | Zea mays             | serine threonine kinase 1. stk1. expressed in mature tassel.   |
| BAA06538.1 | D31737   | Nicotiana tabacum    | protein-serine/threonine kinase.   |
| BAA21132.1 | D88193   | Brassica rapa        | S-receptor kinase. SRK9 (B.c).   |
| BAA06285.1 | D30049   | Brassica rapa        | S-receptor kinase SRK9.  |
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| CAB54520.1 | AJ238845 | Brassica napus       | putative role in cell cycle control. MAP3K epsilon 1 protein kinase. MAP3Ke1.  |
| CAA97692.1 | Z73295   | Catharanthus roseus  | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis. |
| CAA08995.1 | AJ010091 | Brassica napus       | MAP3K alpha 1 protein kinase. MAP3K alpha 1.   |
| BAB40094.1 | AP003210 | Oryza sativa         | putative receptor protein kinase. OSJNBa0010K01.7.   |
| CAB51834.1 | 00069    | Oryza sativa         | 11332.5. contains eukaryotic protein kinase domain PF.   |
| BAB40021.1 | AP003021 | Oryza sativa         | putative wall-associated kinase 2. P0503E05.25. contains EST C24950(S16264).   |
| AAF43496.1 | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.                        |
| AAK11674.1 | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.   |
| AAA62232.1 | U00443   | Brassica napus       | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| BAB40010.1 | AP003021 | Oryza sativa         | putative wall-associated kinase 2. P0503E05.12.  |
| AAC36318.1 | AF053127 | Malus x domestica    | leucine-rich receptor-like protein kinase. LRPKml.   |
| AAC23542.1 | U20948   | Ipomoea trifida      | receptor protein kinase. IRK1.   |
| BAA23676.1 | AB000970 | Brassica rapa        | receptor kinase 1. BcRK1.  |
| BAB18321.1 | AP002865 | Oryza sativa         | putative receptor protein kinase. P0034C11.11.   |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| BAA92836.1     | AB032473 | Brassica oleracea       | S18 S-locus receptor kinase. SRK18.   |
| AAA33915.1     | L27821   | Oryza sativa            | receptor type serine/threonine kinase. protein kinase.  |
| BAB40081.1     | AP003074 | Oryza sativa            | putative receptor protein kinase. OSJNBa0004G10.30.   |
| AAD38286.1     | AC007789 | Oryza sativa            | putative protein kinase. OSJNBa0049B20.13.  |
| BAA92837.1     | AB032474 | Brassica oleracea       | S60 S-locus receptor kinase. SRK60.   |
| SEQ ID NO: 692 |          |                         |   |
| AAK16172.1     | AC079887 | Oryza sativa            | putative glucosyltransferase. OSJNBa0040E01.14.   |
| AAF17077.1     | AF199453 | Sorghum bicolor         | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.   |
| AAF61647.1     | AF190634 | Nicotiana tabacum       | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAA89009.1     | AB027455 | Petunia x hybrida       | anthocyanin 5-O-glucosyltransferase. PH1.   |
| AAF98390.1     | AF287143 | Brassica napus          | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT. |
| AAD21086.1     | AF127218 | Forsythia x intermedia  | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.   |
| BAA93039.1     | AB033758 | Citrus unshiu           | limonoid UDP-glucosyltransferase. LGTase.   |
| BAA83484.1     | AB031274 | Scutellaria baicalensis | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.   |
| BAA12737.1     | D85186   | Gentiana triflora       | UDP-glucose:flavonoid-3-glucosyltransferase.  |
| CAA54612.1     | X77462   | Manihot esculenta       | UTP-glucose glucosyltransferase. CGT5.  |
| BAA19155.1     | AB000623 | Nicotiana tabacum       | glucosyl transferase. JIGT.   |
| BAA89008.1     | AB027454 | Petunia x hybrida       | anthocyanidin 3-O-glucosyltransferase. PGT8.  |
| BAB17060.1     | AP002523 | Oryza sativa            | putative glucosyl transferase. P0013F10.6.  |
| BAA90787.1     | AB038248 | Ipomoea batatas         | UDP glucose: flavonoid 3-O-glucosyltransferase. uf3gt.  |

|            |          |                                 |   |
|------------|----------|---------------------------------|---|
| AAK16178.1 | AC079887 | Oryza sativa                    | putative glucosyltransferase. OSJNBa0040E01.5.  |
| AAK16175.1 | AC079887 | Oryza sativa                    | putative glucosyltransferase. OSJNBa0040E01.15.   |
| AAB36653.1 | U32644   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS5a.   |
| AAB36652.1 | U32643   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS10a.  |
| AAK28303.1 | AF346431 | Nicotiana tabacum               | phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.  |
| AAK28304.1 | AF346432 | Nicotiana tabacum               | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.  |
| BAB17061.1 | AP002523 | Oryza sativa                    | putative glucosyl transferase. P0013F10.7. contains EST C73149(E2992).  |
| CAA30761.1 | X07940   | Zea mays                        | UDPglucose flavonoid glycosyl-transferase. Bz-McC.  |
| AAK16410.1 | AF320086 | Zea mays                        | UDPG-flavonoid 3-O-glucosyl transferase. bz.  |
| CAA31855.1 | X13500   | Zea mays                        | UDPglucose:flavonol 3-O-glucosyltransferase.  |
| BAB17059.1 | AP002523 | Oryza sativa                    | putative glucosyl transferase. P0013F10.5.  |
| BAA36421.1 | AB013596 | Perilla frutescens              | UDP-glucose:anthocynin 5-O-glucosyltransferase. PF3R4.  |
| AAB86473.1 | AF028237 | Ipomoea purpurea                | UDP glucose: flavonoid 3-O-glucosyltransferase. UF3GT-FL1.  |
| CAA30760.1 | X07937   | Zea mays                        | UDPglucose flavonoid glycosyl transferase. Bz-W22.  |
| BAB41021.1 | AB047094 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT1.   |
| BAB41019.1 | AB047092 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT1.   |
| BAB41025.1 | AB047098 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.   |
| AAD55985.1 | AF165148 | Petunia x hybrida               | catalyzes the penultimate step of flavonol glucosylgalactoside biosynthesis from UDP-galactose and flavonol aglycones in petunia pollen. UDP-galactose:flavonol 3-O-galactosyltransferase. F3galtase. |
| BAB41018.1 | AB047091 | Vitis labrusca x Vitis vinifera | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT2. The gene originated in one of the parents V. vinifera cv. Centennial.   |



BAB41017.1 AB047090 *Vitis labrusca* x *Vitis vinifera*  
 UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT1. The gene was derived from one of the parents *V. labruscana* cv. Ishiharawase.

AAK16181.1 AC079887 *Oryza sativa*  
 putative glucosyltransferase. OSJNBa0040E01.16.

BAB41023.1 AB047096 *Vitis vinifera*  
 UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.

CAA59450.1 X85138 *Lycopersicon esculentum*  
 twil. homologous to glucosyltransferases.

CAA54614.1 X77464 *Manihot esculenta*  
 UTP-glucose glucosyltransferase. CGT7.

AAB81682.1 AF000371 *Vitis vinifera*  
 UDP glucose:flavonoid 3-o-glucosyltransferase.

AAB81683.1 AF000372 *Vitis vinifera*  
 UDP glucose:flavonoid 3-o-glucosyltransferase.

BAB41024.1 AB047097 *Vitis vinifera*  
 UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT2.

CAA54558.1 X77369 *Solanum melongena*  
 glycosyl transferase. GT.

BAB41026.1 AB047099 *Vitis vinifera*  
 UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT2.

BAB41020.1 AB047093 *Vitis vinifera*  
 UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT2.

BAB41022.1 AB047095 *Vitis vinifera*  
 UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT2.

BAA19659.1 AB002818 *Perilla frutescens*  
 flavonoid 3-O-glucosyltransferase. UDP glucose.

AAG25643.1 AF303396 *Phaseolus vulgaris*  
 UDP-glucosyltransferase HRA25. putative; defense associated.

SEQ ID NO: 694

BAB40923.1 AB059401 *Oryza sativa*  
 putative selenium binding protein. Os SBP.

SEQ ID NO: 695

AAD26942.1 AF119050 *Datisca glomerata*  
 zinc-finger protein 1. zfp1. DgZFP1.

BAA05079.1 D26086 *Petunia x hybrida*  
 zinc-finger protein.

AAB39638.1 U68763 *Glycine max*  
 putative transcription factor. SCOF-1. scof-1. zinc-finger protein.

CAB77055.1 Y18788 *Medicago sativa*  
 putative TFIIA (or kruppel)-like zinc finger protein.

AAC06243.1 AF053077 *Nicotiana tabacum*  
 transcription factor. osmotic stress-induced zinc-finger protein. zfp.

|   |          |                   |
|---|----------|-------------------|
| BAA05077.1  | D26084   | Petunia x hybrida |
| zinc-finger DNA binding protein.                            |          |                   |
| BAA05076.1  | D26083   | Petunia x hybrida |
| zinc-finger DNA binding protein.                            |          |                   |
| AAB53260.1  | U76554   | Brassica rapa     |
| transcription factor. zinc-finger protein-1. BR42.          |          |                   |
| BAA05078.1  | D26085   | Petunia x hybrida |
| zinc-finger DNA binding protein.                            |          |                   |
| AAB53261.1  | U76555   | Brassica rapa     |
| zinc-finger protein BcZFP1. BcZFP1(3-2z).                   |          |                   |
| AAK01713.1  | AF332876 | Oryza sativa      |
| zinc finger transcription factor ZF1.                       |          |                   |
| BAA21920.1  | AB006598 | Petunia x hybrida |
| ZPT2-11. C2H2 zinc finger protein, 2finger.                 |          |                   |
| BAA21922.1  | AB006600 | Petunia x hybrida |
| ZPT2-13. C2H2 zinc finger protein, 2finger.                 |          |                   |
| BAA19112.1  | AB000453 | Petunia x hybrida |
| PETHy;ZPT3-1. Cys(2) His(2) zinc finger protein, 3 fingers. |          |                   |
| BAA21927.1  | AB006605 | Petunia x hybrida |
| ZPT3-3. C2H2 zinc finger protein, 3 finger.                 |          |                   |
| BAA96071.1  | AB035133 | Petunia x hybrida |
| C2H2 zinc-finger protein ZPT3-3. ZPT3-3.                    |          |                   |
| BAA96070.1  | AB035132 | Petunia x hybrida |
| C2H2 zinc-finger protein ZPT2-10. PETHy;ZPT2-10.            |          |                   |
| BAA21919.1  | AB006597 | Petunia x hybrida |
| ZPT2-10. C2H2 zinc finger protein, 2 finger.                |          |                   |
| BAA19114.1  | AB000455 | Petunia x hybrida |
| PETHy;ZPT4-1. Cys(2) His(2) zinc finger protein, 4 fingers. |          |                   |
| BAA21921.1  | AB006599 | Petunia x hybrida |
| ZPT2-12. C2H2 zinc finger protein, 2 finger.                |          |                   |
| BAA21928.1  | AB006606 | Petunia x hybrida |
| ZPT4-4. C2H2 zinc finger protein, 4 finger.                 |          |                   |
| CAA60828.1  | X87374   | Pisum sativum     |
| putative zinc finger protein.                               |          |                   |
| BAA21925.1  | AB006603 | Petunia x hybrida |
| ZPT2-8. C2H2 zinc finger protein, 2 finger.                 |          |                   |
| BAA19111.1  | AB000452 | Petunia x hybrida |
| PETHy;ZPT2-6. Cys(2) His(2) zinc finger protein, 2 fingers. |          |                   |
| BAA19926.1  | AB000456 | Petunia x hybrida |
| PETHy; ZPT4-2. C2H2 zinc finger protein, 4 finger.          |          |                   |
| BAA21924.1  | AB006602 | Petunia x hybrida |
| ZPT2-7. C2H2 zinc finger protein, 2finger.                  |          |                   |

|  |   |                           |
|--|---|---------------------------|
| BAA21923.1                                       | AB006601  | Petunia x hybrida         |
| ZPT2-14.   | C2H2 zinc finger protein, 2 finger.   |                           |
| BAA21926.1                                       | AB006604  | Petunia x hybrida         |
| ZPT2-9.  | C2H2 zinc finger protein, 2 finger.   |                           |
| BAA19110.1                                       | AB000451  | Petunia x hybrida         |
| PETHy;ZPT2-5.                                    | Cys(2) His(2) zinc finger protein, 2 fingers.   |                           |
| BAA19113.1                                       | AB000454  | Petunia x hybrida         |
| PETHy;ZPT3-2.                                    | Cys(2) His(2) zinc finger protein, 3 fingers.   |                           |
| SEQ ID NO: 702                                   |   |                           |
| AAK29419.1                                       | AF348319  | Zea mays                  |
| TERMINAL EAR1.                                   | tel. RNA-binding protein; 3 putative RRM motifs; similar to Schizosaccharomyces pombe Mei2 and Arabidopsis AML1; tel cDNA sequence is presented in GenBank Accession Number AF047852. |                           |
| CAA57551.1                                       | X82030  | Phaseolus vulgaris        |
| chloroplast RNA binding protein.                 | RNP1.   |                           |
| CAA66479.1                                       | X97905  | Vicia faba                |
| RNA- or ssDNA-binding protein.                   |   |                           |
| CAC01237.1                                       | AJ292767  | Nicotiana plumbaginifolia |
| nuclear RNA binding protein.                     | RNA Binding Protein 45. rbp45.  |                           |
| AAF66823.1                                       | AF190655  | Nicotiana tabacum         |
| poly(A)-binding protein.                         | PABP.   |                           |
| AAK30205.1                                       | AF349964  | Daucus carota             |
| poly(A)-binding protein.                         | Translin1P.   |                           |
| AAG59664.1                                       | AC084319  | Oryza sativa              |
| putative RNA binding protein.                    | OSJNBa0004B24.1.  |                           |
| AAB38974.1                                       | U81318  | Triticum aestivum         |
| poly(A)-binding protein.                         | wheatpab.   |                           |
| CAA81127.1                                       | Z26042  | Anemia phyllitidis        |
| Binding to the poly(A)-tail of eukaryotic mRNAs. | poly(A)-mRNA binding protein.   |                           |
| AAA79045.1                                       | U34742  | Spinacia oleracea         |
| 24 kDa RNA binding protein.                      |   |                           |
| AAC39368.1                                       | AF043297  | Chlamydomonas reinhardtii |
| poly(A) binding protein                          | RB47. PABP.   |                           |
| AAF63202.1                                       | AF240679  | Cucumis sativus           |
| poly(A)-binding protein.                         |   |                           |
| CAA11894.1                                       | AJ224325  | Hordeum vulgare           |
| nucleic acid-binding protein.                    | cp33Hv.   |                           |
| AAF66825.1                                       | AF190657  | Nicotiana tabacum         |
| poly(A)-binding protein.                         | PABP.   |                           |
| CAA06469.1                                       | AJ005286  | Hordeum vulgare           |
| nucleic acid-binding protein.                    | cp31AHv protein.  |                           |
| SEQ ID NO: 709                                   |   |                           |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAB36546.1     | U77940   | Phaseolus vulgaris            | polyubiquitin. expression is regulated by heavy metal stress, UV, virus infection, heat shock and wounding treatment, this suggests that ubiquitin plays a role in plant defense. |
| SEQ ID NO: 714 |          |                               |   |
| CAA61946.1     | X89828   | Pisum sativum                 | fructose-1,6-bisphosphate aldolase.   |
| AAB61592.1     | AF003124 | Mesembryanthemum crystallinum | fructose-biphosphate aldolase.  |
| CAB77243.2     | AJ133146 | Persea americana              | glycolytic enzyme, sixth step in glycolysis. fructose-bisphosphate aldolase. alf.   |
| BAA08845.1     | D50307   | Oryza sativa                  | aldolase C-1.   |
| BAA08830.1     | D50301   | Oryza sativa                  | aldolase C-1.   |
| CAA46649.1     | X65742   | Spinacia oleracea             | fructose-bisphosphate aldolase.   |
| BAA02729.1     | D13512   | Oryza sativa                  | cytoplasmic aldolase.   |
| AAG21429.1     | AF308587 | Fragaria x ananassa           | cytosolic aldolase. SCA1.   |
| CAA37290.1     | X53130   | Oryza sativa                  | fructose-diphosphate aldolase (AA 1-358).   |
| CAA31366.1     | X12872   | Zea mays                      | fructose bisphosphate aldolase.   |
| AAA33435.1     | M16220   | Zea mays                      | aldolase.   |
| CAA61947.1     | X89829   | Pisum sativum                 | fructose-1,6-bisphosphate aldolase.   |
| CAA06308.1     | AJ005041 | Cicer arietinum               | cytosolic fructose-1,6-bisphosphate aldolase.   |
| AAK19324.1     | AF329673 | Dunaliella salina             | fructose-bisphosphate aldolase isoenzyme 1. salt-induced.   |
| CAA71408.1     | Y10380   | Solanum tuberosum             | homologous to plastidic aldolases.  |
| BAA77603.1     | AB027002 | Nicotiana paniculata          | plastidic aldolase.   |
| BAA77604.1     | AB027001 | Nicotiana paniculata          | plastidic aldolase NPALDP1. NpAldP1.  |
| AAA33642.1     | M97476   | Pisum sativum                 | aldolase.   |
| AAA33643.1     | M97477   | Pisum sativum                 | aldolase.   |

|                |          |                                       |   |
|----------------|----------|---------------------------------------|---|
| BAA02730.1     | D13513   | Oryza sativa                          | chloroplastic aldolase.   |
| AAF74220.1     | AF216582 | Avena sativa                          | fructose 1,6-bisphosphate aldolase precursor.   |
| CAA47293.1     | X66814   | Spinacia oleracea                     | fructose-bisphosphate aldolase.   |
| AAK19325.1     | AF329674 | Dunaliella salina                     | fructose-bisphosphate aldolase isoenzyme 2. salt-induced.   |
| CAB46520.1     | AJ243524 | Phleum pratense                       | sixth step in glycolysis. putative fructose-bisphosphate aldolase.  |
| CAA09669.1     | AJ011516 | Scherffelia dubia                     | fructose-bisphosphate aldolase.   |
| CAA49590.1     | X69969   | Chlamydomonas reinhardtii             | fructose-bisphosphate aldolase. ALDCHL.   |
| AAC60574.1     | S72951   | Chloroplast Chlamydomonas reinhardtii | fructosediphosphate aldolase. fructosediphosphate aldolase. This sequence comes from Fig. 4.  |
| AAB70542.1     | AF017362 | Oryza sativa                          | aldolase.   |
| BAA76430.1     | AB025002 | Cicer arietinum                       | fructose-bisphosphate aldolase.   |
| AAD20818.1     | AF107590 | Dendrobium grex Madame Thong-In       | putative fructose-bisphosphate aldolase. otg11.   |
| BAA11395.1     | D78500   | Brassica rapa                         | putative aldolase. Sequence homologous to cytoplasmic aldolase of rice (D13512).  |
| BAA78593.1     | AU066535 | Chlamydomonas sp. HS-5                | fructose-bisphosphate aldolase precursor. NaCl inducible.   |
| CAC34412.1     | Y18576   | Flaveria trinervia                    | fructose-bisphosphate aldolase. alf.  |
| SEQ ID NO: 715 |          |                                       |   |
| AAK07429.1     | AF321287 | Musa acuminata                        | beta-glucosidase.   |
| AAC69619.1     | AF072736 | Pinus contorta                        | beta-glucosidase.   |
| AAF04007.1     | AF163097 | Dalbergia cochinchinensis             | beta-fucosidase beta-glucosidase. dalcocinin 8'-O-beta-glucoside beta-glucosidase precursor. BGLU1. rotenoid beta-glucosidase.  |
| AAA93032.1     | U50201   | Prunus serotina                       | hydrolysis of the cyanogenic glucoside (R)-prunasin. prunasin hydrolase precursor. located in protein bodies of Prunus seeds; encodes 8 putative N-glycosylation sites (N-X-S/T); encodes NEP and ITENG motifs characteristic of the BGA family of beta-glucosidases. |
| BAA78708.1     | AB003089 | Polygonum tinctorium                  | beta-glucosidase.   |

|            |          |                             |  |
|------------|----------|-----------------------------|--|
| AAA91166.1 | U39228   | <i>Prunus avium</i>         | beta-glucosidase.  |
| BAA11831.1 | D83177   | <i>Costus speciosus</i>     | saponin metabolite. furostanol glycoside 26-O-beta-glucosidase (F26G). functional expression in <i>E. coli</i> ; one of the F26G isozymes.   |
| AAF34650.1 | AF221526 | <i>Prunus serotina</i>      | hydrolysis of the cyanogenic glucoside (R)-prunasin. prunasin hydrolase isoform PHA precursor. beta-glucosidase; contains 6 potential N-glycosylation sites (N-X-S/T); glycosyl hydrolase family 1 member. |
| AAG25897.1 | AF170087 | <i>Cucurbita pepo</i>       | silverleaf whitefly-induced protein 3. SLW3. similar to beta-glucosidase.  |
| AAG00614.1 | AF293849 | <i>Secale cereale</i>       | beta-glucosidase.  |
| AAB22162.1 | S35175   | <i>Manihot esculenta</i>    | linamarase. linamarase. beta-glucosidase; Method: conceptual translation with partial peptide sequencing; This sequence comes from Fig. 4.   |
| AAF03675.1 | AF149311 | <i>Rauvolfia serpentina</i> | hydrolyses the glucoalkaloid raucaffricine. raucaffricine-O-beta-D-glucosidase. beta glucosidase; RG; part of the Ajmaline biosynthesis pathway; belongs to family 1 of the glucosyl hydrolases.           |
| AAA87339.1 | L41869   | <i>Hordeum vulgare</i>      | beta-glucosidase. BGQ60. expression specific to starchy endosperm of seed.   |
| AAC49177.1 | U33817   | <i>Sorghum bicolor</i>      | beta-glucosidase, catalyzes the hydrolysis of the cyanogenic beta-glucoside dhurrin. dhurrinase.   |
| CAA64442.1 | X94986   | <i>Manihot esculenta</i>    | beta glucosidase. bglA.  |
| AAD02839.1 | AF082991 | <i>Avena sativa</i>         | beta-D-glucosidase beta subunit precursor. P60b. avenacosidase.  |
| AAD09850.1 | U44087   | <i>Zea mays</i>             | beta-D-glucosidase precursor. glu2. product subunit structure: autodimer of 58.4 kDa monomers Allele: glu2-B73.  |
| AAD10503.1 | U33816   | <i>Zea mays</i>             | functions in defense of young plant parts against pests via the production of hydroxamic acids from hydroxamic acid glucosides. beta-D-glucosidase.  |
| AAB03266.1 | U44773   | <i>Zea mays</i>             | beta-D-glucosidase. glu1. autodimer of 58.4 kD monomers.   |
| AAA65946.1 | U25157   | <i>Zea mays</i>             | functions in defense of young plant parts against pests via the production of hydroxamic acids from hydroxamic acid glucosides. beta-D-glucosidase.  |
| CAA52293.1 | X74217   | <i>Zea mays</i>             | beta-glucosidase. p60.1.   |

|                |          |   |
|----------------|----------|---|
| AAF28800.1     | AF112888 | <i>Catharanthus roseus</i><br>plays a role in secondary metabolism by hydrolyzing strictosidine to cathenamine during indole alkaloid biosynthesis. strictosidine beta-glucosidase. localized in the endoplasmic reticulum.   |
| CAA40057.1     | X56733   | <i>Trifolium repens</i><br>beta-glucosidase. Li.  |
| CAA40058.1     | X56734   | <i>Trifolium repens</i><br>beta-glucosidase. non-cyanogenic.  |
| CAA55196.1     | X78433   | <i>Avena sativa</i><br>beta-D-glucosidase.  |
| CAA79989.2     | Z21977   | <i>Brassica napus</i><br>beta thioglucosidase. myrosinase, thioglucoside glucohydrolase. Myr1.Bn1.  |
| AAB71381.1     | U95298   | <i>Manihot esculenta</i><br>linamarase. pLIN-GEN. beta-glucosidase.   |
| CAA57913.1     | X82577   | <i>Brassica napus</i><br>beta-glucosidase. bgl.   |
| AAB38784.1     | U72154   | <i>Brassica nigra</i><br>beta-glucosidase. psr3.1. PSR3.1; phosphate-starvation responsive enzyme.  |
| AAF34651.1     | AF221527 | <i>Prunus serotina</i><br>putative prunasin hydrolase precursor. beta-glucosidase; glycosyl hydrolase family 1 member.  |
| AAA84906.1     | U28047   | <i>Oryza sativa</i><br>catalyzes the release of either giberellin or cyanogenic substances from their glucoconjugates. beta glucosidase. beta-D-glucoside glucohydrolase; dimer of 60 kDa monomers; localized in the plastid. |
| CAC08209.1     | AJ005950 | <i>Cicer arietinum</i><br>beta-glucosidase.   |
| SEQ ID NO: 718 |          |   |
| AAF61647.1     | AF190634 | <i>Nicotiana tabacum</i><br>UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAA89009.1     | AB027455 | <i>Petunia x hybrida</i><br>anthocyanin 5-O-glucosyltransferase. PH1.   |
| BAA93039.1     | AB033758 | <i>Citrus unshiu</i><br>limonoid UDP-glucosyltransferase. LGTase.   |
| BAA36423.1     | AB013598 | <i>Verbena x hybrida</i><br>UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| BAA36421.1     | AB013596 | <i>Perilla frutescens</i><br>UDP-glucose:anthocynin 5-O-glucosyltransferase. PF3R4.   |
| AAF98390.1     | AF287143 | <i>Brassica napus</i><br>catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT.              |
| BAA36422.1     | AB013597 | <i>Perilla frutescens</i><br>UDP-glucose:anthocyanin 5-O-glucosyltransferase homologue. PF3R6.  |

|            |          |  |  |
|------------|----------|--|--|
| BAB07962.1 | AP002524 | <i>Oryza sativa</i>                    | putative anthocyanin 5-O-glucosyltransferase. P0406H10.16. contains ESTs AU067881(C10481),AU067882(C10481).                          |
| AAD21086.1 | AF127218 | <i>Forsythia x intermedia</i>          | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.                                  |
| AAK16178.1 | AC079887 | <i>Oryza sativa</i>                    | putative glucosyltransferase. OSJNBa0040E01.5.   |
| BAA12737.1 | D85186   | <i>Gentiana triflora</i>               | UDP-glucose:flavonoid-3-glucosyltransferase.   |
| AAK16181.1 | AC079887 | <i>Oryza sativa</i>                    | putative glucosyltransferase. OSJNBa0040E01.16.  |
| AAK16175.1 | AC079887 | <i>Oryza sativa</i>                    | putative glucosyltransferase. OSJNBa0040E01.15.  |
| AAK16172.1 | AC079887 | <i>Oryza sativa</i>                    | putative glucosyltransferase. OSJNBa0040E01.14.  |
| CAA59450.1 | X85138   | <i>Lycopersicon esculentum</i>         | twi1. homologous to glucosyltransferases.  |
| AAF17077.1 | AF199453 | <i>Sorghum bicolor</i>                 | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.  |
| BAA89008.1 | AB027454 | <i>Petunia x hybrida</i>               | anthocyanidin 3-O-glucosyltransferase. PGT8.   |
| AAB81683.1 | AF000372 | <i>Vitis vinifera</i>                  | UDP glucose:flavonoid 3-o-glucosyltransferase.   |
| BAB41017.1 | AB047090 | <i>Vitis labrusca x Vitis vinifera</i> | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT1. The gene was derived from one of the parents V. labruscana cv. Ishiharawase. |
| AAB81682.1 | AF000371 | <i>Vitis vinifera</i>                  | UDP glucose:flavonoid 3-o-glucosyltransferase.   |
| BAB41022.1 | AB047095 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT2.  |
| BAB41020.1 | AB047093 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT2.  |
| BAB41021.1 | AB047094 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT1.  |
| BAB41019.1 | AB047092 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT1.  |
| BAA83484.1 | AB031274 | <i>Scutellaria baicalensis</i>         | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.  |
| BAB41025.1 | AB047098 | <i>Vitis vinifera</i>                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.  |



|                |          |                                 |   |
|----------------|----------|---------------------------------|---|
| BAB41023.1     | AB047096 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.   |
| BAA90787.1     | AB038248 | Ipomoea batatas                 | UDP glucose: flavonoid 3-O-glucosyltransferase. uf3gt.  |
| BAA19659.1     | AB002818 | Perilla frutescens              | flavonoid 3-O-glucosyltransferase. UDP glucose.   |
| BAB41018.1     | AB047091 | Vitis labrusca x Vitis vinifera | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT2. The gene originated in one of the parents V. vinifera cv. Centennial.             |
| AAB36652.1     | U32643   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS10a.  |
| AAK28304.1     | AF346432 | Nicotiana tabacum               | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.  |
| BAB41024.1     | AB047097 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT2.   |
| BAB41026.1     | AB047099 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT2.   |
| CAA31855.1     | X13500   | Zea mays                        | UDPglucose:flavonol 3-O-glucosyltransferase.  |
| AAB86473.1     | AF028237 | Ipomoea purpurea                | UDP glucose: flavonoid 3-O-glucosyltransferase. UF3GT-FL1.  |
| CAA54614.1     | X77464   | Manihot esculenta               | UTP-glucose glucosyltransferase. CGT7.  |
| SEQ ID NO: 719 |          |                                 |   |
| AAC24195.1     | AF020425 | Nicotiana tabacum               | calmodulin binding protein. glutamate decarboxylase isozyme 1. NtGAD1. calcium-calmodulin-dependent enzyme.                               |
| AAK18620.1     | AF352732 | Nicotiana tabacum               | converts glutamate to gamma-aminobutyric acid. glutamate decarboxylase isozyme 3. GAD; GAD3; NtGAD3; calcium/calmodulin-dependent enzyme. |
| AAB40608.1     | U54774   | Nicotiana tabacum               | glutamate decarboxylase. NtGAD1. calmodulin regulated enzyme; calmodulin-binding protein.   |
| AAA33710.1     | L16977   | Petunia x hybrida               | glutamate decarboxylase. gad.   |
| AAA33709.1     | L16797   | Petunia x hybrida               | glutamate decarboxylase. gad.   |
| AAC39483.1     | AF020424 | Nicotiana tabacum               | glutamate decarboxylase isozyme 2. NtGAD2. calcium-calmodulin-dependent enzyme.   |
| BAB32870.1     | AB056062 | Oryza sativa                    | glutamate decarboxylase. GAD.   |
| BAB32868.1     | AB056060 | Oryza sativa                    | glutamate decarboxylase. GAD.   |

|                |          |                                |   |
|----------------|----------|--------------------------------|---|
| CAA56812.1     | X80840   | <i>Lycopersicon esculentum</i> | homology to pyroxidal-5'-phosphate-dependant glutamate decarboxylases; putative start codon.  |
| BAB32871.1     | AB056063 | <i>Oryza sativa</i>            | glutamate decarboxylase. GAD.   |
| BAB32869.1     | AB056061 | <i>Oryza sativa</i>            | glutamate decarboxylase. GAD.   |
| CAA50719.1     | X71900   | <i>Lycopersicon esculentum</i> | histidine decarboxylase. hdc. pyridoxal 5'-phosphate dependant.   |
| SEQ ID NO: 720 |          |                                |   |
| BAA92713.1     | AP001389 | <i>Oryza sativa</i>            | ESTs AU033035(S1515),D39871(S1515) correspond to a region of the predicted gene. Similar to <i>Arabidopsis thaliana</i> chromosome II BAC F22D22 genomic sequence; putative glucan synthase (AC006223). |
| AAD25952.1     | AF085717 | <i>Gossypium hirsutum</i>      | putative callose synthase catalytic subunit. CFL1. cotton FKS1-like protein; similar to <i>Saccharomyces cerevisiae</i> beta-1,3-glucan synthase subunit FKS1.  |
| SEQ ID NO: 721 |          |                                |   |
| CAB55396.1     | AL117264 | <i>Oryza sativa</i>            | zwh12.1. similar to <i>Arabidopsis</i> putative UDP-galactase-4-epimerase (AC007060); Method: conceptual translation with partial peptide sequencing.   |
| CAA06339.1     | AJ005082 | <i>Cyamopsis tetragonoloba</i> | UDP-galactose 4-epimerase.  |
| AAA86532.1     | U31544   | <i>Pisum sativum</i>           | catabolism of galactose to glucose in Leloir pathway, and in galactose synthesis from glucose. UDP-galactose-4-epimerase. galE. galactowaldenase.   |
| CAA06338.1     | AJ005081 | <i>Cyamopsis tetragonoloba</i> | UDP-galactose 4-epimerase.  |
| BAB40967.1     | AB059568 | <i>Pisum sativum</i>           | biosynthesis of UDP-D-xylose. UDP-D-glucuronate carboxy-lyase. uxs1.  |
| SEQ ID NO: 726 |          |                                |   |
| BAA88198.1     | AP000837 | <i>Oryza sativa</i>            | Similar to human dimethylaniline monooxygenase (AC002376).  |
| BAA35120.1     | AB008845 | <i>Oryza sativa</i>            | NADH dependent Glutamate Synthase.  |
| AAB41904.1     | L37606   | <i>Medicago sativa</i>         | NADH-dependent glutamate synthase.  |
| BAA88195.1     | AP000837 | <i>Oryza sativa</i>            | Similar to human dimethylaniline monooxygenase (AC002376).  |

AAB46617.1 L01660 *Medicago sativa*  
the 3 cysteine residues in this region (amino acid residues 1246-1257) may serve to coordinate the [3Fe-4S] cluster that is probably found in this protein. See Knaff et al. (1991) J. Biol. Chem.266:150. NADH-glutamate synthase. NADH-GOGAT; activity increases dramatically during nodule development; the 3 cysteine residues in this region may serve to coordinate the [3Fe-4S] cluster that is probably found in this protein. See Knaff et al. (1991) J. Biol. Chem.266:15080-15084.

## SEQ ID NO: 728

AAG43550.1 AF211532 *Nicotiana tabacum*  
Avr9/Cf-9 rapidly elicited protein 132. ACRE132. similar to RING finger proteins.

BAA96875.1 AB045121 *Oryza sativa*  
RING finger 1. RRF1.

AAK00436.1 AC060755 *Oryza sativa*  
putative zinc finger protein. OSJNBa0003O19.23.

BAA78746.1 AB023482 *Oryza sativa*  
Similar to *Arabidopsis thaliana* RING-H2 finger protein RHX1a mRNA, partial cds.(AF079184).

CAA74911.1 Y14573 *Hordeum vulgare*  
ring finger protein. putative.

AAG46117.1 AC073166 *Oryza sativa*  
putative ring finger protein. OSJNBb0064P21.7.

BAA85438.1 AP000616 *Oryza sativa*  
similar to RING-H2 finger protein RHA1a (AF078683).

BAA77204.1 AB026262 *Cicer arietinum*  
ring finger protein.

## SEQ ID NO: 730

AAA86424.1 U44386 *Lycopersicon esculentum*  
heat shock protein. TFHS1. similar to protein encoded by the arg2 gene in mung bean, encoded by Genbank Accession Number D14411.

## SEQ ID NO: 731

CAA06756.1 AJ005899 *Nicotiana tabacum*  
G subunit. G subunit of Vacuolar-type H<sup>+</sup>-ATPase. vag1.

CAA06757.1 AJ005900 *Nicotiana tabacum*  
Subunit G of vacuolar-type H<sup>+</sup>-ATPase. vag2. vag2.

AAD56039.1 AF184068 *Citrus limon*  
vacuolar membrane ATPase subunit G. LVMA10. V-ATPase VMA10.

## SEQ ID NO: 737

AAF69008.1 AF257779 *Oryza sativa*  
stress-inducible protein. OsSII.

## SEQ ID NO: 739

CAC12996.1 AJ299398 *Medicago truncatula*  
putative auxin import. putative AUX1-like permease. lax2.

AAF21982.1 AF115543 Populus tremula x Populus tremuloides  
AUX1-like protein. PAX1.

AAG17171.1 AF190880 Populus tremula x Populus tremuloides  
putative AUX1-like permease. pax5.

CAC12995.1 AJ299397 Medicago truncatula  
putative auxin import. putative AUX1-like permease. lax1.

CAC12997.1 AJ299399 Medicago truncatula  
putative auxin import. putative AUX1-like permease. lax3.

CAB65535.1 AJ011794 Zea mays  
AUX1 protein. aux1.

SEQ ID NO: 740

CAA65269.1 X96406 Solanum tuberosum  
13-lipoxygenase.

AAB65767.1 U37840 Lycopersicon esculentum  
lipoxygenase. loxD. wound, systemin and methyl-jasmonate induced.

BAA03102.1 D14000 Oryza sativa  
lipoxygenase. lox2osPil.

AAC12951.1 U56406 Hordeum vulgare  
methyljasmonate-inducible lipoxygenase 2.

CAB94852.1 AJ404331 Prunus dulcis  
hydroperoxydation of polyunsaturated fatty acids. lipoxygenase. lox.

AAG21691.1 AY008278 Lycopersicon esculentum  
lipoxygenase. oxido-reductase.

AAA79186.1 U36339 Cucumis sativus  
lipoxygenase.

AAB67858.1 U60200 Solanum tuberosum  
lipoxygenase. POTLX-1. expressed during early tuberization.

CAA64765.1 X95512 Solanum tuberosum  
lipoxygenase.

CAB83038.1 AJ271161 Cucumis sativus  
oxygenase. lipoxygenase-9. lox9.

CAA58859.1 X84040 Nicotiana tabacum  
lipoxygenase. Lox1.

AAB67865.1 U60202 Solanum tuberosum  
lipoxygenase. POTLX-3. expressed in ABA-treated leaves.

CAA55724.1 X79107 Solanum tuberosum  
lipoxygenase. Lox1:St:1.

AAB67860.1 U60201 Solanum tuberosum  
lipoxygenase. POTLX-2. expressed during early tuberization.

CAB65460.1 Y18548 Solanum tuberosum  
lipoxygenase. lox1-St-2.

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAD04258.1 | AF039651 | Solanum tuberosum       | 5-lipoxygenase.  |
| AAB81594.1 | AF019613 | Solanum tuberosum       | lipoxygenase. plox1.   |
| AAA33986.1 | J02795   | Glycine max             | lipoxygenase-1.  |
| AAB67732.1 | U50075   | Glycine max             | lipoxygenase L-5. vlxB.  |
| CAA47717.1 | X67304   | Glycine max             | lipoxygenase.  |
| AAB81595.1 | AF019614 | Solanum tuberosum       | lipoxygenase. plox2.   |
| CAA64766.1 | X95513   | Solanum tuberosum       | lipoxygenase.  |
| AAB31252.1 | S73865   | Solanum tuberosum       | linoleate:oxygen oxidoreductase. linoleate:oxygen oxidoreductase, lipoxygenase, LOX. This sequence comes from Fig. 1; lipoxygenase; LOX. |
| AAA53184.1 | U09026   | Lycopersicon esculentum | lipoxygenase. loxA.  |
| AAB65766.1 | U37839   | Lycopersicon esculentum | lipoxygenase. loxC. expressed during ripening fruit.   |
| CAA65268.1 | X96405   | Solanum tuberosum       | 13-lipoxygenase.   |
| AAF15296.2 | AF204210 | Phaseolus vulgaris      | lipoxygenase. LOX4.  |
| CAA55319.1 | X78581   | Pisum sativum           | lipoxygenase. Lox1:Ps:3.   |
| AAA03728.1 | U04526   | Glycine max             | lipoxygenase.  |
| CAA34906.1 | X17061   | Pisum sativum           | lipoxygenase (AA 1-864).   |
| CAA55318.1 | X78580   | Pisum sativum           | lipoxygenase. Lox1:Ps:2.   |
| BAA03042.1 | D13949   | Glycine max             | lacking. lipoxygenase-2. lox2.   |
| AAA33987.1 | J03211   | Glycine max             | lipoxygenase (EC 1.13.11.12).  |
| AAB71759.1 | U84198   | Pisum sativum           | lipoxygenase. Lox1:Ps:1. expressed in root nodules.  |
| AAB41272.1 | U50081   | Glycine max             | lipoxygenase-3.  |
| CAA39604.1 | X56139   | Glycine max             | lipoxygenase. sc514.   |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAA96817.1     | U26457   | Glycine max                   | lipxygenase. vlxC.  |
| AAG42354.1     | AF234983 | Phaseolus vulgaris            | lipxygenase.  |
| AAG18376.1     | AF283894 | Zantedeschia aethiopica       | lipxygenase. lox2.  |
| AAD39093.1     | AF095895 | Oryza sativa                  | catalyzes the addition of molecular oxygen to fatty acid. lipxygenase. CM-LOX1. |
| CAC04380.1     | AJ293015 | Pisum sativum                 | lipxygenase. lox1:Ps:7.   |
| CAA45088.1     | X63525   | Phaseolus vulgaris            | lipxygenase. loxA.  |
| AAB18970.2     | U76687   | Phaseolus vulgaris            | lipxygenase. PvLOX2.  |
| AAC49159.1     | U36191   | Glycine max                   | linoleate:oxygen oxidoreductase. lipxygenase. lox7.                             |
| AAA03726.1     | U04785   | Glycine max                   | lipxygenase.  |
| CAA45086.1     | X63521   | Phaseolus vulgaris            | lipxygenase.  |
| SEQ ID NO: 741 |          |                               |   |
| AAG28436.1     | AF195029 | Glycine max                   | plasma membrane Ca <sup>2+</sup> -ATPase. SCA2.                                 |
| AAG28435.1     | AF195028 | Glycine max                   | plasma membrane Ca <sup>2+</sup> -ATPase. SCA1.                                 |
| CAA68234.1     | X99972   | Brassica oleracea             | calmodulin-stimulated calcium-ATPase.   |
| AAD31896.1     | AF145478 | Mesembryanthemum crystallinum | calcium ATPase.   |
| BAA90510.2     | AP001111 | Oryza sativa                  | rice EST AU030811, similar to rice Ca <sup>2+</sup> -ATPase (U82966).           |
| AAD11618.1     | AF050496 | Lycopersicon esculentum       | Ca <sup>2+</sup> -ATPase. LCA1B; alternative transcript.                        |
| AAD11617.1     | AF050495 | Lycopersicon esculentum       | Ca <sup>2+</sup> -ATPase. LCA1A; alternative transcript.                        |
| AAA34138.1     | M96324   | Lycopersicon esculentum       | The calcium ATPase is a calcium ion pump. Ca <sup>2+</sup> -ATPase. LCA1.       |
| CAA63790.1     | X93592   | Dunaliella bioculata          | P-type ATPase. cal. calcium pumping; CA1.                                       |
| AAF73985.1     | AF096871 | Zea mays                      | calcium pump. calcium ATPase. cap1.   |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAB58910.1 | U82966   | Oryza sativa                  | Ca <sup>2+</sup> -ATPase.   |
| AAD46188.1 | AF156691 | Nicotiana plumbaginifolia     | plasma membrane proton ATPase. pma9.  |
| AAB17186.1 | U72148   | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase. LHA4. plasma membrane proton pumping ATPase.  |
| CAA47275.1 | X66737   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> -ATPase. pma4.   |
| CAA54045.1 | X76535   | Solanum tuberosum             | H(+)-transporting ATPase. PHA2.   |
| AAD46186.1 | AF156679 | Nicotiana plumbaginifolia     | plasma membrane proton ATPase. pma6.  |
| CAA59800.1 | X85805   | Zea mays                      | H(+)-transporting ATPase. MHA-2.  |
| BAA06629.1 | D31843   | Oryza sativa                  | plasma membrane H <sup>+</sup> -ATPase. OSA2.   |
| AAB35314.2 | S79323   | Vicia faba                    | plasma membrane H(+)-ATPase precursor. plasma membrane H(+)-ATPase. This sequence comes from Fig. 1; conceptual translation presented here differs from translation in publication. |
| CAA59799.1 | X85804   | Phaseolus vulgaris            | H(+)-transporting ATPase. BHA-1.  |
| AAB41898.1 | U84891   | Mesembryanthemum crystallinum | plasma membrane proton pump. H <sup>+</sup> -transporting ATPase. PMA.  |
| AAA34094.1 | M80489   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> ATPase. pma1.  |
| AAB84202.2 | AF029256 | Kosteletzkyia virginica       | plasma membrane proton ATPase. ATP1.  |
| AAA34052.1 | M27888   | Nicotiana plumbaginifolia     | H <sup>+</sup> -translocating ATPase.   |
| AAA34173.1 | M60166   | Lycopersicon esculentum       | H <sup>+</sup> -ATPase. LHA1.   |
| AAA34098.1 | M80490   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> ATPase. pma3.  |
| AAF98344.1 | AF275745 | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase. LHA2. P-type ion pump.  |
| AAD55399.1 | AF179442 | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase isoform LHA2. LHA2.  |
| CAC29436.1 | AJ310524 | Vicia faba                    | P-type H <sup>+</sup> -ATPase. ha5. predominantly expressed in guard cells and flowers.   |
| BAA08134.1 | D45189   | Zostera marina                | plasma membrane H <sup>+</sup> -ATPase. zha1.   |

|                |          |                           |  |
|----------------|----------|---------------------------|--|
| BAA01058.1     | D10207   | Oryza sativa              | H-ATPase. OSA1.  |
| CAB69824.1     | AJ271439 | Prunus persica            | plasma membrane H <sup>+</sup> ATPase. PPA1.   |
| CAA54046.1     | X76536   | Solanum tuberosum         | H(+)-transporting ATPase. PHA1.  |
| AAB60276.1     | U09989   | Zea mays                  | H(+)-transporting ATPase. Mha1.  |
| CAB69823.1     | AJ271438 | Prunus persica            | plasma membrane H <sup>+</sup> ATPase. PPA2.   |
| AAK31799.1     | AY029190 | Lilium longiflorum        | plasma membrane H <sup>+</sup> ATPase. LILHA1.   |
| BAA37150.1     | AB022442 | Vicia faba                | p-type H <sup>+</sup> -ATPase. VHA2.   |
| CAC29435.1     | AJ310523 | Vicia faba                | P-type H <sup>+</sup> -ATPase. vha4. predominantly expressed in flowers.               |
| CAB85495.1     | AJ132892 | Medicago truncatula       | proton pump. H <sup>+</sup> -ATPase. ha1.  |
| CAB85494.1     | AJ132891 | Medicago truncatula       | proton pump. H <sup>+</sup> -ATPase. ha1.  |
| AAD46187.1     | AF156683 | Nicotiana plumbaginifolia | plasma membrane proton ATPase. pma8.   |
| AAD29712.1     | AF140499 | Oryza sativa              | chloroplast envelope calcium ATPase precursor.   |
| AAK32118.1     | AF308816 | Hordeum vulgare           | plasmalemma H <sup>+</sup> -ATPase 1.  |
| AAG01028.1     | AF289025 | Cucumis sativus           | plasma membrane H <sup>+</sup> -ATPase.  |
| AAF97591.1     | AF263917 | Lycopersicon esculentum   | plasma membrane proton ATPase. LHA8.   |
| AAA81348.1     | U38965   | Vicia faba                | p-type H <sup>+</sup> -ATPase. VHA2.   |
| AAA20600.1     | U08984   | Zea mays                  | plasma-membrane H <sup>+</sup> ATPase. Zmpma1.   |
| AAA20601.1     | U08985   | Zea mays                  | plasma-membrane H <sup>+</sup> ATPase. Zmpma1.   |
| AAK32119.1     | AF308817 | Hordeum vulgare           | plasmalemma H <sup>+</sup> -ATPase 2.  |
| SEQ ID NO: 742 |          |                           |  |
| AAD02548.1     | AF049922 | Petunia x hybrida         | PGPS/D6. PGPS/D6. ER lumen protein retaining receptor homolog; putative HDEL receptor. |
| SEQ ID NO: 744 |          |                           |  |



|                |          |                    |   |
|----------------|----------|--------------------|---|
| AAG13424.1     | AC051634 | Oryza sativa       | putative mitochondrial inner membrane protein. OSJNBb0018B10.5.   |
| AAG46068.1     | AC079830 | Oryza sativa       | putative inner mitochondrial membrane protein. OSJNBb0009F04.14.  |
| SEQ ID NO: 746 |          |                    |   |
| BAA32557.1     | AB017159 | Daucus carota      | citrate synthase. DcCS.   |
| AAA82743.1     | U19481   | Citrus maxima      | synthesis of citrate from oxaloacetate and acetylCoA. citrate synthase precursor. cit.  |
| AAG28777.1     | AF302906 | Oryza sativa       | citrate synthase. similar to putative Oryza sativa citrate synthase in GenBank Accession Number AC004521.                                   |
| CAA59008.1     | X84226   | Nicotiana tabacum  | citrate synthase. cit1.   |
| CAA52976.1     | X75082   | Solanum tuberosum  | mitochondrial citrate-synthase. ethanolamine ammonia-lyase.   |
| BAA82390.1     | AP000367 | Oryza sativa       | ESTs C96653(C10531),C96654(C10531),C28571(C61641) correspond to a region of the predicted gene.; Similar to citrate synthetase. (AC004521). |
| CAA59010.1     | X84228   | Beta vulgaris      | citrate (si)-synthase. cit1.  |
| CAA59009.1     | X84227   | Populus x generosa | citrate (si)-synthase. cit1.  |
| BAA07328.1     | D38132   | Cucurbita sp.      | conversion of oxaloacetate to citrate in the glyoxylate cycle. glyoxysomal citrate synthase.  |
| SEQ ID NO: 750 |          |                    |   |
| AAA85365.1     | L42466   | Picea glauca       | ethylene-forming enzyme. EFE.   |
| AAC95363.1     | AF104925 | Solanum chacoense  | 2-oxoglutarate-dependent dioxygenase. SPP2. pollination and fertilization induced gene.   |
| BAA75309.1     | AB023790 | Ipomoea batatas    | flavanone 3-hydroxylase. f3h III.   |
| BAA75308.1     | AB023789 | Ipomoea batatas    | flavanone 3-hydroxylase. f3h II.  |
| BAA75306.1     | AB023787 | Ipomoea batatas    | anthocyanidin synthase. ans II.   |
| AAC48922.1     | U06047   | Vigna radiata      | 1-aminocyclopropane-1-carboxylate oxidase homolog.  |
| SEQ ID NO: 751 |          |                    |   |
| BAB40010.1     | AP003021 | Oryza sativa       | putative wall-associated kinase 2. P0503E05.12.   |
| BAA95893.1     | AP002071 | Oryza sativa       | Similar to Arabidopsis thaliana wak4 gene; wall-associated kinase 4. (AJ009695).  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| BAB40015.1 | AP003021 | Oryza sativa            | putative wall-associated kinase 1. P0503E05.18.  |
| BAA92221.1 | AP001278 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).                     |
| BAA87852.1 | AP000816 | Oryza sativa            | Similar to putative Ser/Thr protein kinase. (AC004218).  |
| BAB40021.1 | AP003021 | Oryza sativa            | putative wall-associated kinase 2. P0503E05.25. contains EST C24950(S16264).   |
| CAB51834.1 | 00069    | Oryza sativa            | 11332.5. contains eukaryotic protein kinase domain PF.   |
| AAB09771.1 | U67422   | Zea mays                | CRINKLY4 precursor. cr4. receptor kinase homolog.  |
| BAB39873.1 | AP002882 | Oryza sativa            | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| AAK11566.1 | AF318490 | Lycopersicon hirsutum   | Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.  |
| BAB39409.1 | AP002901 | Oryza sativa            | putative protein kinase. P0456F08.9. contains EST C23560(R0290).   |
| BAB21240.1 | AP002953 | Oryza sativa            | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).   |
| BAB40022.1 | AP003021 | Oryza sativa            | putative wall-associated kinase 1. P0503E05.26.  |
| CAA97692.1 | Z73295   | Catharanthus roseus     | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis. |
| BAB16871.1 | AP002537 | Oryza sativa            | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).                         |
| AAK11674.1 | AF339747 | Lophopyrum elongatum    | protein kinase. ESI47.   |
| AAF43496.1 | AF131222 | Lophopyrum elongatum    | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.                        |
| BAA90808.1 | AP001168 | Oryza sativa            | Similar to putative receptor-like protein kinase (AL035679).   |
| AAF76313.1 | AF220603 | Lycopersicon esculentum | Pto kinase. LescPth5.  |
| AAB47421.1 | U59316   | Lycopersicon esculentum | serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.                         |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAB47423.1 | U59315   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Pto. Pto. bacterial speck disease resistance in tomato; disease resistance gene.  |
| AAC48914.1 | U02271   | Lycopersicon pimpinellifolium | protein kinase.   |
| AAF76306.1 | AF220602 | Lycopersicon pimpinellifolium | Pto kinase.   |
| AAG59657.1 | AC084319 | Oryza sativa                  | putative protein kinase. OSJNBa0004B24.20.  |
| AAK11567.1 | AF318491 | Lycopersicon hirsutum         | Pto-like protein kinase F. LhirPtoF.  |
| AAK21965.1 | AY028699 | Brassica napus                | receptor protein kinase PERK1.  |
| AAG25966.1 | AF302082 | Nicotiana tabacum             | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.   |
| AAK11569.1 | AF318493 | Lycopersicon hirsutum         | Pto-like protein kinase D. LhirPtoD.  |
| AAF66615.1 | AF142596 | Nicotiana tabacum             | LRR receptor-like protein kinase.   |
| AAG03090.1 | AC073405 | Oryza sativa                  | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| BAA87853.1 | AP000816 | Oryza sativa                  | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAA83373.1 | AP000391 | Oryza sativa                  | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484).  |
| BAA84787.1 | AP000559 | Oryza sativa                  | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484).  |
| CAA74662.1 | Y14286   | Brassica oleracea             | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.  |
| CAA67145.1 | X98520   | Brassica oleracea             | receptor-like kinase. SFR2.   |
| CAA73133.1 | Y12530   | Brassica oleracea             | serine /threonine kinase. ARLK.   |
| AAK11568.1 | AF318492 | Lycopersicon hirsutum         | Pto-like protein kinase B. LhirPtoB.  |
| BAA78764.1 | AB023482 | Oryza sativa                  | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |

|                |          |                       |  |
|----------------|----------|-----------------------|--|
| AAK00425.1     | AC069324 | Oryza sativa          | Putative protein kinase. OSJNBa0071K19.11.   |
| AAK31267.1     | AC079890 | Oryza sativa          | putative protein kinase. OSJNBb0089A17.2.  |
| BAA94509.1     | AB041503 | Populus nigra         | protein kinase 1. PnPK1.   |
| BAA94510.1     | AB041504 | Populus nigra         | protein kinase 2. PnPK2.   |
| SEQ ID NO: 752 |          |                       |  |
| BAB39155.1     | AB048713 | Pisum sativum         | SCARECROW. PsSCR.  |
| AAG13663.1     | AF263457 | Zea mays              | transcription factor. SCARECROW. SCR. ZmSCR.   |
| BAA90816.1     | AP001168 | Oryza sativa          | Similar to SCARECROW (U62798).   |
| AAC98090.1     | AF067400 | Zea mays              | Sc11 protein. Sc11. Scarecrow-like; similar to Zea mays sequence presented in GenBank Accession Number T18310.   |
| BAB39156.1     | AB048714 | Pisum sativum         | SCARECROW. PsSCR.  |
| SEQ ID NO: 753 |          |                       |  |
| AAF21901.1     | AF109392 | Brassica napus        | ligand gated channel-like protein. glutamate receptor homolog.   |
| SEQ ID NO: 757 |          |                       |  |
| CAA92821.1     | Z68504   | Oryza sativa          | 3-hydroxy-3-methylglutaryl-CoA reductase.  |
| AAA33360.1     | M74800   | Hevea brasiliensis    | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. hmgr3.  |
| CAA70440.1     | Y09238   | Zea mays              | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| AAB69727.1     | U72146   | Camptotheca acuminata | 3-hydroxy-3-methylglutaryl coenzyme A reductase. converts HMGC <sub>o</sub> A in mevalonate precursor for isoprenoid compounds; HMGR; hmg2; similar to protein encoded by GenBank Accession Number L10390. |
| AAD08820.1     | U43961   | Oryza sativa          | 3-hydroxy-3-methylglutaryl-CoA reductase. HMGR.  |
| AAA68965.1     | U14624   | Artemisia annua       | 3-hydroxy-3-methylglutaryl coenzyme A reductase. AHM1.   |
| AAD47596.1     | AF142473 | Artemisia annua       | HMG-CoA reductase. HMGR1.  |
| AAD03789.1     | U43711   | Morus alba            | catalyzes the final step in mevalonate pathway. 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR.   |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAA33108.1 | M96068   | Catharanthus roseus     | hydroxymethylglutaryl-CoA reductase. HMGR.   |
| CAA48610.1 | X68651   | Raphanus sativus        | hydroxymethylglutaryl-CoA reductase (NADPH). HMG1.   |
| AAA93498.1 | L01400   | Solanum tuberosum       | convert HMG-CoA into mevalonate. hydroxymethylglutaryl coenzyme A reductase. hmgr. putative.   |
| AAA68966.1 | U14625   | Artemisia annua         | 3-hydroxy-3-methylglutaryl coenzyme A reductase. AHM4.   |
| AAB52552.1 | U51986   | Solanum tuberosum       | HMG-CoA reductase.   |
| CAA48611.1 | X68652   | Raphanus sativus        | hydroxymethylglutaryl-CoA reductase (NADPH). HMG2.   |
| BAA93631.1 | AB022690 | Solanum tuberosum       | 3-hydroxy-3-methylglutaryl coenzyme A reductase. HMG3.   |
| AAB62581.1 | U68072   | Lycopersicon esculentum | 3-hydroxy-3-methylglutaryl CoA reductase 2. HMG2.  |
| AAA34169.1 | M63642   | Lycopersicon esculentum | 3-hydroxy-3-methylglutaryl coenzyme A reductase. hmg2.   |
| AAB69726.1 | U72145   | Camptotheca acuminata   | converts HMGC CoA to mevalonate. 3-hydroxy-3-methylglutaryl coenzyme a reductase. hmg3. HMGR.  |
| AAB52551.1 | U51985   | Solanum tuberosum       | HMG-CoA reductase.   |
| AAA33040.1 | L10390   | Camptotheca acuminata   | 3-hydroxy-3-methylglutaryl coA reductase.  |
| AAD28179.1 | AF110383 | Capsicum annuum         | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR2M. HMGR.   |
| BAB20771.1 | AB041031 | Solanum tuberosum       | 3-hydroxy-3-methylglutaryl coenzyme A reductase. HMG2.   |
| AAB53748.1 | U95816   | Oryza sativa            | 3-hydroxy-3-methylglutaryl-CoA reductase. HMGR.  |
| AAC05088.1 | AF038045 | Gossypium hirsutum      | catalyzes the synthesis of mevalonate from hmg-coA. 3-hydroxy-3-methylglutaryl-coenzyme A reductase 1. hmg1. hmg-coA reductase 1; HMGR1. |
| CAA45181.1 | X63649   | Nicotiana glauca        | catalyses synthesis of mevalonate. 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR. endoplasmic reticulum location.                |
| CAA38469.1 | X54659   | Hevea brasiliensis      | hydroxymethylglutaryl-CoA reductase. HMGR1.  |
| CAA38467.1 | X54657   | Hevea brasiliensis      | hydroxymethylglutaryl-CoA reductase. HMGR1.  |

|            |          |                            |  |
|------------|----------|----------------------------|--|
| AAD38873.1 | AF110382 | Oryza sativa               | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMGR3G. HMG-CoA reductase.  |
| BAA36291.1 | AB021862 | Cucumis melo               | HMG-CoA reductase. Cm-HMGR. putative.  |
| AAB87727.1 | U60452   | Nicotiana tabacum          | hydroxy-methylglutaryl-coenzyme A reductase. HMGR1.  |
| AAC05089.1 | AF038046 | Gossypium hirsutum         | catalyzes the synthesis of mevalonate from hmg-coA. 3-hydroxy-3-methylglutaryl-coenzyme A reductase 2. hmg2. hmg-coA reductase 2; HMGR2.   |
| AAC15475.1 | AF034760 | Tagetes erecta             | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| AAC15476.1 | AF034761 | Tagetes erecta             | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| AAA21720.1 | L28995   | Oryza sativa               | conversion of hydroxymethylglutaryl coenzyme A to mevalonate. 3-hydroxy-3-methylglutaryl coenzyme A reductase. putative.   |
| AAA33358.1 | M74798   | Hevea brasiliensis         | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. hmgr3.  |
| AAC72378.1 | AF096838 | Solanum tuberosum          | 3-hydroxy-3-methylglutaryl coenzyme A reductase.   |
| AAB04043.1 | L40938   | Lycopersicon esculentum    | HMGR CoA reductase. HMGR1.   |
| CAA38468.1 | X54658   | Hevea brasiliensis         | hydroxymethylglutaryl-CoA reductase. HMGR2.  |
| CAA52787.1 | X74783   | Lithospermum erythrorhizon | 3-hydroxy-3-methyl glutaryl coenzyme A reductase. Lehmgr1.   |
| AAD09278.1 | U97683   | Glycine max                | catalyzes the synthesis of mevalonate, the specific precursor of all isoprenoid compounds present in plants. 3-hydroxy-3-methylglutaryl-coenzyme A reductase. HMG-CoA reductase.           |
| AAG43469.1 | AF196964 | Bixa orellana              | catalyzes mevalonate synthesis from hmg-CoA. 3-hydroxy-3-methylglutaryl-coenzyme A reductase.  |
| BAA09705.1 | D63389   | Cucumis sativus            | 3-hydroxy-3-methylglutaryl CoA reductase.  |
| AAB47161.1 | S82272   | Gossypium barbadense       | 3-hydroxy-3-methylglutaryl coenzyme A reductase. /gene="3-hydroxy-3-methylglutaryl coenzyme A reductase,. This sequence comes from Fig. 2; 3-hydroxy-3-methylglutaryl CoA reductase; HMGR. |
| AAA33359.1 | M74799   | Hevea brasiliensis         | 3-hydroxy-3-methylglutaryl-coenzyme A reductase. hmgr3.  |
| AAC37434.1 | L34827   | Solanum tuberosum          | HMG-CoA reductase. hmg1 gene family.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAC37432.1     | L34825   | Solanum tuberosum       | HMG-CoA reductase. hmg1 gene family.   |
| AAC37435.1     | L34828   | Solanum tuberosum       | HMG-CoA reductase. hmg1 gene family.   |
| AAC37433.1     | L34826   | Solanum tuberosum       | HMG-CoA reductase. hmg1 gene family.   |
| AAC37431.1     | L34823   | Solanum tuberosum       | HMG-CoA reductase. hmg1 gene family.   |
| SEQ ID NO: 761 |          |                         |  |
| CAA96512.1     | Z71980   | Malus x domestica       | knotted1-like homeobox protein.  |
| BAA25921.1     | AB004797 | Nicotiana tabacum       | NTH23. homeobox gene.  |
| AAD09582.1     | U76409   | Lycopersicon esculentum | homeobox 1 protein. THox1. homeodomain protein.  |
| AAC49918.1     | AF000142 | Lycopersicon esculentum | class II knotted-like homeodomain protein. LeT12.  |
| BAA08552.1     | D49704   | Oryza sativa            | OSH45. OSH44 transcript homeobox gene.   |
| BAB18585.1     | AB043957 | Ceratopteris richardii  | CRKNOX3. crknox3. knotted1-like homeodomain protein.   |
| AAD00253.1     | U76410   | Lycopersicon esculentum | homeobox 2 protein. THox2. homeodomain protein.  |
| CAA82314.1     | Z29073   | Brassica napus          | homeodomain-containing protein. Bnhdl.   |
| BAA77822.1     | AB007628 | Oryza sativa            | HOS59. homeobox gene.  |
| BAA77823.1     | AB007629 | Oryza sativa            | HOS66. homeobox gene.  |
| AAB81079.1     | AF022390 | Hordeum vulgare         | knotted class 1 homeodomain protein. k. similar to the hooded gene product encoded by GenBank Accession Number X83518; similar to the maize knotted-1 gene product encoded by GenBank Accession Number X61308. |
| AAF32399.1     | AF224499 | Triticum aestivum       | KNOTTED-1-like homeobox protein b. knox1b. KNOX1b.   |
| BAA76750.1     | AB025573 | Nicotiana tabacum       | KN1-type homeobox protein. NTH1.   |
| AAF32400.1     | AF224500 | Triticum aestivum       | KNOTTED-1-like homeobox protein d. knox1d. KNOX1d.   |
| AAC49917.1     | AF000141 | Lycopersicon esculentum | class I knotted-like homeodomain protein. LeT6.  |
| BAA25546.1     | AB004785 | Nicotiana tabacum       | NTH15. homeobox gene.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAF32398.1 | AF224498 | Triticum aestivum       | KNOTTED-1-like homeobox protein a. knox1a. KNOX1.                |
| AAD13611.1 | AF100455 | Zea mays                | knotted class 1 homeodomain protein liguleless3. lg3.            |
| AAC84001.1 | AF063248 | Picea abies             | homeobox protein.  |
| AAC33008.1 | AF080104 | Pisum sativum           | knotted1-like class I homeodomain protein. PsKn1.                |
| AAD00692.1 | U90092   | Picea mariana           | homeobox transcription factor SKN2. knotted1-like homeobox gene. |
| AAC32262.1 | AF063307 | Pisum sativum           | Knox class 1 protein. Hop1.                                      |
| BAA03959.1 | D16507   | Oryza sativa            | homeobox protein. OSH1.  |
| AAG27464.1 | AF308454 | Medicago truncatula     | knotted class I homeodomain KNOX.                                |
| AAD00691.1 | U90091   | Picea mariana           | homeobox transcription factor SKN1. knotted1-like homeobox gene. |
| AAC32817.1 | AF050180 | Oryza sativa            | transcription factor. KNOX class homeodomain protein. Oskn2.     |
| BAA79226.1 | AB028885 | Oryza sativa            | knotted1-type homeobox protein OSH71. OSH71.                     |
| BAA79224.1 | AB028883 | Oryza sativa            | knotted1-type homeobox protein OSH6. OSH6.                       |
| BAA77818.1 | AB007624 | Oryza sativa            | HOS9. homeobox gene.   |
| BAB19772.1 | AP002881 | Oryza sativa            | putative knotted1-type homeobox protein. P0035H10.13.            |
| AAF23753.2 | AF193813 | Brassica oleracea       | shoot meristemless. Stm. homeodomain protein.                    |
| AAD00251.1 | U76407   | Lycopersicon esculentum | knotted 2 protein. TKn2. homeodomain protein.                    |
| CAA96510.1 | Z71978   | Malus x domestica       | kn1-like protein.  |
| BAA31688.1 | AB016071 | Oryza sativa            | OSH15. homeobox gene.  |
| BAA77817.1 | AB007623 | Oryza sativa            | HOS3. homeobox gene.   |
| BAB18582.1 | AB043954 | Ceratopteris richardii  | CRKNOX1. crknox1. knotted1-like homeodomain protein.             |
| BAA76903.1 | AB025713 | Nicotiana tabacum       | homeobox 9. NTH9.  |



AAD00252.1 U76408 *Lycopersicon esculentum*  
knotted 3 protein. TKn3. homeodomain protein.

BAB18584.1 AB043956 *Ceratopteris richardii*  
CRKNOX2. crknox2. knotted1-like homeodomain protein.

CAA96511.1 Z71979 *Malus x domestica*  
kn1-like protein.

AAA20882.1 L13663 *Glycine max*  
SBH1. Sbh1. soybean homeobox-containing gene.

CAB88029.1 AJ276389 *Dendrobium grex Madame Thong-In*  
transcription factor. knotted1-like homeobox protein.

SEQ ID NO: 762

AAB65776.1 U97521 *Vitis vinifera*  
class IV endochitinase. VvChi4A.

AAB65777.1 U97522 *Vitis vinifera*  
class IV endochitinase. VvChi4B.

CAA53626.1 X76041 *Triticum aestivum*  
endochitinase. CHI.

AAG53609.1 AF280437 *Secale cereale*  
31.7 kDa class I endochitinase-antifreeze protein precursor. cht9.

BAA03750.1 D16222 *Oryza sativa*  
endochitinase. Cht-2.

AAA32986.1 M95835 *Brassica napus*  
endochitinase. Ch25.

BAA03749.1 D16221 *Oryza sativa*  
endochitinase. Cht-1.

AAB01895.1 U48687 *Castanea sativa*  
endochitinase.

SEQ ID NO: 763

AAA34065.1 M94135 *Nicotiana tabacum*  
chloroplast carbonic anhydrase.

AAA34057.1 L19255 *Nicotiana tabacum*  
carbonic anhydrase.

AAB65822.1 U55838 *Populus tremula x Populus tremuloides*  
carbonic anhydrase. CA1b. EC 4.2.1.1.

AAC49785.1 U55837 *Populus tremula x Populus tremuloides*  
carbonic anhydrase. CA1a. EC 4.2.1.1.

AAA34026.1 M27295 *Spinacia oleracea*  
carbonic anhydrase precursor.

AAA34027.1 J05403 *Spinacia oleracea*  
carbonic anhydrase (EC 4.2.1.1).

AAA86993.1 U19738 *Flaveria linearis*  
reversible hydration of carbon dioxide. carbonic anhydrase 1.

|            |          |                     |  |
|------------|----------|---------------------|--|
| AAA86942.1 | U08402   | Flaveria brownii    | carbonic anhydrase.  |
| AAA86992.1 | U19737   | Flaveria pringlei   | reversible hydration of carbon dioxide. carbonic anhydrase.  |
| AAA86939.1 | U08398   | Flaveria bidentis   | carbonic anhydrase.  |
| AAD27876.2 | AF139464 | Vigna radiata       | carbonic anhydrase. CipCa1.  |
| AAA33652.1 | M63627   | Pisum sativum       | carbonic anhydrase.  |
| AAD29050.1 | AF132855 | Gossypium hirsutum  | interconversion of CO <sub>2</sub> and HCO <sub>3</sub> <sup>-</sup> . carbonic anhydrase isoform 2. CA2. zinc metalloenzyme; carbonate dehydratase. |
| AAD29049.1 | AF132854 | Gossypium hirsutum  | interconversion of CO <sub>2</sub> and HCO <sub>3</sub> <sup>-</sup> . carbonic anhydrase isoform 1. CA1. zinc metalloenzyme; carbonate dehydratase. |
| AAA86994.1 | U19740   | Flaveria linearis   | reversible hydration of carbon dioxide. carbonic anhydrase 2.  |
| CAB43571.1 | AJ239132 | Glycine max         | hydration of carbon dioxide. carbonic anhydrase. ca1.  |
| CAA63712.1 | X93312   | Medicago sativa     | Carbonic anhydrase. ca1.   |
| AAA86945.1 | U08403   | Zea mays            | carbonic anhydrase.  |
| AAA86944.1 | U08401   | Zea mays            | carbonic anhydrase.  |
| AAC41656.1 | L36959   | Hordeum vulgare     | carbonic anhydrase. putative.  |
| AAA86943.1 | U08404   | Oryza sativa        | carbonic anhydrase. nuclear encoded, localized to chloroplast.   |
| AAD56038.1 | AF182806 | Oryza sativa        | carbonic anhydrase 3. ca3.   |
| BAA31953.1 | AB016283 | Oryza sativa        | carbonic anhydrase.  |
| AAA69027.1 | U19739   | Urochloa panicoides | reversible hydration of carbon dioxide. carbonic anhydrase 2.  |
| AAA69028.1 | U19741   | Urochloa panicoides | reversible hydration of carbon dioxide. carbonic anhydrase 1.  |
| BAA95793.1 | AB009887 | Nicotiana tabacum   | carbonic anhydrase. carbonic anhydrase.  |
| AAF78507.1 | AF195204 | Pyrus pyrifolia     | carbonic anhydrase isoform 1. CA1.   |

|   |        |                           |
|---|--------|---------------------------|
| AAC33484.1                                | U49976 | Coccomyxa sp. PA          |
| beta-type carbonic anhydrase beta-CA1.    |        |                           |
| AAA18560.1                                | M95073 | Zea mays                  |
| putative. silimar to carbonic anhydrases. |        |                           |
| AAB19184.1                                | U41190 | Chlamydomonas reinhardtii |
| carbonic anhydrase precursor. beta-CA2.   |        |                           |
| AAB19183.1                                | U41189 | Chlamydomonas reinhardtii |
| carbonic anhydrase precursor. beta-CA1.   |        |                           |
| AAC49887.1                                | U80804 | Chlamydomonas reinhardtii |
| beta-carbonic anhydrase. ca1. beta-CA1.   |        |                           |
| AAC49888.1                                | U80805 | Chlamydomonas reinhardtii |
| beta-carbonic anhydrase. ca2. beta-CA2.   |        |                           |

SEQ ID NO: 764

|  |          |                   |
|--|----------|-------------------|
| AAC06027.1   | AF052058 | Vigna unguiculata |
| iron storage and mobilization in plants. ferritin subunit cowpea2 precursor. |          |                   |
| AAD50644.1   | AF133814 | Solanum tuberosum |
| ferritin 1. F1.  |          |                   |
| AAB53099.1   | U68217   | Brassica napus    |
| iron binding protein. ferritin. LSC30.                                       |          |                   |
| AAA33959.1   | M64337   | Glycine max       |
| ferritin light chain. ferritin.  |          |                   |
| AAA34016.1   | M72894   | Glycine max       |
| ferritin light chain. SOF-H2.  |          |                   |
| CAA58146.1   | X83076   | Zea mays          |
| ferritin. Fer1.  |          |                   |
| AAB18928.1   | U31648   | Glycine max       |
| iron storage protein. ferritin.  |          |                   |
| CAA65771.1   | X97059   | Medicago sativa   |
| iron storage. ferritin. FER. abscisic acid regulated.                        |          |                   |
| CAA43663.1   | X61391   | Zea mays          |
| ferritin.  |          |                   |
| CAA58147.1   | X83077   | Zea mays          |
| ferritin. Fer2.  |          |                   |
| AAC06026.1   | AF052057 | Vigna unguiculata |
| iron storage and mobilization in plants. ferritin subunit cowpea3 precursor. |          |                   |
| CAA43664.1   | X61392   | Zea mays          |
| ferritin.  |          |                   |
| CAA51786.1   | X73369   | Pisum sativum     |
| ferritin.  |          |                   |
| CAA45763.1   | X64417   | Pisum sativum     |
| ferritin-precursor.  |          |                   |

|                |          |  |
|----------------|----------|--|
| CAA41213.1     | X58274   | Phaseolus vulgaris<br>ferritin. pfe.   |
| AAA33958.1     | M58336   | Glycine max<br>ferritin light chain. SOF-5L.   |
| CAB42587.1     | AJ238628 | Chlorella protothecoides<br>putative ferritin. dee188.                                 |
| BAB17852.1     | AB042612 | Nicotiana tabacum<br>ferritin 1. tob-fer-1. putative.                                  |
| AAC15241.1     | AF028072 | Pinus taeda<br>ferritin.   |
| CAA47983.1     | X67755   | Vigna unguiculata<br>ferritin 2. pfe2.   |
| CAA47982.1     | X67754   | Vigna unguiculata<br>ferritin 1. pfe1.   |
| CAA47984.1     | X67756   | Vigna unguiculata<br>ferritin 5. pfe5.   |
| AAC12282.1     | AF052511 | Glycine max<br>iron storage and mobilization. ferritin 2. soybean 2; cowpea 2 homolog. |
| AAC12281.1     | AF052513 | Glycine max<br>iron storage and mobilization. ferritin 1. soybean 1; cowpea 1 homolog. |
| SEQ ID NO: 765 |          |  |
| AAC36697.1     | AF075579 | Mesembryanthemum crystallinum<br>protein phosphatase-2C. PP2C.                         |
| CAC10358.1     | AJ277086 | Nicotiana tabacum<br>protein phosphatase 2C. PP2C1.                                    |
| CAB90633.1     | AJ277743 | Fagus sylvatica<br>protein phosphatase 2C (PP2C). pp2C1. ABA-induced protein.          |
| AAD17804.1     | AF092431 | Lotus japonicus<br>nodule-enhanced protein phosphatase type 2C. NPP2C1.                |
| CAC10359.1     | AJ277087 | Nicotiana tabacum<br>protein phosphatase 2C. PP2C2.                                    |
| CAC09575.1     | AJ298987 | Fagus sylvatica<br>protein phosphatase 2C (PP2C). pp2Cf1.                              |
| CAA72341.1     | Y11607   | Medicago sativa<br>protein phosphatase 2C. MP2C.                                       |
| CAB61839.1     | AJ242803 | Sporobolus stapfianus<br>putative serine/threonine phosphatase type 2c.                |
| AAD17805.1     | AF092432 | Lotus japonicus<br>protein phosphatase type 2C. PP2C2.                                 |
| AAG43835.1     | AF213455 | Zea mays<br>protein phosphatase type-2C. pp2c-1. PP2C-1.                               |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| AAG46118.1     | AC073166 | Oryza sativa                  | putative protein phosphatase-2C. OSJNBb0064P21.12. tRNA-Lys.  |
| AAG13599.1     | AC051633 | Oryza sativa                  | putative protein phosphatase-2C. OSJNBb0015I11.26.  |
| AAC36698.1     | AF075580 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| BAB12036.1     | AP002820 | Oryza sativa                  | putative protein phosphatase. P0702D12.18.  |
| AAC36700.1     | AF075582 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| AAC36699.1     | AF075581 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| AAK20060.1     | AC025783 | Oryza sativa                  | putative protein phosphatase 2C. OSJNBa0001O14.1.   |
| AAD11430.1     | AF097667 | Mesembryanthemum crystallinum | protein phosphatase 2C homolog. PP2C.   |
| CAB90634.1     | AJ277744 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2C2. ABA and calcium induced protein.  |
| AAC35951.1     | AF079355 | Mesembryanthemum crystallinum | protein phosphatase-2c. PP2C.   |
| AAF19804.1     | AF180355 | Brassica oleracea             | ABI1 protein. ABI1. similar to Arabidopsis thaliana ABI1.   |
| AAB93832.1     | U81960   | Zea mays                      | kinase interaction domain mediates phosphorylation-dependent binding to a subset of active receptor-like protein kinases; type 2C protein phosphatase. kinase associated protein phosphatase. KAPP. |
| AAC26828.1     | AF075603 | Oryza sativa                  | kinase interaction domain mediates phosphorylation-dependent binding to a subset of active receptor-like protein kinases. kinase associated protein phosphatase. kapp. type 2C protein phosphatase. |
| CAC09576.1     | AJ298988 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2Cf2.  |
| SEQ ID NO: 766 |          |                               |   |
| AAG08959.1     | AF122051 | Solanum tuberosum             | tuber-specific and sucrose-responsive element binding factor. TSF. R2R3 MYB class transcription factor.   |
| AAG08960.1     | AF122052 | Solanum tuberosum             | tuber-specific and sucrose-responsive element binding factor. TSF. R2R3 MYB class transcription factor.   |
| AAG08961.1     | AF122053 | Solanum tuberosum             | tuber-specific and sucrose-responsive element binding factor. TSF. R2R3 MYB class transcription factor.   |

|            |          |                       |   |
|------------|----------|-----------------------|---|
| BAA88169.1 | AP000836 | Oryza sativa          | Similar to putative transcription factor (AF062890).  |
| BAA88205.1 | AP000837 | Oryza sativa          | Similar to putative transcription factor (AF062890).  |
| AAF34434.1 | AF172282 | Oryza sativa          | myb-like protein. DUPR11.29.  |
| AAF78890.1 | AF189788 | Hordeum vulgare       | putative c-myb-like transcription factor. MYB3R-1. HvMYB3R-1.   |
| AAF78889.1 | AF189787 | Hordeum vulgare       | putative c-myb-like transcription factor. MYB3R-1. HvMYB3R-1.   |
| AAF67053.1 | AF190304 | Adiantum raddianum    | c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.   |
| AAF67052.1 | AF190303 | Adiantum raddianum    | c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.   |
| AAF78888.1 | AF189786 | Physcomitrella patens | putative c-myb-like transcription factor. MYB3R-1. PpMYB3R-1.   |
| AAF78887.1 | AF189785 | Physcomitrella patens | putative c-myb-like transcription factor. MYB3R-1. PpMYB3R-1.   |
| AAF43043.1 | AF236059 | Papaver rhoeas        | putative Myb-related domain. pmr.   |
| BAA94769.1 | AP001859 | Oryza sativa          | Similar to Arabidopsis thaliana chromosome 4, BAC clone F4D11; putative myb-protein. (AL022537).              |
| AAF67050.1 | AF190301 | Secale cereale        | c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.   |
| AAF67051.1 | AF190302 | Secale cereale        | c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.   |
| CAA78388.1 | Z13998   | Petunia x hybrida     | DNA-binding protein; transcriptional activator. protein 3. myb.Ph3. related to animal myb proto-oncoproteins. |
| BAB39987.1 | AP003020 | Oryza sativa          | putative transcription factor (myb). P0498A12.16. contains ESTs AU097474(S5087),D40175(S1959).                |
| BAB39972.1 | AP003018 | Oryza sativa          | putative transcription factor (myb). OSJNBa0004B13.27. contains ESTs AU097474(S5087),D40175(S1959).           |
| CAA72218.1 | Y11415   | Oryza sativa          | myb.  |
| BAA81731.1 | AB029160 | Glycine max           | GmMYB29A1.  |
| BAA81730.1 | AB029159 | Glycine max           | GmMYB29A1.  |

|            |          |                    |   |
|------------|----------|--------------------|---|
| CAA72217.1 | Y11414   | Oryza sativa       | myb.  |
| BAA81736.1 | AB029165 | Glycine max        | GmMYB29B2.  |
| BAB12688.1 | AP002746 | Oryza sativa       | putative MYB family transcription factor. P0671B11.3. contains ESTs AU082307(E0784),C72014(E0784).  |
| BAA99440.1 | AP002743 | Oryza sativa       | putative MYB family transcription factor. P0710E05.27. contains ESTs AU082307(E0784),C72014(E0784).   |
| AAB41101.1 | U72762   | Nicotiana tabacum  | transcription factor Myb1. myb1. TMV-inducible Myb homolog; contains helix-turn-helix motif; contains redox-sensitive cysteine.               |
| BAA88223.1 | AB028651 | Nicotiana tabacum  | myb-related transcription factor LBM3. lbm3.  |
| BAA23340.1 | D88620   | Oryza sativa       | transfactor. OSMYB4. Osmby4.  |
| BAA93038.1 | AP001552 | Oryza sativa       | EST AU082058(C12976) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana putative transcription factor (AF062916). |
| AAA33067.1 | L04497   | Gossypium hirsutum | MYB A; putative.  |
| BAA81733.2 | AB029162 | Glycine max        | GmMYB29A2.  |
| BAA88222.1 | AB028650 | Nicotiana tabacum  | myb-related transcription factor LBM2. lbm2.  |
| CAB43399.1 | AJ006292 | Antirrhinum majus  | Myb-related transcription factor mixta-like 1. mybml1.  |
| BAA81732.1 | AB029161 | Glycine max        | GmMYB29A2.  |
| AAG28525.1 | AF198498 | Nicotiana tabacum  | anther-specific myb-related protein 2. mybAS2. NtMYBAS2; contains tandem R2, R3 myb domains similar to c-myb family.                          |
| CAA50226.1 | X70881   | Hordeum vulgare    | MybHv33. myb3.  |
| CAA50223.1 | X70878   | Hordeum vulgare    | MybHv33. myb3.  |
| CAA78387.1 | Z13997   | Petunia x hybrida  | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.                                 |
| CAA67000.1 | X98355   | Oryza sativa       | activator of alpha-amylase gene promoter. transcription factor GAMyb. Gam1. Myb-like; expression is regulated by gibberellin.                 |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAK19616.1     | AF336283 | Gossypium hirsutum      | GHMYB25. ghmyb25. similar to myb; contains an unspliced intron.  |
| AAG28526.1     | AF198499 | Nicotiana tabacum       | anther-specific myb-related protein 1. mybAS1. NtMYBAS1; contains N-terminal R2, R3 myb domain repeats similar to c-myb.   |
| CAA78386.1     | Z13996   | Petunia x hybrida       | DNA binding protein; transcriptional activator. protein 1. myb.Ph3. Product related to animal myb proto-oncoproteins. Sequence from nucleotide 992 is not included in clone cPF1 and has been obtained by PCR amplification of cDNA. |
| CAA66952.1     | X98308   | Lycopersicon esculentum | THM18. myb-related transcription factor.   |
| BAA23341.1     | D88621   | Oryza sativa            | transfactor. OSMYB5. Osmyb5.   |
| CAA50221.1     | X70876   | Hordeum vulgare         | MybHv5. myb2.  |
| AAK19611.1     | AF336278 | Gossypium hirsutum      | BNLGHi233. bnlghi6233. similar to myb.   |
| AAC04716.1     | AF034130 | Gossypium hirsutum      | putative MYB-like transcription factor. MYB-like DNA-binding domain protein. Cmy-D. similar to MYB A encoded by GenBank Accession Number L04497.   |
| CAA61021.1     | X87690   | Hordeum vulgare         | transcriptional activator of alpha-amylase gene promoter. GAMyb protein. Gam1.   |
| AAG22863.1     | AY008692 | Hordeum vulgare         | transcription factor GAMyb. Gamyb.   |
| SEQ ID NO: 767 |          |                         |  |
| CAB08111.1     | Z94180   | Lycopersicon esculentum | branched chain alpha-keto acid dehydrogenase E1-alpha subunit.   |
| CAA81558.1     | Z26949   | Solanum tuberosum       | subunit of the mitochondrial pyruvate dehydrogenase complex. E1 alpha subunit of pyruvate dehydrogenase precursor.   |
| AAC72195.1     | AF069911 | Zea mays                | pyruvate dehydrogenase E1 alpha subunit.   |
| AAG43499.1     | AF209924 | Lycopersicon esculentum | pyruvate dehydrogenase.  |
| AAA97411.1     | U51918   | Pisum sativum           | pyruvate dehydrogenase E1 alpha subunit.   |
| CAA10992.1     | AJ222787 | Hordeum vulgare         | alpha-keto acid dehydrogenase-like protein. homology to branched chain alpha-keto acid dehydrogenase E1-alpha subunit.   |
| SEQ ID NO: 768 |          |                         |  |
| AAF64450.1     | AF239928 | Euphorbia esula         | glutathione S-transferase. similar to auxin-inducible GST.   |



---

AAG16758.1 AY007560 *Lycopersicon esculentum*  
putative glutathione S-transferase T3.

---

AAG34803.1 AF243368 *Glycine max*  
glutathione S-transferase GST 13.

---

AAG16756.1 AY007558 *Lycopersicon esculentum*  
putative glutathione S-transferase T1.

---

AAG34796.1 AF243361 *Glycine max*  
glutathione S-transferase GST 6.

---

AAG34809.1 AF243374 *Glycine max*  
glutathione S-transferase GST 19.

---

AAG34797.1 AF243362 *Glycine max*  
glutathione S-transferase GST 7.

---

AAG34807.1 AF243372 *Glycine max*  
glutathione S-transferase GST 17.

---

AAG34798.1 AF243363 *Glycine max*  
glutathione S-transferase GST 8.

---

AAG34804.1 AF243369 *Glycine max*  
glutathione S-transferase GST 14.

---

AAG16759.1 AY007561 *Lycopersicon esculentum*  
putative glutathione S-transferase T4.

---

AAG34801.1 AF243366 *Glycine max*  
glutathione S-transferase GST 11.

---

AAG34810.1 AF243375 *Glycine max*  
glutathione S-transferase GST 20.

---

AAG16757.1 AY007559 *Lycopersicon esculentum*  
putative glutathione S-transferase T2.

---

AAC18566.1 AF048978 *Glycine max*  
2,4-D inducible glutathione S-transferase. GSTa.

---

AAG34808.1 AF243373 *Glycine max*  
glutathione S-transferase GST 18.

---

AAG34800.1 AF243365 *Glycine max*  
glutathione S-transferase GST 10.

---

CAA71784.1 Y10820 *Glycine max*  
glutathione transferase.

---

AAG34844.1 AF244701 *Zea mays*  
glutathione S-transferase GST 36.

---

AAG32472.1 AF309379 *Oryza sativa* subsp. *japonica*  
putative glutathione S-transferase OsGSTU3.

---

AAA68430.1 J03679 *Solanum tuberosum*  
glutathione S-transferase. *gst1*. previously called pathogenesis-related protein; *prp1-1*.

---

CAA04391.1 AJ000923 *Carica papaya*  
glutathione transferase. PGST1.

---

|                |          |  |
|----------------|----------|--|
| AAG34831.1     | AF244688 | <i>Zea mays</i><br>glutathione S-transferase GST 23.   |
| CAA09187.1     | AJ010448 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1a.   |
| CAA09188.1     | AJ010449 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1b.   |
| AAG34802.1     | AF243367 | <i>Glycine max</i><br>glutathione S-transferase GST 12.  |
| AAG34805.1     | AF243370 | <i>Glycine max</i><br>glutathione S-transferase GST 15.  |
| AAG34832.1     | AF244689 | <i>Zea mays</i><br>glutathione S-transferase GST 24.   |
| AAG34837.1     | AF244694 | <i>Zea mays</i><br>glutathione S-transferase GST 29.   |
| AAG34836.1     | AF244693 | <i>Zea mays</i><br>glutathione S-transferase GST 28.   |
| AAG32471.1     | AF309378 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTU4.  |
| AAG34849.1     | AF244706 | <i>Zea mays</i><br>glutathione S-transferase GST 41.   |
| CAC24549.1     | AJ296343 | <i>Cichorium intybus</i> x <i>Cichorium endivia</i><br>glutathione S-transferase. chi-GST1. auxin-induced GST.   |
| AAC32118.1     | AF051214 | <i>Picea mariana</i><br>probable glutathione S-transferase. Sb18. similar to <i>Nicotiana tabacum</i> probable glutathione S-transferase encoded by GenBank Accession Number X56266. |
| AAG34795.1     | AF243360 | <i>Glycine max</i><br>glutathione S-transferase GST 5.   |
| AAG34841.1     | AF244698 | <i>Zea mays</i><br>glutathione S-transferase GST 33.   |
| AAF29773.1     | AF159229 | <i>Gossypium hirsutum</i><br>glutathione S-transferase. GST.   |
| SEQ ID NO: 769 |          |  |
| AAG34797.1     | AF243362 | <i>Glycine max</i><br>glutathione S-transferase GST 7.   |
| AAG34798.1     | AF243363 | <i>Glycine max</i><br>glutathione S-transferase GST 8.   |
| AAG34803.1     | AF243368 | <i>Glycine max</i><br>glutathione S-transferase GST 13.  |
| AAG16758.1     | AY007560 | <i>Lycopersicon esculentum</i><br>putative glutathione S-transferase T3.   |
| AAF64450.1     | AF239928 | <i>Euphorbia esula</i><br>glutathione S-transferase. similar to auxin-inducible GST.   |

|            |          |   |
|------------|----------|---|
| AAG34801.1 | AF243366 | Glycine max<br>glutathione S-transferase GST 11.  |
| AAG34804.1 | AF243369 | Glycine max<br>glutathione S-transferase GST 14.  |
| AAG34796.1 | AF243361 | Glycine max<br>glutathione S-transferase GST 6.   |
| AAG34809.1 | AF243374 | Glycine max<br>glutathione S-transferase GST 19.  |
| AAG16759.1 | AY007561 | Lycopersicon esculentum<br>putative glutathione S-transferase T4.   |
| AAG16757.1 | AY007559 | Lycopersicon esculentum<br>putative glutathione S-transferase T2.   |
| AAG16756.1 | AY007558 | Lycopersicon esculentum<br>putative glutathione S-transferase T1.   |
| AAG34807.1 | AF243372 | Glycine max<br>glutathione S-transferase GST 17.  |
| AAG34810.1 | AF243375 | Glycine max<br>glutathione S-transferase GST 20.  |
| AAG34844.1 | AF244701 | Zea mays<br>glutathione S-transferase GST 36.   |
| AAG34831.1 | AF244688 | Zea mays<br>glutathione S-transferase GST 23.   |
| AAC18566.1 | AF048978 | Glycine max<br>2,4-D inducible glutathione S-transferase. GSTa.   |
| AAG34832.1 | AF244689 | Zea mays<br>glutathione S-transferase GST 24.   |
| AAG34808.1 | AF243373 | Glycine max<br>glutathione S-transferase GST 18.  |
| AAG34837.1 | AF244694 | Zea mays<br>glutathione S-transferase GST 29.   |
| AAG34800.1 | AF243365 | Glycine max<br>glutathione S-transferase GST 10.  |
| AAG34836.1 | AF244693 | Zea mays<br>glutathione S-transferase GST 28.   |
| CAA04391.1 | AJ000923 | Carica papaya<br>glutathione transferase. PGST1.  |
| CAA71784.1 | Y10820   | Glycine max<br>glutathione transferase.   |
| AAG34849.1 | AF244706 | Zea mays<br>glutathione S-transferase GST 41.   |
| AAA68430.1 | J03679   | Solanum tuberosum<br>glutathione S-transferase. gst1. previously called pathogenesis-related protein; prp1-1. |

|            |          |  |
|------------|----------|--|
| AAG34802.1 | AF243367 | Glycine max<br>glutathione S-transferase GST 12.   |
| CAC24549.1 | AJ296343 | Cichorium intybus x Cichorium endivia<br>glutathione S-transferase. chi-GST1. auxin-induced GST. |
| CAA09187.1 | AJ010448 | Alopecurus myosuroides<br>glutathione transferase. GST1a.  |
| AAF22518.1 | AF118925 | Papaver somniferum<br>glutathione S-transferase 2. GST2.   |
| AAG32471.1 | AF309378 | Oryza sativa subsp. japonica<br>putative glutathione S-transferase OsGSTU4.                      |
| CAA09188.1 | AJ010449 | Alopecurus myosuroides<br>glutathione transferase. GST1b.  |
| AAF22517.1 | AF118924 | Papaver somniferum<br>glutathione S-transferase 1. GST1.   |
| AAF22647.1 | AF193439 | Lycopersicon esculentum<br>glutathione S-transferase/peroxidase. BI-GST/GPX.                     |
| AAG32473.1 | AF309380 | Oryza sativa subsp. japonica<br>putative glutathione S-transferase OsGSTU2.                      |
| AAG32472.1 | AF309379 | Oryza sativa subsp. japonica<br>putative glutathione S-transferase OsGSTU3.                      |

SEQ ID NO: 771

|            |          |  |
|------------|----------|--|
| AAG46118.1 | AC073166 | Oryza sativa<br>putative protein phosphatase-2C. OSJNBb0064P21.12. tRNA-Lys. |
| AAG13599.1 | AC051633 | Oryza sativa<br>putative protein phosphatase-2C. OSJNBb0015I11.26.           |
| BAB12036.1 | AP002820 | Oryza sativa<br>putative protein phosphatase. P0702D12.18.                   |
| AAC36698.1 | AF075580 | Mesembryanthemum crystallinum<br>protein phosphatase-2C. PP2C.               |
| CAA72341.1 | Y11607   | Medicago sativa<br>protein phosphatase 2C. MP2C.                             |
| AAG43835.1 | AF213455 | Zea mays<br>protein phosphatase type-2C. pp2c-1. PP2C-1.                     |
| CAB61839.1 | AJ242803 | Sporobolus stapfianus<br>putative serine/threonine phosphatase type 2c.      |
| AAD17804.1 | AF092431 | Lotus japonicus<br>nodule-enhanced protein phosphatase type 2C. NPP2C1.      |
| AAD17805.1 | AF092432 | Lotus japonicus<br>protein phosphatase type 2C. PP2C2.                       |
| AAC36697.1 | AF075579 | Mesembryanthemum crystallinum<br>protein phosphatase-2C. PP2C.               |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| CAB90633.1     | AJ277743 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2C1. ABA-induced protein.  |
| CAC10358.1     | AJ277086 | Nicotiana tabacum             | protein phosphatase 2C. PP2C1.  |
| CAC10359.1     | AJ277087 | Nicotiana tabacum             | protein phosphatase 2C. PP2C2.  |
| AAC36700.1     | AF075582 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| CAC09575.1     | AJ298987 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2Cf1.  |
| AAK20060.1     | AC025783 | Oryza sativa                  | putative protein phosphatase 2C. OSJNBa0001O14.1.   |
| CAB90634.1     | AJ277744 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2C2. ABA and calcium induced protein.  |
| AAC35951.1     | AF079355 | Mesembryanthemum crystallinum | protein phosphatase-2c. PP2C.   |
| AAD11430.1     | AF097667 | Mesembryanthemum crystallinum | protein phosphatase 2C homolog. PP2C.   |
| AAB93832.1     | U81960   | Zea mays                      | kinase interaction domain mediates phosphorylation-dependent binding to a subset of active receptor-like protein kinases; type 2C protein phosphatase. kinase associated protein phosphatase. KAPP. |
| AAC26828.1     | AF075603 | Oryza sativa                  | kinase interaction domain mediates phosphorylation-dependent binding to a subset of active receptor-like protein kinases. kinase associated protein phosphatase. kapp. type 2C protein phosphatase. |
| AAC36699.1     | AF075581 | Mesembryanthemum crystallinum | protein phosphatase-2C. PP2C.   |
| CAC09576.1     | AJ298988 | Fagus sylvatica               | protein phosphatase 2C (PP2C). pp2Cf2.  |
| SEQ ID NO: 777 |          |                               |   |
| AAD21872.1     | AF078082 | Phaseolus vulgaris            | receptor-like protein kinase homolog RK20-1.  |
| AAB93834.1     | U82481   | Zea mays                      | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.  |
| CAA74661.1     | Y14285   | Brassica oleracea             | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.  |
| AAD52097.1     | AF088885 | Nicotiana tabacum             | receptor-like kinase CHRK1. Chrkl.  |
| CAA73134.1     | Y12531   | Brassica oleracea             | serine/threonine kinase. BRLK.  |

|            |          |                             |  |
|------------|----------|-----------------------------|--|
| CAB41879.1 | Y18260   | Brassica oleracea           | SRK15 protein. SRK15. receptor-like kinase.  |
| CAB41878.1 | Y18259   | Brassica oleracea           | SRK5 protein. SRK5. receptor-like kinase.  |
| AAA33008.1 | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| CAB89179.1 | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| CAA67145.1 | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.  |
| AAA33000.1 | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| CAA73133.1 | Y12530   | Brassica oleracea           | serine /threonine kinase. ARLK.  |
| AAA62232.1 | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| AAC23542.1 | U20948   | Ipomoea trifida             | receptor protein kinase. IRK1.   |
| CAA74662.1 | Y14286   | Brassica oleracea           | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| BAA92837.1 | AB032474 | Brassica oleracea           | S60 S-locus receptor kinase. SRK60.  |
| BAA07577.2 | D38564   | Brassica rapa               | receptor protein kinase SRK12.   |
| BAA23676.1 | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.  |
| CAA79355.1 | Z18921   | Brassica oleracea           | S-receptor kinase-like protein.  |
| BAB21001.1 | AB054061 | Brassica rapa               | S locus receptor kinase. SRK22.  |
| BAA21132.1 | D88193   | Brassica rapa               | S-receptor kinase. SRK9 (B.c).   |
| BAA06285.1 | D30049   | Brassica rapa               | S-receptor kinase SRK9.  |
| BAA92836.1 | AB032473 | Brassica oleracea           | S18 S-locus receptor kinase. SRK18.  |
| BAA07576.1 | D38563   | Brassica rapa               | receptor protein kinase SRK8.  |
| BAB18292.1 | AP002860 | Oryza sativa                | putative receptor-like protein kinase. P0409B08.19.  |
| AAK21965.1 | AY028699 | Brassica napus              | receptor protein kinase PERK1.   |

|                |          |                            |  |
|----------------|----------|----------------------------|--|
| BAA92954.1     | AP001551 | Oryza sativa               | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).  |
| BAB39873.1     | AP002882 | Oryza sativa               | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| AAK00425.1     | AC069324 | Oryza sativa               | Putative protein kinase. OSJNBa0071K19.11.   |
| BAB39435.1     | AP003338 | Oryza sativa               | putative receptor kinase. OJ1212_B09.2.  |
| BAB07904.1     | AP002835 | Oryza sativa               | putative S-receptor kinase. P0417G05.12.   |
| BAA94518.1     | AP001800 | Oryza sativa               | Similar to Arabidopsis thaliana chromosome 2 section 111 of 255; putative receptor-like protein kinase (AC002392).               |
| AAG16628.1     | AY007545 | Brassica napus             | protein serine/threonine kinase BNK1.  |
| BAB39409.1     | AP002901 | Oryza sativa               | putative protein kinase. P0456F08.9. contains EST C23560(R0290).   |
| BAB16871.1     | AP002537 | Oryza sativa               | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).                       |
| AAF91324.1     | AF244890 | Glycine max                | receptor-like protein kinase 3. RLK3. GmRLK3.  |
| BAA82556.1     | AB030083 | Populus nigra              | lectin-like protein kinase. PnLPK.   |
| AAF91323.1     | AF244889 | Glycine max                | receptor-like protein kinase 2. RLK2. GmRLK2.  |
| AAF91322.1     | AF244888 | Glycine max                | receptor-like protein kinase 1. RLK1. GmRLK1.  |
| SEQ ID NO: 779 |          |                            |  |
| AAK31284.1     | AC079890 | Oryza sativa               | putative quinone oxidoreductase. OSJNBb0089A17.10.   |
| BAA78050.1     | AB027757 | Cicer arietinum            | NADPH oxidoreductase homolog.  |
| BAA83082.1     | AB030704 | Lithospermum erythrorhizon | LEDI-4 protein. LEDI-4. preferentially expressed in darkness; putative NADPH quinone oxidoreductase; similar to zeta-crystallin. |
| AAG53944.1     | AF304461 | Triphysaria versicolor     | quinone-oxidoreductase QR1. TvQR1.   |
| SEQ ID NO: 780 |          |                            |  |
| AAK17067.1     | AF254558 | Oryza sativa               | NAC6. NAC6.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| BAA89800.1     | AB028185 | Oryza sativa            | OsNAC6 protein. OsNAC6.   |
| BAA89799.1     | AB028184 | Oryza sativa            | OsNAC5 protein. OsNAC5.   |
| BAA89798.1     | AB028183 | Oryza sativa            | OsNAC4 protein. OsNAC4.   |
| BAA89797.1     | AB028182 | Oryza sativa            | OsNAC3 protein. OsNAC3.   |
| BAA89801.1     | AB028186 | Oryza sativa            | OsNAC7 protein. OsNAC7.   |
| AAF68626.1     | AF254124 | Medicago truncatula     | NAC1. NAC1. NAC domain containing protein.  |
| BAA89802.1     | AB028187 | Oryza sativa            | OsNAC8 protein. OsNAC8.   |
| BAA78417.1     | AB021178 | Nicotiana tabacum       | NAC-domain protein. TERN. elicitor-responsive gene.                                   |
| SEQ ID NO: 783 |          |                         |   |
| CAA54390.1     | X77134   | Brassica napus          | acyl-CoA binding protein.   |
| CAA70200.1     | Y08996   | Ricinus communis        | acyl-CoA-binding protein.   |
| CAB56693.1     | AJ249833 | Digitalis lanata        | binds medium- and long-chain acyl-coa esters. Acyl-CoA binding protein (ACBP). acbp3. |
| CAB56694.1     | AJ249834 | Digitalis lanata        | binds medium- and long-chain acyl-coa esters. Acyl-CoA binding protein (ACBP). acbp4. |
| AAB67736.1     | U35015   | Gossypium hirsutum      | acyl-CoA-binding protein.   |
| AAB86851.1     | AF031541 | Fritillaria agrestis    | acyl-CoA-binding protein. acabp.  |
| SEQ ID NO: 784 |          |                         |   |
| CAA58994.1     | X84208   | Sinapis alba            | trypsin inhibitor 2. mti-2.   |
| CAA76116.1     | Y16190   | Sinapis alba            | trypsin inhibitor 2. mti-2.   |
| SEQ ID NO: 785 |          |                         |   |
| AAF66242.1     | AF243180 | Lycopersicon esculentum | dicyanin. binuclear blue copper protein; contains two stellacyanins linked together.  |



|                |          |                                |  |
|----------------|----------|--------------------------------|--|
| AAC32421.1     | U65511   | Cucumis sativus                | putative oxygen activation and/or lignin formation. stellacyanin. member of a strictly plant-specific family of mononuclear blue copper proteins known as phytoeyanins, which are further classified into three distinct subfamilies: Uclacyanins, Stellacyanins, and Plantacyanins; member of the subfamily of Stellacyanins; similar to Rhus vernicifera stellacyanin: SwissProt Accession Number P00302; similar to umecyanin: SwissProt Accession Number P42849; similar to mavicyanin: SwissProt Accession Number P80728; similar to BCB encoded by GenBank Accession Number Z15058; similar to F18A8.9, encoded by GenBank Accession Number AC003105; similar to F7F1.27 encoded by GenBank Accession Number AC004669; similar to F9D12.16 encoded by GenBank Accession Number AF077407; glycoprotein. |
| CAA80963.1     | Z25471   | Pisum sativum                  | blue copper protein.   |
| AAD10251.1     | AF031195 | Triticum aestivum              | blue copper-binding protein homolog. S85.  |
| AAC64163.1     | AF093537 | Zea mays                       | blue copper protein. similar to pea blue copper protein in GenBank Accession Number Z25471.  |
| CAA10134.1     | AJ012693 | Cicer arietinum                | basic blue copper protein.   |
| AAF66243.1     | AF243181 | Lycopersicon esculentum        | plantacyanin. naturally occurring cupredoxin with a Val residue in the position of the axial ligand Met; member of the plantacyanin subfamily of a strictly plant-specific family of mononuclear blue copper proteins known as phytoeyanins.   |
| CAB65280.1     | AJ248323 | Medicago sativa subsp. x varia | basic blue protein. babl.  |
| AAC32448.1     | U76296   | Spinacia oleracea              | plantacyanin. member of a strictly plant-specific family of mononuclear blue copper proteins known as phytoeyanins, which are further classified into three distinct subfamilies: Uclacyanins, Stellacyanins, and Plantacyanins; member of the subfamily of Plantacyanins; non-glycosylated.   |
| SEQ ID NO: 793 |          |                                |  |
| BAA81862.1     | AB026295 | Oryza sativa                   | Similar to leucoanthocyanidin dioxygenase.(AI440611).  |
| AAB39995.1     | U82432   | Dianthus caryophyllus          | anthocyanidin synthase. allele: S; 2-oxoglutarat-dependent dioxygenase.  |
| BAA36554.1     | AB011796 | Citrus unshiu                  | flavonol synthase. CitFLS.   |
| AAD56580.1     | AF184273 | Daucus carota                  | leucoanthocyanidin dioxygenase 1. LDOX1. 2-oxoglutarate dependent dioxygenase.   |
| AAD56581.1     | AF184274 | Daucus carota                  | leucoanthocyanidin dioxygenase 2. LDOX. 2-oxoglutarate dependent dioxygenase.  |
| CAA50498.1     | X71360   | Malus sp.                      | anthocyanidin hydroxylase. apple equivalent to 'Candi' from Antirrhinum majus.   |

|                |          |                                 |   |
|----------------|----------|---------------------------------|---|
| AAD26205.1     | AF117269 | Malus x domestica               | anthocyanidin synthase. ANS.  |
| AAB82287.1     | AF026058 | Matthiola incana                | anthocyanidin synthase.   |
| CAA80264.1     | Z22543   | Petunia x hybrida               | flavonol synthase.  |
| AAF64168.1     | AF240764 | Eustoma grandiflorum            | flavonol synthase. fls.   |
| BAA20143.1     | AB003779 | Perilla frutescens              | leucoanthocyanidin dioxygenase.   |
| AAB66560.1     | AF015885 | Callistephus chinensis          | anthocyanidin synthase.   |
| BAB21477.1     | AB044091 | Torenia fournieri               | anthocyanidin synthase. 2-oxoglutarate dependent oxygenase.   |
| CAA63092.1     | X92178   | Solanum tuberosum               | flavonol synthase.  |
| AAD26261.1     | AF119095 | Malus x domestica               | flavonol synthase. FLS.   |
| CAA53580.1     | X75966   | Vitis vinifera                  | leucoanthocyanidin dioxygenase. LDOX.   |
| BAA75305.1     | AB023786 | Ipomoea batatas                 | anthocyanidin synthase. ans I.  |
| AAB84049.1     | AF028602 | Ipomoea purpurea                | anthocyanidin synthase. ANS-FL1.  |
| BAA75306.1     | AB023787 | Ipomoea batatas                 | anthocyanidin synthase. ans II.   |
| CAA73094.1     | Y12489   | Forsythia x intermedia          | anthocyanidin synthase.   |
| CAA69252.1     | Y07955   | Oryza sativa                    | anthocyanidin synthase. ANS.  |
| SEQ ID NO: 794 |          |                                 |   |
| AAD10204.1     | AF030260 | Vicia sativa                    | CYP94A1. vagh111. cytochrome P450 fatty acid hydroxylase; Method: conceptual translation with partial peptide sequencing. |
| AAG33645.1     | AF092917 | Vicia sativa                    | cytochrome P450-dependent fatty acid hydroxylase. CYP94A2.  |
| AAG17470.1     | AF123609 | Triticum aestivum               | cytochrome P450.  |
| BAA99523.1     | AP002484 | Oryza sativa                    | putative cytochrome P450. P0489A01.14.  |
| AAK31592.1     | AY029178 | Brassica rapa subsp. pekinensis | cytochrome P450. mf-CYP450. possible relevance to male-sterility.   |

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAA99522.1 | AP002484 | Oryza sativa         | putative cytochrome P450. P0489A01.13.  |
| BAA83370.1 | AP000391 | Oryza sativa         | ESTs AU056036(S20239),C72753(E2173), AU056035(S20239) correspond to a region of the predicted gene.; Similar to putative cytochrome P-450 (AC003680). |
| CAB41474.1 | AJ238402 | Catharanthus roseus  | cytochrome P450. CYP96C1.   |
| AAB94586.1 | AF022457 | Glycine max          | CYP97B2p. CYP97B2. cytochrome P450 monooxygenase.   |
| AAK20054.1 | AC025783 | Oryza sativa         | putative cytochrome P450 monooxygenase. OSJNBa0001014.16.   |
| AAK38086.1 | AF321862 | Lolium rigidum       | putative cytochrome P450.   |
| AAK38085.1 | AF321861 | Lolium rigidum       | putative cytochrome P450.   |
| AAB94588.1 | AF022459 | Glycine max          | CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.   |
| CAA89260.1 | Z49263   | Pisum sativum        | cytochrome P450.  |
| AAG09208.1 | AF175278 | Pisum sativum        | wound-inducible P450 hydroxylase. CYP82A1.  |
| AAC49188.2 | U29333   | Pisum sativum        | cytochrome P450 monooxygenase. CYP82. new cytochrome P450 family.   |
| BAB19083.1 | AP002744 | Oryza sativa         | putative cytochrome P450. P0006C01.25. contains ESTs AU081507(C12518),C26520(C12518).   |
| BAB19104.1 | AP002839 | Oryza sativa         | putative cytochrome P450. P0688A04.10. contains ESTs AU081507(C12518),C26520(C12518).   |
| AAK38092.1 | AF321868 | Lolium rigidum       | putative cytochrome P450.   |
| CAA04117.1 | AJ000478 | Helianthus tuberosus | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-race).   |
| AAK38091.1 | AF321867 | Lolium rigidum       | putative cytochrome P450.   |
| CAA04116.1 | AJ000477 | Helianthus tuberosus | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.   |
| BAA22423.1 | AB001380 | Glycyrrhiza echinata | cytochrome P450. CYP93B1.   |
| CAA71876.1 | Y10982   | Glycine max          | putative cytochrome P450.   |

|            |          |                              |  |
|------------|----------|------------------------------|--|
| BAB39252.1 | AP002968 | Oryza sativa                 | putative cytochrome P450. P0416G11.1.  |
| AAA33106.1 | L10081   | Catharanthus roseus          | cytochrome P-450 protein. CYP72. putative; CYP72 protein.  |
| AAA17746.1 | L19075   | Catharanthus roseus          | cytochrome P450. CYP72C. putative.   |
| BAA74466.1 | AB022733 | Glycyrrhiza echinata         | cytochrome P450. CYP Ge-51.  |
| BAB19121.1 | AP002839 | Oryza sativa                 | putative cytochrome P450. P0688A04.28.   |
| AAA17732.1 | L19074   | Catharanthus roseus          | cytochrome P450. CYP72B.   |
| BAA93634.1 | AB025016 | Lotus japonicus              | cytochrome P450.   |
| AAB05376.3 | U35226   | Nicotiana plumbaginifolia    | putative cytochrome P-450.   |
| AAB61965.1 | U48435   | Solanum chacoense            | putative cytochrome P450.  |
| AAF27282.1 | AF122821 | Capsicum annuum              | cytochrome P450. PepCYP.   |
| CAB50768.1 | AJ243804 | Cicer arietinum              | putative isoflavone synthase. cytochrome P450. cyp93C3.  |
| AAC34853.1 | AF082028 | Hemerocallis hybrid cultivar | putative cyt P450-containing fatty acid hydroxylase. senescence-associated protein 3. SA3. mRNA accumulates in senescing petals. |
| CAB43505.1 | AJ239051 | Cicer arietinum              | cytochrome P450. cyp81E2.  |
| BAB21156.1 | AP002899 | Oryza sativa                 | putative cytochrome P450. P0456A01.12.   |
| CAA50648.1 | X71657   | Solanum melongena            | P450 hydroxylase.  |
| BAB19112.1 | AP002839 | Oryza sativa                 | putative cytochrome P450. P0688A04.18. contains ESTs AU067870(C10320),AU067869(C10320).  |
| BAB19091.1 | AP002744 | Oryza sativa                 | putative cytochrome P450. P0006C01.33. contains ESTs AU067870(C10320),AU067869(C10320).  |
| BAA84072.1 | AB028152 | Torenia hybrida              | flavone synthase II. cytochrome P450. TFNS5.   |
| BAA76380.1 | AB023636 | Glycyrrhiza echinata         | cytochrome P450. CYP Ge-8.   |
| CAA72208.1 | Y11404   | Zea mays                     | cytochrome p450. cyp71c2.  |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| CAA57423.1     | X81829   | Zea mays                | cytochrome P450. CYP71C2. family CYP71, subfamily CYP71C.  |
| BAB12433.1     | AB025030 | Coptis japonica         | p450.  |
| SEQ ID NO: 798 |          |                         |  |
| BAB21205.1     | AP002913 | Oryza sativa            | nucleoid DNA-binding protein cnd41-like protein. P0480E02.11. contains ESTs AU166073(E31027),AU029516(E31027).   |
| BAA22813.1     | D26015   | Nicotiana tabacum       | aspartic protease activity. CND41, chloroplast nucleoid DNA binding protein. cnd41.  |
| SEQ ID NO: 804 |          |                         |  |
| AAD46491.1     | AF135014 | Zea mays                | dihydrolipoamide S-acetyltransferase. pyruvate dehydrogenase complex E2 subunit.   |
| SEQ ID NO: 805 |          |                         |  |
| BAA85412.1     | AP000615 | Oryza sativa            | ESTs AU065232(E60855),C23624(S1554), AU078241(E60855) correspond to a region of the predicted gene.; similar to putative adenylate kinase. (AC005896). |
| BAA01181.1     | D10335   | Oryza sativa            | adenylate kinase-b. Adk-b.   |
| BAA01180.1     | D10334   | Oryza sativa            | adenylate kinase-a. Adk-a.   |
| BAA94761.1     | AB041773 | Oryza sativa            | adenylate kinase. Adk-a.   |
| AAB68604.1     | U82330   | Prunus armeniaca        | adenylate kinase homolog.  |
| AAF23372.1     | AF187063 | Oryza sativa            | catalyzes the transfer of phosphate from ATP to UMP or CMP to form ADP and UDP or CDP. UMP/CMP kinase b. ura6.   |
| AAF23371.1     | AF187062 | Oryza sativa            | catalyzes the transfer of phosphate from ATP to UMP or CMP to form ADP and UDP or CDP. UMP/CMP kinase a. ura6.   |
| AAD41679.1     | AF086603 | Ceratopteris richardii  | adenylate kinase. ADK1.  |
| BAA85443.1     | AP000616 | Oryza sativa            | similar to UMP/CMP kinase (AF000147).  |
| SEQ ID NO: 808 |          |                         |  |
| CAA41774.1     | X59046   | Oryza sativa            | sucrose-UDP glucosyltransferase (isoenzyme 2). RSs2.   |
| BAA89049.1     | AB029401 | Citrus unshiu           | sucrose synthase. CitSUS1-2.   |
| AAA34196.1     | L19762   | Lycopersicon esculentum | sucrose synthase.  |

|  |          |                         |
|--|----------|-------------------------|
| BAA88905.1   | AB022092 | Citrus unshiu           |
| sucrose synthase. CitSUS1.   |          |                         |
| AAD28641.1   | U73588   | Gossypium hirsutum      |
| sucrose synthase.  |          |                         |
| CAA49428.1   | X69773   | Vicia faba              |
| sucrose synthase. VfSucs.  |          |                         |
| AAC37346.1   | M97551   | Vicia faba              |
| cleavage of sucrose. UDP-glucose:D-fructose-2-glucosyltransferase. putative. |          |                         |
| CAA09681.1   | AJ011535 | Lycopersicon esculentum |
| sucrose synthase. sus2.  |          |                         |
| AAA97572.1   | U24088   | Solanum tuberosum       |
| sucrose synthase.  |          |                         |
| CAA09593.1   | AJ011319 | Lycopersicon esculentum |
| sucrose synthase. sus3.  |          |                         |
| CAB40794.1   | AJ131943 | Medicago truncatula     |
| sucrose synthase. sucS1.   |          |                         |
| AAC17867.1   | AF049487 | Medicago sativa         |
| sucrose hydrolysis. sucrose synthase.  |          |                         |
| CAB40795.1   | AJ131964 | Medicago truncatula     |
| sucrose synthase. sucS1.   |          |                         |
| CAA65640.1   | X96939   | Tulipa gesneriana       |
| sucrose-synthase 21.   |          |                         |
| AAA97571.1   | U24087   | Solanum tuberosum       |
| sucrose synthase.  |          |                         |
| CAA63122.1   | X92378   | Alnus glutinosa         |
| sucrose synthase. sus1.  |          |                         |
| AAA33514.1   | L22296   | Zea mays                |
| UDP-glucose:D-fructose 2-glucosyl-transferase. Sus1.                         |          |                         |
| CAA65639.1   | X96938   | Tulipa gesneriana       |
| sucrose-synthase 1.  |          |                         |
| AAC41682.1   | L03366   | Oryza sativa            |
| sucrose synthase 3. RSs3.  |          |                         |
| CAA75793.1   | Y15802   | Hordeum vulgare         |
| sucrose synthase 2. Ss2.   |          |                         |
| CAA49551.1   | X69931   | Hordeum vulgare         |
| sucrose synthase. Ss2.   |          |                         |
| CAA76056.1   | Y16090   | Daucus carota           |
| sucrose synthase isoform I. Susy*Dc1.  |          |                         |
| CAA53081.1   | X75332   | Daucus carota           |
| sucrose synthase.  |          |                         |
| AAA33515.1   | L33244   | Zea mays                |
| sucrose synthase 2. Sus1.  |          |                         |

|            |          |                            |   |
|------------|----------|----------------------------|---|
| BAB20799.1 | AB045710 | Pyrus pyrifolia            | sucrose synthase 1. PypSUS1.                          |
| CAA03935.1 | AJ000153 | Triticum aestivum          | sucrose synthase type 2.                              |
| AAC39323.1 | AF030231 | Glycine max                | sucrose synthase. SS. nodulin-100.                    |
| BAA01108.1 | D10266   | Vigna radiata              | sucrose synthase. vss1.                               |
| CAA09910.1 | AJ012080 | Pisum sativum              | sucrose synthase.                                     |
| AAC28107.1 | AF079851 | Pisum sativum              | nodule-enhanced sucrose synthase. ness.               |
| CAC32462.1 | AJ311496 | Pisum sativum              | sucrose metabolism. sucrose synthase isoform 3. sus3. |
| CAA57881.1 | X82504   | Chenopodium rubrum         | sucrose synthase. CSS1.                               |
| CAA26229.1 | X02382   | Zea mays                   | sucrose synthase.                                     |
| CAA26247.1 | X02400   | Zea mays                   | sucrose synthase.                                     |
| CAA46017.1 | X64770   | Oryza sativa               | sucrose synthase. RSs1.                               |
| CAB38022.1 | AJ132000 | Craterostigma plantagineum | sucrose metabolism. sucrose synthase. Ss2.            |
| CAA78747.1 | Z15028   | Oryza sativa               | sucrose synthase.                                     |
| AAF85966.1 | AF263384 | Saccharum officinarum      | sucrose synthase-1.                                   |
| CAA46701.1 | X65871   | Hordeum vulgare            | sucrose synthase.                                     |
| CAA04543.1 | AJ001117 | Triticum aestivum          | sucrose synthase type I. Ss1.                         |
| BAA88904.1 | AB022091 | Citrus unshiu              | sucrose synthase. CitSUSA.                            |
| BAA88981.1 | AB025778 | Citrus unshiu              | sucrose synthase. CitSUSA-2.                          |
| CAA04512.1 | AJ001071 | Pisum sativum              | second sucrose synthase.                              |
| CAA76057.1 | Y16091   | Daucus carota              | sucrose synthase isoform II. Susy*Dc2.                |
| CAB38021.1 | AJ131999 | Craterostigma plantagineum | sucrose metabolism. sucrose synthase. Ss1.            |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| CAA57499.1     | X81974   | Beta vulgaris                 | sucrose synthase. SBSS1.                                  |
| CAA47264.1     | X66728   | Hordeum vulgare               | sucrose synthase.   |
| SEQ ID NO: 809 |          |                               |   |
| AAB69317.1     | AF012861 | Petroselinum crispum          | plastidic glucose-6-phosphate dehydrogenase. pG6PDH.      |
| AAF87216.1     | AF231351 | Nicotiana tabacum             | plastidic glucose 6-phosphate dehydrogenase. G6PDHP2.     |
| CAA67782.1     | X99405   | Nicotiana tabacum             | glucose-6-phosphate dehydrogenase. G6PD.                  |
| CAB52708.1     | AJ010712 | Solanum tuberosum             | glucose-6-phosphate 1-dehydrogenase. g6pd.                |
| CAB52685.1     | AJ132346 | Dunaliella bioculata          | plastidic glucose-6-phosphate dehydrogenase. g6PD.        |
| CAA58775.1     | X83923   | Solanum tuberosum             | glucose-6-phosphate dehydrogenase.                        |
| CAA03941.1     | AJ000184 | Spinacia oleracea             | Glucose-6-phosphate dehydrogenase. G6PD.                  |
| CAA03939.1     | AJ000182 | Spinacia oleracea             | Glucose-6-phosphate dehydrogenase. G6PD.                  |
| CAA04994.1     | AJ001772 | Nicotiana tabacum             | glucose-6-phosphate dehydrogenase. TCG18.                 |
| CAA03940.1     | AJ000183 | Spinacia oleracea             | Glucose-6-phosphate dehydrogenase. G6PD.                  |
| AAD11426.1     | AF097663 | Mesembryanthemum crystallinum | cytoplasmic glucose-6-phosphate 1-dehydrogenase. G6PD.    |
| AAB41552.1     | U18238   | Medicago sativa subsp. sativa | glucose-6-phosphate dehydrogenase.                        |
| CAA52442.1     | X74421   | Solanum tuberosum             | glucose-6-phosphate 1-dehydrogenase. g6pdh. homotetramer. |
| AAB69318.1     | AF012862 | Petroselinum crispum          | cytosolic glucose-6-phosphate dehydrogenase 1. cG6PDH1.   |
| AAB69319.1     | AF012863 | Petroselinum crispum          | cytosolic glucose-6-phosphate dehydrogenase 2. cG6PDH2.   |
| CAA04992.1     | AJ001769 | Nicotiana tabacum             | glucose-6-phosphate dehydrogenase. TCG6.                  |
| CAA04993.1     | AJ001770 | Nicotiana tabacum             | glucose-6-phosphate dehydrogenase. TCG9.                  |
| BAA97662.1     | AB029454 | Triticum aestivum             | glucose-6-phosphate dehydrogenase. g6pdh.                 |



|                |          |                           |   |
|----------------|----------|---------------------------|---|
| BAA97663.1     | AB029455 | Triticum aestivum         | glucose-6-phosphate dehydrogenase. g6pdh.   |
| BAA97664.1     | AB029456 | Triticum aestivum         | glucose-6-phosphate dehydrogenase. g6pdh.   |
| AAG23802.1     | AF260736 | Cucurbita pepo            | plastidic glucose-6-phosphate dehydrogenase.  |
| CAB66330.1     | AJ279688 | Betula pendula            | glucose-6-phosphate dehydrogenase. g6pd.  |
| BAA82155.1     | AB011441 | Triticum aestivum         | glucose-6-phosphate dehydrogenase. WESR5. salt-stress responding gene.  |
| CAA06200.1     | AJ004900 | Glycine max               | pentose phosphate pathway oxidoreductase generating NADPH. glucose-6-phosphate-dehydrogenase.                               |
| SEQ ID NO: 812 |          |                           |   |
| BAA08910.1     | D50407   | Cucumis sativus           | glutamyl-tRNA reductase. hemA.  |
| AAD16897.1     | AF105221 | Glycine max               | converts glutamyl-tRNA to glutamate 1-semialdehyde. glutamyl-tRNA reductase precursor. gtr1.                                |
| BAA11091.1     | D67088   | Cucumis sativus           | glutamyl-tRNA reductase. hemA2.   |
| BAA25003.1     | AB011416 | Oryza sativa              | glutamyl-tRNA reductase.  |
| AAG13620.1     | AC078840 | Oryza sativa              | putative glutamyl-tRNA reductase. OSJNBb0073N24.1.  |
| CAA60054.1     | X86101   | Hordeum vulgare           | aminolevulinate synthesis (chlorophyll synthesis). glutamyl tRNA reductase. hemA 1. 1st isoform.                            |
| CAA63140.1     | X92403   | Hordeum vulgare           | glutamyl-tRNA reductase. hemA1. isoform I.  |
| BAA25167.1     | D88382   | Hordeum vulgare           | glutamyl-tRNA reductase. hemA1. isoform 1.  |
| CAA60055.1     | X86102   | Hordeum vulgare           | aminolevulinate synthesis (chlorophyll synthesis). glutamyl tRNA reductase. hemA 2. 2nd isoform.                            |
| BAA25168.1     | D88383   | Hordeum vulgare           | glutamyl-tRNA reductase. hemA3. isoform 3.  |
| AAG41962.1     | AF305613 | Chlamydomonas reinhardtii | glutamyl-tRNA reductase precursor. HemA. pGtr.  |
| AAG02480.1     | AF294753 | Hordeum vulgare           | converts glutamyl-tRNA to glutamate 1-semialdehyde in the chlorophyll biosynthetic pathway. glutamyl-tRNA reductase. hemA2. |

|                |          |                              |   |
|----------------|----------|------------------------------|---|
| AAG02479.1     | AF294752 | Hordeum vulgare              | converts glutamyl-tRNA to glutamate 1-semialdehyde in the chlorophyll biosynthetic pathway. glutamyl-tRNA reductase. hemA1. |
| SEQ ID NO: 813 |          |                              |   |
| BAA82556.1     | AB030083 | Populus nigra                | lectin-like protein kinase. PnLPK.  |
| AAD21872.1     | AF078082 | Phaseolus vulgaris           | receptor-like protein kinase homolog RK20-1.  |
| AAF43408.1     | AF230515 | Oryza sativa subsp. japonica | serine/threonine protein kinase. YK35.  |
| CAB51480.1     | Y14600   | Sorghum bicolor              | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.  |
| CAA73134.1     | Y12531   | Brassica oleracea            | serine/threonine kinase. BRLK.  |
| AAB93834.1     | U82481   | Zea mays                     | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.                        |
| AAK00425.1     | AC069324 | Oryza sativa                 | Putative protein kinase. OSJNBa0071K19.11.  |
| BAA92954.1     | AP001551 | Oryza sativa                 | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).   |
| BAB39873.1     | AP002882 | Oryza sativa                 | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).                                       |
| BAB19337.1     | AP003044 | Oryza sativa                 | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).                                      |
| AAK21965.1     | AY028699 | Brassica napus               | receptor protein kinase PERK1.  |
| BAA92953.1     | AP001551 | Oryza sativa                 | Similar to Arabidopsis thaliana chromosome 4 BAC clone F10M6 ; S-receptor kinase -like protein. (AL021811).                 |
| BAB07906.1     | AP002835 | Oryza sativa                 | putative S-receptor kinase. P0417G05.14.  |
| BAA94516.1     | AP001800 | Oryza sativa                 | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| AAD52097.1     | AF088885 | Nicotiana tabacum            | receptor-like kinase CHRK1. Chrk1.  |
| BAB21240.1     | AP002953 | Oryza sativa                 | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| AAC23542.1     | U20948   | Ipomoea trifida              | receptor protein kinase. IRK1.  |

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAB18292.1 | AP002860 | Oryza sativa         | putative receptor-like protein kinase. P0409B08.19.   |
| AAG16628.1 | AY007545 | Brassica napus       | protein serine/threonine kinase BNK1.   |
| AAA33915.1 | L27821   | Oryza sativa         | receptor type serine/threonine kinase. protein kinase.  |
| BAB03429.1 | AP002817 | Oryza sativa         | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587). |
| BAB07999.1 | AP002525 | Oryza sativa         | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAA94509.1 | AB041503 | Populus nigra        | protein kinase 1. PnPK1.  |
| AAD46420.1 | AF100771 | Hordeum vulgare      | receptor-like kinase. Hv3ARK. similar to wheat ARK1AS.  |
| BAA23676.1 | AB000970 | Brassica rapa        | receptor kinase 1. BcRK1.   |
| AAD38286.1 | AC007789 | Oryza sativa         | putative protein kinase. OSJNBa0049B20.13.  |
| BAB40081.1 | AP003074 | Oryza sativa         | putative receptor protein kinase. OSJNBa0004G10.30.   |
| BAB18321.1 | AP002865 | Oryza sativa         | putative receptor protein kinase. P0034C11.11.  |
| BAA94517.1 | AP001800 | Oryza sativa         | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| BAA21132.1 | D88193   | Brassica rapa        | S-receptor kinase. SRK9 (B.c).  |
| BAA06285.1 | D30049   | Brassica rapa        | S-receptor kinase SRK9.   |
| BAB07905.1 | AP002835 | Oryza sativa         | putative S-receptor kinase. P0417G05.13.  |
| BAA94529.2 | AP001800 | Oryza sativa         | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| AAB61708.1 | U93048   | Daucus carota        | somatic embryogenesis receptor-like kinase. SERK.   |
| CAA79355.1 | Z18921   | Brassica oleracea    | S-receptor kinase-like protein.   |
| AAK11674.1 | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| AAF43496.1 | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |

|                |          |                             |  |
|----------------|----------|-----------------------------|--|
| BAA94528.1     | AP001800 | Oryza sativa                | Similar to Arabidopsis thaliana chromosome 2 BAC T20K24; putative receptor-like protein kinase (AC002392). |
| BAA94510.1     | AB041504 | Populus nigra               | protein kinase 2. PnPK2.   |
| AAA33000.1     | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| CAA67145.1     | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.  |
| AAB47421.1     | U59316   | Lycopersicon esculentum     | serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato. |
| BAA07577.2     | D38564   | Brassica rapa               | receptor protein kinase SRK12.   |
| AAA33008.1     | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| CAB89179.1     | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| BAA92837.1     | AB032474 | Brassica oleracea           | S60 S-locus receptor kinase. SRK60.  |
| SEQ ID NO: 814 |          |                             |  |
| AAF23903.1     | AF194416 | Oryza sativa                | MAP kinase homolog. MAPK2. RMAPK2.   |
| AAD52659.1     | AF177392 | Oryza sativa                | blast and wounding induced mitogen-activated protein kinase. BWMK1. BWMK1 MAP kinase.                      |
| AAF23902.1     | AF194415 | Oryza sativa                | MAP kinase homolog. MAPK1. RMAPK1.   |
| AAD28617.1     | AF129087 | Medicago sativa             | mitogen-activated protein kinase homologue. TDY1.  |
| CAB61750.1     | AJ275316 | Cicer arietinum             | MAP kinase protein.  |
| AAB57843.1     | U96716   | Selaginella lepidophylla    | MAP kinase-like protein. sdhn-6r.  |
| AAF65766.1     | AF242308 | Euphorbia esula             | mitogen-activated protein kinase. regulated by tyrosine and threonine phosphorylation.                     |
| BAB18271.1     | AB035141 | Chlamydomonas reinhardtii   | mitogen-activated protein kinase. CrMPK2.  |
| CAA58761.1     | X83880   | Nicotiana tabacum           | p45Ntf4 serine/threonine protein kinase. ntf4.   |
| CAA47099.1     | X66469   | Medicago sativa             | MAP Kinase. MSK7.  |

|            |          |                      |   |
|------------|----------|----------------------|---|
| AAB41548.1 | L07042   | Medicago sativa      | autophosphorylating serine/threonine protein kinase. MAP kinase. MsERK1.      |
| AAB58396.1 | U94192   | Nicotiana tabacum    | salicylic acid-activated MAP kinase. NtSIPK.                                  |
| CAA50036.1 | X70703   | Pisum sativum        | MAP kinase homologue. PSMAPKIN.   |
| AAF73236.1 | AF153061 | Pisum sativum        | MAP kinase 3. Mapk3. PsMAPK3.   |
| BAB32406.1 | AB055515 | Nicotiana tabacum    | NRK1 MAPK. nrk1. A tobacco MAPK that is phosphorylated and activated by NQK1. |
| CAA58760.1 | X83879   | Nicotiana tabacum    | p43Ntf6 serine/threonine protein kinase. ntf6.                                |
| CAA57721.1 | X82270   | Medicago sativa      | protein kinase. MMK4.   |
| AAF81420.1 | AF247136 | Capsicum annuum      | MAP kinase 2. MK2. CAMK2; wound, UV-C, and cold-inducible expression.         |
| AAD37790.1 | AF149424 | Ipomoea batatas      | MAP kinase.   |
| AAG40580.1 | AF216316 | Oryza sativa         | MAP kinase 2. protein kinase; MAP2.   |
| CAB37188.1 | AJ224336 | Medicago sativa      | MAP kinase. MMK3.   |
| AAF61238.1 | AF241166 | Oryza sativa         | MAP kinase MAPK2.   |
| AAG40581.1 | AF216317 | Oryza sativa         | MAP kinase 3. protein kinase; MAP3.   |
| CAB61889.1 | AJ251330 | Oryza sativa         | protein kinase. MAPK4 protein. mapk4.   |
| CAA73323.1 | Y12785   | Petroselinum crispum | MAP kinase I.   |
| CAC13967.1 | AJ250311 | Oryza sativa         | protein kinase. MAPK2 protein. mapk2.   |
| CAA56314.1 | X79993   | Avena sativa         | MAP KINASE. Asmap1.   |
| CAA49592.1 | X69971   | Nicotiana tabacum    | serine/threonine protein kinase. NTF3.  |
| CAA58466.1 | X83440   | Petunia x hybrida    | MAP/ERK kinase 1. MEK1.   |
| AAK01710.1 | AF332873 | Oryza sativa         | MAP kinase BIMK1.   |
| AAG40579.1 | AF216315 | Oryza sativa         | MAP kinase 1. protein kinase; MAP1.   |

|                |          |                   |  |
|----------------|----------|-------------------|--|
| CAA57719.1     | X82268   | Medicago sativa   | protein kinase. MMK2.  |
| AAC28850.1     | AF079318 | Triticum aestivum | protein kinase. MAP kinase homolog. WCK-1.   |
| AAD32204.1     | AF134730 | Prunus armeniaca  | putative mitogen-activated protein kinase MAPK. MAP kinase.  |
| BAA74734.1     | AB016802 | Zea mays          | MAP kinase 5. ZmMPK5.  |
| AAF73257.1     | AF154329 | Pisum sativum     | MAP kinase PsMAPK2. Mapk2.   |
| BAA09600.1     | D61377   | Nicotiana tabacum | WIPK. MAP (mitogen-activated protein) kinase.  |
| AAF81419.1     | AF247135 | Capsicum annuum   | MAP kinase 1. MK1. wound and UV-C inducible expression.  |
| BAA74733.1     | AB016801 | Zea mays          | MAP kinase 4. ZmMPK4.  |
| CAA05328.1     | AJ002314 | Nicotiana tabacum | serine/threonine protein kinase. shaggy-like kinase 111. NSK 111.  |
| CAA05329.1     | AJ002315 | Nicotiana tabacum | shaggy-like kinase 59. NSK 59.   |
| CAA11861.1     | AJ224164 | Petunia x hybrida | shaggy kinase 6. PSK6.   |
| CAA58595.1     | X83620   | Petunia x hybrida | Petunia Shaggy kinase 6. PSK6.   |
| CAA11862.1     | AJ224165 | Petunia x hybrida | shaggy kinase 7. PSK7.   |
| CAA58594.1     | X83619   | Petunia x hybrida | Petunia Shaggy kinase 4. PSK4.   |
| AAA92823.1     | U18365   | Brassica napus    | cyclin dependent protein kinase homolog; similar to moth bean p34cdc2 protein, PIR<br>Accession Number JQ2243.   |
| BAA92214.1     | AP001278 | Oryza sativa      | ESTs C22403(C50132),C22404(C50132) correspond to a region of the predicted gene.<br>Similar to Arabidopsis thaliana shaggy related protein kinase ASK-gamma. (P43289). |
| CAA67554.1     | X99100   | Trifolium repens  | protein kinase. trK.   |
| SEQ ID NO: 816 |          |                   |  |
| CAA65065.1     | X95759   | Solanum tuberosum | glycogen (starch) synthase.  |
| CAA64173.1     | X94400   | Solanum tuberosum | soluble-starch-synthase. SSSIII.   |

|            |          |                           |  |
|------------|----------|---------------------------|--|
| AAC14014.1 | AF023159 | Zea mays                  | starch synthase DULL1. dull1. similar to potato starch synthase SSIII; likely to be the maize starch synthase defined biochemically as SSII. |
| CAB40374.1 | AJ225088 | Vigna unguiculata         | ADP-glucose-starch glucosyltransferase. Starch synthase isoform SS III.  |
| AAF88000.1 | AF258609 | Aegilops tauschii         | starch synthase III.   |
| AAF87999.1 | AF258608 | Triticum aestivum         | starch synthase III. wSSIII.   |
| CAB40375.1 | AJ006752 | Vigna unguiculata         | ADP-glucose starch glucosyltransferase. starch synthase, isoform V.  |
| AAC14015.1 | AF023160 | Zea mays                  | starch synthase DULL1. dull1. similar to potato starch synthase SSIII; like to be the maize starch synthase defined biochemically as SSII.   |
| AAC17971.2 | AF026422 | Chlamydomonas reinhardtii | soluble starch synthase. ADP-glucose:alpha-1, 4-D-glucan-4-alpha-D-glucosyltransferase.  |
| AAD13342.1 | AF019297 | Zea mays                  | starch synthase isoform zSTSII-2. zSSIIb.  |
| CAB86618.1 | AJ269502 | Triticum aestivum         | transfers the glucosyl unit of ADP glucose to the non-reducing end of an alpha 1,4. starch synthase IIa-1. wSs2a-1.                          |
| CAA61269.1 | X88790   | Pisum sativum             | glycogen (starch) synthase.  |
| CAB96626.1 | AJ269503 | Triticum aestivum         | transfers the glucosyl unit of ADP glucose to the non-reducing end of an alpha 1,4. starch synthase IIa-2. wSs2a-2.                          |
| CAB96627.1 | AJ269504 | Triticum aestivum         | transfers the glucosyl unit of ADP glucose to the non-reducing end of an alpha 1,4. starch synthase IIa-3. wSs2a-3.                          |
| CAA71442.1 | Y10416   | Solanum tuberosum         | soluble starch (bacterial glycogen) synthase. SS I.  |
| AAD53263.1 | AF155217 | Triticum aestivum         | starch synthase IIa.   |
| AAF37876.1 | AF234163 | Hordeum vulgare           | starch synthase I. SSI.  |
| CAB99209.1 | AJ292521 | Triticum aestivum         | essential for starch synthesis. starch synthase I-1. wSsI-1.   |
| AAD54661.1 | AF091803 | Triticum aestivum         | starch synthase I.   |
| AAB17085.1 | U66377   | Triticum aestivum         | starch synthase. TaSS. EC 2.4.1.11.  |
| CAB99210.1 | AJ292522 | Triticum aestivum         | essential for starch synthesis. starch synthase I-2. wSsI-2.   |

|            |          |   |
|------------|----------|---|
| AAF03557.1 | AF091802 | <i>Aegilops tauschii</i><br>starch synthase I.  |
| AAD13341.1 | AF019296 | <i>Zea mays</i><br>starch synthase isoform zSTSII-1. zSSIIa.  |
| AAC17969.2 | AF026420 | <i>Chlamydomonas reinhardtii</i><br>ADP-Glucose:alpha-1, 4-D-glucan-4-alpha-D-glucosyltransferase. granule-bound starch synthase I precursor. STA2. GBSSI.                                    |
| BAA82346.1 | AB029546 | <i>Phaseolus vulgaris</i><br>granule-bound starch synthase I. GBSSI.  |
| CAA37732.1 | X53694   | <i>Oryza sativa</i><br>starch synthase.   |
| CAA52273.1 | X74160   | <i>Manihot esculenta</i><br>starch (bacterial glycogen) synthase. GBSS.   |
| AAF72561.1 | AF141954 | <i>Oryza sativa</i><br>granule-bound starch synthase. Waxy.   |
| CAA46294.1 | X65183   | <i>Oryza sativa</i><br>glycogen (starch) synthase. waxy gene. starch granule enzyme.  |
| AAF72562.1 | AF141955 | <i>Oryza sativa</i><br>granule-bound starch synthase. Waxy.   |
| CAA44065.1 | X62134   | <i>Oryza sativa</i><br>starch biosynthesis. starch (bacterial glycogen) synthase. Wx.   |
| AAB02197.1 | U48227   | <i>Triticum aestivum</i><br>soluble starch synthase.  |
| AAF13168.1 | AF173900 | <i>Manihot esculenta</i><br>granule bound starch synthase II precursor. GBSSII. MEGBSSII.   |
| CAA45472.1 | X64108   | <i>Oryza sativa</i><br>starch granule-bound starch synthase. waxy.  |
| AAC61675.2 | AF031162 | <i>Oryza sativa</i><br>granule-bound starch synthase. Waxy.   |
| AAC70779.1 | AF097922 | <i>Astragalus membranaceus</i><br>granule-bound glycogen (starch) synthase. GBSS.   |
| CAA06958.1 | AJ006293 | <i>Antirrhinum majus</i><br>granule-bound starch synthase. waxy.  |
| AAC19119.1 | AF068834 | <i>Ipomoea batatas</i><br>starch synthase.  |
| AAD49850.1 | AF165890 | <i>Oryza sativa</i> subsp. japonica<br>soluble starch synthase.   |
| BAA81848.1 | AB026295 | <i>Oryza sativa</i><br>ESTs AU075322(C11109),D22430(C11109) correspond to a region of the predicted gene.;<br>Rice gene for soluble starch synthase (SSS1), complete cds (exon1-15).(D38221). |
| BAA03739.1 | D16202   | <i>Oryza sativa</i><br>soluble starch synthase precursor.   |



|                |          |                           |   |
|----------------|----------|---------------------------|---|
| CAA61268.1     | X88789   | Pisum sativum             | glycogen (starch) synthase.   |
| AAA86423.1     | U44126   | Ipomoea batatas           | starch synthase. SPSS67.  |
| AAF14233.1     | AF109395 | Triticum aestivum         | granule-bound starch synthase GBSSII.   |
| CAA41359.1     | X58453   | Solanum tuberosum         | glycogen (starch) synthase. amf. waxy protein, granule-bound starch synthase. |
| AAG43519.1     | AF210699 | Perilla frutescens        | granule-bound starch synthase. GBSSI. waxy protein.                           |
| SEQ ID NO: 819 |          |                           |   |
| BAA13032.1     | D86180   | Pisum sativum             | phosphoribosylanthranilate transferase. PAT1.                                 |
| SEQ ID NO: 822 |          |                           |   |
| AAB86850.1     | AF031540 | Fritillaria agrestis      | cytochrome C. cytC.   |
| AAC84135.1     | AF101422 | Cichorium intybus         | cytochrome.   |
| BAA02159.1     | D12634   | Oryza sativa              | 'cytochrome C'.   |
| AAA63515.1     | M63704   | Oryza sativa              | cytochrome c. Cc-1.   |
| AAA92712.1     | L77113   | Helianthus annuus         | cytochrome c. cytcl. putative.  |
| AAB70265.1     | AF017367 | Oryza sativa              | cytochrome C.   |
| AAA33084.1     | M35173   | Chlamydomonas reinhardtii | apocytochrome c (cyc).  |
| CAB16954.1     | Z99829   | Chlamydomonas reinhardtii | cytochrome c. CYC1.   |
| CAA79708.1     | Z21499   | Stellaria longipes        | mitochondrial cytochrome c.   |
| SEQ ID NO: 823 |          |                           |   |
| BAA02159.1     | D12634   | Oryza sativa              | 'cytochrome C'.   |
| AAA63515.1     | M63704   | Oryza sativa              | cytochrome c. Cc-1.   |
| AAB86850.1     | AF031540 | Fritillaria agrestis      | cytochrome C. cytC.   |
| AAC84135.1     | AF101422 | Cichorium intybus         | cytochrome.   |

|                |          |                                   |   |
|----------------|----------|-----------------------------------|---|
| AAA92712.1     | L77113   | Helianthus annuus                 | cytochrome c. cytcl. putative.  |
| AAB70265.1     | AF017367 | Oryza sativa                      | cytochrome C.   |
| AAA33084.1     | M35173   | Chlamydomonas reinhardtii         | apocytochrome c (cyc).  |
| CAB16954.1     | Z99829   | Chlamydomonas reinhardtii         | cytochrome c. CYC1.   |
| CAA79708.1     | Z21499   | Stellaria longipes                | mitochondrial cytochrome c.   |
| SEQ ID NO: 825 |          |                                   |   |
| BAB17113.1     | AP002866 | Oryza sativa                      | putative white protein; ATP-binding cassette transporter. P0410E01.34.                                    |
| AAF43869.1     | AF166114 | Chloroplast Mesostigma viride     | probable transport protein. cysA.   |
| BAA90508.1     | AP001111 | Oryza sativa                      | similar to ABC transporter of Arabidopsis thaliana (AC004697).  |
| AAD54843.1     | AF137379 | Chloroplast Nephroselmis olivacea | probable transport protein. cysA.   |
| BAA57907.1     | AB001684 | Chlorella vulgaris                | sulfate transport system permease protein. cysA.  |
| BAA90507.1     | AP001111 | Oryza sativa                      | similar to ABC transporter of Arabidopsis thaliana (AC004697).  |
| BAB40032.1     | AP003046 | Oryza sativa                      | putative ABC transporter. P0445D12.3.   |
| AAG49003.1     | AY013246 | Hordeum vulgare                   | putative ABC transporter. 635P2.4b; GC splice donor confirmed by cDNA alignment and comparative sequence. |
| BAB21275.1     | AP002844 | Oryza sativa                      | putative ABC transporter protein. P0410E03.6.   |
| AAG45492.1     | AY013245 | Oryza sativa                      | 36I5.4. putative ABC transporter; GC splice donor confirmed by cDNA alignment and comparative sequence.   |
| AAG49002.1     | AY013246 | Hordeum vulgare                   | putative ABC transporter. 635P2.4a; GC splice donor confirmed by cDNA and comparative sequencing.         |
| BAB21276.1     | AP002844 | Oryza sativa                      | putative ABC transporter protein. P0410E03.7. contains EST D22472(C1173).                                 |
| AAD10836.1     | U52079   | Solanum tuberosum                 | P-glycoprotein. pmdr1. binds ATP; ATPase; transporter; transmembrane protein.                             |
| BAB21279.1     | AP002844 | Oryza sativa                      | putative ABC transporter protein. P0410E03.10. contains ESTs AU065360(R3463),AU101680(R3463).             |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAB21273.1     | AP002844 | Oryza sativa            | putative ABC transporter protein. P0410E03.4.  |
| BAA83352.1     | AP000391 | Oryza sativa            | ESTs AU067992(C11433),AU077424(C11433) correspond to a region of the predicted gene.; Similar to ABC transporter-7 (U43892). |
| BAA96612.1     | AP002482 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome 2, BAC F14M4 ; putative ABC transporter (AC004411).                               |
| SEQ ID NO: 827 |          |                         |  |
| AAG34803.1     | AF243368 | Glycine max             | glutathione S-transferase GST 13.  |
| AAG16758.1     | AY007560 | Lycopersicon esculentum | putative glutathione S-transferase T3.   |
| AAG34798.1     | AF243363 | Glycine max             | glutathione S-transferase GST 8.   |
| AAF64450.1     | AF239928 | Euphorbia esula         | glutathione S-transferase. similar to auxin-inducible GST.   |
| AAG34807.1     | AF243372 | Glycine max             | glutathione S-transferase GST 17.  |
| AAG34796.1     | AF243361 | Glycine max             | glutathione S-transferase GST 6.   |
| AAG16759.1     | AY007561 | Lycopersicon esculentum | putative glutathione S-transferase T4.   |
| AAG34797.1     | AF243362 | Glycine max             | glutathione S-transferase GST 7.   |
| AAG34801.1     | AF243366 | Glycine max             | glutathione S-transferase GST 11.  |
| AAG34804.1     | AF243369 | Glycine max             | glutathione S-transferase GST 14.  |
| AAG34809.1     | AF243374 | Glycine max             | glutathione S-transferase GST 19.  |
| AAG34808.1     | AF243373 | Glycine max             | glutathione S-transferase GST 18.  |
| AAG34810.1     | AF243375 | Glycine max             | glutathione S-transferase GST 20.  |
| AAG16757.1     | AY007559 | Lycopersicon esculentum | putative glutathione S-transferase T2.   |
| AAG16756.1     | AY007558 | Lycopersicon esculentum | putative glutathione S-transferase T1.   |
| AAG34844.1     | AF244701 | Zea mays                | glutathione S-transferase GST 36.  |
| AAG34805.1     | AF243370 | Glycine max             | glutathione S-transferase GST 15.  |

|            |          |                              |   |
|------------|----------|------------------------------|---|
| AAG34831.1 | AF244688 | Zea mays                     | glutathione S-transferase GST 23.   |
| AAG34832.1 | AF244689 | Zea mays                     | glutathione S-transferase GST 24.   |
| AAG34849.1 | AF244706 | Zea mays                     | glutathione S-transferase GST 41.   |
| AAG34802.1 | AF243367 | Glycine max                  | glutathione S-transferase GST 12.   |
| CAA09187.1 | AJ010448 | Alopecurus myosuroides       | glutathione transferase. GST1a.   |
| AAG34829.1 | AF244686 | Zea mays                     | glutathione S-transferase GST 21.   |
| CAA09188.1 | AJ010449 | Alopecurus myosuroides       | glutathione transferase. GST1b.   |
| AAA68430.1 | J03679   | Solanum tuberosum            | glutathione S-transferase. gst1. previously called pathogenesis-related protein; prp1-1.  |
| AAG34836.1 | AF244693 | Zea mays                     | glutathione S-transferase GST 28.   |
| AAG34837.1 | AF244694 | Zea mays                     | glutathione S-transferase GST 29.   |
| AAG34800.1 | AF243365 | Glycine max                  | glutathione S-transferase GST 10.   |
| AAC18566.1 | AF048978 | Glycine max                  | 2,4-D inducible glutathione S-transferase. GSTa.  |
| AAC32118.1 | AF051214 | Picea mariana                | probable glutathione S-transferase. Sb18. similar to Nicotiana tabacum probable glutathione S-transferase encoded by GenBank Accession Number X56266. |
| AAF22517.1 | AF118924 | Papaver somniferum           | glutathione S-transferase 1. GST1.  |
| AAF22518.1 | AF118925 | Papaver somniferum           | glutathione S-transferase 2. GST2.  |
| AAG32471.1 | AF309378 | Oryza sativa subsp. japonica | putative glutathione S-transferase OsGSTU4.   |
| AAG34806.1 | AF243371 | Glycine max                  | glutathione S-transferase GST 16.   |
| CAA04391.1 | AJ000923 | Carica papaya                | glutathione transferase. PGST1.   |
| AAG32472.1 | AF309379 | Oryza sativa subsp. japonica | putative glutathione S-transferase OsGSTU3.   |
| AAG34833.1 | AF244690 | Zea mays                     | glutathione S-transferase GST 25.   |
| CAA71784.1 | Y10820   | Glycine max                  | glutathione transferase.  |

---

AAG34847.1 AF244704 Zea mays  
glutathione S-transferase GST 39.

---

AAF22519.1 AF118926 Papaver somniferum  
glutathione S-transferase 3. GST3.

---

SEQ ID NO: 828

---

AAG34803.1 AF243368 Glycine max  
glutathione S-transferase GST 13.

---

AAG34797.1 AF243362 Glycine max  
glutathione S-transferase GST 7.

---

AAG34798.1 AF243363 Glycine max  
glutathione S-transferase GST 8.

---

AAF64450.1 AF239928 Euphorbia esula  
glutathione S-transferase. similar to auxin-inducible GST.

---

AAG16758.1 AY007560 Lycopersicon esculentum  
putative glutathione S-transferase T3.

---

AAG34796.1 AF243361 Glycine max  
glutathione S-transferase GST 6.

---

AAG16759.1 AY007561 Lycopersicon esculentum  
putative glutathione S-transferase T4.

---

AAG34801.1 AF243366 Glycine max  
glutathione S-transferase GST 11.

---

AAG34804.1 AF243369 Glycine max  
glutathione S-transferase GST 14.

---

AAG34807.1 AF243372 Glycine max  
glutathione S-transferase GST 17.

---

AAG34809.1 AF243374 Glycine max  
glutathione S-transferase GST 19.

---

AAG34810.1 AF243375 Glycine max  
glutathione S-transferase GST 20.

---

AAG16757.1 AY007559 Lycopersicon esculentum  
putative glutathione S-transferase T2.

---

AAG16756.1 AY007558 Lycopersicon esculentum  
putative glutathione S-transferase T1.

---

AAG34802.1 AF243367 Glycine max  
glutathione S-transferase GST 12.

---

AAG34808.1 AF243373 Glycine max  
glutathione S-transferase GST 18.

---

AAG34844.1 AF244701 Zea mays  
glutathione S-transferase GST 36.

---

CAA09188.1 AJ010449 Alopecurus myosuroides  
glutathione transferase. GST1b.

---

|                |          |  |
|----------------|----------|--|
| CAA09187.1     | AJ010448 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1a.   |
| AAG32472.1     | AF309379 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTU3.  |
| AAA68430.1     | J03679   | <i>Solanum tuberosum</i><br>glutathione S-transferase. <i>gst1</i> . previously called pathogenesis-related protein; <i>prp1-1</i> .   |
| AAG34837.1     | AF244694 | <i>Zea mays</i><br>glutathione S-transferase GST 29.   |
| AAG34800.1     | AF243365 | <i>Glycine max</i><br>glutathione S-transferase GST 10.  |
| AAG34831.1     | AF244688 | <i>Zea mays</i><br>glutathione S-transferase GST 23.   |
| AAC32118.1     | AF051214 | <i>Picea mariana</i><br>probable glutathione S-transferase. Sb18. similar to <i>Nicotiana tabacum</i> probable glutathione S-transferase encoded by GenBank Accession Number X56266. |
| AAG34805.1     | AF243370 | <i>Glycine max</i><br>glutathione S-transferase GST 15.  |
| AAC18566.1     | AF048978 | <i>Glycine max</i><br>2,4-D inducible glutathione S-transferase. GSTa.   |
| AAG34829.1     | AF244686 | <i>Zea mays</i><br>glutathione S-transferase GST 21.   |
| CAA04391.1     | AJ000923 | <i>Carica papaya</i><br>glutathione transferase. PGST1.  |
| CAA71784.1     | Y10820   | <i>Glycine max</i><br>glutathione transferase.   |
| AAG34795.1     | AF243360 | <i>Glycine max</i><br>glutathione S-transferase GST 5.   |
| AAG34836.1     | AF244693 | <i>Zea mays</i><br>glutathione S-transferase GST 28.   |
| AAG34832.1     | AF244689 | <i>Zea mays</i><br>glutathione S-transferase GST 24.   |
| AAG34833.1     | AF244690 | <i>Zea mays</i><br>glutathione S-transferase GST 25.   |
| AAG34849.1     | AF244706 | <i>Zea mays</i><br>glutathione S-transferase GST 41.   |
| AAG34806.1     | AF243371 | <i>Glycine max</i><br>glutathione S-transferase GST 16.  |
| CAA09189.1     | AJ010450 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1c.   |
| SEQ ID NO: 829 |          |  |
| AAG34803.1     | AF243368 | <i>Glycine max</i><br>glutathione S-transferase GST 13.  |

|            |          |  |
|------------|----------|--|
| AAF64450.1 | AF239928 | <i>Euphorbia esula</i><br>glutathione S-transferase. similar to auxin-inducible GST. |
| AAG16758.1 | AY007560 | <i>Lycopersicon esculentum</i><br>putative glutathione S-transferase T3.             |
| AAG34798.1 | AF243363 | <i>Glycine max</i><br>glutathione S-transferase GST 8.                               |
| AAG34801.1 | AF243366 | <i>Glycine max</i><br>glutathione S-transferase GST 11.                              |
| AAG34797.1 | AF243362 | <i>Glycine max</i><br>glutathione S-transferase GST 7.                               |
| AAG34796.1 | AF243361 | <i>Glycine max</i><br>glutathione S-transferase GST 6.                               |
| AAG34807.1 | AF243372 | <i>Glycine max</i><br>glutathione S-transferase GST 17.                              |
| AAG16759.1 | AY007561 | <i>Lycopersicon esculentum</i><br>putative glutathione S-transferase T4.             |
| AAG34804.1 | AF243369 | <i>Glycine max</i><br>glutathione S-transferase GST 14.                              |
| AAG34810.1 | AF243375 | <i>Glycine max</i><br>glutathione S-transferase GST 20.                              |
| AAG34809.1 | AF243374 | <i>Glycine max</i><br>glutathione S-transferase GST 19.                              |
| AAG16757.1 | AY007559 | <i>Lycopersicon esculentum</i><br>putative glutathione S-transferase T2.             |
| AAG16756.1 | AY007558 | <i>Lycopersicon esculentum</i><br>putative glutathione S-transferase T1.             |
| AAG34805.1 | AF243370 | <i>Glycine max</i><br>glutathione S-transferase GST 15.                              |
| AAC18566.1 | AF048978 | <i>Glycine max</i><br>2,4-D inducible glutathione S-transferase. GSTa.               |
| AAG34808.1 | AF243373 | <i>Glycine max</i><br>glutathione S-transferase GST 18.                              |
| AAG34800.1 | AF243365 | <i>Glycine max</i><br>glutathione S-transferase GST 10.                              |
| AAG34829.1 | AF244686 | <i>Zea mays</i><br>glutathione S-transferase GST 21.                                 |
| AAG34802.1 | AF243367 | <i>Glycine max</i><br>glutathione S-transferase GST 12.                              |
| AAG34837.1 | AF244694 | <i>Zea mays</i><br>glutathione S-transferase GST 29.                                 |
| CAA09187.1 | AJ010448 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1a.                     |

|                |          |  |
|----------------|----------|--|
| CAA09188.1     | AJ010449 | <i>Alopecurus myosuroides</i><br>glutathione transferase. GST1b.   |
| AAG34849.1     | AF244706 | <i>Zea mays</i><br>glutathione S-transferase GST 41.   |
| AAG34844.1     | AF244701 | <i>Zea mays</i><br>glutathione S-transferase GST 36.   |
| AAG34806.1     | AF243371 | <i>Glycine max</i><br>glutathione S-transferase GST 16.  |
| CAA71784.1     | Y10820   | <i>Glycine max</i><br>glutathione transferase.   |
| AAA68430.1     | J03679   | <i>Solanum tuberosum</i><br>glutathione S-transferase. gst1. previously called pathogenesis-related protein; prp1-1.   |
| CAA04391.1     | AJ000923 | <i>Carica papaya</i><br>glutathione transferase. PGST1.  |
| AAG34836.1     | AF244693 | <i>Zea mays</i><br>glutathione S-transferase GST 28.   |
| AAG34831.1     | AF244688 | <i>Zea mays</i><br>glutathione S-transferase GST 23.   |
| AAG34847.1     | AF244704 | <i>Zea mays</i><br>glutathione S-transferase GST 39.   |
| AAC32118.1     | AF051214 | <i>Picea mariana</i><br>probable glutathione S-transferase. Sb18. similar to <i>Nicotiana tabacum</i> probable glutathione S-transferase encoded by GenBank Accession Number X56266. |
| AAF29773.1     | AF159229 | <i>Gossypium hirsutum</i><br>glutathione S-transferase. GST.   |
| AAG32472.1     | AF309379 | <i>Oryza sativa</i> subsp. <i>japonica</i><br>putative glutathione S-transferase OsGSTU3.  |
| AAG41204.1     | AF321437 | <i>Suaeda maritima</i><br>glutathione transferase.   |
| CAC24549.1     | AJ296343 | <i>Cichorium intybus</i> x <i>Cichorium endivia</i><br>glutathione S-transferase. chi-GST1. auxin-induced GST.   |
| SEQ ID NO: 830 |          |  |
| AAD37699.1     | AF145730 | <i>Oryza sativa</i><br>homeodomain leucine zipper protein. Oshox6. transcription factor.   |
| BAA93461.1     | AB028073 | <i>Physcomitrella patens</i><br>homeobox protein PpHB2. PpHB2. homeodomain-leucine zipper gene.  |
| AAF01765.1     | AF184278 | <i>Glycine max</i><br>homeodomain-leucine zipper protein 57. Hdl57. transcription factor.  |
| AAF01764.2     | AF184277 | <i>Glycine max</i><br>homeodomain-leucine zipper protein 56. Hdl56. transcription factor.  |
| BAA93466.1     | AB028078 | <i>Physcomitrella patens</i><br>homeobox protein PpHB7. PpHB7. homeodomain-leucine zipper gene.  |



|            |          |   |   |
|------------|----------|---|---|
| CAB67118.1 | Y17306   | <i>Lycopersicon esculentum</i>                | homeodomain protein. h52.   |
| AAF73482.1 | AF268422 | <i>Brassica rapa</i> subsp. <i>pekinensis</i> | hb-6-like protein. transcription factor; similar to <i>Arabidopsis thaliana</i> hb-6 protein.             |
| AAD37697.1 | AF145728 | <i>Oryza sativa</i>                           | homeodomain leucine zipper protein. Oshox4. transcription factor.   |
| BAA21017.1 | D26578   | <i>Daucus carota</i>                          | transcriptional regulator. DNA-binding protein. homeodomain at nt 761-940; leucine zipper at nt 941-1048. |
| BAA05624.1 | D26575   | <i>Daucus carota</i>                          | transcriptional regulator. DNA-binding protein. homeodomain at nt 520-699; leucine zipper at nt 700-805.  |
| BAA93460.1 | AB028072 | <i>Physcomitrella patens</i>                  | homeobox protein PpHB1. PpHB1. homeodomain-leucine zipper gene.   |
| BAA93464.1 | AB028076 | <i>Physcomitrella patens</i>                  | homeobox protein PpHB5. PpHB5. homeodomain-leucine zipper gene.   |
| BAA93467.1 | AB028079 | <i>Physcomitrella patens</i>                  | homeobox protein PpHB8. PpHB8. homeodomain-leucine zipper gene.   |
| BAA93465.1 | AB028077 | <i>Physcomitrella patens</i>                  | homeobox protein PpHB6. PpHB6. homeodomain-leucine zipper gene.   |
| BAA05625.1 | D26576   | <i>Daucus carota</i>                          | transcriptional regulator. DNA-binding protein. homeodomain at nt 300-479; leucine zipper at nt 480-587.  |
| BAA93468.1 | AB028080 | <i>Physcomitrella patens</i>                  | homeobox protein PpHB9. PpHB9. homeodomain-leucine zipper gene.   |
| BAA05623.1 | D26574   | <i>Daucus carota</i>                          | transcriptional regulator. DNA-binding protein. homeodomain at nt 498-677; leucine zipper at nt 678-785.  |
| BAA05622.1 | D26573   | <i>Daucus carota</i>                          | transcriptional regulator. DNA-binding protein. homeodomain at nt 585-764; leucine zipper at nt 765-851.  |
| AAD37698.1 | AF145729 | <i>Oryza sativa</i>                           | homeodomain leucine zipper protein. Oshox5. transcription factor.   |
| CAA64221.1 | X94449   | <i>Pimpinella brachycarpa</i>                 | transcription activator. homeobox-leucine zipper protein. PHZ4.   |
| CAA64152.1 | X94375   | <i>Pimpinella brachycarpa</i>                 | transcription activator. homeobox-leucine zipper protein.   |
| CAA64491.1 | X95193   | <i>Pimpinella brachycarpa</i>                 | transcription activator. homeobox-leucine zipper protein.   |
| BAA93463.1 | AB028075 | <i>Physcomitrella patens</i>                  | homeobox protein PpHB4. PpHB4. homeodomain-leucine zipper gene.   |
| CAA06728.1 | AJ005833 | <i>Craterostigma plantagineum</i>             | transcription factor. homeodomain leucine zipper protein. hb-2.   |

|                |          |                            |  |
|----------------|----------|----------------------------|--|
| AAD37695.1     | AF145726 | Oryza sativa               | homeodomain leucine zipper protein. Oshox2. transcription factor.  |
| CAA65456.2     | X96681   | Oryza sativa               | transcription factor. DNA-binding protein. Oshox1. homeodomain leucine zipper gene.  |
| AAF19980.1     | AF211193 | Oryza sativa               | homeodomain-leucine zipper transcription factor. Hox1. hox1.   |
| AAK31270.1     | AC079890 | Oryza sativa               | homeodomain leucine zipper protein hox1. OSJNBb0089A17.12.   |
| CAA63222.1     | X92489   | Glycine max                | transcription activator. homeobox-leucine zipper protein.  |
| CAA06717.1     | AJ005820 | Craterostigma plantagineum | transcription factor. homeodomain leucine zipper protein. hb-1.  |
| AAA79778.1     | L48485   | Helianthus annuus          | homeodomain protein. putative.   |
| SEQ ID NO: 831 |          |                            |  |
| CAA06334.1     | AJ005077 | Lycopersicon esculentum    | protein kinase. TCTR2 protein. TCTR2.  |
| AAG31141.1     | AF305911 | Oryza sativa               | EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.  |
| AAG31142.1     | AF305912 | Hordeum vulgare            | EDR1. EDR1. MAP kinase kinase kinase; similar to Arabidopsis thaliana EDR1.  |
| AAK30005.1     | AY029067 | Rosa hybrid cultivar       | CTR2 protein kinase.   |
| AAD46406.1     | AF096250 | Lycopersicon esculentum    | ethylene-responsive protein kinase TCTR1. ER50. serine/threonine kinase; similar to Arabidopsis thaliana negative regulator of the ethylene response pathway encoded by GenBank Accession Number L08789. |
| CAA73722.1     | Y13273   | Lycopersicon esculentum    | putative protein kinase.   |
| AAD10057.1     | AF110519 | Lycopersicon esculentum    | ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1v.   |
| AAD10056.1     | AF110518 | Lycopersicon esculentum    | ethylene-inducible CTR1-like protein kinase. protein kinase homolog; ethylene and fruit ripening inducible CTR1-like protein kinase; TCTR1.  |
| AAA34002.1     | M67449   | Glycine max                | protein kinase. PK6.   |
| AAK11734.1     | AY027437 | Arachis hypogaea           | serine/threonine/tyrosine kinase.  |
| BAB16918.1     | AP002863 | Oryza sativa               | putative protein kinase. P0005A05.22.  |
| CAC09580.1     | AJ298992 | Fagus sylvatica            | Abscisic acid (ABA) and calcium induced protein kinase. protein kinase (PK). pk1.  |

|            |          |                      |  |
|------------|----------|----------------------|--|
| CAA97692.1 | Z73295   | Catharanthus roseus  | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.     |
| AAF59906.1 | AF197947 | Glycine max          | receptor protein kinase-like protein. CLV1B.   |
| AAF59905.1 | AF197946 | Glycine max          | receptor protein kinase-like protein. CLV1A.   |
| AAK00425.1 | AC069324 | Oryza sativa         | Putative protein kinase. OSJNBa0071K19.11.   |
| BAA87852.1 | AP000816 | Oryza sativa         | Similar to putative Ser/Thr protein kinase. (AC004218).  |
| BAB40094.1 | AP003210 | Oryza sativa         | putative receptor protein kinase. OSJNBa0010K01.7.   |
| BAA92221.1 | AP001278 | Oryza sativa         | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).                         |
| CAA08995.1 | AJ010091 | Brassica napus       | MAP3K alpha 1 protein kinase. MAP3K alpha 1.   |
| BAB21240.1 | AP002953 | Oryza sativa         | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).   |
| CAB51834.1 | 00069    | Oryza sativa         | l1332.5. contains eukaryotic protein kinase domain PF.   |
| CAA08997.1 | AJ010093 | Brassica napus       | MAP3K beta 1 protein kinase. MAP3K beta 1.   |
| CAB54520.1 | AJ238845 | Brassica napus       | putative role in cell cycle control. MAP3K epsilon 1 protein kinase. MAP3Ke1.  |
| AAF34436.1 | AF172282 | Oryza sativa         | similar to mitogen-activated protein kinases. DUPR11.32.   |
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| AAG25966.1 | AF302082 | Nicotiana tabacum    | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.                  |
| BAA06538.1 | D31737   | Nicotiana tabacum    | protein-serine/threonine kinase.   |
| AAF76189.1 | AF271206 | Rosa hybrid cultivar | CTR1-like protein kinase. Raf-like protein kinase.   |
| BAA84787.1 | AP000559 | Oryza sativa         | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |

BAA83373.1 AP000391 *Oryza sativa*  
ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484).

AAF66615.1 AF142596 *Nicotiana tabacum*  
LRR receptor-like protein kinase.

AAF91322.1 AF244888 *Glycine max*  
receptor-like protein kinase 1. RLK1. GmRLK1.

AAD21872.1 AF078082 *Phaseolus vulgaris*  
receptor-like protein kinase homolog RK20-1.

CAA61510.1 X89226 *Oryza sativa*  
leucine-rich repeat/receptor protein kinase. lrk2.

BAA87853.1 AP000816 *Oryza sativa*  
EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).

AAF91323.1 AF244889 *Glycine max*  
receptor-like protein kinase 2. RLK2. GmRLK2.

AAF91324.1 AF244890 *Glycine max*  
receptor-like protein kinase 3. RLK3. GmRLK3.

AAF43394.1 AF230501 *Oryza sativa* subsp. *japonica*  
serine/threonine protein kinase. YK1.

AAK16409.1 AF320086 *Zea mays*  
serine threonine kinase 1. stk1. expressed in mature tassel.

AAK21965.1 AY028699 *Brassica napus*  
receptor protein kinase PERK1.

BAB39437.1 AP003338 *Oryza sativa*  
receptor-like kinase. OJ1212\_B09.6.

AAK11568.1 AF318492 *Lycopersicon hirsutum*  
Pto-like protein kinase B. LhirPtoB.

SEQ ID NO: 832

AAF35901.1 AF230332 *Zinnia elegans*  
expansin 2.

CAC19184.1 AJ291817 *Cicer arietinum*  
expansin.

AAG13982.1 AF297521 *Prunus avium*  
expansin 1. Exp1. PruavExp1.

BAB19676.1 AB029083 *Prunus persica*  
expansin. PchExp1.

AAC33529.1 U93167 *Prunus armeniaca*  
expansin. PA-Exp1.

AAC33530.1 AF038815 *Prunus armeniaca*  
expansin. Exp2.

AAD47901.1 AF085330 *Pinus taeda*  
expansin.

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAB37746.1 | U30382   | Cucumis sativus         | expansin S1 precursor. Cs-EXP1. similar to pollen allergen Lol pI, Lolium perenne, Swiss-Prot Accession Number P14946; former gene name CuExS1; expansin-29 (Ex29) protein.  |
| AAF21101.1 | AF159563 | Fragaria x ananassa     | expansin. Exp2. ripening regulated.  |
| AAB40634.1 | U64890   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAB40637.1 | U64893   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAB40635.1 | U64891   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| CAB43197.1 | AJ239068 | Lycopersicon esculentum | cell wall loosening enzyme. expansin2. exp2.   |
| AAB40636.1 | U64892   | Pinus taeda             | expansin. similar to Arabidopsis expansin encoded by GenBank Accession Numbers U30476, U30478, U30480 and U30481, to Cucumis sativus expansin encoded by GenBank Accession Numbers U30382 and U30460, and to rice expansin encoded by GenBank Accession Numbers U30477 and U30479. |
| AAC64201.1 | AF096776 | Lycopersicon esculentum | expansin. LeEXP2.  |
| AAD49956.1 | AF167360 | Rumex palustris         | expansin. EXP1.  |
| AAC96081.1 | AF049354 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP5. cell wall protein.  |
| AAC39512.1 | AF043284 | Gossypium hirsutum      | expansin. GhEX1. contains N-terminal signal peptide.   |
| AAB81662.1 | U85246   | Oryza sativa            | expansin. Os-EXP4.   |
| AAG13983.1 | AF297522 | Prunus avium            | expansin 2. Exp2. PruavExp2.   |
| AAF32409.1 | AF230276 | Triphysaria versicolor  | alpha-expansin 3.  |
| AAG32921.1 | AF184233 | Lycopersicon esculentum | expansin. Exp10.   |

|            |          |   |
|------------|----------|---|
| BAB32732.1 | AB049406 | <i>Eustoma grandiflorum</i><br>expansin. Eg Expansin.   |
| AAF32411.1 | AF230278 | <i>Triphysaria versicolor</i><br>alpha-expansin 1.  |
| AAF35902.1 | AF230333 | <i>Zinnia elegans</i><br>expansin 3.  |
| AAB38074.1 | U30477   | <i>Oryza sativa</i><br>induces extension (creep) in plant cell walls. expansin Os-EXP2. Os-EXP2. former gene name RiExB.  |
| AAC96080.1 | AF049353 | <i>Nicotiana tabacum</i><br>involved in acid-growth response. alpha-expansin precursor. Nt-EXP4. cell wall protein.   |
| AAF17570.1 | AF202119 | <i>Marsilea quadrifolia</i><br>alpha-expansin. EXP1. Mq-EXP1.   |
| CAC06433.1 | AJ276007 | <i>Festuca pratensis</i><br>expansin. exp2.   |
| AAD13633.1 | AF059489 | <i>Lycopersicon esculentum</i><br>expansin precursor. Exp5.   |
| CAC19183.1 | AJ291816 | <i>Cicer arietinum</i><br>expansin.   |
| AAF62181.1 | AF247163 | <i>Oryza sativa</i><br>alpha-expansin OsEXP6. cell wall loosening factor; expressed in internodes and leaves.   |
| AAF62180.1 | AF247162 | <i>Oryza sativa</i><br>alpha-expansin OsEXP5. cell wall loosening factor; expressed in internodes, leaves, coleoptiles, and roots.  |
| CAB46492.1 | AJ243340 | <i>Lycopersicon esculentum</i><br>expansin9. exp9.  |
| BAA88200.1 | AP000837 | <i>Oryza sativa</i><br>EST AU078708(E60526) corresponds to a region of the predicted gene. Similar to expansin (U85246).  |
| AAF32410.1 | AF230277 | <i>Triphysaria versicolor</i><br>alpha-expansin 2.  |
| AAB37749.1 | U30460   | <i>Cucumis sativus</i><br>expansin S2 precursor. Cs-EXP2. similar to pollen allergen Lol pI, <i>Lolium perenne</i> , Swiss-Prot Accession Number P14946; former gene name CuExS2; expansin-30 (Ex30) protein. |
| CAA04385.1 | AJ000885 | <i>Brassica napus</i><br>Cell wall extension in plants. Expansin.   |
| AAF17571.1 | AF202120 | <i>Regnellidium diphyllum</i><br>alpha-expansin. EXP1. Rd-EXP1.   |
| AAD13632.1 | AF059488 | <i>Lycopersicon esculentum</i><br>expansin precursor. Exp4.   |
| CAA06271.2 | AJ004997 | <i>Lycopersicon esculentum</i><br>expansin18. exp18.  |

|                |          |                               |  |
|----------------|----------|-------------------------------|--|
| AAC63088.1     | U82123   | Lycopersicon esculentum       | expansin. LeEXP1. fruit ripening regulated expansin.                                       |
| AAC96077.1     | AF049350 | Nicotiana tabacum             | involved in acid-growth response. alpha-expansin precursor. Nt-EXP1. cell wall protein.    |
| AAF62182.1     | AF247164 | Oryza sativa                  | alpha-expansin OsEXP7. cell wall loosening factor; expressed in internodes and leaves.     |
| CAC18802.1     | AJ289154 | Glycine max                   | expansion of cell walls. expansin. dd2/63.   |
| AAC96078.1     | AF049351 | Nicotiana tabacum             | involved in acid-growth response. alpha-expansin precursor. Nt-EXP2. cell wall protein.    |
| AAG01875.1     | AF291659 | Striga asiatica               | alpha-expansin 3. Exp3.  |
| CAA69105.1     | Y07782   | Oryza sativa                  | expansin. RiExA.   |
| AAC96079.1     | AF049352 | Nicotiana tabacum             | involved in acid-growth response. alpha-expansin precursor. Nt-EXP3. cell wall protein.    |
| SEQ ID NO: 833 |          |                               |  |
| AAA34030.1     | J03492   | Spinacia oleracea             | glycolate oxidase (EC 1.1.3.15).   |
| AAB40396.1     | U80071   | Mesembryanthemum crystallinum | glycolate oxidase. GOX.  |
| BAA03131.1     | D14044   | Cucurbita sp.                 | glycolate oxidase.   |
| AAB82143.1     | AF022740 | Oryza sativa                  | glycolate oxidase. GOX.  |
| CAA63482.1     | X92888   | Lycopersicon esculentum       | conversion of glycolate to glyoxylate + H <sub>2</sub> O <sub>2</sub> . glycolate oxidase. |
| AAC32392.1     | AF082874 | Medicago sativa               | glycolate oxidase.   |
| AAC33509.1     | U62485   | Nicotiana tabacum             | photorespiration. glycolate oxidase. GLO.  |
| AAF03097.1     | AF162196 | Lactuca sativa                | glycolate oxidase.   |
| SEQ ID NO: 838 |          |                               |  |
| CAA06770.1     | AJ005928 | Brassica napus                | squalene epoxidase homologue. Sqp1;2.  |
| CAA06773.1     | AJ005931 | Brassica napus                | squalene epoxidase homologue. Sqp1;1.  |
| BAA24448.1     | AB003516 | Panax ginseng                 | squalene epoxidase.  |
| CAA06223.1     | AJ004923 | Lycopersicon esculentum       | Squalene epoxidase. ERG.   |

## SEQ ID NO: 840

BAB12686.1 AP002746 *Oryza sativa*  
putative pyrophosphate-dependent phosphofructo-1-kinase. P0671B11.1. contains ESTs  
AU068014(C11507),C28532(C61484),AU090544(C61415).

BAA99438.1 AP002743 *Oryza sativa*  
putative pyrophosphate-dependent phosphofructo-1-kinase. P0710E05.25. contains ESTs  
AU068014(C11507),C28532(C61484),AU090544(C61415).

AAB88875.1 U93272 *Prunus armeniaca*  
pyrophosphate-dependent phosphofructo-1-kinase.

CAA83683.1 Z32850 *Ricinus communis*  
pyrophosphate-dependent phosphofructokinase beta subunit.

AAC67587.1 AF095521 *Citrus x paradisi*  
pyrophosphate-dependent phosphofructokinase alpha subunit. PPi-PFKa.

AAA63452.1 M55191 *Solanum tuberosum*  
pyrophosphate-fructose 6-phosphate 1-phosphotransferase beta-subunit.

AAC67586.1 AF095520 *Citrus x paradisi*  
pyrophosphate-dependent phosphofructokinase beta subunit. PPi-PFKb. PFP.

AAA63451.1 M55190 *Solanum tuberosum*  
pyrophosphate-fructose 6-phosphate 1-phosphotransferase alpha-subunit.

CAA83682.1 Z32849 *Ricinus communis*  
pyrophosphate-dependent phosphofructokinase alpha subunit.

## SEQ ID NO: 841

AAG60182.1 AC084763 *Oryza sativa*  
putative ethylene-responsive element binding protein. OSJNBa0027P10.12.

AAK31279.1 AC079890 *Oryza sativa*  
putative ethylene-responsive element binding protein. OSJNBb0089A17.16.

AAG43545.1 AF211527 *Nicotiana tabacum*  
Avr9/Cf-9 rapidly elicited protein 1. ACRE1. similar to EREBP transcription factors.

AAF63205.1 AF245119 *Mesembryanthemum crystallinum*  
AP2-related transcription factor. CDBP. stress induced transcription factor.

BAA07321.1 D38123 *Nicotiana tabacum*  
ERF1. ethylene-responsive transcription factor.

BAA97122.1 AB016264 *Nicotiana sylvestris*  
ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf2.

CAB93940.1 AJ238740 *Catharanthus roseus*  
putative transcription factor. AP2-domain DNA-binding protein. orca2.

BAA87068.1 AB035270 *Matricaria chamomilla*  
ethylene-responsive element binding protein1 homolog. McEREBP1.

BAA97124.1 AB016266 *Nicotiana sylvestris*  
ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf4.



|            |          |                     |  |
|------------|----------|---------------------|--|
| CAB96900.1 | AJ251250 | Catharanthus roseus | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| CAB96899.1 | AJ251249 | Catharanthus roseus | transcription factor. AP2-domain DNA-binding protein. orca3.   |
| AAC62619.1 | AF057373 | Nicotiana tabacum   | transcription factor. ethylene response element binding protein 1. EREBP1.   |
| BAA97123.1 | AB016265 | Nicotiana glauca    | ERF (EREBP); ethylene-responsive element binding factor for basic PR (Pathogenesis-related) gene of higher plant. ethylene-responsive element binding factor. nserf3.                |
| AAC24587.1 | AF071893 | Prunus armeniaca    | AP2 domain containing protein. AP2DCP.   |
| CAC12822.1 | AJ299252 | Nicotiana tabacum   | AP2 domain-containing transcription factor. ap2.   |
| AAF76898.1 | AF274033 | Atriplex hortensis  | apetala2 domain-containing protein.  |
| BAA94514.2 | AP001800 | Oryza sativa        | Similar to Arabidopsis thaliana chromosome 4, BAC clone F9D16; putative Ap2 domain protein (AL035394).   |
| AAC14323.1 | AF058827 | Nicotiana tabacum   | TSI1. Tsi1. contains putative AP2 DNA-binding domain; similar to Pti6.   |
| AAD00708.1 | U91857   | Stylosanthes hamata | ethylene-responsive element binding protein homolog. similar to EREBP1, -2, -3 and -4 proteins encoded by GenBank Accession Numbers D38123, D38126, D38124, and D38125 respectively. |
| BAA76734.1 | AB024575 | Nicotiana tabacum   | ethylene responsive element binding factor.  |
| BAB03248.1 | AB037183 | Oryza sativa        | ERF protein transcriptional repressor. ethylene responsive element binding factor3. osERF3.  |
| BAB16083.1 | AB036883 | Oryza sativa        | transcriptional repressor. osERF3. osERF3. ERF protein family ERF3 associated repression domain.   |
| AAF23899.1 | AF193803 | Oryza sativa        | transcription factor EREBP1. EREBP/AP2-like transcription factor.  |
| AAF05606.1 | AF190770 | Oryza sativa        | EREBP-like protein. tsh1. TSH1; induced by ethylene.   |
| CAB93939.1 | AJ238739 | Catharanthus roseus | putative transcription factor. AP2-domain DNA-binding protein. orca1.  |
| BAA78738.1 | AB023482 | Oryza sativa        | EST AU055776(S20048) corresponds to a region of the predicted gene.; Similar to Arabidopsis thaliana AP2 domain containing protein RAP2.10 mRNA, partial cds.(AF003103).             |
| AAG43548.1 | AF211530 | Nicotiana tabacum   | Avr9/Cf-9 rapidly elicited protein 111A. ACRE111A. similar to EREBP transcription factors.   |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| AAG43549.1     | AF211531 | Nicotiana tabacum         | Avr9/Cf-9 rapidly elicited protein 111B. ACRE111B. similar to EREBP transcription factors.  |
| BAA99376.1     | AP002526 | Oryza sativa              | ESTs AU093391(E60370),AU091593(C60458), AU093392(E60370) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana BAC F21J9; AP2 domain protein. (AC000103).   |
| AAK31271.1     | AC079890 | Oryza sativa              | putative transcriptional factor. OSJNBb0089A17.22.  |
| AAK01089.1     | AF298231 | Hordeum vulgare           | CBF3-like protein BCBF3. BCBF3. AP2 domain protein; DRE binding factor.   |
| AAG59618.1     | AF239616 | Hordeum vulgare           | CRT/DRE-binding factor. CBF.  |
| AAC49567.1     | U41466   | Zea mays                  | Glossy15. Glossy15. AP2 DNA-binding domain protein; similar to the Arabidopsis floral homeotic gene APETALA2; maize AP2-domain regulator of leaf epidermal traits; homeotic regulator of leaf epidermal cell identity; allele: W64A; putative transcription factor. |
| SEQ ID NO: 842 |          |                           |   |
| AAC32034.1     | AF023472 | Hordeum vulgare           | peptide transporter. ptr1. PTR1; integral membrane protein.   |
| BAB40113.1     | AP003311 | Oryza sativa              | putative peptide transport protein. P0024G09.4. contains ESTs D40448(S2437),C71800(E0368),AU102190(E2393), AU055921(S20154),AU102191(E2393),AU055922(S20154), C98524(E0368),AU097146(S2437).  |
| BAB16458.1     | AP002483 | Oryza sativa              | putative peptide transport protein. P0019D06.16. contains ESTs D40448(S2437),C71800(E0368),AU102190(E2393), AU055921(S20154),AU102191(E2393),AU055922(S20154), C98524(E0368),AU097146(S2437).   |
| AAD01600.1     | AF016713 | Lycopersicon esculentum   | LeOPT1. LeOPT1. oligopeptide transporter.   |
| AAF07875.1     | AF140606 | Oryza sativa              | nitrate transporter. NRT1.  |
| AAF20002.1     | AF213936 | Prunus dulcis             | amino acid/peptide transporter. PTR2. similar to transporters of nitrogenous compounds.   |
| AAG46153.1     | AC018727 | Oryza sativa              | putative peptide transporter. OSJNBa0056G17.8.  |
| CAC00544.1     | AJ277084 | Nicotiana plumbaginifolia | ion transport. putative low-affinity nitrate transporter. nrt1.1.   |
| CAC00545.1     | AJ277085 | Nicotiana plumbaginifolia | ion transport. putative low-affinity nitrate transporter. nrt1.2.   |
| AAA80582.1     | U17987   | Brassica napus            | putative nitrate transporter. RCH2 protein.   |

---

CAC07206.1 AJ278966 *Brassica napus*  
Low-affinity nitrate transporter, nitrate transporter, nrt1.

---

AAG21898.1 AC026815 *Oryza sativa*  
putative peptide transport protein. OSJNBa0079L16.13.

---

BAB19758.1 AB052786 *Glycine max*  
putative nitrate transporter NRT1-3. NRT1-3.

---

AAK15441.1 AC037426 *Oryza sativa*  
putative nitrate transporter. OSJNBb0014I11.9.

---

AAG21906.1 AC026815 *Oryza sativa*  
putative peptide transport protein. OSJNBa0079L16.9.

---

BAB19760.1 AB052788 *Glycine max*  
nitrate transporter NRT1-5. NRT1-5.

---

AAG46154.1 AC018727 *Oryza sativa*  
putative peptide transporter. OSJNBa0056G17.27.

---

BAB19757.1 AB052785 *Glycine max*  
nitrate transporter NRT1-2. NRT1-2.

---

BAB19756.1 AB052784 *Glycine max*  
nitrate transporter NRT1-1. NRT1-1.

---

BAB16322.1 AP002818 *Oryza sativa*  
putative peptide transporter-like protein. P0436E04.4.

---

AAB69642.1 AF000392 *Lotus japonicus*  
peptide transporter. LjNOD65.

---

CAA93316.1 Z69370 *Cucumis sativus*  
nitrite transporter. NiTR1.

---

AAD16016.1 AF080545 *Nepenthes alata*  
peptide transporter. PTR1.

---

BAB19759.1 AB052787 *Glycine max*  
putative nitrate transporter NRT1-4. NRT1-4.

---

AAD42860.1 AF154930 *Prunus dulcis*  
transporter-like protein. TLP1.

---

SEQ ID NO: 847

---

CAA61980.1 X89890 *Bidens pilosa*  
Calmodulin.

---

AAF73157.1 AF150059 *Brassica napus*  
calmodulin. CaM1. involved in seed germination.

---

BAA87825.1 AP000815 *Oryza sativa*  
ESTs AU030013(E50493),AU081341(E50493) correspond to a region of the predicted gene.  
Similar to O.sativa gene encoding calmodulin. (Z12828).

---

CAA67054.1 X98404 *Capsicum annuum*  
calmodulin-2.

---

AAA87347.1 M88307 *Brassica juncea*  
calmodulin.

---

|            |          |                    |   |
|------------|----------|--------------------|---|
| AAA33397.1 | L18912   | Lilium longiflorum | calcium binding protein, signal transduction. calmodulin. putative.   |
| AAG27432.1 | AF295637 | Elaeis guineensis  | calmodulin.   |
| CAA42423.1 | X59751   | Daucus carota      | calmodulin. Ccam-1.   |
| AAG11418.1 | AF292108 | Prunus avium       | calmodulin.   |
| AAA92681.1 | U13882   | Pisum sativum      | calcium-binding protein. calmodulin.  |
| AAB46588.1 | U83402   | Capsicum annuum    | calmodulin.   |
| AAA33706.1 | M80836   | Petunia x hybrida  | calmodulin. CAM81.  |
| AAF65511.1 | AF108889 | Capsicum annuum    | calmodulin.   |
| CAA43143.1 | X60738   | Malus x domestica  | Calmodulin. CaM.  |
| AAA19571.1 | U10150   | Brassica napus     | calcium binding. calmodulin. bcml.  |
| CAA78301.1 | Z12839   | Lilium longiflorum | calcium binding protein, signal transduction. calmodulin.   |
| BAA88540.1 | AP000969 | Oryza sativa       | ESTs AU081349(E61253), D41425(S3918) correspond to a region of the predicted gene. Similar to calmodulin. (AF042840). |
| AAB36130.1 | S81594   | Vigna radiata      | auxin-regulated calmodulin. auxin-regulated calmodulin, arCaM. This sequence comes from Fig. 1; arCaM.                |
| AAC36059.1 | AF042840 | Oryza sativa       | calmodulin. CaM1.   |
| AAA33901.1 | L18913   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin. putative.   |
| AAA33900.1 | L18914   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin.   |
| AAA34237.1 | L20691   | Vigna radiata      | calmodulin.   |
| CAA78288.1 | Z12828   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin.   |
| AAA32938.1 | M27303   | Hordeum vulgare    | calmodulin.   |
| CAA78287.1 | Z12827   | Oryza sativa       | calcium binding protein, signal transduction. calmodulin.   |

|            |          |   |
|------------|----------|---|
| AAC49587.1 | U49105   | Triticum aestivum<br>calmodulin TaCaM4-1. calcium-binding protein.  |
| AAC49586.1 | U49104   | Triticum aestivum<br>calmodulin TaCaM3-3. calcium-binding protein.  |
| AAC49583.1 | U48692   | Triticum aestivum<br>calmodulin TaCaM2-3. calcium-binding protein.  |
| AAC49585.1 | U49103   | Triticum aestivum<br>calmodulin TaCaM3-2. calcium-binding protein.  |
| AAC49584.1 | U48693   | Triticum aestivum<br>calmodulin TaCaM3-1. calcium-binding protein.  |
| AAC49582.1 | U48691   | Triticum aestivum<br>calmodulin TaCaM2-2. calcium-binding protein.  |
| AAC49580.1 | U48689   | Triticum aestivum<br>calmodulin TaCaM1-3. calcium-binding protein.  |
| AAC49579.1 | U48688   | Triticum aestivum<br>calmodulin TaCaM1-2. calcium binding protein.  |
| AAC49578.1 | U48242   | Triticum aestivum<br>calmodulin TaCaM1-1. calcium-binding.  |
| AAA03580.1 | L01431   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-2. putative.  |
| AAC36058.1 | AF042839 | Oryza sativa<br>calmodulin. CaM2.   |
| AAA85156.1 | U20296   | Solanum tuberosum<br>calcium-binding protein. calmodulin.   |
| AAA34014.1 | L01432   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-3. putative.  |
| AAD10244.1 | AF030032 | Phaseolus vulgaris<br>calmodulin. CaM. EF-hand protein; calcium-dependent protein; functions in calcium signal transduction pathways. |
| CAA36644.1 | X52398   | Medicago sativa<br>calmodulin (AA 1-149).   |
| AAA85155.1 | U20294   | Solanum tuberosum<br>calcium-binding protein. calmodulin.   |
| AAB68399.1 | U79736   | Helianthus annuus<br>calmodulin. HaCaM.   |
| AAA62351.1 | U20295   | Solanum tuberosum<br>calcium-binding protein. calmodulin.   |
| AAA34238.1 | L20507   | Vigna radiata<br>calmodulin.  |
| AAA85157.1 | U20297   | Solanum tuberosum<br>calcium-binding protein. calmodulin.   |
| AAA34013.1 | L01430   | Glycine max<br>calcium-binding regulatory protein. calmodulin. SCaM-1. putative.  |

|                |          |                            |  |
|----------------|----------|----------------------------|--|
| AAA33705.1     | M80831   | Petunia x hybrida          | calmodulin-related protein. CAM53.   |
| CAA74307.1     | Y13974   | Zea mays                   | calmodulin.  |
| CAA54583.1     | X77397   | Zea mays                   | calmodulin. CaM2.  |
| SEQ ID NO: 848 |          |                            |  |
| CAA06486.1     | AJ005340 | Linum usitatissimum        | IAA amidohydrolase. homolog.   |
| SEQ ID NO: 850 |          |                            |  |
| BAB17350.1     | AP002747 | Oryza sativa               | putative nodulin. P0698G03.34. contains ESTs D39891(S1543),D41717(S4395),AU033037(S1543).  |
| BAA85440.1     | AP000616 | Oryza sativa               | ESTs AU055729(S20023),AU055730(S20023) correspond to a region of the predicted gene.; similar to Medicago nodulin N21-like protein (AC004218). |
| CAB53493.1     | AJ245900 | Oryza sativa               | CAA303720.1 protein. q3037.20. Similar to Medicago nodulin N21 (MtN21).  |
| SEQ ID NO: 852 |          |                            |  |
| AAD16018.1     | AF081514 | Taxus canadensis           | prenyltransferase. geranylgeranyl diphosphate synthase. geranylgeranyl pyrophosphate synthase.   |
| SEQ ID NO: 853 |          |                            |  |
| BAB32588.1     | AB055807 | Momordica charantia        | inhibitor against trypsin. bgit.   |
| AAA34180.1     | J05094   | Lycopersicon peruvianum    | proteinase inhibitor I precursor.  |
| AAA34198.1     | M59427   | Lycopersicon peruvianum    | proteinase inhibitor I. proteinase inhibitor I.  |
| CAB61327.1     | AJ132473 | Amaranthus hypochondriacus | Proteinase inhibition. trypsin inhibitor.  |
| AAA60745.1     | J04099   | Lycopersicon esculentum    | proteinase inhibitor I. ER1.   |
| CAA78269.1     | Z12623   | Nicotiana tabacum          | Putative precursor of serine proteinase inhibitor type I. Pre-pro-proteinase inhibitor I.  |
| CAA47461.1     | X67076   | Nicotiana tabacum          | inhibitor of microbial serine proteinases (major isoform). TIMPa.  |
| CAA78265.1     | Z12619   | Nicotiana tabacum          | precursor for serine proteinase inhibitor I. Pre-pro-proteinase inhibitor I.   |
| CAA47460.1     | X67075   | Nicotiana tabacum          | inhibitor of microbial serine proteinases (minor isoform). TIMPb.  |
| AAA34067.1     | M74102   | Nicotiana glauca           | pre-pro-proteinase inhibitor I.  |

|                |          |   |  |
|----------------|----------|---|--|
| AAC49603.1     | U30861   | Solanum tuberosum                         | serine proteinase inhibitor. wound-inducible proteinase inhibitor I.   |
| BAA02823.1     | D13662   | Nicotiana glauca X Nicotiana langsdorffii | genetic tumor-related proteinase inhibitor I precursor. GTI.   |
| AAA34199.1     | K03290   | Lycopersicon esculentum                   | wound-induced proteinase inhibitor I prepropeptide.  |
| AAA34200.1     | M13938   | Lycopersicon esculentum                   | proteinase inhibitor I. PIIF.  |
| AAA69780.1     | L06137   | Solanum tuberosum                         | proteinase inhibitor I. pin1. putative.  |
| AAA72133.1     | L06985   | Solanum tuberosum                         | proteinase inhibitor I. pin1. The 'a' of the first atg is missing.   |
| CAA78259.1     | Z12611   | Solanum tuberosum                         | proteinase inhibitor I.  |
| AAA69781.1     | L06606   | Solanum tuberosum                         | proteinase inhibitor I. precursor.   |
| CAA48136.1     | X67950   | Solanum tuberosum                         | protease inhibitor I. pin1.  |
| CAA47907.1     | X67675   | Solanum tuberosum                         | proteinase inhibitor I. pin1.  |
| CAB71340.1     | AJ250663 | Hordeum vulgare                           | putative proteinase inhibitor. bci-7. similarity to subtilisin/chymotrypsin inhibitor.                                   |
| CAA57677.1     | X82187   | Zea mays                                  | serine proteinase inhibitor. substilin /chymotrypsin-like inhibitor. pis7.   |
| CAA55588.1     | X78988   | Zea mays                                  | proteinase inhibitor. MPI.   |
| CAA49593.1     | X69972   | Zea mays                                  | proteinase inhibitor. MPI.   |
| AAA33816.1     | M17108   | Solanum tuberosum                         | proteinase inhibitor I. precursor.   |
| CAA57307.1     | X81647   | Cucurbita maxima                          | Pumpkin fruit trypsin inhibitor. pfiAF4.   |
| CAA57203.1     | X81447   | Cucurbita maxima                          | Pumpkin Fruit Chymotrypsin Inhibitor. pfiBM7.  |
| SEQ ID NO: 859 |          |   |  |
| AAC34855.1     | AF082030 | Hemerocallis hybrid cultivar              | senescence-associated protein 5. SA5. mRNA accumulates in senescing petals and accumulation is induced by exogenous ABA. |
| AAG13616.1     | AC078840 | Oryza sativa                              | putative senescence-associated protein. OSJNBb0073N24.21.  |
| SEQ ID NO: 864 |          |   |  |
| AAF62403.1     | AF212183 | Nicotiana tabacum                         | harpin inducing protein. hin1. similar to hin1 protein.  |

CAA68848.1 Y07563 *Nicotiana tabacum*  
activated during hypersensitive response. hin1.

AAB97367.1 AF039532 *Oryza sativa*  
harpin induced gene 1 homolog. Hin1.

SEQ ID NO: 871

AAC61839.1 AF025430 *Papaver somniferum*  
berberine bridge enzyme. bbe1. (S)-reticuline:oxygen oxidoreductase (methylene bridge forming).

AAB20352.1 S65550 *Eschscholzia californica*  
(S)-reticuline:oxygen oxidoreductase (methylene-bridge-forming). /gene="(S)-reticuline:oxygen oxidoreductase. This sequence comes from Fig 2; berberine bridge enzyme.

AAC39358.1 AF005655 *Eschscholzia californica*  
oxidizes the N-methyl group of (S)-reticuline to the berberine bridge carbon C-8 of (S)-scoulerine in berberine and benzophenanthridine alkaloid biosynthesis. berberine bridge enzyme. bbe1. covalently-bound FAD-dependent oxidase; elicitor-inducible.

AAD17487.1 AF049347 *Berberis stolonifera*  
Catalyzes the oxidative cyclization of the N-methyl group of (S)-reticuline into the berberine bridge carbon (C-8) of (S)-scoulerine. berberine bridge enzyme. bbe1. Covalently flavinylated oxidase of isoquinoline alkaloid biosynthesis in plants.

SEQ ID NO: 872

AAF98369.1 AF158253 *Nicotiana tabacum*  
patatin-like protein 3. PAT3. NtPat3.

CAA73328.1 Y12793 *Cucumis sativus*  
mobilization of fat during seed germination. patatin-like protein.

CAA11042.1 AJ223039 *Hevea brasiliensis*  
latex allergen. sequence similarity to patatins.

AAF25553.1 AF113546 *Hevea brasiliensis*  
latex protein allergen Hev b 7. putative PLA2; similar to *Solanum tubulin* patatin encoded by GenBank Accession Number X03932.

CAA11041.1 AJ223038 *Hevea brasiliensis*  
latex allergen. with sequence similarity to patatins.

AAC27724.1 U80598 *Hevea brasiliensis*  
latex patatin homolog. putative PLA2; latex protein allergen; similar to *Solanum tubulin* patatin encoded by GenBank Accession Number X03932.

AAK27797.1 AF318315 *Vigna unguiculata*  
patatin-like protein.

AAK18751.1 AF193067 *Vigna unguiculata*  
patatin-like protein.

AAB08428.1 U68484 *Nicotiana tabacum*  
patatin homolog.

AAD22170.1 AF061282 *Sorghum bicolor*  
patatin-like protein.



|                |          |                    |  |
|----------------|----------|--------------------|--|
| AAF98368.1     | AF158027 | Nicotiana tabacum  | patatin-like protein 1. PAT1. NtPat1.  |
| AAD22169.1     | AF061282 | Sorghum bicolor    | patatin-like protein.  |
| CAA81735.1     | Z27221   | Solanum tuberosum  | patatin.   |
| CAA31575.1     | X13178   | Solanum tuberosum  | patatin B2 (AA 1 - 386).   |
| AAA33819.1     | M18880   | Solanum tuberosum  | patatin.   |
| CAA31576.1     | X13179   | Solanum tuberosum  | patatin B1 (377 AA) (1 is 3rd base in codon).  |
| CAA27588.1     | X03956   | Solanum tuberosum  | patatin.   |
| AAA33828.1     | M21879   | Solanum tuberosum  | patatin.   |
| CAA27571.1     | X03932   | Solanum tuberosum  | patatin.   |
| CAA25592.1     | X01125   | Solanum tuberosum  | patatin.   |
| AAA66198.1     | U09331   | Solanum brevidens  | patatin precursor.   |
| AAB08427.1     | U68483   | Nicotiana tabacum  | patatin homolog.   |
| AAF98370.1     | AF158254 | Nicotiana tabacum  | patatin-like protein 2. PAT2.  |
| AAD22149.1     | AF061282 | Sorghum bicolor    | patatin-like protein. similar to the EST sequences E0496(panicle at flowering stage), R2382 (root), R2382 (root), S4036 (shoot), S3728 (shoot), S13457 (green shoot).  |
| SEQ ID NO: 875 |          |                    |  |
| BAA93022.1     | AP001552 | Oryza sativa       | ESTs C74776(E51022),C26123(C116681) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana cultivar Landsberg extra-large G-protein (AF060942). |
| SEQ ID NO: 876 |          |                    |  |
| BAA87853.1     | AP000816 | Oryza sativa       | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).                                   |
| AAK21965.1     | AY028699 | Brassica napus     | receptor protein kinase PERK1.   |
| AAD21872.1     | AF078082 | Phaseolus vulgaris | receptor-like protein kinase homolog RK20-1.   |
| AAK00425.1     | AC069324 | Oryza sativa       | Putative protein kinase. OSJNBa0071K19.11.   |

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).                                 |
| AAA33915.1 | L27821   | Oryza sativa         | receptor type serine/threonine kinase. protein kinase.  |
| BAA82556.1 | AB030083 | Populus nigra        | lectin-like protein kinase. PnLPK.  |
| AAK11674.1 | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| AAF43496.1 | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.           |
| AAG59657.1 | AC084319 | Oryza sativa         | putative protein kinase. OSJNBa0004B24.20.  |
| AAG03090.1 | AC073405 | Oryza sativa         | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).                                      |
| AAB93834.1 | U82481   | Zea mays             | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.                  |
| AAF66615.1 | AF142596 | Nicotiana tabacum    | LRR receptor-like protein kinase.   |
| AAG25966.1 | AF302082 | Nicotiana tabacum    | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment. |
| AAB09771.1 | U67422   | Zea mays             | CRINKLY4 precursor. cr4. receptor kinase homolog.   |
| AAF34428.1 | AF172282 | Oryza sativa         | receptor-like protein kinase. DUPR11.18.  |
| BAB07906.1 | AP002835 | Oryza sativa         | putative S-receptor kinase. P0417G05.14.  |
| AAC23542.1 | U20948   | Ipomoea trifida      | receptor protein kinase. IRK1.  |
| BAA94516.1 | AP001800 | Oryza sativa         | Similar to Zea mays S-domain receptor-like protein kinase (AJ010166).   |
| BAA87852.1 | AP000816 | Oryza sativa         | Similar to putative Ser/Thr protein kinase. (AC004218).   |
| BAA92221.1 | AP001278 | Oryza sativa         | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).        |
| CAA73134.1 | Y12531   | Brassica oleracea    | serine/threonine kinase. BRLK.  |
| BAA92954.1 | AP001551 | Oryza sativa         | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).   |

|            |          |                             |  |
|------------|----------|-----------------------------|--|
| AAG16628.1 | AY007545 | Brassica napus              | protein serine/threonine kinase BNK1.  |
| CAB51834.1 | 00069    | Oryza sativa                | l1332.5. contains eukaryotic protein kinase domain PF.   |
| BAA06538.1 | D31737   | Nicotiana tabacum           | protein-serine/threonine kinase.   |
| CAA73133.1 | Y12530   | Brassica oleracea           | serine /threonine kinase. ARLK.  |
| CAB89179.1 | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| AAA33008.1 | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| BAA92837.1 | AB032474 | Brassica oleracea           | S60 S-locus receptor kinase. SRK60.  |
| AAA33000.1 | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| CAA79355.1 | Z18921   | Brassica oleracea           | S-receptor kinase-like protein.  |
| CAA67145.1 | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.  |
| CAA74661.1 | Y14285   | Brassica oleracea           | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.     |
| AAA62232.1 | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| BAA23676.1 | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.  |
| BAB07904.1 | AP002835 | Oryza sativa                | putative S-receptor kinase. P0417G05.12.   |
| BAA94518.1 | AP001800 | Oryza sativa                | Similar to Arabidopsis thaliana chromosome 2 section 111 of 255; putative receptor-like protein kinase (AC002392). |
| BAA07577.2 | D38564   | Brassica rapa               | receptor protein kinase SRK12.   |
| CAA74662.1 | Y14286   | Brassica oleracea           | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.   |
| BAA07576.1 | D38563   | Brassica rapa               | receptor protein kinase SRK8.  |
| BAB07999.1 | AP002525 | Oryza sativa                | putative protein kinase. P0462H08.22. contains EST C22619(S11214).   |
| BAB07905.1 | AP002835 | Oryza sativa                | putative S-receptor kinase. P0417G05.13.   |

|                |          |                               |   |
|----------------|----------|-------------------------------|---|
| BAB03429.1     | AP002817 | Oryza sativa                  | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587). |
| SEQ ID NO: 883 |          |                               |   |
| BAA90510.2     | AP001111 | Oryza sativa                  | rice EST AU030811, similar to rice Ca <sup>2+</sup> -ATPase (U82966).   |
| AAF73985.1     | AF096871 | Zea mays                      | calcium pump. calcium ATPase. cap1.   |
| CAA63790.1     | X93592   | Dunaliella bioculata          | P-type ATPase. cal. calcium pumping; CA1.   |
| AAD11618.1     | AF050496 | Lycopersicon esculentum       | Ca <sup>2+</sup> -ATPase. LCA1B; alternative transcript.  |
| AAA34138.1     | M96324   | Lycopersicon esculentum       | The calcium ATPase is a calcium ion pump. Ca <sup>2+</sup> -ATPase. LCA1.   |
| AAB58910.1     | U82966   | Oryza sativa                  | Ca <sup>2+</sup> -ATPase.   |
| AAD11617.1     | AF050495 | Lycopersicon esculentum       | Ca <sup>2+</sup> -ATPase. LCA1A; alternative transcript.  |
| AAG28435.1     | AF195028 | Glycine max                   | plasma membrane Ca <sup>2+</sup> -ATPase. SCA1.   |
| AAG28436.1     | AF195029 | Glycine max                   | plasma membrane Ca <sup>2+</sup> -ATPase. SCA2.   |
| CAA68234.1     | X99972   | Brassica oleracea             | calmodulin-stimulated calcium-ATPase.   |
| AAD31896.1     | AF145478 | Mesembryanthemum crystallinum | calcium ATPase.   |
| AAB60276.1     | U09989   | Zea mays                      | H <sup>(+)</sup> -transporting ATPase. Mha1.  |
| CAB69824.1     | AJ271439 | Prunus persica                | plasma membrane H <sup>+</sup> ATPase. PPA1.  |
| AAD46187.1     | AF156683 | Nicotiana glauca              | plasma membrane proton ATPase. pma8.  |
| BAA01058.1     | D10207   | Oryza sativa                  | H-ATPase. OSA1.   |
| AAB49042.1     | U54690   | Dunaliella acidophila         | plasma membrane proton ATPase. dha1. DaDHA1; proton pump.   |
| AAA34173.1     | M60166   | Lycopersicon esculentum       | H <sup>+</sup> -ATPase. LHA1.   |
| CAA52107.1     | X73901   | Dunaliella bioculata          | plasma membrane ATPase. pma1.   |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAB35314.2 | S79323   | Vicia faba                    | plasma membrane H(+)-ATPase precursor, plasma membrane H(+)-ATPase. This sequence comes from Fig. 1; conceptual translation presented here differs from translation in publication. |
| BAA06629.1 | D31843   | Oryza sativa                  | plasma membrane H <sup>+</sup> -ATPase. OSA2.   |
| AAA34094.1 | M80489   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> ATPase. pma1.  |
| BAA08134.1 | D45189   | Zostera marina                | plasma membrane H <sup>+</sup> -ATPase. zha1.   |
| CAB85494.1 | AJ132891 | Medicago truncatula           | proton pump. H <sup>+</sup> -ATPase. ha1.   |
| CAB85495.1 | AJ132892 | Medicago truncatula           | proton pump. H <sup>+</sup> -ATPase. ha1.   |
| AAB84202.2 | AF029256 | Kosteletzkya virginica        | plasma membrane proton ATPase. ATP1.  |
| CAA47275.1 | X66737   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> -ATPase. pma4.   |
| CAA54045.1 | X76535   | Solanum tuberosum             | H(+)-transporting ATPase. PHA2.   |
| AAD46186.1 | AF156679 | Nicotiana plumbaginifolia     | plasma membrane proton ATPase. pma6.  |
| AAB17186.1 | U72148   | Lycopersicon esculentum       | plasma membrane H <sup>+</sup> -ATPase. LHA4, plasma membrane proton pumping ATPase.  |
| CAB69823.1 | AJ271438 | Prunus persica                | plasma membrane H <sup>+</sup> ATPase. PPA2.  |
| AAB41898.1 | U84891   | Mesembryanthemum crystallinum | plasma membrane proton pump. H <sup>+</sup> -transporting ATPase. PMA.  |
| CAC29436.1 | AJ310524 | Vicia faba                    | P-type H <sup>+</sup> -ATPase. ha5, predominantly expressed in guard cells and flowers.   |
| BAA37150.1 | AB022442 | Vicia faba                    | p-type H <sup>+</sup> -ATPase. VHA2.  |
| CAA59800.1 | X85805   | Zea mays                      | H(+)-transporting ATPase. MHA-2.  |
| CAA59799.1 | X85804   | Phaseolus vulgaris            | H(+)-transporting ATPase. BHA-1.  |
| CAC29435.1 | AJ310523 | Vicia faba                    | P-type H <sup>+</sup> -ATPase. vha4, predominantly expressed in flowers.  |
| AAD46188.1 | AF156691 | Nicotiana plumbaginifolia     | plasma membrane proton ATPase. pma9.  |
| AAA34099.1 | M80491   | Nicotiana plumbaginifolia     | plasma membrane H <sup>+</sup> ATPase. pma3.  |

AAA34052.1 M27888 *Nicotiana plumbaginifolia*  
H<sup>+</sup>-translocating ATPase.

CAA54046.1 X76536 *Solanum tuberosum*  
H(+)-transporting ATPase. PHA1.

AAA34098.1 M80490 *Nicotiana plumbaginifolia*  
plasma membrane H<sup>+</sup> ATPase. pma3.

AAD55399.1 AF179442 *Lycopersicon esculentum*  
plasma membrane H<sup>+</sup>-ATPase isoform LHA2. LHA2.

AAF98344.1 AF275745 *Lycopersicon esculentum*  
plasma membrane H<sup>+</sup>-ATPase. LHA2. P-type ion pump.

AAG01028.1 AF289025 *Cucumis sativus*  
plasma membrane H<sup>+</sup>-ATPase.

AAK31799.1 AY029190 *Lilium longiflorum*  
plasma membrane H<sup>+</sup> ATPase. LILHA1.

AAA81348.1 U38965 *Vicia faba*  
p-type H<sup>+</sup>-ATPase. VHA2.

AAK32119.1 AF308817 *Hordeum vulgare*  
plasmalemma H<sup>+</sup>-ATPase 2.

AAK32118.1 AF308816 *Hordeum vulgare*  
plasmalemma H<sup>+</sup>-ATPase 1.

AAA20600.1 U08984 *Zea mays*  
plasma-membrane H<sup>+</sup> ATPase. Zmpma1.

SEQ ID NO: 884

AAD21872.1 AF078082 *Phaseolus vulgaris*  
receptor-like protein kinase homolog RK20-1.

AAB93834.1 U82481 *Zea mays*  
KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.

CAA73134.1 Y12531 *Brassica oleracea*  
serine/threonine kinase. BRLK.

AAC23542.1 U20948 *Ipomoea trifida*  
receptor protein kinase. IRK1.

CAA67145.1 X98520 *Brassica oleracea*  
receptor-like kinase. SFR2.

CAA73133.1 Y12530 *Brassica oleracea*  
serine /threonine kinase. ARLK.

BAA23676.1 AB000970 *Brassica rapa*  
receptor kinase 1. BcRK1.

CAB41879.1 Y18260 *Brassica oleracea*  
SRK15 protein. SRK15. receptor-like kinase.

CAB41878.1 Y18259 *Brassica oleracea*  
SRK5 protein. SRK5. receptor-like kinase.

|            |          |                             |  |
|------------|----------|-----------------------------|--|
| CAA74662.1 | Y14286   | Brassica oleracea           | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413. |
| CAA74661.1 | Y14285   | Brassica oleracea           | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554.   |
| BAA06285.1 | D30049   | Brassica rapa               | S-receptor kinase SRK9.  |
| BAA21132.1 | D88193   | Brassica rapa               | S-receptor kinase. SRK9 (B.c).   |
| BAA92836.1 | AB032473 | Brassica oleracea           | S18 S-locus receptor kinase. SRK18.  |
| CAA79355.1 | Z18921   | Brassica oleracea           | S-receptor kinase-like protein.  |
| AAA33000.1 | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| AAA62232.1 | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| AAA33008.1 | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| CAB89179.1 | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| BAB18292.1 | AP002860 | Oryza sativa                | putative receptor-like protein kinase. P0409B08.19.  |
| AAD52097.1 | AF088885 | Nicotiana tabacum           | receptor-like kinase CHRK1. Chrk1.   |
| BAA92837.1 | AB032474 | Brassica oleracea           | S60 S-locus receptor kinase. SRK60.  |
| BAB21001.1 | AB054061 | Brassica rapa               | S locus receptor kinase. SRK22.  |
| BAA07576.1 | D38563   | Brassica rapa               | receptor protein kinase SRK8.  |
| BAA07577.2 | D38564   | Brassica rapa               | receptor protein kinase SRK12.   |
| AAK02023.1 | AC074283 | Oryza sativa                | Putative protein kinase-like. OSJNBa0087H07.5.   |
| CAA79324.1 | Z18884   | Brassica oleracea           | S-receptor kinase related protein.   |
| BAB16871.1 | AP002537 | Oryza sativa                | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).       |

BAB39873.1 AP002882 *Oryza sativa*  
putative protein kinase. P0439B06.8. contains ESTs  
AU056701(S20808),AU056702(S20808).

AAK21965.1 AY028699 *Brassica napus*  
receptor protein kinase PERK1.

BAA87853.1 AP000816 *Oryza sativa*  
EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative  
NAK-like Ser/Thr protein kinase. (AF001308).

CAB51836.1 AJ243961 *Oryza sativa*  
Putitive Ser/Thr protein kinase. 11332.7.

AAA33915.1 L27821 *Oryza sativa*  
receptor type serine/threonine kinase. protein kinase.

BAB21240.1 AP002953 *Oryza sativa*  
Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).

AAK00425.1 AC069324 *Oryza sativa*  
Putative protein kinase. OSJNBa0071K19.11.

BAA92954.1 AP001551 *Oryza sativa*  
Similar to *Oryza sativa* protein kinase (OSPK10) mRNA. (L27821).

AAF66615.1 AF142596 *Nicotiana tabacum*  
LRR receptor-like protein kinase.

BAB17139.1 AP002867 *Oryza sativa*  
putative receptor kinase. P0463F06.31.

SEQ ID NO: 885

CAA80358.1 Z22645 *Solanum tuberosum*  
cleavage of sucrose to glucose and fructose. beta-fructofuranosidase.

CAA79676.1 Z21486 *Solanum tuberosum*  
cleavage of sucrose to glucose and fructose. beta-fructofuranosidase.

CAA57428.1 X81834 *Nicotiana tabacum*  
beta-fructofuranosidase. Ntbfruc1. beta-fructosidase.

CAA49162.1 X69321 *Daucus carota*  
beta-fructofuranosidase. Inval\*DC1.

CAA57389.1 X81792 *Chenopodium rubrum*  
beta-fructofuranosidase. CIN1.

AAC17166.1 AF063246 *Pisum sativum*  
hydrolyzes sucrose to glucose and fructose. cell wall invertase. bfruct1. isoform Pcl-2; the  
Pcl-1 isoform is in the file with GenBank Accession Number X85327; beta-  
fructofuranosidase.

CAA59677.1 X85327 *Pisum sativum*  
hydrplyze sucrose into fructose and glucose. invertase. bfruct1. beta-fructofuranosidase.

AAD02263.1 AF043346 *Zea mays*  
sucrose hydrolysis. cell wall invertase. incw3. Incw3; beta-fructofuranosidase.

CAA84526.1 Z35162 *Vicia faba*  
hydrolyze sucrose. beta-fructofuranosidase; cell wall invertase I; fructosidase. VFCWINV1.



|                |          |                    |   |
|----------------|----------|--------------------|---|
| AAC96065.1     | AF030420 | Triticum aestivum  | hydrolyzes sucrose. cell wall invertase. IVR1. hydrolase; beta-fructofuranosidase; fructosidase.  |
| CAA53099.1     | X75353   | Daucus carota      | beta-fructofuranosidase.  |
| AAB68679.1     | U92438   | Phaseolus vulgaris | soluble acid invertase. PVSAI; potential vacuolar targeted enzyme; beta-fructofuranosidase.   |
| CAA89992.1     | Z49831   | Vicia faba         | hydrolyze sucrose. vacuolar invertase; beta-fructofuranosidase. VFVCINV.  |
| CAA53097.1     | X75351   | Daucus carota      | beta-fructofuranosidase.  |
| CAA77267.1     | Y18707   | Daucus carota      | beta-fructofuranosidase, isoform I. Inv*Dc4. soluble acid invertase.  |
| CAA53098.1     | X75352   | Daucus carota      | beta-fructofuranosidase.  |
| CAA77266.1     | Y18706   | Daucus carota      | beta-fructofuranosidase, isoform II. Inv*Dc5. soluble acid invertase.   |
| AAC96066.1     | AF030421 | Triticum aestivum  | hydrolyzes sucrose. cell wall invertase. IVR3. hydrolase; beta-fructofuranosidase; fructosidase.  |
| AAG36943.1     | AF274299 | Brassica oleracea  | cleaves sucrose into glucose and fructose at acid pH optima. acid invertase AI7-3. sucrose hydrolysing enzyme; beta-fructofuranosidase. |
| SEQ ID NO: 887 |          |                    |   |
| AAD10836.1     | U52079   | Solanum tuberosum  | P-glycoprotein. pmdr1. binds ATP; ATPase; transporter; transmembrane protein.   |
| BAA96612.1     | AP002482 | Oryza sativa       | Similar to Arabidopsis thaliana chromosome 2, BAC F14M4 ; putative ABC transporter (AC004411).  |
| AAG49002.1     | AY013246 | Hordeum vulgare    | putative ABC transporter. 635P2.4a; GC splice donor confirmed by cDNA and comparative sequencing.                                       |
| AAG45492.1     | AY013245 | Oryza sativa       | 36I5.4. putative ABC transporter; GC splice donor confirmed by cDNA alignment and comparative sequence.                                 |
| BAA83352.1     | AP000391 | Oryza sativa       | ESTs AU067992(C11433),AU077424(C11433) correspond to a region of the predicted gene.; Similar to ABC transporter-7 (U43892).            |
| AAG49003.1     | AY013246 | Hordeum vulgare    | putative ABC transporter. 635P2.4b; GC splice donor confirmed by cDNA alignment and comparative sequence.                               |
| BAB17113.1     | AP002866 | Oryza sativa       | putative white protein; ATP-binding cassette transporter. P0410E01.34.  |

|                |          |                             |  |
|----------------|----------|-----------------------------|--|
| BAA90508.1     | AP001111 | Oryza sativa                | similar to ABC transporter of Arabidopsis thaliana (AC004697).   |
| BAA90507.1     | AP001111 | Oryza sativa                | similar to ABC transporter of Arabidopsis thaliana (AC004697).   |
| BAB16495.1     | AP002861 | Oryza sativa                | putative ABC transporter ATP-binding protein. P0665D10.21.   |
| BAB21276.1     | AP002844 | Oryza sativa                | putative ABC transporter protein. P0410E03.7. contains EST D22472(C1173).                                      |
| BAB21275.1     | AP002844 | Oryza sativa                | putative ABC transporter protein. P0410E03.6.  |
| BAB21273.1     | AP002844 | Oryza sativa                | putative ABC transporter protein. P0410E03.4.  |
| BAB40032.1     | AP003046 | Oryza sativa                | putative ABC transporter. P0445D12.3.  |
| SEQ ID NO: 888 |          |                             |  |
| AAD21872.1     | AF078082 | Phaseolus vulgaris          | receptor-like protein kinase homolog RK20-1.   |
| CAA73134.1     | Y12531   | Brassica oleracea           | serine/threonine kinase. BRLK.   |
| AAB93834.1     | U82481   | Zea mays                    | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.           |
| AAC23542.1     | U20948   | Ipomoea trifida             | receptor protein kinase. IRK1.   |
| AAA33000.1     | M76647   | Brassica oleracea           | receptor protein kinase. SKR6.   |
| CAB89179.1     | AJ245479 | Brassica napus subsp. napus | ser /thr kinase. S-locus receptor kinase. srk.   |
| AAA33008.1     | M97667   | Brassica napus              | serine/threonine kinase receptor.  |
| CAA74661.1     | Y14285   | Brassica oleracea           | SFR1. extracellular S domain: 8-1342; transmembrane domain: 1343-1411; intracellular kinase domain: 1412-2554. |
| CAA67145.1     | X98520   | Brassica oleracea           | receptor-like kinase. SFR2.  |
| AAA62232.1     | U00443   | Brassica napus              | S-receptor kinase. protein contains an immunoglobulin-like domain.   |
| CAA73133.1     | Y12530   | Brassica oleracea           | serine /threonine kinase. ARLK.  |
| BAA23676.1     | AB000970 | Brassica rapa               | receptor kinase 1. BcRK1.  |
| BAA92836.1     | AB032473 | Brassica oleracea           | S18 S-locus receptor kinase. SRK18.  |

|            |          |                   |  |
|------------|----------|-------------------|--|
| CAA74662.1 | Y14286   | Brassica oleracea | SFR3. extracellular S domain: 123-1346; transmembrane domain: 1347-1412; intracellular kinase domain: from 1413.                     |
| CAA79355.1 | Z18921   | Brassica oleracea | S-receptor kinase-like protein.  |
| CAB41879.1 | Y18260   | Brassica oleracea | SRK15 protein. SRK15. receptor-like kinase.  |
| BAA06285.1 | D30049   | Brassica rapa     | S-receptor kinase SRK9.  |
| BAA21132.1 | D88193   | Brassica rapa     | S-receptor kinase. SRK9 (B.c).   |
| CAB41878.1 | Y18259   | Brassica oleracea | SRK5 protein. SRK5. receptor-like kinase.  |
| BAA92837.1 | AB032474 | Brassica oleracea | S60 S-locus receptor kinase. SRK60.  |
| BAA07577.2 | D38564   | Brassica rapa     | receptor protein kinase SRK12.   |
| BAB21001.1 | AB054061 | Brassica rapa     | S locus receptor kinase. SRK22.  |
| BAA07576.1 | D38563   | Brassica rapa     | receptor protein kinase SRK8.  |
| AAD52097.1 | AF088885 | Nicotiana tabacum | receptor-like kinase CHRK1. Chrk1.   |
| AAK02023.1 | AC074283 | Oryza sativa      | Putative protein kinase-like. OSJNBa0087H07.5.   |
| BAB18292.1 | AP002860 | Oryza sativa      | putative receptor-like protein kinase. P0409B08.19.  |
| BAA87853.1 | AP000816 | Oryza sativa      | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308). |
| AAK21965.1 | AY028699 | Brassica napus    | receptor protein kinase PERK1.   |
| BAB21240.1 | AP002953 | Oryza sativa      | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).   |
| BAB39873.1 | AP002882 | Oryza sativa      | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |
| BAB17331.1 | AP002747 | Oryza sativa      | putative receptor kinase. P0698G03.12.   |
| BAB17139.1 | AP002867 | Oryza sativa      | putative receptor kinase. P0463F06.31.   |
| AAK00425.1 | AC069324 | Oryza sativa      | Putative protein kinase. OSJNBa0071K19.11.   |

---

BAB16871.1 AP002537 *Oryza sativa*  
putative protein kinase APK1A *Arabidopsis thaliana*. P0001B06.24. contains ESTs  
C22608(R3192), D25110(R3192).

---

BAA92954.1 AP001551 *Oryza sativa*  
Similar to *Oryza sativa* protein kinase (OSPK10) mRNA. (L27821).

---

CAB51836.1 AJ243961 *Oryza sativa*  
Putative Ser/Thr protein kinase. I1332.7.

---

AAF78016.1 AF238472 *Oryza sativa*  
receptor-like kinase. RLG15. protein kinase.

---

BAB39451.1 AP003338 *Oryza sativa*  
putative receptor kinase. OJ1212\_B09.24.

---

SEQ ID NO: 890

---

AAB47181.1 S82324 *Zea mays*  
/gene="calcium/calmodulin-dependent protein kinase. This sequence comes from Fig. 1.

---

BAA22410.1 D38452 *Zea mays*  
calcium-dependent protein kinase-related kinase.

---

BAA12692.1 D84508 *Zea mays*  
CDPK-related protein kinase. Does not require calcium for its activity.

---

AAG01179.1 AF289237 *Zea mays*  
calcium/calmodulin dependent protein kinase MCK2. MCK2.

---

CAA58750.1 X83869 *Daucus carota*  
CDPK-related protein kinase. CRK (or PK421).

---

BAA12691.1 D84507 *Zea mays*  
CDPK-related protein kinase. Does not require calcium for its activity (by similarity).

---

AAC24961.1 AF009337 *Tradescantia virginiana*  
CDPK-related protein kinase. CRK1.

---

AAF23901.2 AF194414 *Oryza sativa*  
calcium-dependent protein kinase. CDPK5. OsCDPK5.

---

AAF23900.1 AF194413 *Oryza sativa*  
calcium-dependent protein kinase. CDPK1. OsCDPK1.

---

AAC78558.1 AF030879 *Solanum tuberosum*  
protein kinase CPK1.

---

AAD17800.1 AF090835 *Mesembryanthemum crystallinum*  
Ca<sup>2+</sup>-dependent protein kinase. CPK1. serine/threonine protein kinase.

---

AAB49984.1 U90262 *Cucurbita pepo*  
calcium-dependent calmodulin-independent protein kinase CDPK. cpCPK1. serine/threonine  
protein kinase that is activated by direct binding of calcium.

---

CAA07481.1 AJ007366 *Zea mays*  
calcium-dependent protein kinase.

---

BAB21081.1 AP002819 *Oryza sativa*  
putative calcium-dependent protein kinase. P0501G01.10.

---

BAA12715.1 D85039 *Zea mays*  
calcium-dependent protein kinase.

---

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAC25423.1 | AF072908 | Nicotiana tabacum       | calcium-dependent protein kinase. CDPK1.                               |
| CAA39936.1 | X56599   | Daucus carota           | calcium- dependent protein kinase. DcPK431.                            |
| CAA57157.1 | X81394   | Oryza sativa            | calcium-dependent protein kinase. OSCPK2.                              |
| AAF21062.1 | AF216527 | Dunaliella tertiolecta  | calcium-dependent protein kinase. CPK1; CDPK.                          |
| BAA81751.1 | AB017517 | Marchantia polymorpha   | calcium-dependent protein kinase. CDPK1. CDPK-B; alternative splicing. |
| BAA81749.1 | AB017515 | Marchantia polymorpha   | calcium-dependent protein kinase. CDPK1. CDPK-B; alternative splicing. |
| AAB88537.1 | AF035944 | Fragaria x ananassa     | calcium-dependent protein kinase. MAX17.                               |
| BAA81750.1 | AB017516 | Marchantia polymorpha   | calcium-dependent protein kinase. CDPK1. CDPK-A; alternative splicing. |
| BAA81748.1 | AB017515 | Marchantia polymorpha   | calcium-dependent protein kinase. CDPK1. CDPK-A; alternative splicing. |
| AAA69507.1 | U28376   | Zea mays                | calcium-dependent protein kinase. MZECDPK2.                            |
| AAA61682.1 | L27484   | Zea mays                | calcium-dependent protein kinase. CDPK.                                |
| AAB80693.1 | U69174   | Glycine max             | calmodulin-like domain protein kinase isoenzyme gamma. CDPK gamma.     |
| BAA13232.1 | D87042   | Zea mays                | Calcium-dependent protein kinase.                                      |
| BAA13440.1 | D87707   | Ipomoea batatas         | calcium dependent protein kinase. CDPK.                                |
| CAA89202.1 | Z49233   | Chlamydomonas eugametos | calcium-stimulated protein kinase.                                     |
| AAB70706.1 | U82087   | Tortula ruralis         | calmodulin-like domain protein kinase. TrCPK1.                         |
| AAD28192.2 | AF115406 | Solanum tuberosum       | calcium-dependent protein kinase. CDPK; catalytic domain.              |
| AAC49405.1 | U08140   | Vigna radiata           | calcium dependent protein kinase. CDPK.                                |
| BAA12338.1 | D84408   | Zea mays                | calcium dependent protein kinase. ZmCDPK1.                             |
| CAA65500.1 | X96723   | Medicago sativa         | protein kinase. CDPK.  |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| BAA85396.1     | AP000615 | Oryza sativa            | ESTs C22369(C12239),C22370(C12239), AU057852(S21844),AU057853(S21844) correspond to a region of the predicted gene.; similar to calcium dependent protein kinase. (AF048691). |
| AAA33443.1     | L15390   | Zea mays                | calcium-dependent protein kinase. CDPK.   |
| AAB80692.1     | U69173   | Glycine max             | calmodulin-like domain protein kinase isoenzyme beta. CDPK beta.  |
| CAA57156.1     | X81393   | Oryza sativa            | calcium-dependent protein kinase. OSCPKII.  |
| AAC05270.1     | AF048691 | Oryza sativa            | calcium dependent protein kinase. CDPK12.   |
| AAK26164.1     | AY027885 | Cucumis sativus         | calcium-dependent calmodulin-independent protein kinase 5. CDPK5. CsCDPK5.  |
| AAG46110.1     | AC073166 | Oryza sativa            | calcium-dependent protein kinase. OSJNBb0064P21.2.  |
| BAA02698.1     | D13436   | Oryza sativa            | calcium-dependent protein kinase. spk.  |
| BAA90814.1     | AP001168 | Oryza sativa            | ESTs AU030197(E50746),AU030196(E50746) correspond to a region of the predicted gene.; Similar to calcium-dependent calmodulin-independent protein kinase CDPK (U90262).       |
| AAF19401.1     | AF203479 | Glycine max             | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                          |
| CAB46228.1     | Y18055   | Arachis hypogaea        | calcium dependent protein kinase. CDPK.   |
| AAF19403.1     | AF203481 | Lycopersicon esculentum | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                          |
| AAF19402.1     | AF203480 | Lycopersicon esculentum | phosphoenolpyruvate carboxylase kinase. protein kinase; member of Ca <sup>2+</sup> /CaM kinase family; lacks the autoinhibitory region and EF hands.                          |
| SEQ ID NO: 892 |          |                         |   |
| AAF20931.1     | AF206721 | Brassica juncea         | ascorbate oxidase.  |
| BAA07734.1     | D43624   | Nicotiana tabacum       | ascorbate oxidase precursor.  |
| AAA33119.1     | J04494   | Cucumis sativus         | ascorbate oxidase precursor (EC 1.10.3.3).  |
| CAA75577.1     | Y15295   | Medicago truncatula     | L-ascorbate oxidase. MtN23.   |
| AAF35911.2     | AF233594 | Cucumis melo            | ascorbate oxidase AO4. multicopper oxidase.   |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| CAA39300.1 | X55779   | Cucurbita sp.           | ascorbate oxidase.  |
| BAA09528.1 | D55677   | Cucurbita maxima        | ascorbate oxidase. AAO.   |
| AAF35910.1 | AF233593 | Cucumis melo            | ascorbate oxidase AO1. multicopper oxidase.   |
| CAA71275.1 | Y10226   | Cucumis melo            | L-ascorbate oxidase. ao3.   |
| AAF20932.1 | AF206722 | Brassica juncea         | ascorbate oxidase.  |
| AAF20933.1 | AF206723 | Brassica juncea         | ascorbate oxidase.  |
| CAA71273.1 | Y10224   | Cucumis melo            | L-ascorbate oxidase. ao1.   |
| CAA71274.1 | Y10225   | Cucumis melo            | L-ascorbate oxidase. ao1.   |
| BAA20520.1 | AB004799 | Oryza sativa            | ascorbate oxidase.  |
| AAF33751.1 | AF202460 | Capsicum annuum         | ascorbic acid oxidase.  |
| AAB17193.1 | U73105   | Liriodendron tulipifera | monolignol polymerization; lignin biosynthesis. laccase. LAC2-3. diphenol oxidase; blue copper oxidase. |
| AAB17191.1 | U73103   | Liriodendron tulipifera | monolignol polymerization; lignin biosynthesis. laccase. LAC2-1. diphenol oxidase; blue copper oxidase. |
| AAC49536.1 | U43542   | Nicotiana tabacum       | diphenol oxidase. laccase.  |
| AAB17194.1 | U73106   | Liriodendron tulipifera | monolignol polymerization; lignin biosynthesis. laccase. LAC2-4. diphenol oxidase; blue copper oxidase. |
| AAB17192.1 | U73104   | Liriodendron tulipifera | monolignol polymerization; lignin biosynthesis. laccase. LAC2-2. diphenol oxidase; blue copper oxidase. |
| AAC04576.1 | AF047697 | Oryza sativa            | p-diphenol oxidase. putative high-pI laccase.   |
| AAB09228.1 | U12757   | Acer pseudoplatanus     | monolignol polymerization; lignin biosynthesis. laccase. diphenol oxidase.                              |
| CAA45554.1 | X64257   | Brassica napus          | Bp10. protein homologous to ascorbate oxidase.  |
| AAC49538.1 | U45243   | Nicotiana tabacum       | diphenol oxidase. laccase.  |

|                |          |                              |  |
|----------------|----------|------------------------------|--|
| AAC49537.1     | U43543   | Nicotiana tabacum            | diphenol oxidase. laccase.   |
| AAD02557.1     | AF049931 | Petunia x hybrida            | PGPS/NH15. PGPS/NH15. ascorbate oxidase homolog.   |
| SEQ ID NO: 894 |          |                              |  |
| CAB43505.1     | AJ239051 | Cicer arietinum              | cytochrome P450. cyp81E2.  |
| BAA74465.1     | AB022732 | Glycyrrhiza echinata         | cytochrome P450. CYP Ge-31.  |
| BAA93634.1     | AB025016 | Lotus japonicus              | cytochrome P450.   |
| BAA22422.1     | AB001379 | Glycyrrhiza echinata         | cytochrome P450. CYP81E1.  |
| CAA04117.1     | AJ000478 | Helianthus tuberosus         | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-race).                                    |
| CAB41490.1     | AJ238439 | Cicer arietinum              | cytochrome P450 monooxygenase. cyp81E3v2.  |
| CAA04116.1     | AJ000477 | Helianthus tuberosus         | fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.  |
| CAA10067.1     | AJ012581 | Cicer arietinum              | cytochrome P450. cyp81E3.  |
| AAK38079.1     | AF321855 | Lolium rigidum               | putative cytochrome P450.  |
| AAK38080.1     | AF321856 | Lolium rigidum               | putative cytochrome P450.  |
| AAK38081.1     | AF321857 | Lolium rigidum               | putative cytochrome P450.  |
| AAD56282.1     | AF155332 | Petunia x hybrida            | flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.   |
| AAC34853.1     | AF082028 | Hemerocallis hybrid cultivar | putative cyt P450-containing fatty acid hydroxylase. senescence-associated protein 3. SA3. mRNA accumulates in senescing petals. |
| AAG09208.1     | AF175278 | Pisum sativum                | wound-inducible P450 hydroxylase. CYP82A1.   |
| BAA12159.1     | D83968   | Glycine max                  | Cytochrome P-450 (CYP93A1).  |
| CAA71515.1     | Y10491   | Glycine max                  | putative cytochrome P450.  |
| CAA71876.1     | Y10982   | Glycine max                  | putative cytochrome P450.  |
| CAA65580.1     | X96784   | Nicotiana tabacum            | cytochrome P450. hsr515.   |



|                |          |  |
|----------------|----------|--|
| AAC49188.2     | U29333   | <i>Pisum sativum</i><br>cytochrome P450 monooxygenase. CYP82. new cytochrome P450 family.  |
| AAG44132.1     | AF218296 | <i>Pisum sativum</i><br>cytochrome P450. P450 isolog.  |
| CAA71516.1     | Y10492   | <i>Glycine max</i><br>putative cytochrome P450.  |
| CAA64635.1     | X95342   | <i>Nicotiana tabacum</i><br>cytochrome P450. hsr515. hypersensitivity-related gene.  |
| AAC39454.1     | AF014802 | <i>Eschscholzia californica</i><br>(S)-N-methylcoclaurine 3'-hydroxylase. CYP82B1. cytochrome P-450-dependent monooxygenase; methyl jasmonate-inducible cytochrome P-450-dependent, homologous to wound-inducible CYP82A1 of <i>Pisum sativum</i> GenBank Accession Number U29333. |
| AAB94590.1     | AF022461 | <i>Glycine max</i><br>CYP82C1p. CYP82C1. cytochrome P450 monooxygenase.  |
| AAA32913.1     | M32885   | <i>Persea americana</i><br>cytochrome P-450LXXIA1 (cyp71A1).   |
| CAA71877.1     | Y10983   | <i>Glycine max</i><br>putative cytochrome P450.  |
| BAA84072.1     | AB028152 | <i>Torenia hybrida</i><br>flavone synthase II. cytochrome P450. TFNS5.   |
| BAA13076.1     | D86351   | <i>Glycine max</i><br>cytochrome P-450 (CYP93A2).  |
| AAG34695.1     | AF313492 | <i>Matthiola incana</i><br>putative cytochrome P450.   |
| BAA84071.1     | AB028151 | <i>Antirrhinum majus</i><br>flavone synthase II. cytochrome P450. AFNS2.   |
| AAC32274.1     | AF081575 | <i>Petunia x hybrida</i><br>flavonoid 3',5'-hydroxylase. Hfl. P450 enzyme.   |
| BAA35080.1     | AB015762 | <i>Nicotiana tabacum</i><br>putative cytochrome P450. CYP82E1.   |
| CAA50155.1     | X70824   | <i>Solanum melongena</i><br>flavonoid hydroxylase (P450). CYP75.   |
| CAA70575.1     | Y09423   | <i>Nepeta racemosa</i><br>cytochrome P450. CYP71A5.  |
| BAA92894.1     | AB006790 | <i>Petunia x hybrida</i><br>cytochrome P450. IMT-2.  |
| CAA71513.1     | Y10489   | <i>Glycine max</i><br>putative cytochrome P450.  |
| AAB94587.1     | AF022458 | <i>Glycine max</i><br>CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.  |
| SEQ ID NO: 895 |          |  |

|            |          |                      |   |
|------------|----------|----------------------|---|
| BAA78764.1 | AB023482 | Oryza sativa         | ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAF43496.1 | AF131222 | Lophopyrum elongatum | protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and ABA treatment.   |
| AAK11674.1 | AF339747 | Lophopyrum elongatum | protein kinase. ESI47.  |
| AAG16628.1 | AY007545 | Brassica napus       | protein serine/threonine kinase BNK1.   |
| BAA94510.1 | AB041504 | Populus nigra        | protein kinase 2. PnPK2.  |
| BAA94509.1 | AB041503 | Populus nigra        | protein kinase 1. PnPK1.  |
| BAA87853.1 | AP000816 | Oryza sativa         | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB07999.1 | AP002525 | Oryza sativa         | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB03429.1 | AP002817 | Oryza sativa         | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587).             |
| BAB16871.1 | AP002537 | Oryza sativa         | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).  |
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| BAB39409.1 | AP002901 | Oryza sativa         | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| AAK00425.1 | AC069324 | Oryza sativa         | Putative protein kinase. OSJNBa0071K19.11.  |
| BAB21240.1 | AP002953 | Oryza sativa         | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| AAK21965.1 | AY028699 | Brassica napus       | receptor protein kinase PERK1.  |
| AAG59657.1 | AC084319 | Oryza sativa         | putative protein kinase. OSJNBa0004B24.20.  |
| AAG03090.1 | AC073405 | Oryza sativa         | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| CAB51834.1 | 00069    | Oryza sativa         | 11332.5. contains eukaryotic protein kinase domain PF.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAC61805.1 | U28007   | Lycopersicon esculentum | serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.   |
| AAF91336.1 | AF249317 | Glycine max             | Pti1 kinase-like protein. Pti1a. protein kinase.   |
| AAF91337.1 | AF249318 | Glycine max             | Pti1 kinase-like protein. Pti1b. protein kinase.   |
| BAB21241.1 | AP002953 | Oryza sativa            | Putative Pto kinase interactor 1. P0426D06.21. contains ESTs AU108280(E0721),D48017(S13927).                                       |
| AAB09771.1 | U67422   | Zea mays                | CRINKLY4 precursor. cr4. receptor kinase homolog.  |
| CAA97692.1 | Z73295   | Catharanthus roseus     | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis. |
| BAA92221.1 | AP001278 | Oryza sativa            | Similar to Arabidopsis thaliana chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).                     |
| BAA87852.1 | AP000816 | Oryza sativa            | Similar to putative Ser/Thr protein kinase. (AC004218).  |
| AAB61708.1 | U93048   | Daucus carota           | somatic embryogenesis receptor-like kinase. SERK.  |
| AAG25966.1 | AF302082 | Nicotiana tabacum       | cytokinin-regulated kinase 1. CRK1. protein kinase; transcript abundance decreases rapidly after cytokinin treatment.              |
| BAA90808.1 | AP001168 | Oryza sativa            | Similar to putative receptor-like protein kinase (AL035679).   |
| AAF76313.1 | AF220603 | Lycopersicon esculentum | Pto kinase. LescPth5.  |
| AAB47421.1 | U59316   | Lycopersicon esculentum | serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.                         |
| AAC23542.1 | U20948   | Ipomoea trifida         | receptor protein kinase. IRK1.   |
| BAB19337.1 | AP003044 | Oryza sativa            | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).   |
| AAC27894.1 | AF023164 | Zea mays                | leucine-rich repeat transmembrane protein kinase 1. ltk1.  |
| AAF66615.1 | AF142596 | Nicotiana tabacum       | LRR receptor-like protein kinase.  |
| CAB51480.1 | Y14600   | Sorghum bicolor         | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.   |

|                |          |                       |   |
|----------------|----------|-----------------------|---|
| BAA92954.1     | AP001551 | Oryza sativa          | Similar to Oryza sativa protein kinase (OSPK10) mRNA. (L27821).   |
| BAA92953.1     | AP001551 | Oryza sativa          | Similar to Arabidopsis thaliana chromosome 4 BAC clone F10M6 ; S-receptor kinase -like protein. (AL021811).   |
| AAK11566.1     | AF318490 | Lycopersicon hirsutum | Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.   |
| AAK11567.1     | AF318491 | Lycopersicon hirsutum | Pto-like protein kinase F. LhirPtoF.  |
| BAA82556.1     | AB030083 | Populus nigra         | lectin-like protein kinase. PnLPK.  |
| CAA67145.1     | X98520   | Brassica oleracea     | receptor-like kinase. SFR2.   |
| SEQ ID NO: 899 |          |                       |   |
| AAF61647.1     | AF190634 | Nicotiana tabacum     | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.   |
| BAA89009.1     | AB027455 | Petunia x hybrida     | anthocyanin 5-O-glucosyltransferase. PH1.   |
| AAA59054.1     | L34847   | Zea mays              | conjugation of the growth hormone indole-3-acetic acid (IAA). IAA-glu synthetase. iaglu.  |
| BAA93039.1     | AB033758 | Citrus unshiu         | limonoid UDP-glucosyltransferase. LGTase.   |
| BAA36423.1     | AB013598 | Verbena x hybrida     | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| BAA36421.1     | AB013596 | Perilla frutescens    | UDP-glucose:anthocynin 5-O-glucosyltransferase. PF3R4.  |
| AAF98390.1     | AF287143 | Brassica napus        | catalyzes the transfer of glucose from UDP-glucose to sinapate and some other hydroxycinnamates (4-coumarate, caffeate, ferulate). UDP-glucose:sinapate glucosyltransferase. SGT1. SGT. |
| BAA36422.1     | AB013597 | Perilla frutescens    | UDP-glucose:anthocyanin 5-O-glucosyltransferase homologue. PF3R6.   |
| BAB07962.1     | AP002524 | Oryza sativa          | putative anthocyanin 5-O-glucosyltransferase. P0406H10.16. contains ESTs AU067881(C10481),AU067882(C10481).   |
| AAK16175.1     | AC079887 | Oryza sativa          | putative glucosyltransferase. OSJNBa0040E01.15.   |
| AAK16181.1     | AC079887 | Oryza sativa          | putative glucosyltransferase. OSJNBa0040E01.16.   |
| AAK16178.1     | AC079887 | Oryza sativa          | putative glucosyltransferase. OSJNBa0040E01.5.  |
| AAG25643.1     | AF303396 | Phaseolus vulgaris    | UDP-glucosyltransferase HRA25. putative; defense associated.  |

|            |          |                                 |  |
|------------|----------|---------------------------------|--|
| AAK16172.1 | AC079887 | Oryza sativa                    | putative glucosyltransferase. OSJNBa0040E01.14.  |
| BAA83484.1 | AB031274 | Scutellaria baicalensis         | UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.  |
| AAB36652.1 | U32643   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS10a.   |
| AAK28304.1 | AF346432 | Nicotiana tabacum               | phenylpropanoid:glucosyltransferase 2. togt2. glucosyltransferase.   |
| AAD21086.1 | AF127218 | Forsythia x intermedia          | adds glucose residue to position 3 of flavonoid compounds. flavonoid 3-O-glucosyltransferase. UFGT.                                  |
| AAB36653.1 | U32644   | Nicotiana tabacum               | immediate-early salicylate-induced glucosyltransferase. IS5a.  |
| BAA12737.1 | D85186   | Gentiana triflora               | UDP-glucose:flavonoid-3-glucosyltransferase.   |
| AAK28303.1 | AF346431 | Nicotiana tabacum               | phenylpropanoid:glucosyltransferase 1. togt1. glucosyltransferase.   |
| AAF17077.1 | AF199453 | Sorghum bicolor                 | UDP-glucose glucosyltransferase. UDP-glucose:p-hydroxymandelonitrile-o-glucosyltransferase.  |
| AAK16180.1 | AC079887 | Oryza sativa                    | putative glucosyltransferase. OSJNBa0040E01.21.  |
| CAA59450.1 | X85138   | Lycopersicon esculentum         | twil. homologous to glucosyltransferases.  |
| BAB41017.1 | AB047090 | Vitis labrusca x Vitis vinifera | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT1. The gene was derived from one of the parents V. labruscana cv. Ishiharawase. |
| BAB41026.1 | AB047099 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT2.  |
| BAB41024.1 | AB047097 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT2.  |
| BAB41022.1 | AB047095 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT2.  |
| BAB41020.1 | AB047093 | Vitis vinifera                  | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT2.  |
| BAA89008.1 | AB027454 | Petunia x hybrida               | anthocyanidin 3-O-glucosyltransferase. PGT8.   |
| BAB41018.1 | AB047091 | Vitis labrusca x Vitis vinifera | UDP-glucose:flavonoid 3-O-glucosyltransferase. KyUFGT2. The gene originated in one of the parents V. vinifera cv. Centennial.        |
| CAB56231.1 | Y18871   | Dorotheanthus bellidiformis     | betanidin-5-O-glucosyltransferase.   |

|                |          |                         |   |
|----------------|----------|-------------------------|---|
| AAB81683.1     | AF000372 | Vitis vinifera          | UDP glucose:flavonoid 3-o-glucosyltransferase.  |
| BAB41019.1     | AB047092 | Vitis vinifera          | UDP-glucose:flavonoid 3-O-glucosyltransferase. ItUFGT1.   |
| BAB41025.1     | AB047098 | Vitis vinifera          | UDP-glucose:flavonoid 3-O-glucosyltransferase. FIUFGT1.   |
| BAB41023.1     | AB047096 | Vitis vinifera          | UDP-glucose:flavonoid 3-O-glucosyltransferase. AIUFGT1.   |
| BAB41021.1     | AB047094 | Vitis vinifera          | UDP-glucose:flavonoid 3-O-glucosyltransferase. RuUFGT1.   |
| AAB81682.1     | AF000371 | Vitis vinifera          | UDP glucose:flavonoid 3-o-glucosyltransferase.  |
| BAA19659.1     | AB002818 | Perilla frutescens      | flavonoid 3-O-glucosyltransferase. UDP glucose.   |
| BAA90787.1     | AB038248 | Ipomoea batatas         | UDP glucose: flavonoid 3-O-glucosyltransferase. uf3gt.  |
| AAD04166.1     | AF101972 | Phaseolus lunatus       | catalyzes formation of O-glucosylzeatin from zeatin and UDPG; catalyzes formation of O-xylosylzeatin from zeatin and UDPX. zeatin O-glucosyltransferase. ZOG1. cytokinin O-glucosyltransferase. |
| AAB86473.1     | AF028237 | Ipomoea purpurea        | UDP glucose: flavonoid 3-O-glucosyltransferase. UF3GT-FL1.  |
| SEQ ID NO: 900 |          |                         |   |
| AAA33975.1     | M11395   | Glycine max             | small heat shock protein.   |
| CAA41547.1     | X58711   | Medicago sativa         | heat shock protein.   |
| AAB03893.1     | M11318   | Glycine max             | 17.5 kd heat shock protein Gmhsp17.6L.  |
| AAA33672.1     | M33899   | Pisum sativum           | 18.1 kDa heat shock protein (hsp18.1).  |
| CAA25578.1     | X01104   | Glycine max             | heat shock protein 6871 (aa 1-153).   |
| CAB55634.2     | AJ237596 | Helianthus annuus       | 17.9 kDa heat-shock protein. hsp17.9.   |
| BAA33062.1     | AB017273 | Cuscuta japonica        | low-molecular-weight heat shock protein. CJHSP17.   |
| AAA33974.1     | M11317   | Glycine max             | 17.6 kd heat shock protein Gmhsp17.6L.  |
| AAD30454.1     | AF123257 | Lycopersicon esculentum | 17.6 kD class I small heat shock protein. HSP17.6.  |
| AAD30452.1     | AF123255 | Lycopersicon esculentum | 17.7 kD class I small heat shock protein. HSP17.7.  |

|            |          |                         |  |
|------------|----------|-------------------------|--|
| CAA39603.1 | X56138   | Lycopersicon esculentum | small heat shock protein (class I).  |
| AAD30453.1 | AF123256 | Lycopersicon esculentum | 17.8 kD class I small heat shock protein. HSP17.8.                               |
| CAA37847.1 | X53851   | Daucus carota           | heat shock protein.  |
| AAA33671.1 | M33900   | Pisum sativum           | 17.9 kDa heat shock protein (hsp17.9).   |
| AAF34133.1 | AF161179 | Malus x domestica       | low molecular weight heat shock protein. Hsp1.                                   |
| CAA41546.1 | X58710   | Medicago sativa         | heat shock protein.  |
| AAB63310.1 | U46544   | Helianthus annuus       | 18.6 kDa heat-shock protein. Class I low-molecular-weight heat-shock protein.    |
| AAB63311.1 | U46545   | Helianthus annuus       | 17.7 kDa heat shock protein. Class I low-molecular-weight heat-shock protein.    |
| CAB08441.1 | Z95153   | Helianthus annuus       | 17.6 kD class I small heat-shock protein HSP17.6. Ha hsp17.6.                    |
| CAA42222.1 | X59701   | Helianthus annuus       | 17.6 kDa heat shock protein.   |
| AAC39360.1 | U63631   | Fragaria x ananassa     | LMW heat shock protein.  |
| AAA33910.1 | M80939   | Oryza sativa            | 16.9 kDa heat shock protein.   |
| CAA37848.1 | X53852   | Daucus carota           | heat shock protein.  |
| AAA33909.1 | M80938   | Oryza sativa            | 16.9 kDa heat shock protein.   |
| CAA43210.1 | X60820   | Oryza sativa            | 16.9 KD low molecular weight heat shock protein.                                 |
| CAA37864.1 | X53870   | Chenopodium rubrum      | heat-shock protein.  |
| AAD49336.1 | AF166277 | Nicotiana tabacum       | low molecular weight heat-shock protein. LHS-1. TLHS-1.                          |
| AAA61632.1 | U08601   | Papaver somniferum      | low molecular weight heat-shock protein.   |
| AAB72109.1 | AF022217 | Brassica rapa           | low molecular weight heat-shock protein. BcHSP17.6. 17.6 kDa; cytosolic class I. |
| CAA63902.1 | X94192   | Pennisetum glaucum      | heat shock protein 16.9. hsp16.9.  |
| AAB39856.1 | U81385   | Oryza sativa            | heat shock protein. Oshsp16.9C. class I, low molecular mass.                     |

|                |          |                       |   |
|----------------|----------|-----------------------|---|
| CAA46641.1     | X65725   | Zea mays              | heat shock protein 17.2. Zmhs17.2. Class I low molecular weight heat shock protein.                     |
| CAA08908.1     | AJ009880 | Castanea sativa       | molecular chaperone. cytosolic class I small heat-shock protein HSP17.5. hsp17.5.                       |
| CAB93512.1     | AJ243565 | Brassica oleracea     | putative class I small heat shock protein. HSP17.7-a protein. hsp17.7-a.                                |
| CAA63903.1     | X94193   | Pennisetum glaucum    | heat shock protein 17.9. hsp17.9.   |
| CAB36910.1     | AJ000691 | Quercus suber         | stress protein chaperone. heat shock protein 17.4. hsp17.   |
| CAA63901.1     | X94191   | Pennisetum glaucum    | heat shock protein 17.0. hsp17.0.   |
| AAC78392.1     | U83669   | Oryza sativa          | low molecular mass heat shock protein Oshsp17.3. OSHSP17.3. class I LMMHSP.                             |
| BAA02160.1     | D12635   | Oryza sativa          | 'low molecular weight heat shock protein'.  |
| AAC78393.1     | U83670   | Oryza sativa          | low molecular mass heat shock protein Oshsp18.0. OSHSP18.0. class I LMMHSP.                             |
| CAA31785.1     | X13431   | Triticum aestivum     | put. heat shock protein (AA 1 -151).  |
| CAA63570.1     | X92983   | Pseudotsuga menziesii | low molecular weight heat-shock protein.  |
| AAC78394.1     | U83671   | Oryza sativa          | low molecular mass heat shock protein Oshsp17.7. OSHSP17.7. class I LMMHSP.                             |
| CAA63571.1     | X92984   | Pseudotsuga menziesii | low molecular weight heat-shock protein.  |
| SEQ ID NO: 901 |          |                       |   |
| AAG08959.1     | AF122051 | Solanum tuberosum     | tuber-specific and sucrose-responsive element binding factor. TSF. R2R3 MYB class transcription factor. |
| AAG08960.1     | AF122052 | Solanum tuberosum     | tuber-specific and sucrose-responsive element binding factor. TSF. R2R3 MYB class transcription factor. |
| AAG08961.1     | AF122053 | Solanum tuberosum     | tuber-specific and sucrose-responsive element binding factor. TSF. R2R3 MYB class transcription factor. |
| AAF67053.1     | AF190304 | Adiantum raddianum    | c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.                                   |
| AAF67052.1     | AF190303 | Adiantum raddianum    | c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.                                   |
| AAF34434.1     | AF172282 | Oryza sativa          | myb-like protein. DUPR11.29.  |



|            |          |  |
|------------|----------|--|
| AAF78888.1 | AF189786 | Physcomitrella patens<br>putative c-myb-like transcription factor. MYB3R-1. PpMYB3R-1.   |
| AAF78887.1 | AF189785 | Physcomitrella patens<br>putative c-myb-like transcription factor. MYB3R-1. PpMYB3R-1.   |
| AAF43043.1 | AF236059 | Papaver rhoeas<br>putative Myb-related domain. pmr.  |
| BAA94769.1 | AP001859 | Oryza sativa<br>Similar to Arabidopsis thaliana chromosome 4, BAC clone F4D11; putative myb-protein.<br>(AL022537).                          |
| AAF78890.1 | AF189788 | Hordeum vulgare<br>putative c-myb-like transcription factor. MYB3R-1. HvMYB3R-1.   |
| AAF78889.1 | AF189787 | Hordeum vulgare<br>putative c-myb-like transcription factor. MYB3R-1. HvMYB3R-1.   |
| BAA88169.1 | AP000836 | Oryza sativa<br>Similar to putative transcription factor (AF062890).   |
| BAA88205.1 | AP000837 | Oryza sativa<br>Similar to putative transcription factor (AF062890).   |
| CAA78388.1 | Z13998   | Petunia x hybrida<br>DNA-binding protein; transcriptional activator. protein 3. myb.Ph3. related to animal myb<br>proto-oncoproteins.        |
| AAF67051.1 | AF190302 | Secale cereale<br>c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.  |
| AAF67050.1 | AF190301 | Secale cereale<br>c-myb-like transcription factor. MYB3R-1. contains three MYB repeats.  |
| AAG28525.1 | AF198498 | Nicotiana tabacum<br>anther-specific myb-related protein 2. myBAS2. NtMYBAS2; contains tandem R2, R3 myb<br>domains similar to c-myb family. |
| BAB12688.1 | AP002746 | Oryza sativa<br>putative MYB family transcription factor. P0671B11.3. contains ESTs<br>AU082307(E0784),C72014(E0784).                        |
| BAA99440.1 | AP002743 | Oryza sativa<br>putative MYB family transcription factor. P0710E05.27. contains ESTs<br>AU082307(E0784),C72014(E0784).                       |
| AAK19619.1 | AF336286 | Gossypium hirsutum<br>GHMYB9. ghmyb9. similar to myb.  |
| CAA64615.1 | X95297   | Lycopersicon esculentum<br>transcription factor. THM1. myb-related.  |
| BAA81730.1 | AB029159 | Glycine max<br>GmMYB29A1.  |
| CAA50224.1 | X70879   | Hordeum vulgare<br>MybHv1. myb1.   |
| CAA50222.1 | X70877   | Hordeum vulgare<br>MybHv1. myb1.   |

|            |          |                    |   |
|------------|----------|--------------------|---|
| CAB40189.1 | AJ133638 | Avena sativa       | transcriptional activator. myb protein. gamyb.  |
| AAG28526.1 | AF198499 | Nicotiana tabacum  | anther-specific myb-related protein 1. myBAS1. NtMYBAS1; contains N-terminal R2, R3 myb domain repeats similar to c-myb.      |
| AAD31395.1 | AF114162 | Lolium temulentum  | gibberellin MYB transcription factor. GAMyb. R2/R3-MYB.   |
| BAA81732.1 | AB029161 | Glycine max        | GmMYB29A2.  |
| BAA81731.1 | AB029160 | Glycine max        | GmMYB29A1.  |
| CAA61021.1 | X87690   | Hordeum vulgare    | transcriptional activator of alpha-amylase gene promoter. GAMyb protein. Gam1.  |
| AAG22863.1 | AY008692 | Hordeum vulgare    | transcription factor GAMyb. Gamyb.  |
| BAA96421.1 | AB044084 | Triticum aestivum  | transcription activator for gibberellin response. GAMyb protein. Ta-GAMyb.  |
| BAA81733.2 | AB029162 | Glycine max        | GmMYB29A2.  |
| BAA81736.1 | AB029165 | Glycine max        | GmMYB29B2.  |
| AAA33067.1 | L04497   | Gossypium hirsutum | MYB A; putative.  |
| BAB39972.1 | AP003018 | Oryza sativa       | putative transcription factor (myb). OSJNBa0004B13.27. contains ESTs AU097474(S5087),D40175(S1959).                           |
| CAA72218.1 | Y11415   | Oryza sativa       | myb.  |
| CAA67000.1 | X98355   | Oryza sativa       | activator of alpha-amylase gene promoter. transcription factor GAMyb. Gam1. Myb-like; expression is regulated by gibberellin. |
| BAB39987.1 | AP003020 | Oryza sativa       | putative transcription factor (myb). P0498A12.16. contains ESTs AU097474(S5087),D40175(S1959).                                |
| CAA78387.1 | Z13997   | Petunia x hybrida  | DNA-binding protein, transcriptional activator. protein 2. myb.Ph2. related to animal myb proto-oncoproteins.                 |
| CAA72217.1 | Y11414   | Oryza sativa       | myb.  |
| BAA23341.1 | D88621   | Oryza sativa       | transfactor. OSMYB5. Osmyb5.  |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAA93038.1     | AP001552 | Oryza sativa            | EST AU082058(C12976) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana putative transcription factor (AF062916).  |
| CAA64614.1     | X95296   | Lycopersicon esculentum | transcription factor. THM27. myb-related.  |
| AAK19616.1     | AF336283 | Gossypium hirsutum      | GHMYB25. ghmyb25. similar to myb; contains an unspliced intron.  |
| CAA78386.1     | Z13996   | Petunia x hybrida       | DNA binding protein; transcriptional activator. protein 1. myb.Ph3. Product related to animal myb proto-oncoproteins. Sequence from nucleotide 992 is not included in clone cPF1 and has been obtained by PCR amplification of cDNA. |
| CAA67575.1     | X99134   | Lycopersicon esculentum | transcription factor. THM6. myb-related.   |
| BAA88222.1     | AB028650 | Nicotiana tabacum       | myb-related transcription factor LBM2. lbm2.   |
| SEQ ID NO: 902 |          |                         |  |
| CAA57773.1     | X82329   | Arachis hypogaea        | chitinase (class II). chi2;1.  |
| AAD54935.1     | AF141373 | Petroselinum crispum    | random hydrolysis of 1,4-beta-acetamido-2-deoxy-D-glucoside linkages in chitin. chitinase precursor. Chi2-1. class II.   |
| AAD54936.1     | AF141374 | Petroselinum crispum    | random hydrolysis of 1,4-beta-acetamido-2-deoxy-D-glucoside linkages in chitin. chitinase precursor. Chi2-2. class II.   |
| AAF00131.1     | AF147091 | Fragaria x ananassa     | chitin degradation. class II chitinase. Chi2-1.  |
| AAC95376.1     | AF105426 | Cynodon dactylon        | chitinase. Cht2.   |
| BAA95846.1     | AP002070 | Oryza sativa            | Similar to Oryza sativa clone RGCH7 chitinase. (AF013581).   |
| AAA32986.1     | M95835   | Brassica napus          | endochitinase. Ch25.   |
| AAF69783.1     | AF135143 | Arabis lemmonii         | class I chitinase.   |
| AAF69775.1     | AF135135 | Arabis drummondii       | class I chitinase.   |
| AAF69792.1     | AF135152 | Arabis parishii         | class I chitinase.   |
| AAC95375.1     | AF105425 | Cynodon dactylon        | chitinase. Cht1.   |
| AAF69785.1     | AF135145 | Arabis lignifera        | class I chitinase.   |

|            |          |  |
|------------|----------|--|
| AAF69770.1 | AF135130 | <i>Arabis holboellii</i><br>class I chitinase.                         |
| AAF69781.1 | AF135141 | <i>Arabis gunnisoniana</i><br>class I chitinase.                       |
| AAF69777.1 | AF135137 | <i>Arabis fecunda</i><br>class I chitinase.                            |
| AAF69790.1 | AF135150 | <i>Arabis microphylla</i><br>class I chitinase.                        |
| AAF69787.1 | AF135147 | <i>Arabis lignifera</i><br>class I chitinase.                          |
| AAF69772.1 | AF135132 | <i>Arabis gunnisoniana</i><br>class I chitinase.                       |
| AAF69782.1 | AF135142 | <i>Halimolobos perplexa</i> var. <i>perplexa</i><br>class I chitinase. |
| AAF69784.1 | AF135144 | <i>Arabis lemmonii</i><br>class I chitinase.                           |
| AAF69788.1 | AF135148 | <i>Arabis lyallii</i><br>class I chitinase.                            |
| BAA03750.1 | D16222   | <i>Oryza sativa</i><br>endochitinase. Cht-2.                           |
| AAF69776.1 | AF135136 | <i>Arabis fecunda</i><br>class I chitinase.                            |
| CAA40107.1 | X56787   | <i>Oryza sativa</i><br>chitinase.                                      |
| BAB13369.1 | AB048531 | <i>Psophocarpus tetragonolobus</i><br>class I chitinase.               |
| AAF69778.1 | AF135138 | <i>Arabis glabra</i><br>class I chitinase.                             |
| AAF69786.1 | AF135146 | <i>Arabis lignifera</i><br>class I chitinase.                          |
| BAA82826.1 | AB023464 | <i>Arabis gemmifera</i><br>basic endochitinase. ChiB.                  |
| AAF69773.1 | AF135133 | <i>Arabis blepharophylla</i><br>class I chitinase.                     |
| AAF69791.1 | AF135151 | <i>Arabis microphylla</i><br>class I chitinase.                        |
| AAF69793.1 | AF135153 | <i>Arabis parishii</i><br>class I chitinase.                           |
| CAA39535.1 | X56063   | <i>Oryza sativa</i><br>chitinase.                                      |
| BAA03749.1 | D16221   | <i>Oryza sativa</i><br>endochitinase. Cht-1.                           |

|                |          |  |
|----------------|----------|--|
| AAF69789.1     | AF135149 | Arabis microphylla<br>class I chitinase.   |
| CAA71402.1     | Y10373   | Medicago truncatula<br>chitinase.  |
| AAC16010.1     | AF061805 | Elaeagnus umbellata<br>acidic chitinase.   |
| CAA53626.1     | X76041   | Triticum aestivum<br>endochitinase. CHI.   |
| BAA33971.1     | AB008892 | Nicotiana tabacum<br>chitinase 134. Chn134.  |
| CAA47921.1     | X67693   | Solanum tuberosum<br>chitinase. SK2. endochitinase.  |
| AAF69780.1     | AF135140 | Arabis glabra<br>class I chitinase.  |
| BAB18519.1     | AB051578 | Secale cereale<br>seed chitinase-a. rsca.  |
| AAA51377.1     | L37289   | Oryza sativa<br>chitinolytic activity, antifungal activity. chitinase.   |
| AAB41324.1     | U83591   | Medicago sativa<br>class I chitinase.  |
| CAC17793.1     | AJ301671 | Nicotiana sylvestris<br>hydrolysis of chitin. endochitinase. chnb. class I chitinase.  |
| AAB41325.1     | U83592   | Medicago sativa<br>class I chitinase.  |
| AAB23263.1     | S43926   | Phaseolus vulgaris<br>chitinase. CH5B. This sequence comes from Fig. 1.  |
| AAA33756.1     | M13968   | Phaseolus vulgaris<br>chitinase (EC 3.2.1.14).   |
| CAA35945.1     | X51599   | Nicotiana tabacum<br>chitinase. CHN50.   |
| AAA34070.1     | M15173   | Nicotiana tabacum<br>endochitinase precursor (EC 3.2.1.14).  |
| CAA45822.1     | X64519   | Nicotiana tabacum<br>chitinase B class I. CHN200.  |
| SEQ ID NO: 903 |          |  |
| BAA78764.1     | AB023482 | Oryza sativa<br>ESTs C98382(C2985),D22444(C11129) correspond to a region of the predicted gene.; Similar<br>to Arabidopsis thaliana APK1 gene for protein tyrosine-serine-threonine kinase.(D12522). |
| AAF43496.1     | AF131222 | Lophopyrum elongatum<br>protein serine/threonine kinase. ESI47. induced in roots by salt stress, osmotic stress, and<br>ABA treatment.   |
| AAK11674.1     | AF339747 | Lophopyrum elongatum<br>protein kinase. ESI47.   |

|            |          |                               |   |
|------------|----------|-------------------------------|---|
| AAG16628.1 | AY007545 | Brassica napus                | protein serine/threonine kinase BNK1.   |
| BAA94509.1 | AB041503 | Populus nigra                 | protein kinase 1. PnPK1.  |
| BAA87853.1 | AP000816 | Oryza sativa                  | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).                                  |
| BAA94510.1 | AB041504 | Populus nigra                 | protein kinase 2. PnPK2.  |
| BAB16871.1 | AP002537 | Oryza sativa                  | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).  |
| BAB03429.1 | AP002817 | Oryza sativa                  | EST C22619(S11214) corresponds to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 2, BAC clone T17D12; putative protein kinase (AC006587). |
| BAB07999.1 | AP002525 | Oryza sativa                  | putative protein kinase. P0462H08.22. contains EST C22619(S11214).  |
| BAB39409.1 | AP002901 | Oryza sativa                  | putative protein kinase. P0456F08.9. contains EST C23560(R0290).  |
| BAB21240.1 | AP002953 | Oryza sativa                  | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| AAK00425.1 | AC069324 | Oryza sativa                  | Putative protein kinase. OSJNBa0071K19.11.  |
| CAB51834.1 | 00069    | Oryza sativa                  | 11332.5. contains eukaryotic protein kinase domain PF.  |
| BAB39873.1 | AP002882 | Oryza sativa                  | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).   |
| AAK21965.1 | AY028699 | Brassica napus                | receptor protein kinase PERK1.  |
| AAG03090.1 | AC073405 | Oryza sativa                  | Similar to an Arabidopsis somatic embryogenesis receptor-like kinase (AC007504).  |
| BAB19337.1 | AP003044 | Oryza sativa                  | putative protein kinase. P0038C05.10. contains ESTs AU056335(S20481),AU056336(S20481).  |
| BAA90808.1 | AP001168 | Oryza sativa                  | Similar to putative receptor-like protein kinase (AL035679).  |
| AAG59657.1 | AC084319 | Oryza sativa                  | putative protein kinase. OSJNBa0004B24.20.  |
| AAB47424.1 | U59317   | Lycopersicon pimpinellifolium | serine/threonine protein kinase Fen. Fen. fenthion sensitivity gene from tomato.  |

|            |          |  |
|------------|----------|--|
| AAC48932.1 | U13923   | <i>Lycopersicon pimpinellifolium</i><br>Fen. putative serine/threonine protein kinase; similar to product encoded by <i>Lycopersicon pimpinellifolium</i> Pto gene, GenBank Accession Number U02271; Fen is a member of the Pto gene family. |
| AAF76307.1 | AF220602 | <i>Lycopersicon pimpinellifolium</i><br>Fen kinase.  |
| BAA92221.1 | AP001278 | <i>Oryza sativa</i><br>Similar to <i>Arabidopsis thaliana</i> chromosome II BAC F12L6 genomic sequence, putative protein kinase. (AC004218).   |
| BAA87852.1 | AP000816 | <i>Oryza sativa</i><br>Similar to putative Ser/Thr protein kinase. (AC004218).   |
| AAB61708.1 | U93048   | <i>Daucus carota</i><br>somatic embryogenesis receptor-like kinase. SERK.  |
| AAK11566.1 | AF318490 | <i>Lycopersicon hirsutum</i><br>Pto-like protein kinase E. LhirPtoE. confers resistance to bacterial speck disease.  |
| AAK11568.1 | AF318492 | <i>Lycopersicon hirsutum</i><br>Pto-like protein kinase B. LhirPtoB.   |
| AAD21872.1 | AF078082 | <i>Phaseolus vulgaris</i><br>receptor-like protein kinase homolog RK20-1.  |
| AAF66615.1 | AF142596 | <i>Nicotiana tabacum</i><br>LRR receptor-like protein kinase.  |
| AAC23542.1 | U20948   | <i>Ipomoea trifida</i><br>receptor protein kinase. IRK1.   |
| AAF91337.1 | AF249318 | <i>Glycine max</i><br>PtiI kinase-like protein. PtiIb. protein kinase.   |
| AAC27894.1 | AF023164 | <i>Zea mays</i><br>leucine-rich repeat transmembrane protein kinase 1. ltk1.   |
| AAF76314.1 | AF220603 | <i>Lycopersicon esculentum</i><br>Fen kinase. Lescfen.   |
| AAB47422.1 | U59318   | <i>Lycopersicon esculentum</i><br>serine/threonine protein kinase Fen. fen. allele of Fen, fenthion sensitivity gene from tomato.  |
| AAF76313.1 | AF220603 | <i>Lycopersicon esculentum</i><br>Pto kinase. LescPth5.  |
| AAB47421.1 | U59316   | <i>Lycopersicon esculentum</i><br>serine/threonine protein kinase Pto. pto. allele of Pto bacterial speck disease resistance gene in tomato.   |
| AAF91336.1 | AF249317 | <i>Glycine max</i><br>PtiI kinase-like protein. PtiIa. protein kinase.   |
| AAK11567.1 | AF318491 | <i>Lycopersicon hirsutum</i><br>Pto-like protein kinase F. LhirPtoF.   |
| AAC61805.1 | U28007   | <i>Lycopersicon esculentum</i><br>serine/threonine protein kinase. Pto kinase interactor 1. Pti1. Pti1 kinase.   |

|                |          |                                    |   |
|----------------|----------|------------------------------------|---|
| AAB93834.1     | U82481   | Zea mays                           | KI domain interacting kinase 1. KIK1. receptor-like protein kinase; serine/threonine protein kinase.                                      |
| CAA97692.1     | Z73295   | Catharanthus roseus                | receptor-like protein kinase. CRPK1. Autophosphorylation predominantly on Thr, less on Ser. Mechanism: autophosphorylation in cis.        |
| CAA73134.1     | Y12531   | Brassica oleracea                  | serine/threonine kinase. BRLK.  |
| AAF76306.1     | AF220602 | Lycopersicon pimpinellifolium      | Pto kinase.   |
| AAB47423.1     | U59315   | Lycopersicon pimpinellifolium      | serine/threonine protein kinase Pto. Pto. bacterial speck disease resistance in tomato; disease resistance gene.                          |
| BAB21241.1     | AP002953 | Oryza sativa                       | Putative Pto kinase interactor 1. P0426D06.21. contains ESTs AU108280(E0721),D48017(S13927).  |
| AAC48914.1     | U02271   | Lycopersicon pimpinellifolium      | protein kinase.   |
| SEQ ID NO: 904 |          |                                    |   |
| AAF65545.1     | AF233894 | Perilla citriodora                 | limonene synthase.  |
| AAG31437.1     | AF241792 | Perilla frutescens                 | limonene synthase.  |
| AAK06663.1     | AF317695 | Perilla frutescens var. frutescens | limonene synthase.  |
| AAG31435.1     | AF241790 | Perilla citriodora                 | limonene synthase.  |
| BAA21629.1     | AB005744 | Perilla frutescens                 | catalyzing the cyclization of geranyl pyrophosphate to 1-limonene. 1-limonene synthase. gPFLC. similar to perilla PFLC-1:DDBJ Acc#D49368. |
| AAG31438.1     | AF241793 | Perilla frutescens                 | limonene synthase.  |
| BAA08367.1     | D49368   | Perilla frutescens                 | limonene cyclase.   |
| AAC37366.1     | L13459   | Mentha spicata                     | 4S-limonene synthase.   |
| AAD50304.1     | AF175323 | Mentha longifolia                  | limonene synthase. monoterpene synthase.  |
| AAG01140.1     | AF282875 | Schizonepeta tenuifolia            | (+)-4R-limonene synthase.   |
| AAF21053.1     | AF212433 | Capsicum annuum                    | UV-induced sesquiterpene cyclase. SC2.  |



|            |          |                         |  |
|------------|----------|-------------------------|--|
| BAA82141.1 | AB023816 | Solanum tuberosum       | vetispiradiene synthase. PVS4. potato sesquiterpene cyclase.   |
| AAF74977.1 | AF270425 | Gossypium hirsutum      | (E,E)-farnesyl diphosphate cyclase. (+)-delta-cadinene synthase. cdn2. sesquiterpene cyclase.                            |
| BAA82092.1 | AB022598 | Solanum tuberosum       | vetispiradiene synthase. PVS1. potato sesquiterpene cyclase.   |
| BAA82109.1 | AB022720 | Solanum tuberosum       | vetispiradiene synthase. PVS3. potato sesquiterpene cyclase; PVS3.   |
| AAG09949.1 | AF171216 | Lycopersicon esculentum | vetispiradiene synthase. LEVS2. sesquiterpene cyclase.   |
| BAA82108.1 | AB022719 | Solanum tuberosum       | vetispiradiene synthase. PVS2. potato sesquiterpene cyclase; PVS2.   |
| AAC12784.1 | U88318   | Gossypium hirsutum      | (E,E)-farnesyl diphosphate cyclizing. (+)-delta-cadinene synthase. cdn1. sesquiterpene cyclase; delta-cadinene synthase. |
| AAK15641.1 | AF326117 | Capsicum annuum         | sesquiterpene cyclase. PSC2.   |
| AAC61260.1 | AF061285 | Capsicum annuum         | sesquiterpene cyclase. UV induced.   |
| AAG24640.2 | AF304444 | Artemisia annua         | sesquiterpene cyclase.   |
| CAC12731.1 | AJ271792 | Artemisia annua         | putative sesquiterpene cyclase. cASC125.   |

## SEQ ID NO: 905

|            |          |                         |  |
|------------|----------|-------------------------|--|
| AAF64450.1 | AF239928 | Euphorbia esula         | glutathione S-transferase. similar to auxin-inducible GST. |
| AAG34803.1 | AF243368 | Glycine max             | glutathione S-transferase GST 13.                          |
| AAG16759.1 | AY007561 | Lycopersicon esculentum | putative glutathione S-transferase T4.                     |
| AAG34798.1 | AF243363 | Glycine max             | glutathione S-transferase GST 8.                           |
| AAG34797.1 | AF243362 | Glycine max             | glutathione S-transferase GST 7.                           |
| AAG16758.1 | AY007560 | Lycopersicon esculentum | putative glutathione S-transferase T3.                     |
| AAG34809.1 | AF243374 | Glycine max             | glutathione S-transferase GST 19.                          |
| AAG34801.1 | AF243366 | Glycine max             | glutathione S-transferase GST 11.                          |
| AAG34796.1 | AF243361 | Glycine max             | glutathione S-transferase GST 6.                           |

---

|            |          |   |
|------------|----------|---|
| AAG34807.1 | AF243372 | Glycine max<br>glutathione S-transferase GST 17.  |
| AAG16757.1 | AY007559 | Lycopersicon esculentum<br>putative glutathione S-transferase T2.   |
| AAG16756.1 | AY007558 | Lycopersicon esculentum<br>putative glutathione S-transferase T1.   |
| AAG34804.1 | AF243369 | Glycine max<br>glutathione S-transferase GST 14.  |
| AAG34810.1 | AF243375 | Glycine max<br>glutathione S-transferase GST 20.  |
| AAG34808.1 | AF243373 | Glycine max<br>glutathione S-transferase GST 18.  |
| AAG34805.1 | AF243370 | Glycine max<br>glutathione S-transferase GST 15.  |
| AAG34844.1 | AF244701 | Zea mays<br>glutathione S-transferase GST 36.   |
| AAG34829.1 | AF244686 | Zea mays<br>glutathione S-transferase GST 21.   |
| AAG34831.1 | AF244688 | Zea mays<br>glutathione S-transferase GST 23.   |
| AAA68430.1 | J03679   | Solanum tuberosum<br>glutathione S-transferase. gst1. previously called pathogenesis-related protein; prp1-1.   |
| AAC18566.1 | AF048978 | Glycine max<br>2,4-D inducible glutathione S-transferase. GSTa.   |
| AAG34832.1 | AF244689 | Zea mays<br>glutathione S-transferase GST 24.   |
| AAG34800.1 | AF243365 | Glycine max<br>glutathione S-transferase GST 10.  |
| AAG34849.1 | AF244706 | Zea mays<br>glutathione S-transferase GST 41.   |
| AAG34837.1 | AF244694 | Zea mays<br>glutathione S-transferase GST 29.   |
| AAC32118.1 | AF051214 | Picea mariana<br>probable glutathione S-transferase. Sb18. similar to Nicotiana tabacum probable glutathione<br>S-transferase encoded by GenBank Accession Number X56266. |
| AAG41204.1 | AF321437 | Suaeda maritima<br>glutathione transferase.   |
| AAG34802.1 | AF243367 | Glycine max<br>glutathione S-transferase GST 12.  |
| AAF29773.1 | AF159229 | Gossypium hirsutum<br>glutathione S-transferase. GST.   |
| AAG32472.1 | AF309379 | Oryza sativa subsp. japonica<br>putative glutathione S-transferase OsGSTU3.   |

---

|                |          |                                       |   |
|----------------|----------|---------------------------------------|---|
| AAG34836.1     | AF244693 | Zea mays                              | glutathione S-transferase GST 28.   |
| CAA04391.1     | AJ000923 | Carica papaya                         | glutathione transferase. PGST1.   |
| CAA71784.1     | Y10820   | Glycine max                           | glutathione transferase.  |
| CAA09187.1     | AJ010448 | Alopecurus myosuroides                | glutathione transferase. GST1a.   |
| AAG34795.1     | AF243360 | Glycine max                           | glutathione S-transferase GST 5.  |
| CAA09188.1     | AJ010449 | Alopecurus myosuroides                | glutathione transferase. GST1b.   |
| AAG34806.1     | AF243371 | Glycine max                           | glutathione S-transferase GST 16.   |
| CAC24549.1     | AJ296343 | Cichorium intybus x Cichorium endivia | glutathione S-transferase. chi-GST1. auxin-induced GST.                                     |
| AAC28101.1     | AF079511 | Mesembryanthemum crystallinum         | glutathione S-transferase.  |
| SEQ ID NO: 907 |          |                                       |   |
| AAA33280.1     | L20475   | Datura stramonium                     | 29kDa protein; high homology to aa sequence of tropinone reductases.                        |
| AAA33281.1     | L20473   | Datura stramonium                     | catalyses a stereospecific reduction of tropinone to tropine. tropinone reductase-I.        |
| CAC34420.1     | AJ307584 | Solanum tuberosum                     | tropine forming tropinone reductase. tropinone reductase I. TRI.                            |
| BAA85844.1     | AB026544 | Hyoscyamus niger                      | tropinone reductase-I. tr1.   |
| BAA13547.1     | D88156   | Hyoscyamus niger                      | tropinone reductase-I.  |
| AAA33282.1     | L20474   | Datura stramonium                     | catalyses a stereospecific reduction of tropinone to pseudotropine. tropinone reductase-II. |
| AAB09776.1     | L20485   | Hyoscyamus niger                      | tropinone reductase-II.   |
| BAA85845.1     | AB026545 | Hyoscyamus niger                      | tropinone reductase-II. tr2.  |
| CAB52307.1     | AJ245634 | Solanum tuberosum                     | pseudotropine forming tropinone reductase. tropinone reductase II. TRII.                    |
| CAC19810.1     | AJ292343 | Solanum tuberosum                     | pseudotropine forming tropinone reductase. tropinone reductase II. trII.                    |
| CAA45866.1     | X64566   | Cuphea lanceolata                     | beta-ketoacyl-ACP reductase. Clkr27. 3-oxoacyl-[acyl-carrier protein] reductase.            |

|                |          |                        |  |
|----------------|----------|------------------------|--|
| CAA45793.1     | X64463   | Brassica napus         | 3-oxoacyl-[acyl-carrier protein] reductase.  |
| AAB82767.1     | U89509   | Zea mays               | cuticular wax biosynthesis. b-keto acyl reductase. glossy8.  |
| AAB82766.1     | U89510   | Hordeum vulgare        | cuticular wax biosynthesis. b-keto acyl reductase. glossy8.  |
| CAA74176.1     | Y13861   | Nicotiana tabacum      | enoyl-ACP reductase. enr-T2.   |
| AAB20114.2     | S60064   | Brassica napus         | enoyl-acyl carrier protein reductase. This sequence comes from fig3; conceptual translation differs from published sequence. |
| CAA74177.1     | Y13862   | Nicotiana tabacum      | enoyl-ACP reductase. enr-T1.   |
| AAC78100.1     | AF093628 | Oryza sativa           | protochlorophyllide reductase homolog.   |
| CAA05879.1     | AJ003124 | Petunia x hybrida      | enoyl-ACP reductase. pte.  |
| CAA64729.1     | X95462   | Brassica napus         | enoyl reductase.   |
| CAA05816.1     | AJ003025 | Oryza sativa           | enoyl-ACP reductase.   |
| BAA99570.1     | AB036823 | Chlorella vulgaris     | oxidoreductase. oxi.   |
| SEQ ID NO: 909 |          |                        |  |
| BAA82107.1     | AB022693 | Nicotiana tabacum      | transcription factor. NtWRKY1.   |
| AAC31956.1     | AF080595 | Pimpinella brachycarpa | zinc finger protein. ZFP1. WRKY1.  |
| AAD55974.1     | AF121353 | Petroselinum crispum   | zinc-finger type transcription factor WRKY1. WRKY1.  |
| BAA77383.1     | AB020590 | Nicotiana tabacum      | transcription factor NtWRKY2.  |
| AAC49527.1     | U48831   | Petroselinum crispum   | WRKY1. contains two WRKY domains; WRKY-type DNA-binding protein; sequence-specific DNA-binding protein.                      |
| AAD32677.1     | AF140554 | Avena sativa           | DNA-binding protein WRKY1. wrky1. putative transcription factor.   |
| CAA88326.1     | Z48429   | Avena fatua            | binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein.   |
| AAD16139.1     | AF096299 | Nicotiana tabacum      | DNA-binding protein 2. WRKY2. transcription factor.  |
| BAA86031.1     | AB026890 | Nicotiana tabacum      | transcription factor NtWRKY4.  |

|            |          |                      |  |
|------------|----------|----------------------|--|
| AAC37515.1 | L44134   | Cucumis sativus      | SPF1-like DNA-binding protein.   |
| AAK16171.1 | AC079887 | Oryza sativa         | putative DNA-binding protein. OSJNBa0040E01.10.  |
| AAF23898.1 | AF193802 | Oryza sativa         | zinc finger transcription factor WRKY1.  |
| AAD16138.1 | AF096298 | Nicotiana tabacum    | DNA-binding protein 1. WRKY1. transcription factor.  |
| AAC49529.1 | U58540   | Petroselinum crispum | WRKY2. Contains two WRKY domains; WRKY-type DNA-binding protein.   |
| CAB97004.1 | AJ278507 | Solanum tuberosum    | putative transcription factor. WRKY DNA binding protein. WRKY1.  |
| BAB19075.1 | AP002744 | Oryza sativa         | putative DNA-binding protein homolog. P0006C01.17.   |
| BAB19096.1 | AP002839 | Oryza sativa         | putative DNA-binding protein homolog. P0688A04.2.  |
| AAK16170.1 | AC079887 | Oryza sativa         | putative DNA binding protein. OSJNBa0040E01.4.   |
| BAB40073.1 | AP003074 | Oryza sativa         | putative WRKY DNA binding protein. OSJNBa0004G10.20. contains EST C26525(C12525).                                    |
| AAD38283.1 | AC007789 | Oryza sativa         | putative WRKY DNA binding protein. OSJNBa0049B20.9.  |
| BAB18313.1 | AP002865 | Oryza sativa         | putative WRKY DNA binding protein. P0034C11.1. contains EST C26525(C12525).  |
| CAA88331.1 | Z48431   | Avena fatua          | binds conserved cis-element from cereal alpha-Amy2 promoters. DNA-binding protein.                                   |
| AAC49528.1 | U56834   | Petroselinum crispum | DNA-binding. WRKY3. WRKY-type DNA-binding protein.   |
| AAG35658.1 | AF204925 | Petroselinum crispum | transcription factor WRKY4. WRKY4. binds to W box (TTGACC) elements.   |
| BAB16432.1 | AB041520 | Nicotiana tabacum    | WRKY transcription factor Nt-SubD48. Nt-SubD48.  |
| AAD32676.1 | AF140553 | Avena sativa         | DNA-binding protein WRKY3. wrky3. putative transcription factor.   |
| AAG46150.1 | AC018727 | Oryza sativa         | putative DNA-binding protein. OSJNBa0056G17.18.  |
| BAA77358.1 | AB020023 | Nicotiana tabacum    | WRKY domain Zn-finger type DNA-binding protein. DNA-binding protein NtWRKY3.   |
| AAG35659.1 | AF204926 | Petroselinum crispum | transcription factor WRKY5. WRKY5. binds to W box (TTGACC) elements.   |
| AAD27591.1 | AF121354 | Petroselinum crispum | binds sequence specifically to W Boxes (TTGACC). transcription factor. WRKY3. sequence specific DNA-binding protein. |

|                |          |   |
|----------------|----------|---|
| CAB66338.1     | AJ279697 | Betula pendula<br>wrky-type DNA binding protein. wrky.  |
| AAF61864.1     | AF193771 | Nicotiana tabacum<br>DNA-binding protein 4. WRKY4. transcription factor.  |
| BAA87069.1     | AB035271 | Matricaria chamomilla<br>elicitor-induced DNA-binding protein homolog. McWRKY1.   |
| AAF61863.1     | AF193770 | Nicotiana tabacum<br>DNA-binding protein 3. WRKY3. transcription factor.  |
| SEQ ID NO: 910 |          |   |
| AAC09420.1     | M68929   | Mitochondrion Marchantia polymorpha<br>rps14.   |
| CAA33994.1     | X15901   | Plastid Oryza sativa<br>ribosomal protein S14. rps14.   |
| SEQ ID NO: 911 |          |   |
| CAA55128.1     | X78325   | Nicotiana tabacum<br>chitinase/lysozyme. Pz.  |
| CAA54373.1     | X77110   | Nicotiana tabacum<br>chitinase, class V. chi-V.   |
| CAA54374.1     | X77111   | Nicotiana tabacum<br>chitinase, class V. chi-V.   |
| SEQ ID NO: 912 |          |   |
| CAB55394.1     | AL117264 | Oryza sativa<br>zwh0010.1. similar to Arabidopsis germin-like protein 6 (AF032976); Method: conceptual translation with partial peptide sequencing. |
| BAA78563.1     | AB024338 | Atriplex lentiformis<br>germin-like protein.  |
| AAB97470.1     | AF042489 | Oryza sativa<br>germin-like protein 16. glp16.  |
| AAA33030.1     | M93041   | Mesembryanthemum crystallinum<br>germin-like protein. germin-like protein.  |
| CAB65371.1     | AJ250834 | Pisum sativum<br>germin-like protein. ger2b. 2nd variant of the clone PsGER2.   |
| CAB55559.1     | AJ237943 | Triticum aestivum<br>germin-like protein. glp2b.  |
| CAB65370.1     | AJ250833 | Pisum sativum<br>germin-like protein. ger2a. 1st variant of this clone.   |
| CAB55558.1     | AJ237942 | Triticum aestivum<br>germin-like protein. glp2a.  |
| AAD43971.1     | AF141878 | Oryza sativa<br>germin-like protein 1 precursor. RGLP1. similar to barley epidermis/papilla-specific oxalate oxidase-like protein.                  |

|            |          |                   |  |
|------------|----------|-------------------|--|
| AAD43973.1 | AF141880 | Oryza sativa      | germin-like protein 1 precursor. RGLP1. similar to barley epidermis/papilla-specific oxalate oxidase-like protein. |
| AAC04837.1 | AF032976 | Oryza sativa      | germin-like protein 6. GER6. similar to wheat and barley oxalate oxidase.  |
| AAG00425.1 | AF250933 | Hordeum vulgare   | germin A. GerA. apoplastic protein; contains prepeptide for targeting into the cell wall.                          |
| AAD43972.1 | AF141879 | Oryza sativa      | germin-like protein 2 precursor. RGLP2. similar to barley epidermis/papilla-specific oxalate oxidase-like protein. |
| CAA63659.1 | X93171   | Hordeum vulgare   | oxalate oxidase-like protein or germin-like protein.   |
| AAC04833.1 | AF032972 | Oryza sativa      | germin-like protein 2. GER2. similar to wheat and barley oxalate oxidase.  |
| AAC04834.1 | AF032973 | Oryza sativa      | germin-like protein 3. GER3. similar to wheat and barley oxalate oxidase.  |
| AAC04832.1 | AF032971 | Oryza sativa      | germin-like protein 1. GER1. similar to wheat and barley oxalate oxidase.  |
| AAG00426.1 | AF250934 | Hordeum vulgare   | germin B. GerB. apoplastic protein; contains prepeptide for targeting into the cell wall.                          |
| AAG00427.1 | AF250935 | Hordeum vulgare   | germin F. GerF. apoplastic protein; contains prepeptide for targeting into the cell wall.                          |
| AAF34811.1 | AF005084 | Triticum aestivum | oxalate oxidase. up-regulated by aluminum.   |
| AAC04835.1 | AF032974 | Oryza sativa      | germin-like protein 4. GER4. similar to wheat and barley oxalate oxidase.  |
| BAB39965.1 | AP003018 | Oryza sativa      | probable germin protein 4. OSJNBa0004B13.19. contains ESTs AU101991(S4037),AU070167(R0031).                        |
| BAB39980.1 | AP003020 | Oryza sativa      | probable germin protein 4. P0498A12.8. contains ESTs AU101991(S4037),AU070167(R0031).                              |
| AAG00428.1 | AF250936 | Hordeum vulgare   | germin D. GerD. apoplastic protein; contains prepeptide for targeting into the cell wall.                          |
| AAA20245.1 | U01963   | Hordeum vulgare   | germin subunit.  |
| CAC19429.1 | AJ291825 | Lolium perenne    | oxalate oxidase. oxO1.   |
| AAA34270.1 | M63223   | Triticum aestivum | germin. germin 9f-2.8.   |
| AAA34268.1 | M21962   | Triticum aestivum | germin protein precursor.  |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| CAA74595.1     | Y14203   | Hordeum vulgare           | oxalate oxidase.  |
| AAG00429.1     | AF250937 | Hordeum vulgare           | germin E. GerE. apoplastic protein.                                       |
| CAB65369.1     | AJ250832 | Pisum sativum             | germin-like protein. ger1.  |
| AAA32959.1     | L15737   | Hordeum vulgare           | oxalate oxidase. This is the sequence of the complete mature peptide.     |
| CAA71052.1     | Y09917   | Triticum aestivum         | germin homolog. pSBGer3.  |
| AAA34271.1     | M63224   | Triticum aestivum         | germin. germin 9f-3.8.  |
| CAA71050.1     | Y09915   | Triticum aestivum         | germin homolog. pSBGer1.  |
| BAB18339.1     | AP002865 | Oryza sativa              | putative germin protein. P0034C11.30. contains EST C97263(C53484).        |
| AAD38298.1     | AC007789 | Oryza sativa              | putative oxalate oxidase (germin protein). OSJNBa0049B20.25.              |
| BAA86880.1     | AB028454 | Barbula unguiculata       | germin-like protein.  |
| AAF03355.1     | AF132671 | Nicotiana plumbaginifolia | nectarin I precursor. NEC1. germin-like protein.                          |
| BAA25197.1     | AB012138 | Lycopersicon esculentum   | adaptation to Mn-deficiency. germin-like protein. Mdip1.                  |
| AAC78470.1     | AF067731 | Solanum tuberosum         | germin-like protein. OXAOXA. similar to oxalate oxidase.                  |
| AAC25777.1     | AF072694 | Oryza sativa              | germin-like protein 7. GER7. similar to wheat and barley oxalate oxidase. |
| AAC99473.1     | AF039201 | Pinus caribaea            | germin-like protein. PcGER1.  |
| CAA71051.1     | Y09916   | Triticum aestivum         | germin homolog. pSBGer2.  |
| AAC05146.1     | AF049065 | Pinus radiata             | germin-like protein. PRGer1.  |
| CAC34417.1     | AJ311624 | Pisum sativum             | Germin-like protein. glp3.  |
| SEQ ID NO: 915 |          |                           |   |
| BAB19363.1     | AP002542 | Oryza sativa              | putative beta-1,3-glucanase. P0679C08.2.                                  |
| AAD10386.1     | U72255   | Oryza sativa              | beta-1,3-glucanase precursor. Gns9.                                       |



|            |          |                                  |   |
|------------|----------|----------------------------------|---|
| CAB85903.1 | AJ251646 | <i>Pisum sativum</i>             | hydrolysis of beta-1,3 conjugated glucans. beta-1,3 glucanase. gns2.  |
| AAA90953.1 | U30323   | <i>Triticum aestivum</i>         | beta 1,3-glucanase. Glc1.   |
| BAA89481.1 | AB029462 | <i>Salix gilgiana</i>            | beta-1,3-glucanase. SgGN1.  |
| BAB40807.1 | AB052291 | <i>Pyrus pyrifolia</i>           | catalyzing the hydrolysis of 1,3-beta-glucosyl linkages. endo-1,3-beta-glucanase-like protein. bgn-1. Amino acid alignment of the protein(BGN-1) encoded by pear pollen bgn-1 with barley endo-1,3-beta-glucanase(GII. accession number: pdb/1GHS-B/2.3/2/306/N/)(40% identity) and their hydrophobic cluster analysis(HCA)(a overall HCA homology score of 87.1%) showed that it was most likely that the bgn-1 encoded a endo-1,3-beta-glucanase. A higher identity(59.3%) was found between BGN-1 and a putative pea endo-1,3-beta-glucanase (accession number: dad/AJ251646-1). |
| AAB82772.2 | AF001523 | <i>Musa acuminata</i>            | beta-1, 3-glucanase. similar to beta-1, 3-glucanase.  |
| CAA49513.1 | X69887   | <i>Brassica napus</i>            | beta-1,3-glucanase homologue.   |
| AAF08679.1 | AF004838 | <i>Musa acuminata</i>            | beta-1,3-glucanase.   |
| CAA82271.1 | Z28697   | <i>Nicotiana tabacum</i>         | beta-1,3-glucanase.   |
| AAD10384.1 | U72253   | <i>Oryza sativa</i>              | beta-1,3-glucanase precursor. Gns7.   |
| CAB71021.1 | AJ271598 | <i>Hieracium piloselloides</i>   | putative role in callose degradation. putative beta-1,3-glucanase. gluc.  |
| CAA30261.1 | X07280   | <i>Nicotiana plumbaginifolia</i> | beta-glucanase.   |
| AAA51643.1 | M23120   | <i>Nicotiana plumbaginifolia</i> | beta-glucanase precursor.   |
| AAA87456.1 | U22147   | <i>Hevea brasiliensis</i>        | beta-1,3-glucanase. HGN1. hydrolytic enzyme.  |
| AAA34078.1 | M63634   | <i>Nicotiana plumbaginifolia</i> | regulator of beta(1,3)-glucanase. beta(1,3)-glucanase regulator.  |
| AAD10385.1 | U72254   | <i>Oryza sativa</i>              | beta-1,3-glucanase precursor. Gns8.   |
| CAB38443.1 | AJ133470 | <i>Hevea brasiliensis</i>        | beta-1,3-glucanase. hgn1.   |
| AAB03501.1 | U41323   | <i>Glycine max</i>               | beta-1,3-glucanase. SGN1.   |
| AAA32939.1 | M62907   | <i>Hordeum vulgare</i>           | hydrolysis of beta-(1-3)-glucan. (1-3)-beta-glucanase. cBGL32.  |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAD33881.1     | AF141654 | Nicotiana tabacum       | beta-1,3-glucanase. GGL4.                        |
| AAA03617.1     | M80604   | Lycopersicon esculentum | beta-1,3-glucanase.                              |
| BAA77786.1     | AB027431 | Oryza sativa            | beta-1,3-glucanase.                              |
| BAA77787.1     | AB027432 | Oryza sativa            | beta-1,3-glucanase.                              |
| CAB91554.1     | AJ277900 | Vitis vinifera          | beta 1-3 glucanase. gl.                          |
| AAC14399.1     | AF030771 | Hordeum vulgare         | beta-1,3-glucanase 2. BGL32.                     |
| CAA03908.1     | AJ000081 | Citrus sinensis         | glucan hydrolase. beta-1,3-glucanase. gns1.      |
| AAG24921.1     | AF311749 | Hevea brasiliensis      | beta-1,3-glucanase.                              |
| AAA33946.1     | M37753   | Glycine max             | beta-1,3-endoglucanase (EC 3.2.1.39).            |
| AAA63542.1     | M59443   | Nicotiana tabacum       | acidic beta-1,3-glucanase. glucanase.            |
| AAD28732.1     | AF112965 | Triticum aestivum       | beta-1,3-glucanase precursor. Glb3.              |
| AAD10381.1     | U72250   | Oryza sativa            | beta-1,3-glucanase precursor. Gns4.              |
| CAA57255.1     | X81560   | Nicotiana tabacum       | (1-)-beta-glucanase. Sp41a.                      |
| AAD33880.1     | AF141653 | Nicotiana tabacum       | beta-1,3-glucanase. GGL1.                        |
| AAB86541.1     | AF030166 | Oryza sativa            | glucanase. glu1.                                 |
| AAA63539.1     | M60402   | Nicotiana tabacum       | glucan beta-1,3-glucanase. glucanase GLA.        |
| AAA34053.1     | M60464   | Nicotiana tabacum       | beta-1,3-glucanase.                              |
| AAD10380.1     | U72249   | Oryza sativa            | beta-1,3-glucanase precursor. Gns3.              |
| AAA63540.1     | M60403   | Nicotiana tabacum       | glucan-1,3-beta-glucosidase. glucanase GLB.      |
| SEQ ID NO: 916 |          |                         |  |
| AAA85440.1     | U32624   | Sorghum bicolor         | cytochrome P-450. CYP79. P450TYR; N-hydroxylase. |

|            |          |                      |  |
|------------|----------|----------------------|--|
| AAF27289.1 | AF140613 | Manihot esculenta    | N-hydroxylating cytochrome P450. CYP79D1.  |
| AAF27290.1 | AF140614 | Manihot esculenta    | N-hydroxylating cytochrome P450. CYP79D2.  |
| AAD03415.1 | AF069494 | Sinapis alba         | converts tyrosine to para-hydrophenylacetaldoxime in para-hydroxybenzylglucosinolate biosynthesis. cytochrome P450. CYP79B1. |
| AAG59648.1 | AC084319 | Oryza sativa         | putative cytochrome p450tyr. OSJNBa0004B24.15.   |
| AAF66543.1 | AF140609 | Triglochin maritimum | cytochrome P450 CYP79E1. CYP79E1.  |
| AAF66544.1 | AF140610 | Triglochin maritimum | cytochrome P450 CYP79E2. CYP79E2.  |
| BAA92894.1 | AB006790 | Petunia x hybrida    | cytochrome P450. IMT-2.  |
| AAB94587.1 | AF022458 | Glycine max          | CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.  |
| AAC32274.1 | AF081575 | Petunia x hybrida    | flavonoid 3',5'-hydroxylase. Hfl. P450 enzyme.   |
| AAD56282.1 | AF155332 | Petunia x hybrida    | flavonoid 3'-hydroxylase. Htl. cytochrome P450; CYP75B2.   |
| CAA64635.1 | X95342   | Nicotiana tabacum    | cytochrome P450. hsr515. hypersensitivity-related gene.  |
| CAA50155.1 | X70824   | Solanum melongena    | flavonoid hydroxylase (P450). CYP75.   |
| AAG44132.1 | AF218296 | Pisum sativum        | cytochrome P450. P450 isolog.  |
| AAA32913.1 | M32885   | Persea americana     | cytochrome P-450LXXIA1 (cyp71A1).  |
| BAA84071.1 | AB028151 | Antirrhinum majus    | flavone synthase II. cytochrome P450. AFNS2.   |
| AAD38930.1 | AF135485 | Glycine max          | cytochrome P450 monooxygenase CYP93D1. CYP93E1.  |
| AAB17562.1 | U72654   | Eustoma grandiflorum | flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.  |
| AAG34695.1 | AF313492 | Matthiola incana     | putative cytochrome P450.  |
| CAA65580.1 | X96784   | Nicotiana tabacum    | cytochrome P450. hsr515.   |
| BAA93634.1 | AB025016 | Lotus japonicus      | cytochrome P450.   |
| CAA71513.1 | Y10489   | Glycine max          | putative cytochrome P450.  |

|                |          |  |
|----------------|----------|--|
| AAB94590.1     | AF022461 | Glycine max<br>CYP82C1p. CYP82C1. cytochrome P450 monooxygenase.   |
| CAA71876.1     | Y10982   | Glycine max<br>putative cytochrome P450.   |
| BAA12159.1     | D83968   | Glycine max<br>Cytochrome P-450 (CYP93A1).   |
| AAG14962.1     | AF214008 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H2.   |
| BAA74465.1     | AB022732 | Glycyrrhiza echinata<br>cytochrome P450. CYP Ge-31.  |
| BAA22422.1     | AB001379 | Glycyrrhiza echinata<br>cytochrome P450. CYP81E1.  |
| AAG14961.1     | AF214007 | Brassica napus<br>cytochrome p450-dependent monooxygenase. BNF5H1.   |
| CAA04117.1     | AJ000478 | Helianthus tuberosus<br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-<br>race).                            |
| CAA04116.1     | AJ000477 | Helianthus tuberosus<br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.  |
| AAB94593.1     | AF022464 | Glycine max<br>CYP77A3p. CYP77A3. cytochrome P450 monooxygenase.   |
| BAA13076.1     | D86351   | Glycine max<br>cytochrome P-450 (CYP93A2).   |
| CAA50647.1     | X71656   | Solanum melongena<br>P450 hydroxylase.   |
| SEQ ID NO: 918 |          |  |
| CAA06223.1     | AJ004923 | Lycopersicon esculentum<br>Squalene epoxidase. ERG.  |
| SEQ ID NO: 919 |          |  |
| AAF71823.1     | AF153276 | Populus tremula x Populus tremuloides<br>pumilio domain-containing protein PPD1. PPD1. similar to Drosophila pumilio protein RNA-<br>binding domain. |
| SEQ ID NO: 920 |          |  |
| BAA82393.1     | AP000367 | Oryza sativa<br>EST C96716(C10608) corresponds to a region of the predicted gene.; Similar to putative<br>receptor protein kinase. (AC002334).       |
| CAC20842.1     | AJ250467 | Pinus sylvestris<br>receptor protein kinase. upk.  |
| AAB36558.1     | U77888   | Ipomoea nil<br>receptor-like protein kinase INRPK1. inrpkl. leucine-rich repeat.   |
| AAF91324.1     | AF244890 | Glycine max<br>receptor-like protein kinase 3. RLK3. GmRLK3.   |

|            |          |                      |  |
|------------|----------|----------------------|--|
| BAB40094.1 | AP003210 | Oryza sativa         | putative receptor protein kinase. OSJNBa0010K01.7.   |
| AAF91322.1 | AF244888 | Glycine max          | receptor-like protein kinase 1. RLK1. GmRLK1.  |
| AAC36318.1 | AF053127 | Malus x domestica    | leucine-rich receptor-like protein kinase. LRPKml.   |
| BAA84787.1 | AP000559 | Oryza sativa         | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| BAA83373.1 | AP000391 | Oryza sativa         | ESTs C22657(S0014),C22656(S0014) correspond to a region of the predicted gene.; Similar to receptor protein kinase, ERECTA (AC004484). |
| AAF59905.1 | AF197946 | Glycine max          | receptor protein kinase-like protein. CLV1A.   |
| AAK27806.1 | AC022457 | Oryza sativa         | putative protein kinase. OSJNBa0006L06.21.   |
| AAF91323.1 | AF244889 | Glycine max          | receptor-like protein kinase 2. RLK2. GmRLK2.  |
| AAF34426.1 | AF172282 | Oryza sativa         | leucine rich repeat containing protein kinase. DUPR11.16.  |
| AAK27817.1 | AC022457 | Oryza sativa         | putative protein kinase. OSJNBa0006L06.16.   |
| CAA61510.1 | X89226   | Oryza sativa         | leucine-rich repeat/receptor protein kinase. lrk2.   |
| BAB03621.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.21.   |
| AAB82755.1 | U72725   | Oryza longistaminata | receptor kinase-like protein. Xa21 gene family member A1; downstream of microsatellite region; disease resistance gene family member.  |
| BAB03629.1 | AP002522 | Oryza sativa         | putative protein kinase Xa21. P0009G03.30.   |
| AAB61708.1 | U93048   | Daucus carota        | somatic embryogenesis receptor-like kinase. SERK.  |
| AAF66615.1 | AF142596 | Nicotiana tabacum    | LRR receptor-like protein kinase.  |
| AAB82756.1 | U72724   | Oryza sativa         | receptor kinase-like protein. Xa21 gene family member E.   |
| CAB51480.1 | Y14600   | Sorghum bicolor      | putative protein serine /threonine kinase. RLK1. accumulates in mesophyll cells.   |
| BAB39873.1 | AP002882 | Oryza sativa         | putative protein kinase. P0439B06.8. contains ESTs AU056701(S20808),AU056702(S20808).  |

|                |          |                            |   |
|----------------|----------|----------------------------|---|
| BAB18321.1     | AP002865 | Oryza sativa               | putative receptor protein kinase. P0034C11.11.  |
| BAB40081.1     | AP003074 | Oryza sativa               | putative receptor protein kinase. OSJNBa0004G10.30.   |
| AAD38286.1     | AC007789 | Oryza sativa               | putative protein kinase. OSJNBa0049B20.13.  |
| BAB21240.1     | AP002953 | Oryza sativa               | Putative protein kinase. P0426D06.20. contains ESTs C22359(C11461),C22360(C11461).  |
| AAK21965.1     | AY028699 | Brassica napus             | receptor protein kinase PERK1.  |
| BAA87853.1     | AP000816 | Oryza sativa               | EST AU030604(E51294) corresponds to a region of the predicted gene. Similar to putative NAK-like Ser/Thr protein kinase. (AF001308).  |
| BAB16871.1     | AP002537 | Oryza sativa               | putative protein kinase APK1AArabidopsis thaliana. P0001B06.24. contains ESTs C22608(R3192),D25110(R3192).  |
| AAC27894.1     | AF023164 | Zea mays                   | leucine-rich repeat transmembrane protein kinase 1. ltk1.   |
| AAK00425.1     | AC069324 | Oryza sativa               | Putative protein kinase. OSJNBa0071K19.11.  |
| AAG59657.1     | AC084319 | Oryza sativa               | putative protein kinase. OSJNBa0004B24.20.  |
| BAA94519.1     | AP001800 | Oryza sativa               | ESTs AU032341(R3918),AU071016(R10613) correspond to a region of the predicted gene. Similar to Arabidopsis thaliana chromosome 4, BAC F9D16; putative receptor kinase (AL035394). |
| BAB07903.1     | AP002835 | Oryza sativa               | putative receptor kinase. P0417G05.10. contains ESTs AU032341(R3918),AU071016(R10613).  |
| SEQ ID NO: 921 |          |                            |   |
| BAA96250.1     | AB027617 | Pyrus pyrifolia            | UDP-glucose pyrophosphorylase.  |
| BAA25917.1     | AB013353 | Pyrus pyrifolia            | UDP-glucose pyrophosphorylase.  |
| CAA62689.1     | X91347   | Hordeum vulgare            | UDP-glucose pyrophosphorylase.  |
| AAF62555.1     | AF249880 | Oryza sativa subsp. indica | UDP-glucose pyrophosphorylase. UDPGase.   |
| SEQ ID NO: 922 |          |                            |   |
| AAD03415.1     | AF069494 | Sinapis alba               | converts tyrosine to para-hydroxyphenylacetaldoxime in para-hydroxybenzylglucosinolate biosynthesis. cytochrome P450. CYP79B1.  |

|            |          |  |
|------------|----------|--|
| AAA85440.1 | U32624   | <i>Sorghum bicolor</i><br>cytochrome P-450. CYP79. P450TYR; N-hydroxylase.   |
| AAF27289.1 | AF140613 | <i>Manihot esculenta</i><br>N-hydroxylating cytochrome P450. CYP79D1.  |
| AAF27290.1 | AF140614 | <i>Manihot esculenta</i><br>N-hydroxylating cytochrome P450. CYP79D2.  |
| AAG59648.1 | AC084319 | <i>Oryza sativa</i><br>putative cytochrome p450tyr. OSJNBa0004B24.15.  |
| AAF66543.1 | AF140609 | <i>Triglochin maritimum</i><br>cytochrome P450 CYP79E1. CYP79E1.   |
| AAF66544.1 | AF140610 | <i>Triglochin maritimum</i><br>cytochrome P450 CYP79E2. CYP79E2.   |
| BAA92894.1 | AB006790 | <i>Petunia x hybrida</i><br>cytochrome P450. IMT-2.  |
| AAD56282.1 | AF155332 | <i>Petunia x hybrida</i><br>flavonoid 3'-hydroxylase. Ht1. cytochrome P450; CYP75B2.   |
| CAA50155.1 | X70824   | <i>Solanum melongena</i><br>flavonoid hydroxylase (P450). CYP75.   |
| AAC32274.1 | AF081575 | <i>Petunia x hybrida</i><br>flavonoid 3',5'-hydroxylase. Hf1. P450 enzyme.   |
| AAG34695.1 | AF313492 | <i>Matthiola incana</i><br>putative cytochrome P450.   |
| AAA32913.1 | M32885   | <i>Persea americana</i><br>cytochrome P-450LXXIA1 (cyp71A1).   |
| CAA64635.1 | X95342   | <i>Nicotiana tabacum</i><br>cytochrome P450. hsr515. hypersensitivity-related gene.  |
| CAA65580.1 | X96784   | <i>Nicotiana tabacum</i><br>cytochrome P450. hsr515.   |
| AAB94587.1 | AF022458 | <i>Glycine max</i><br>CYP98A2p. CYP98A2. cytochrome P450 monooxygenase.  |
| AAB17562.1 | U72654   | <i>Eustoma grandiflorum</i><br>flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.   |
| AAG09208.1 | AF175278 | <i>Pisum sativum</i><br>wound-inducible P450 hydroxylase. CYP82A1.   |
| AAD37433.1 | AF150881 | <i>Lycopersicon esculentum x Lycopersicon peruvianum</i><br>catalyzes the hydroxylation of ferulic acid to 5-hydroxyferulic acid. ferulate-5-hydroxylase. CYP84. cytochrome P450-dependent monooxygenase; F5H; FAH1. |
| CAA04117.1 | AJ000478 | <i>Helianthus tuberosus</i><br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1l. chimeric sequence (from 5'-race).   |
| CAA04116.1 | AJ000477 | <i>Helianthus tuberosus</i><br>fatty acid in-chain hydroxylase. cytochrome P450. CYP81B1c.   |

|            |          |                   |   |
|------------|----------|-------------------|---|
| CAB43505.1 | AJ239051 | Cicer arietinum   | cytochrome P450. cyp81E2.   |
| AAC49188.2 | U29333   | Pisum sativum     | cytochrome P450 monooxygenase. CYP82. new cytochrome P450 family. |
| AAG14963.1 | AF214009 | Brassica napus    | cytochrome p450-dependent monooxygenase. BNF5H3.                  |
| AAD38930.1 | AF135485 | Glycine max       | cytochrome P450 monooxygenase CYP93D1. CYP93E1.                   |
| AAG14962.1 | AF214008 | Brassica napus    | cytochrome p450-dependent monooxygenase. BNF5H2.                  |
| AAG44132.1 | AF218296 | Pisum sativum     | cytochrome P450. P450 isolog.                                     |
| AAB94588.1 | AF022459 | Glycine max       | CYP71D10p. CYP71D10. cytochrome P450 monooxygenase.               |
| AAG14961.1 | AF214007 | Brassica napus    | cytochrome p450-dependent monooxygenase. BNF5H1.                  |
| AAB94590.1 | AF022461 | Glycine max       | CYP82C1p. CYP82C1. cytochrome P450 monooxygenase.                 |
| CAA71513.1 | Y10489   | Glycine max       | putative cytochrome P450.   |
| BAA84071.1 | AB028151 | Antirrhinum majus | flavone synthase II. cytochrome P450. AFNS2.                      |
| CAA50442.1 | X71130   | Petunia x hybrida | P450 hydroxylase. PET 1.  |

## SEQ ID NO: 924

|            |          |                       |  |
|------------|----------|-----------------------|--|
| BAB20581.1 | AB042268 | Zea mays              | response regulator 6. ZmRR6.                             |
| AAK14395.1 | AF339732 | Dianthus caryophyllus | response regulator protein. RR.                          |
| BAB20580.1 | AB042267 | Zea mays              | response regulator 5. ZmRR5.                             |
| BAB20579.1 | AB042261 | Zea mays              | response regulator 4. ZmRR4.                             |
| BAA85113.1 | AB031012 | Zea mays              | response regulator 2. ZmRR2. cytokinin-inducible.        |
| BAA82873.1 | AB024291 | Zea mays              | response regulator. ZmRR2.                               |
| BAA75253.1 | AB004882 | Zea mays              | response regulator. ZmCip1, cytokinin-inducible protein. |
| BAA85112.1 | AB031011 | Zea mays              | response regulator 1. ZmRR1. cytokinin-inducible.        |



|                |          |                   |  |
|----------------|----------|-------------------|--|
| BAB17300.1     | AB042260 | Zea mays          | response regulator. ZmRR3.   |
| BAB20582.1     | AB042269 | Zea mays          | response regulator 7. ZmRR7.   |
| BAB41137.1     | AB060130 | Zea mays          | response regulator 8. ZmRR8.   |
| SEQ ID NO: 925 |          |                   |  |
| AAA33280.1     | L20475   | Datura stramonium | 29kDa protein; high homology to aa sequence of tropinone reductases.   |
| AAA33281.1     | L20473   | Datura stramonium | catalyses a stereospecific reduction of tropinone to tropine. tropinone reductase-I.   |
| BAA85844.1     | AB026544 | Hyoscyamus niger  | tropinone reductase-I. tr1.  |
| CAC34420.1     | AJ307584 | Solanum tuberosum | tropine forming tropinone reductase. tropinone reductase I. TRI.   |
| BAA13547.1     | D88156   | Hyoscyamus niger  | tropinone reductase-I.   |
| AAA33282.1     | L20474   | Datura stramonium | catalyses a stereospecific reduction of tropinone to pseudotropine. tropinone reductase-II.                                  |
| BAA85845.1     | AB026545 | Hyoscyamus niger  | tropinone reductase-II. tr2.   |
| AAB09776.1     | L20485   | Hyoscyamus niger  | tropinone reductase-II.  |
| CAB52307.1     | AJ245634 | Solanum tuberosum | pseudotropine forming tropinone reductase. tropinone reductase II. TRII.   |
| CAC19810.1     | AJ292343 | Solanum tuberosum | pseudotropine forming tropinone reductase. tropinone reductase II. trII.   |
| CAA45866.1     | X64566   | Cuphea lanceolata | beta-ketoacyl-ACP reductase. Clkr27. 3-oxoacyl-[acyl-carrier protein] reductase.   |
| CAA45793.1     | X64463   | Brassica napus    | 3-oxoacyl-[acyl-carrier protein] reductase.  |
| AAB82767.1     | U89509   | Zea mays          | cuticular wax biosynthesis. b-keto acyl reductase. glossy8.  |
| AAB82766.1     | U89510   | Hordeum vulgare   | cuticular wax biosynthesis. b-keto acyl reductase. glossy8.  |
| AAB20114.2     | S60064   | Brassica napus    | enoyl-acyl carrier protein reductase. This sequence comes from fig3; conceptual translation differs from published sequence. |
| CAA74176.1     | Y13861   | Nicotiana tabacum | enoyl-ACP reductase. enr-T2.   |
| AAB82764.1     | U89511   | Allium porrum     | cuticular wax biosynthesis. b-keto acyl reductase. glossy8.  |

AAC78100.1 AF093628 *Oryza sativa*  
protochlorophyllide reductase homolog.

CAA74177.1 Y13862 *Nicotiana tabacum*  
enoyl-ACP reductase. enr-T1.

SEQ ID NO: 927

AAD43046.1 AF124045 *Sorghum bicolor*  
GTP-binding protein-like. 110K5.8. terminal exon, defined by homology to maize ESTs.

SEQ ID NO: 928

CAA09619.1 AJ011418 *Lycopersicon esculentum*  
ubiquitin activating enzyme.

AAC32140.1 AF051239 *Picea mariana*  
probable ubiquitin activating enzyme 2. Sb53. similar to *Arabidopsis thaliana* ubiquitin  
activating enzyme 2 encoded by GenBank Accession Number U40566.

AAA34308.1 M55604 *Triticum aestivum*  
ubiquitin-activating enzyme E1. UBA1.

AAA34265.1 M90663 *Triticum aestivum*  
ubiquitin activating enzyme. E1.

AAA34266.1 M90664 *Triticum aestivum*  
ubiquitin activating enzyme. E1.

CAA71762.1 Y10804 *Nicotiana tabacum*  
Ubiquitin activating enzyme E1. UBA1.

SEQ ID NO: 929

CAA55693.1 X79086 *Zea mays*  
initiator-binding protein. IBP2.

CAC19789.1 AJ251686 *Catharanthus roseus*  
putative transcription factor. MYB-like DNA-binding protein. bpf-1.

CAA55691.1 X79085 *Zea mays*  
initiator binding protein. IBP1.

AAF97508.1 AF242298 *Oryza sativa*  
telomere binding protein-1. TBP1.

SEQ ID NO: 930

AAK15447.1 AC037426 *Oryza sativa*  
putative flavin-containing monooxygenase. OSJNBb0014I11.12.

AAK15439.1 AC037426 *Oryza sativa*  
putative flavin-containing monooxygenase. OSJNBb0014I11.8.

BAA88198.1 AP000837 *Oryza sativa*  
Similar to human dimethylaniline monooxygenase (AC002376).

BAA88195.1 AP000837 *Oryza sativa*  
Similar to human dimethylaniline monooxygenase (AC002376).

BAB32703.1 AP002902 *Oryza sativa*  
dimethylaniline monooxygenase-like protein. P0492F05.5. contains EST C72123(E1029).

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| BAB07916.2     | AP002835 | Oryza sativa            | dimethylaniline monooxygenase-like protein. P0417G05.24. contains EST C72123(E1029).   |
| SEQ ID NO: 931 |          |                         |  |
| AAG38521.1     | AF283536 | Citrus x paradisi       | cystatin-like protein. cystein proteinase inhibitor.   |
| AAA97905.1     | U51853   | Glycine max             | cysteine proteinase inhibitor.   |
| CAA79954.1     | Z21954   | Vigna unguiculata       | cysteine proteinase inhibitor.   |
| AAB66355.1     | U54702   | Oryza sativa            | thiol proteinase inhibitor; cysteine proteinase inhibitor. oryzacystatin. has potential for insect resistance in rice and for cancer therapeutics. |
| AAB24010.1     | S49967   | Oryza                   | oryzacystatin. oryzacystatin. cysteine protease inhibitor; This sequence comes from Fig. 1.  |
| CAA60610.1     | X87126   | Zea mays                | cysteine proteinase inhibitor. zmc-II.   |
| BAA09666.1     | D63342   | Zea mays                | cysteine proteinase inhibitor. gCC.  |
| AAA32672.1     | L16624   | Ambrosia artemisiifolia | cystatin proteinase inhibitor.   |
| BAA07327.1     | D38130   | Zea mays                | inhibition against cysteine proteinases. cystatin II.  |
| AAA97907.1     | U51855   | Glycine max             | cysteine proteinase inhibitor.   |
| BAB18766.1     | AB038392 | Triticum aestivum       | cysteine proteinase inhibitor. WC61. cystatin.   |
| BAB18767.1     | AB038393 | Triticum aestivum       | cysteine proteinase inhibitor. WC92. cystatin.   |
| CAA11899.1     | AJ224331 | Castanea sativa         | cysteine proteinase inhibitor. cystatin.   |
| AAB71505.1     | U82220   | Pyrus communis          | cysteine protease inhibitor.   |
| AAA79239.1     | L48182   | Brassica rapa           | cysteine proteinase inhibitor. N-terminal deletion clone; putative.  |
| CAA60634.1     | X87168   | Sorghum bicolor         | cysteine proteinase inhibitor. CPII.   |
| AAA68150.1     | L42819   | Brassica rapa           | cysteine protenase inhibitor. N-terminal deletion clone; putative.   |
| BAB18765.1     | AB038391 | Triticum aestivum       | cysteine proteinase inhibitor. gWC2. cystatin.   |
| BAB18768.1     | AB038394 | Triticum aestivum       | cysteine proteinase inhibitor. WC83. cystatin.   |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAC37479.1 | L41355   | Brassica rapa           | cysteine proteinase inhibitor.  |
| BAA28867.1 | AB014760 | Cucumis sativus         | root-specific cystein protease inhibitor. cystein proteinase inhibitor.   |
| AAD33907.1 | AF143677 | Artemisia vulgaris      | cysteine proteinase inhibitor. CPI. pollen cystatin.  |
| BAA89582.1 | AP001073 | Oryza sativa            | ESTs AU067919(C10906),AU067918(C10906) correspond to a region of the predicted gene. Similar to cysteine proteinase inhibitor (D31700). |
| CAA72790.1 | Y12068   | Hordeum vulgare         | cysteine proteinase inhibitor. CPI.   |
| AAA96316.1 | U51119   | Brassica rapa           | cysteine proteinase inhibitor. BCPI-2.  |
| AAK15090.1 | AF240007 | Sesamum indicum         | cystatin. cysteine proteinase inhibitor.  |
| AAF23127.1 | AF198389 | Lycopersicon esculentum | cysteine proteinase inhibitor. cystatin. STC. expression induced by arachidonic acid and gamma linolenic acid.                          |
| BAA19610.1 | D64115   | Glycine max             | cysteine proteinase inhibitor. cystatin.  |
| BAA19608.1 | D31700   | Glycine max             | cysteine proteinase inhibitor. cystatin.  |
| CAA89697.1 | Z49697   | Ricinus communis        | cysteine proteinase inhibitor.  |
| AAF72202.1 | AF265551 | Manihot esculenta       | cysteine protease inhibitor.  |
| AAA97906.1 | U51854   | Glycine max             | cysteine proteinase inhibitor.  |
| CAA50437.1 | X71124   | Carica papaya           | cysteine proteinase inhibitor (cystatin).   |
| AAF23126.1 | AF198388 | Lycopersicon esculentum | cysteine proteinase inhibitor. cystatin. LTC. expression induced by arachidonic acid and gamma linolenic acid.                          |
| AAD13812.1 | AF117334 | Ipomoea batatas         | cysteine proteinase inhibitor. cystatin.  |
| AAF64480.1 | AF241536 | Ipomoea batatas         | cysteine protease inhibitor. CYSPI.   |
| AAK30004.1 | AY028994 | Dianthus caryophyllus   | cysteine proteinase inhibitor. DC-CPI.  |
| BAB18769.1 | AB038395 | Triticum aestivum       | cysteine proteinase inhibitor. WC81. cystatin.  |
| AAC32853.1 | AF083253 | Lycopersicon esculentum | cysteine protease inhibitor.  |

| SEQ ID NO: 938 |          |   |
|----------------|----------|---|
| CAA06486.1     | AJ005340 | Linum usitatissimum<br>IAA amidohydrolase. homolog.   |
| SEQ ID NO: 940 |          |   |
| BAB17350.1     | AP002747 | Oryza sativa<br>putative nodulin. P0698G03.34. contains ESTs<br>D39891(S1543),D41717(S4395),AU033037(S1543).  |
| BAA85440.1     | AP000616 | Oryza sativa<br>ESTs AU055729(S20023),AU055730(S20023) correspond to a region of the predicted gene.;<br>similar to Medicago nodulin N21-like protein (AC004218). |
| CAB53493.1     | AJ245900 | Oryza sativa<br>CAA303720.1 protein. q3037.20. Similar to Medicago nodulin N21 (MtN21).   |
| SEQ ID NO: 941 |          |   |
| BAA21921.1     | AB006599 | Petunia x hybrida<br>ZPT2-12. C2H2 zinc finger protein, 2 finger.   |
| BAA21922.1     | AB006600 | Petunia x hybrida<br>ZPT2-13. C2H2 zinc finger protein, 2finger.  |
| BAA19110.1     | AB000451 | Petunia x hybrida<br>PETHy;ZPT2-5. Cys(2) His(2) zinc finger protein, 2 fingers.  |
| BAA21923.1     | AB006601 | Petunia x hybrida<br>ZPT2-14. C2H2 zinc finger protein, 2 finger.   |
| BAA21925.1     | AB006603 | Petunia x hybrida<br>ZPT2-8. C2H2 zinc finger protein, 2 finger.  |
| BAA21924.1     | AB006602 | Petunia x hybrida<br>ZPT2-7. C2H2 zinc finger protein, 2finger.   |
| BAA21920.1     | AB006598 | Petunia x hybrida<br>ZPT2-11. C2H2 zinc finger protein, 2finger.  |
| CAA60828.1     | X87374   | Pisum sativum<br>putative zinc finger protein.  |
| BAA19111.1     | AB000452 | Petunia x hybrida<br>PETHy;ZPT2-6. Cys(2) His(2) zinc finger protein, 2 fingers.  |
| BAA21926.1     | AB006604 | Petunia x hybrida<br>ZPT2-9. C2H2 zinc finger protein, 2 finger.  |
| BAA21927.1     | AB006605 | Petunia x hybrida<br>ZPT3-3. C2H2 zinc finger protein, 3 finger.  |
| BAA96071.1     | AB035133 | Petunia x hybrida<br>C2H2 zinc-finger protein ZPT3-3. ZPT3-3.   |
| BAA21919.1     | AB006597 | Petunia x hybrida<br>ZPT2-10. C2H2 zinc finger protein, 2 finger.   |
| BAA96070.1     | AB035132 | Petunia x hybrida<br>C2H2 zinc-finger protein ZPT2-10. PETHy;ZPT2-10.   |
| CAB77055.1     | Y18788   | Medicago sativa<br>putative TFIIIA (or kruppel)-like zinc finger protein.   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAB53260.1     | U76554   | Brassica rapa           | transcription factor. zinc-finger protein-1. BR42.   |
| AAB53261.1     | U76555   | Brassica rapa           | zinc-finger protein BcZFP1. BcZFP1(3-2z).  |
| BAA19114.1     | AB000455 | Petunia x hybrida       | PETHy;ZPT4-1. Cys(2) His(2) zinc finger protein, 4 fingers.  |
| BAA05078.1     | D26085   | Petunia x hybrida       | zinc-finger DNA binding protein.   |
| AAD26942.1     | AF119050 | Datisca glomerata       | zinc-finger protein 1. zfp1. DgZFP1.   |
| BAA05077.1     | D26084   | Petunia x hybrida       | zinc-finger DNA binding protein.   |
| AAB39638.1     | U68763   | Glycine max             | putative transcription factor. SCOF-1. scof-1. zinc-finger protein.  |
| AAC06243.1     | AF053077 | Nicotiana tabacum       | transcription factor. osmotic stress-induced zinc-finger protein. zfp.   |
| BAA19112.1     | AB000453 | Petunia x hybrida       | PETHy;ZPT3-1. Cys(2) His(2) zinc finger protein, 3 fingers.  |
| AAK01713.1     | AF332876 | Oryza sativa            | zinc finger transcription factor ZF1.  |
| BAA05076.1     | D26083   | Petunia x hybrida       | zinc-finger DNA binding protein.   |
| BAA05079.1     | D26086   | Petunia x hybrida       | zinc-finger protein.   |
| BAA21928.1     | AB006606 | Petunia x hybrida       | ZPT4-4. C2H2 zinc finger protein, 4 finger.  |
| BAA19926.1     | AB000456 | Petunia x hybrida       | PETHy; ZPT4-2. C2H2 zinc finger protein, 4 finger.   |
| BAA19113.1     | AB000454 | Petunia x hybrida       | PETHy;ZPT3-2. Cys(2) His(2) zinc finger protein, 3 fingers.  |
| SEQ ID NO: 945 |          |                         |  |
| AAG10793.1     | AF296158 | Citrus unshiu           | beta-carotene hydroxylase. CHX1.   |
| AAG33636.1     | AF315289 | Citrus unshiu           | beta-carotene hydroxylase. CHX2. similar to beta-carotene hydroxylase of Citrus unshiu encoded by GenBank Accession Number AF296158. |
| AAG10430.1     | AF251018 | Tagetes erecta          | beta hydroxylase.  |
| CAB55625.1     | Y14809   | Lycopersicon esculentum | beta-carotene hydroxylase. CrtR-b1.  |
| CAA70888.1     | Y09722   | Capsicum annuum         | beta-carotene hydroxylase 2.   |

|                |          |                           |   |
|----------------|----------|---------------------------|---|
| CAB55626.1     | Y14810   | Lycopersicon esculentum   | beta-carotene hydroxylase. CrtR-b2.   |
| CAA70427.1     | Y09225   | Capsicum annuum           | beta-carotene hydrolase.  |
| CAC06712.1     | AJ278882 | Narcissus pseudonarcissus | synthesis of zeaxanthin. beta-carotene hydroxylase.   |
| AAD54243.1     | AF162276 | Haematococcus pluvialis   | carotenoid hydroxylase.   |
| SEQ ID NO: 946 |          |                           |   |
| AAC18914.1     | U94748   | Petunia x hybrida         | AN11. An11. No functional information available. Protein is involved in transcriptional regulation of anthocyanin biosynthesis in petunia. Protein contains five WD 40 repeats; WD 40 repeat protein. |
| SEQ ID NO: 948 |          |                           |   |
| AAG52887.1     | AF333386 | Nicotiana tabacum         | beta-expansin-like protein. PPAL. pollen allergen-like protein.   |
| AAF72986.1     | AF261273 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB5.  |
| AAF72990.1     | AF261277 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB9. putative group-1 pollen allergen.  |
| BAB20817.1     | AB051899 | Atriplex lentiformis      | beta-expansin. A1-EXP1.   |
| AAF72984.1     | AF261271 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB3.  |
| AAK15453.1     | AC037426 | Oryza sativa              | beta-expansin EXPB3. OSJNBb0014I11.1.   |
| AAF72991.1     | AF261278 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB10. putative group-1 pollen allergen.   |
| AAK15442.1     | AC037426 | Oryza sativa              | beta-expansin EXPB6. OSJNBb0014I11.3.   |
| AAF72987.1     | AF261274 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB6.  |
| AAF72983.1     | AF261270 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB1. putative group-1 pollen allergen Ory sl.   |
| AAF72988.1     | AF261275 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB7.  |
| AAF72989.2     | AF261276 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB8.  |
| AAF72985.1     | AF261272 | Oryza sativa              | putative cell wall loosening activity. beta-expansin. EXPB4.  |

|            |          |                         |   |
|------------|----------|-------------------------|---|
| AAK15440.1 | AC037426 | Oryza sativa            | beta-expansin EXPB2. OSJNBb0014I11.2.   |
| AAB61710.1 | U95968   | Oryza sativa            | beta-expansin. EXPB2. cell wall loosening protein.  |
| AAB37749.1 | U30460   | Cucumis sativus         | expansin S2 precursor. Cs-EXP2. similar to pollen allergen Lol pI, Lolium perenne, Swiss-Prot Accession Number P14946; former gene name CuExS2; expansin-30 (Ex30) protein. |
| AAD38296.1 | AC007789 | Oryza sativa            | putative expansin. OSJNBa0049B20.23.  |
| BAB18336.1 | AP002865 | Oryza sativa            | putative expansin. P0034C11.27.   |
| CAC19183.1 | AJ291816 | Cicer arietinum         | expansin.   |
| AAG13983.1 | AF297522 | Prunus avium            | expansin 2. Exp2. PruavExp2.  |
| AAG01875.1 | AF291659 | Striga asiatica         | alpha-expansin 3. Exp3.   |
| AAG32920.1 | AF184232 | Lycopersicon esculentum | expansin. Exp8.   |
| AAC96077.1 | AF049350 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP1. cell wall protein.   |
| AAC96078.1 | AF049351 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP2. cell wall protein.   |
| AAF35902.1 | AF230333 | Zinnia elegans          | expansin 3.   |
| BAA88200.1 | AP000837 | Oryza sativa            | EST AU078708(E60526) corresponds to a region of the predicted gene. Similar to expansin (U85246).   |
| AAC39512.1 | AF043284 | Gossypium hirsutum      | expansin. GhEX1. contains N-terminal signal peptide.  |
| AAD38297.1 | AC007789 | Oryza sativa            | putative expansin. OSJNBa0049B20.24.  |
| AAF32411.1 | AF230278 | Triphysaria versicolor  | alpha-expansin 1.   |
| BAB18338.1 | AP002865 | Oryza sativa            | putative expansin. P0034C11.29.   |
| AAG32921.1 | AF184233 | Lycopersicon esculentum | expansin. Exp10.  |
| CAC19184.1 | AJ291817 | Cicer arietinum         | expansin.   |
| AAC96079.1 | AF049352 | Nicotiana tabacum       | involved in acid-growth response. alpha-expansin precursor. Nt-EXP3. cell wall protein.   |

SEQ ID NO: 950



|            |          |                                |   |
|------------|----------|--------------------------------|---|
| CAA52213.1 | X74115   | <i>Picea abies</i>             | short-chain alcohol dehydrogenase.  |
| AAC35342.1 | AF072449 | <i>Ipomoea trifida</i>         | short-chain alcohol dehydrogenase. SSP. self-incompatibility-locus specific stigma protein.   |
| AAC35340.1 | AF072447 | <i>Ipomoea trifida</i>         | short-chain alcohol dehydrogenase. SSP. self-incompatibility-locus specific stigma protein.   |
| CAA11153.1 | AJ223177 | <i>Nicotiana tabacum</i>       | short chain alcohol dehydrogenase.  |
| CAA11154.1 | AJ223178 | <i>Nicotiana tabacum</i>       | short chain alcohol dehydrogenase. SCANT.   |
| AAK29646.1 | AF349916 | <i>Solanum tuberosum</i>       | putative short-chain type alcohol dehydrogenase. GAN; similar to tomato Leert10 and maize Ts2.  |
| AAC37345.1 | L20621   | <i>Zea mays</i>                | alcohol dehydrogenase. short chain.   |
| AAC35341.1 | AF072448 | <i>Ipomoea trifida</i>         | short-chain alcohol dehydrogenase. SSP. self-incompatibility-locus specific stigma protein.   |
| AAC35343.1 | AF072450 | <i>Ipomoea trifida</i>         | short-chain alcohol dehydrogenase. SSP. self-incompatibility-locus specific stigma protein.   |
| AAB57737.1 | U89270   | <i>Tripsacum dactyloides</i>   | short-chain alcohol dehydrogenase. gynomonoecious sex form 1. similar to <i>Zea mays</i> tasselseed 2: SwissProt Accession Number P50160.   |
| AAF89645.1 | AF169018 | <i>Glycine max</i>             | seed maturation protein PM34. PM34. similar to bacterial glucose and ribitol dehydrogenase.   |
| AAF04253.1 | AF097651 | <i>Pisum sativum</i>           | short-chain alcohol dehydrogenase SAD-C. sadC. contains the entire nucleotide binding motif of 3(alpha), 20(beta)-hydroxysteroid dehydrogenases, GXXXXXXGXXG(A)XGXXXXA (Ghosh et al., 1991, Proc. Natl. Acad. Sci. USA 88, 10064-10068).  |
| AAF04193.1 | AF053638 | <i>Pisum sativum</i>           | short-chain alcohol dehydrogenase. sadA. contains the entire nucleotide binding motif of the 3(alfa),20(beta)-hydroxysteroid dehydrogenase: GXXXXXXGXXG(A)XGXXXXA (Ghosh et al. (1991) Proc. Natl. Acad. Sci. USA 88, 10064-10068); similar to <i>Lycopersicon esculentum</i> product encoded by GenBank Accession Number U21801 and <i>Streptomyces hydrogenans</i> steroid alcohol dehydrogenase. |
| AAB57738.1 | U89271   | <i>Tripsacum dactyloides</i>   | short-chain alcohol dehydrogenase. gynomonoecious sex form 1. similar to <i>Zea mays</i> tasselseed 2: SwissProt Accession Number P50160.   |
| CAB91875.1 | AJ277945 | <i>Lycopersicon esculentum</i> | putative alcohol dehydrogenase. yfe37.  |
| AAF04194.1 | AF053639 | <i>Pisum sativum</i>           | short-chain alcohol dehydrogenase. sadB. contains the Prosite pattern no. PS00061 for short-chain alcohol dehydrogenases; contains a deletion in the nucleotide binding motif of the 3(alfa),20(beta)-hydroxysteroid dehydrogenase: GXXXXXXGXXG(A)XGXXXXA (Ghosh et al. (1991) Proc. Natl. Acad. Sci. USA 88, 10064-10068).   |

|                |          |                         |  |
|----------------|----------|-------------------------|--|
| AAB00109.1     | U21801   | Lycopersicon esculentum | alcohol dehydrogenase homolog. GAD3. mRNA is suppressed in the presence of gibberellin; similar to nonmetallo-short-chain alcohol dehydrogenases, PIR Accession Number A47542. |
| SEQ ID NO: 952 |          |                         |  |
| CAB63264.1     | AJ251808 | Lotus japonicus         | calcium-binding protein. cbp1.   |
| AAG43547.1     | AF211529 | Nicotiana tabacum       | Avr9/Cf-9 rapidly elicited protein 31. ACRE31. similar to Solanum tuberosum CAST calcium binding protein encoded by GenBank Accession Number L02830.                           |
| AAA34014.1     | L01432   | Glycine max             | calcium-binding regulatory protein. calmodulin. SCaM-3. putative.  |
| AAA34013.1     | L01430   | Glycine max             | calcium-binding regulatory protein. calmodulin. SCaM-1. putative.  |
| AAA92681.1     | U13882   | Pisum sativum           | calcium-binding protein. calmodulin.   |
| CAA78301.1     | Z12839   | Lilium longiflorum      | calcium binding protein, signal transduction. calmodulin.  |
| AAA33397.1     | L18912   | Lilium longiflorum      | calcium binding protein, signal transduction. calmodulin. putative.  |
| AAA19571.1     | U10150   | Brassica napus          | calcium binding. calmodulin. bcm1.   |
| AAA85157.1     | U20297   | Solanum tuberosum       | calcium-binding protein. calmodulin.   |
| AAA85156.1     | U20296   | Solanum tuberosum       | calcium-binding protein. calmodulin.   |
| AAA62351.1     | U20295   | Solanum tuberosum       | calcium-binding protein. calmodulin.   |
| AAA85155.1     | U20294   | Solanum tuberosum       | calcium-binding protein. calmodulin.   |
| AAC49587.1     | U49105   | Triticum aestivum       | calmodulin TaCaM4-1. calcium-binding protein.  |
| AAC49586.1     | U49104   | Triticum aestivum       | calmodulin TaCaM3-3. calcium-binding protein.  |
| AAC49585.1     | U49103   | Triticum aestivum       | calmodulin TaCaM3-2. calcium-binding protein.  |
| AAC49584.1     | U48693   | Triticum aestivum       | calmodulin TaCaM3-1. calcium-binding protein.  |
| AAC49580.1     | U48689   | Triticum aestivum       | calmodulin TaCaM1-3. calcium-binding protein.  |
| AAC49579.1     | U48688   | Triticum aestivum       | calmodulin TaCaM1-2. calcium binding protein.  |
| AAC49578.1     | U48242   | Triticum aestivum       | calmodulin TaCaM1-1. calcium-binding.  |

---

|   |        |              |
|---|--------|--------------|
| CAA78287.1  | Z12827 | Oryza sativa |
| calcium binding protein, signal transduction. calmodulin. |        |              |

---

|   |        |             |
|---|--------|-------------|
| AAA03580.1  | L01431 | Glycine max |
| calcium-binding regulatory protein. calmodulin. SCaM-2. putative. |        |             |

---

|   |        |              |
|---|--------|--------------|
| AAA33901.1  | L18913 | Oryza sativa |
| calcium binding protein, signal transduction. calmodulin. putative. |        |              |

---

|   |        |             |
|---|--------|-------------|
| AAA34015.1  | L01433 | Glycine max |
| calcium-binding regulatory protein. calmodulin. SCaM-4. putative. |        |             |

---

**Table 24** Plant Open Reading Frames (ORFs) orthologous to specific *Chenopodium* ORFs

| SEQ ID NO:1956 |  |
|----------------|--|
| CAB51903.1     | AJ242807 <i>Brassica napus</i><br>endo-1,4-beta-D-glucanase. Cell6. cellulase.   |
| BAA94257.1     | AB040769 <i>Hordeum vulgare</i><br>endo-1,4-beta-glucanase Cell. Cell.   |
| AAC49704.1     | U78526 <i>Lycopersicon esculentum</i><br>endo-1,4-beta-glucanase. Cel3.  |
| SEQ ID NO:1957 |  |
| BAB21273.1     | AP002844 <i>Oryza sativa</i><br>putative ABC transporter protein. P0410E03.4.  |
| BAB21275.1     | AP002844 <i>Oryza sativa</i><br>putative ABC transporter protein. P0410E03.6.  |
| BAB21276.1     | AP002844 <i>Oryza sativa</i><br>putative ABC transporter protein. P0410E03.7. contains EST<br>D22472(C1173).   |
| CAA94437.1     | Z70524 <i>Spirodela polyrrhiza</i><br>multidrug resistance protein. PDR5-like ABC transporter.   |
| BAB21279.1     | AP002844 <i>Oryza sativa</i><br>putative ABC transporter protein. P0410E03.10. contains ESTs<br>AU065360(R3463), AU101680(R3463).  |
| CAA03960.1     | AJ000234 <i>Hordeum vulgare</i><br>partial sequence, homology to PDR5-like ABC transporter.  |
| SEQ ID NO:1960 |  |
| AAD51778.1     | AF116858 <i>Phaseolus vulgaris</i><br>utilizes UDPX as the sugar donor and catalyzes the formation<br>of O-xylosylzeatin from zeatin. zeatin O-xylosyltransferase. ZOX1.   |
| AAD04166.1     | AF101972 <i>Phaseolus lunatus</i><br>catalyzes formation of O-glucosylzeatin from zeatin and<br>UDPG; catalyzes formation of O-xylosylzeatin from zeatin and UDPX. zeatin<br>O-glucosyltransferase. ZOG1. cytokinin O-glucosyltransferase. |
| BAA36410.1     | AB012114 <i>Vigna mungo</i><br>UDP-glucose:flavonoid glycosyltransferase. UFGlyT.  |
| BAB17061.1     | AP002523 <i>Oryza sativa</i><br>putative glucosyl transferase. P0013F10.7. contains EST<br>C73149(E2992).  |

---

BAA36412.1 AB012116Vigna mungo  
UDP-glucose:flavonoid glycosyltransferase. UFGlyT.

---

AAK28303.1 AF346431Nicotiana tabacum  
phenylpropanoid:glucosyltransferase 1. togt1.  
glucosyltransferase.

---

BAB17059.1 AP002523Oryza sativa  
putative glucosyl transferase. P0013F10.5.

---

AAF17551.1 AF198453Glycine max  
UDP-glucose:flavonoid glycosyltransferase. UFGlyT.

---

AAB36653.1 U32644 Nicotiana tabacum  
immediate-early salicylate-induced glucosyltransferase.  
IS5a.

---

CAB88666.1 AJ400861Cicer arietinum  
flavonoid glycosyltransferase. putative UDP-glucose.

---

AAB36652.1 U32643 Nicotiana tabacum  
immediate-early salicylate-induced glucosyltransferase.  
IS10a.

---

CAA54612.1 X77462 Manihot esculenta  
UTP-glucose glucosyltransferase. CGT5.

---

BAB17060.1 AP002523Oryza sativa  
putative glucosyl transferase. P0013F10.6.

---

AAK28304.1 AF346432Nicotiana tabacum  
phenylpropanoid:glucosyltransferase 2. togt2.  
glucosyltransferase.

---

CAA59450.1 X85138 Lycopersicon esculentum  
twi1. homologous to glucosyltransferases.

---

BAA83484.1 AB031274Scutellaria baicalensis  
UDP-glucose: flavonoid 7-O-glucosyltransferase. ufgt.

---

CAA54611.1 X77461 Manihot esculenta  
UTP-glucose glucosyltransferase. CGT2.

---

BAA36411.1 AB012115Vigna mungo  
UDP-glucose:flavonoid glycosyltransferase. UFGlyT.

---

CAC35167.1 AJ310148Rauvolfia serpentina  
arbutin synthase. as.

---

BAB17182.1 AP002843Oryza sativa  
putative UTP-glucose glucosyltransferase. P0407B12.19.

---

|            |          |                                    |   |
|------------|----------|------------------------------------|---|
| CAB56231.1 | Y18871   | <i>Dorotheanthus bellidiformis</i> | betanidin-5-O-glucosyltransferase.  |
| BAB17176.1 | AP002843 | <i>Oryza sativa</i>                | putative UTP-glucose glucosyltransferase. P0407B12.13.  |
| AAK16172.1 | AC079887 | <i>Oryza sativa</i>                | putative glucosyltransferase. OSJNBa0040E01.14.   |
| CAA54613.1 | X77463   | <i>Manihot esculenta</i>           | UTP-glucose glucosyltransferase. CGT6.  |
| AAF61647.1 | AF190634 | <i>Nicotiana tabacum</i>           | UDP-glucose:salicylic acid glucosyltransferase. SA-GTase.                                       |
| BAA93039.1 | AB033758 | <i>Citrus unshiu</i>               | limonoid UDP-glucosyltransferase. LGTase.   |
| CAA54610.1 | X77460   | <i>Manihot esculenta</i>           | UTP-glucose glucosyltransferase. CGT4.  |
| AAA59054.1 | L34847   | <i>Zea mays</i>                    | conjugation of the growth hormone indole-3-acetic acid (IAA). IAA-glu synthetase. iaglu.        |
| BAA36423.1 | AB013598 | <i>Verbena x hybrida</i>           | UDP-glucose:anthocyanin 5-O-glucosyltransferase. HGT8.  |
| AAF17077.1 | AF199453 | <i>Sorghum bicolor</i>             | UDP-glucose glucosyltransferase.<br>UDP-glucose:p-hydroxymandelonitrile-o- glucosyltransferase. |
| CAA54558.1 | X77369   | <i>Solanum melongena</i>           | glycosyl transferase. GT.   |
| AAK16175.1 | AC079887 | <i>Oryza sativa</i>                | putative glucosyltransferase. OSJNBa0040E01.15.   |
| CAA81057.1 | Z25802   | <i>Petunia x hybrida</i>           | UDP rhamnose: anthocyanidin-3-glucoside<br>rhamnosyltransferase.                                |
| CAA50377.1 | X71060   | <i>Petunia x hybrida</i>           | anthocyanin: rhamnosyltransferase. rt.  |
| CAA50376.1 | X71059   | <i>Petunia x hybrida</i>           | anthocyanin 3 glucoside: rhamnosyltransferase. rt.  |
| AAK16181.1 | AC079887 | <i>Oryza sativa</i>                | putative glucosyltransferase. OSJNBa0040E01.16.   |
| AAK16178.1 | AC079887 | <i>Oryza sativa</i>                | putative glucosyltransferase. OSJNBa0040E01.5.  |

---

AAD21086.1 AF127218 *Forsythia x intermedia*  
adds glucose residue to position 3 of flavonoid compounds.  
flavonoid 3-O-glucosyltransferase. UFGT.

---

BAA89008.1 AB027454 *Petunia x hybrida*  
anthocyanidin 3-O-glucosyltransferase. PGT8.

---

**SEQ ID NO:1962**

---

CAA65580.1 X96784 *Nicotiana tabacum*  
cytochrome P450. hsr515.

---

CAA64635.1 X95342 *Nicotiana tabacum*  
cytochrome P450. hsr515. hypersensitivity-related gene.

---

AAG44132.1 AF218296 *Pisum sativum*  
cytochrome P450. P450 isolog.

---

AAG49299.1 AF313489 *Callistephus chinensis*  
flavonoid 3',5'-hydroxylase.

---

BAA03438.1 D14588 *Petunia x hybrida*  
flavonoid-3',5'-hydroxylase. Hfl.

---

AAC32274.1 AF081575 *Petunia x hybrida*  
flavonoid 3',5'-hydroxylase. Hfl. P450 enzyme.

---

CAA50442.1 X71130 *Petunia x hybrida*  
P450 hydroxylase. PET 1.

---

CAA80266.1 Z22545 *Petunia x hybrida*  
flavonoid 3',5'-hydroxylase.

---

AAG49315.1 AF315465 *Pelargonium x hortorum*  
flavonoid 3'-hydroxylase.

---

AAG49298.1 AF313488 *Callistephus chinensis*  
putative flavonoid 3'-hydroxylase.

---

AAB17562.1 U72654 *Eustoma grandiflorum*  
flavonoid 3'5'-hydroxylase. F3'5'H. cytochrome P450; CYP75.

---

BAB20076.1 AB012925 *Torenia hybrida*  
flavonoid 3',5'-hydroxylase. F3'5'H.

---

AAG14961.1 AF214007 *Brassica napus*  
cytochrome p450-dependent monooxygenase. BNF5H1.

---

CAC26920.1 AJ295586 *Arabidopsis lyrata* subsp. *petraea*  
ferulate-5-hydroxylase. fah1.

---

AAG14962.1 AF214008 *Brassica napus*  
cytochrome p450-dependent monooxygenase. BNF5H2.

---

## References

- Aarts et al., Proc Natl Acad Sci 95:10306 (1998).
- Alonso et al., Science, 284:2148 (1999)
- Altschul et al., J. Mol. Biol. 215:403 (1990).
- 5 Altschul et al., Nucleic Acids Res. 25:3389 (1997).
- An et al., EMBO J. 4:277 (1985).
- Aoyama et al., N-H Plant Journal, 11:605 (1997).
- Bachem et al., Plant J., 9:745 (1996).
- Bailey and Elkan In: Proceedings of the second international Conference on  
 10 Intelligent Systems for Molecular Biology (Altman, R., ed), pp 28-36, AAAI  
 press (1994).
- Ballas et al., Nucleic Acids Res. 17:7891 (1989).
- Balzi et al., J. Biol. Chem., 269:2206 (1994).
- Bansal et al., Proc. Natl. Acad. Sci. USA, 89:3654 (1992).
- 15 Batzer et al., Nucleic Acid Res., 19:5081 (1991).
- Beachy and Murakishi, Phytopathology, 61:877 (1971).
- Beals et al., Plant Cell, 9:1527 (1997).
- Belanger et al., Genetics, 129:863 (1991).
- Bevan et al., Nature, 304:184 (1983).
- 20 Blochinger & Diggelmann, Mol Cell Biol, 4:2929.
- Bourouis et al., EMBO J., 2:1099 (1983).
- Boyes et al., Proc Natl Acad Sci USA, 95:15849 (1998).
- Byrne et al. Plant Cell Tissue and Organ Culture, 8:3 (1987).
- Campbell and Gowri, Plant Physiol., 92:1 (1990).
- 25 Canto and Palukaitis, Virology, 265:74 (1999).
- Cao et al., Plant Cell, 6:1583 (1994).
- Cao et al., Cell, 88:57 (1997).
- Century et al., Science, 278:1963 (1997).
- Chandler et al., Plant Cell, 1:1175 (1989).
- 30 Christou et al., Biotechnology, 9:957 (1991).
- Christou et al., Plant Physiol., 87:671 (1988).
- Conklin and Last, Plant Physiol., 109:203 (1995).
- Cooper et al., Virology, 206:307 (1995)



- Cooper et al., Virology, 216:208 (1996).
- Cordero et al., Plant J., 6:141 (1994).
- Corpet et al., Nucleic Acids Res. 16:10881 (1988).
- Cramer et al., Nature Biotech., 15:436 (1997).
- 5 Cramer et al., Nature, 391:288 (1998).
- Creelman et al., Annu Rev Plant Physiol Plant Mol Biol 48:355 (1997).
- Crossway et al., BioTechniques, 4:320 (1986).
- Czako et al., Mol. Gen. Genet. 23 5 (1), 33-40 (1992).
- Czernic et al., Plant Mol. Biol., 31:255 (1996).
- 10 Datta et al., Bio/Technology, 8, 736 (1990).
- Dayhoff et al., Atlas of Protein Sequence and Structure, Natl. Biomed. Res. Found., Washington, C.D. (1978).
- De Blaere et al., Meth. Enzymol. 143:277 (1987).
- de Framond, FEBS, 290:103 (1991).
- 15 Delaney et al., Proc. Natl. Acad. Sci. USA 92:6602 (1995).
- Della-Cioppa et al., Plant Physiology, 84:965 (1987).
- De Oliveira et al., Microbios., 76:213 (1993).
- Dennis et al., Nucleic Acids Res., 12:3983 (1984).
- Dong et al., Curr Opin Plant Biol 1:316 (1998).
- 20 Dunigan and Madlener, Virology, 207:460 (1995).
- Durrant et al., The Plant Cell, 12:963 (2000).
- Dzelkalns et al., Plant Cell, 5:855 (1993).
- Eisen et al., Trends in Plant Sci. 95:14863 (2000).
- Ellis and Jones, Curr Opin Plant Bio 1:288 (1998).
- 25 Elroy-Stein et al., PNAS USA, 86:6126 (1989).
- English, et al., Plant Cell 8:179 (1996).
- Eulgem et al., Trends in Plant Sci. 5:199 (2000).
- Falk et al., Proc Natl Acad Sci USA 96:3292 (1999).
- Feys et al., Plant Cell 6:751 (1994).
- 30 Franken et al., EMBO J., 10:2605 (1991).
- Fromm et al., Bio/Technology 8:833 (1990).
- Gallie et al., Molecular Biology of RNA, 237 (1989).
- Gallie et al., Nucl. Acids Res., 15:8693 (1987).

- Gan et al., Science (1995), 270 (5244), 1986-8).
- Gatz Current Opinion in Biotechnology, 7:168 (1996).
- Gatz, C., Annu. Rev. Plant Physiol. Plant Mol. Biol., 48:89 (1997).
- Gelfand, eds., PCR Strategies (Academic Press, New York (1995)).
- 5 Glazebrook, J., Curr. Opin. Plant Biology, 2:280 (1999).
- Glazebrook et al., Genetics 143:973 (1996).
- Glazebrook et al., Proc. Natl. Acad. Sci. USA 91:8955 (1994).
- Gordon-Kamm et al., Plant Cell, 2, 603 (1990).
- Graham et al., Biochem. Biophys. Res. Comm., 101:1164 (1981).
- 10 Graham et al., J. Biol. Chem., 260:6555 (1985).
- Graham et al., J. Biol. Chem., 260:6561 (1985).
- Guerineau et al., Mol. Gen. Genet. 262:141 (1991).
- Gulyas and Farkas, Phytopath. Z., 91:182 (1978)
- Hammand-Kosack and Jones, Ann. Rev. Plant Physiol. Plant Mol. Biol., 48:575 \
- 15 1997).
- Heinlein et al., Science, 270:1983 (1995).
- Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA, 89:10915 (1989).
- Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA, 89:10915 (1992).
- Hiei et al., Plant J. 6:271 (1994).
- 20 Higgins et al., Gene 73:237 (1988).
- Higgins et al., CABIOS 5:151 (1989).
- Hinchee et al., Biotechnology, 6:915 (1988).
- Hoekema, In: The Binary Plant Vector System. Offset-drukkerij Kanters B.V.
- Horvath and Chua, Plant Mol. Biol., 31:1061 (1996).
- 25 Huang et al., CABIOS 8:155 (1992).
- Hudspeth & Grula, Plant Molec. Biol., 12:579 (1989).
- Huffman et al., J. Cell. Biochem., 17B: Abstract.
- Hunt et al., Mol. Plant-Microbe Int. 9:261 (1997).
- Ingelbrecht et al., Plant Cell, 1:671 (1989).
- 30 Innis et al., eds., PCR Protocols: A Guide to Methods and Applications (Academic Press, New York (1995).
- Innis and Gelfand, eds., PCR Methods Manual (Academic Press, New York) (1999).

- Jobling et al., Nature, 325:622 (1987).
- John et al., Proc. Natl. Acad. Sci. USA 89(13):5769 (1992).
- Jones et al., Adv Bot Res 24:89 (1997).
- Joshi et al., Nucleic Acid Res. 15:9627 (1987).
- 5 Joshi, Nucl. Acid Res., 15, 6643 (197).
- Karlin and Altschul, Proc. Natl. Acad. Sci. USA 87:2264 (1990).
- Karlin and Altschul, Proc. Natl. Acad. Sci. USA 90:5873 (1993).
- Keller et al., Genes Dev., 3:1639 (1989).
- Klein et al., Bio/Technology, 6:559 (1988).
- 10 Klein et al., Nature (London) 327:70 (1987).
- Klein et al., Plant Physiol., 91:440 (1988).
- Klein et al., Proc. Natl. Acad. Sci. USA, 85:4305 (1988).
- Kohler et al., Plant Mol. Biol., 29:1293 (1995).
- Knauf, et al., Genetic Analysis of Host Range Expression by *Agrobacterium* In:
- 15 Molecular Genetics of the Bacteria-Plant Interaction, Puhler, A. ed., Springer-Verlag, New York, 245 (1983).
- Komari, Plant Cell Reports, 9:303 (1990).
- Kozziel et al., Biotechnology, 11:194 (1993).
- Kridl et al., Seed Science Research, 1:209 (1991).
- 20 Kriz et al., Mol. Gen. Genet., 207:90 (1987).
- Kunkel, Proc. Natl. Acad. Sci. USA, 82:488 (1985).
- Kunkel et al., Methods in Enzymol., 154:367 (1987).
- Lange et al., Plant Sci., 142:133 (1999).
- Langridge et al., Cell, 34:1015 (1983).
- 25 Lashbrook et al., Plant Cell, 6:1485 (1994).
- Leister et al., P.N.A.S., (USA), 95:370 (1998).
- Leister et al., Proc. Natl. Acad. Sci. USA, 95:370 (1998).
- Liang et al., Science, 257:967 (1992).
- Lindstrom et al., Dev. Genet., 11:160 (1990).
- 30 Lommel et al., Virology, 81:382 (1991).
- Ly et al., Science, 287:2486 (2000).
- Macejak et al., Nature, 353:90 (1991).
- Mansson et al., Gen. Genet., 200:356 (1985).

- Martin and Paz-Ares, Trends in Genetics 13:67 (1997).
- Martinez et al., J. Mol. Biol., 208:551 (1989).
- McBride et al., Proc. Natl. Acad. Sci. USA, 91:7301 (1994).
- McCabe et al., Bio/Technology, 6:923 (1988).
- 5 McDowell et al., Plant J. 22:523 (2000).
- McDowell et al., Plant Cell 10:1861 (1998).
- McNellis et al., Plant J. 1998, 14:247-257.
- Meinkoth and Wahl, Anal. Biochem., 138:267 (1984).
- Messing & Vierra, Gene, 19:259 (1982).
- 10 Meyers et al., Plant J. 20:317 (1999).
- Mogen et al., Plant Cell 2:1261 (1990).
- Moore et al., J. Mol. Biol., 272:336 (1997).
- Munroe et al., Gene 91:151 (1990).
- Murray et al., Nucleic Acids Res., 17:477 (1989).
- 15 Myers and Miller, CABIOS 4:11 (1988).
- Needleman and Wunsch, J. Mol. Biol. 48:443 (1970).
- Odell et al., Nature, 313:810 (1985).
- Odell et al., Nature, 313:810 (1985).
- Ohtsuka et al., J. Biol. Chem., 260:2605 (1985).
- 20 Okamuro et al., Biochemistry of Plants, 15:1 (1989).
- Otsuki et al., Virology, 50:45 (1972)
- Paszkowski et al., EMBO J., 3:2717 (1984).
- Pacciotti et al. Bio/Technology 3:241 (1985).
- Padgett and Beachy, Plant Cell, 5:577 (1993).
- 25 Park et al., J. Plant Biol. 38(4):365 (1985).
- Pearson et al., Meth. Mol. Biol. 24:307 (1994).
- Pearson and Lipman, Proc. Natl. Acad. Sci. 85:2444 (1988).
- Penninck et al., Plant Cell 8:2309 (1996).
- Perlak et al., Proc. Natl. Acad. Sci. USA, 88:3324 (1991).
- 30 Pieterse et al., Plant Cell, 10:1571 (1998).
- Pieterse et al., Plant Cell, 8:1225 (1996).
- Proudfoot, Cell 64:671 (1991).
- Quigley et al., J. Mol. Evol., 29:412 (1989).

- Ralston et al., Genetics, 119:185 (1988).
- Reina et al., Nucleic Acids Res., 18:6425 (1990).
- Reina et al., Nucleic Acids Res., 18:7449 (1990).
- Riggs et al., Proc. Natl. Acad. Sci. USA, 83:5602 (1986).
- 5     Rochester et al., (1986).
- Rommens et al., Plant Cell, 7:1537 (1995).
- Ronald, Curr Opin Plant Biol 1:294 (1998).
- Rossolini et al., Mol. Cell. Probes, 8:91 (1994).
- Roth et al., Nature Biotechnology, 16:939 (1998).
- 10     Ruiz et al., Plant Cell 10:937 (1998).
- Ryals et al., Plant Cell 8:1809 (1996).
- Ryals et al., Plant Cell 9:425 (1997)
- Sambrook et al., Molecular Cloning: A Laboratory Manual (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York) (1989).
- 15     Sanchez-Fernandez et al., Mol. Gen. Genet., 258:655 (1998).
- Sanfacon et al., Genes Dev. 5:141 (1991).
- Sanford et al., Particulate Science and Technology, 5:27 (1987).
- Schernthaner et al., EMBO J., 7:1249 (1988).
- Schindler et al., Plant Cell 4:1309 (1992).
- 20     Schmitz and Rao, Virology, 248:323 (1996)
- Schwob et al., Plant J., 4:423 (1993).
- Shah et al., Mol. Plant-Microbe Interact. 10:69 (1997).
- Shimamoto et al., Nature, 338:274 (1989).
- Shirasu et al., Plant Cell 9:261 (1997).
- 25     Shulaev et al., Plant Cell 7:1691 (1995).
- Simpson, Plant Mol. Biol., 19:699 (1985).
- Skuzeski et al., Plant Molec. Biol., 15:65 (1990).
- Slater et al., Plant Mol. Biol., 5:137 (1985).
- Smart and Fleming, J. Biol. Chem., 271:19351 (1996).
- 30     Smith et al., Adv. Appl. Math. 2:482 (1981).
- Smith et al., Planta 168:94 (1986).
- Song et al., Science, 270:1804 (1995).
- Spencer et al., Theor Appl Genet, 79:625 (1990).

- Staswick et al., Proc. Natl. Acad. Sci. USA 89:6837 (1992).
- Staub et al., EMBO J., 12:601 (1993).
- Staub et al., Plant Cell, 4:39 (1992).
- Stemmer, Nature, 370:389 (1994).
- 5 Stemmer, Proc. Natl. Acad. Sci. USA, 91:10747 (1994).
- Sukhapinda et al. Plant Mol. Biol. 8:209 (1987).
- Sullivan et al., Mol. Gen. Genet., 215:431 (1989).
- Svab et al., Proc. Natl. Acad. Sci. USA, 87:8526 (1990).
- Svab et al., Proc. Natl. Acad. Sci. USA, 90:913 (1993).
- 10 Thomma et al., Plant Physiol., 121:1093 (1999).
- Thomma et al., Proc. Natl. Acad. sci. U.S.A., 85:15107 (1998).
- Tijssen, Laboratory Techniques in Biochemistry and Molecular Biology-  
Hybridization with Nucleic Acid Probes, part I chapter 2 "Overview of  
principles of hybridization and the strategy of nucleic acid probe assays"
- 15 Elsevier, New York (1993).
- Turner et al., Molecular Biotechnology, 3:225 (1995).
- VanTunen et al., EMBO J., 7:1257 (1988).
- Vasil et al., Biotechnology, 11:1553 (1993).
- Verduin, J. Gen. Virol., 38:571 (1978).
- 20 Vernooij et al., Plant Cell 6:959 (1994).
- Visedo et al., Physiologia Plantarum, 78:218 (1990).
- Vodkin, Prog. Clin. Biol. Res., 138:87 (1983).
- Vogel et al., EMBO J., 11:157 (1992).
- Walker and Gastra, eds., Techniques in Molecular Biology, MacMillan
- 25 Publishing Company, New York (1983).
- Wandelt et al., Nucleic Acids Res., 17:2354 (1989).
- Waterman, M.S. Introduction to Computational Biology: Maps, sequences and  
genomes. Chapman & Hall. London (1995).
- Weeks et al., Plant Physiol., 102:1077 (1993).
- 30 Weissinger et al., Annual Rev. Genet., 22:421 (1988).
- Wenzler et al., Plant Mol. Biol., 13:347 (1989).
- Weymann et al., Plant Cell 7:2013 (1995).
- White et al., Nucl Acids Res, 18:1062 (1990).

- Whitham et al., P.N.A.S., (USA), 93:8776 (1996).
- Whitham et al., Cell, 78:1101 (1994).
- Willitset al., Mol Plant-Microbe Interact 11:795 (1998).
- Xie et al., Science 280:1091 (1998).
- 5 Yamamoto et al., Nucleic Acids Res., 18:7449 (1990).
- Yang et al., Plant Mol. Biol., 38:1201 (1998).
- Yang & Klessig, Proc. Natl. Acad. Sci. USA 93:14972 (2000).
- Yu et al., Proc. Natl. Acad. Sci. USA 95:7819 (1998).
- Zhang et al., Proc. Natl. Acad. Sci. USA, 94:4504 (1997).
- 10 Zhou et al., EMBO J., 16:3207 (1997).
- Zhou et al., Plant Cell 8:2235 (1996).
- Zhou et al., Plant Cell 10:1021 (1998).

- 15 All publications, patents and patent applications are incorporated herein by reference.
- While in the foregoing specification this invention has been described in relation to certain preferred embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled in the art that the invention is susceptible to additional embodiments and that certain of the details described herein may be varied
- 20 considerably without departing from the basic principles of the invention.

**WHAT IS CLAIMED IS:**

1. A method to identify at least one gene involved in plant resistance or response to infection by at least one pathogen comprising:
  - 5 a) determining or detecting plant gene expression in an incompatible interaction; and
  - b) identifying at least one gene whose expression is significantly altered in the incompatible interaction relative to expression of the at least one gene in an uninfected plant, in a mutant plant that does not express a gene associated with response to infection by a pathogen, or in a corresponding compatible interaction.
- 10 2. A method to identify at least one gene involved in plant resistance or response to infection by at least one pathogen comprising:
  - a) determining or detecting plant gene expression in a compatible interaction; and
  - 15 b) identifying at least one gene whose expression is significantly altered in the compatible interaction relative to expression of the at least one gene in an uninfected plant, in a mutant plant that does not express a gene associated with response to infection by a pathogen, or in a corresponding incompatible interaction.
- 20 3. The method of claim 1 or 2 wherein the compatible interaction is between a plant having a resistance gene and a pathogen lacking a corresponding avirulence gene, a plant lacking a resistance gene to a pathogen having a corresponding avirulence gene, or a plant lacking a resistance gene and a pathogen lacking a corresponding avirulence gene.
- 25 4. The method of claim 1 or 2 wherein expression of the at least one gene is upregulated in response to infection.
5. The method of claim 1 or 2 wherein expression of the at least one gene is downregulated in response to infection.
- 30 6. The method of claim 1 or 2 wherein the at least one gene encodes a polypeptide that is substantially similar to a polypeptide encoded by an open reading frame comprising one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737.



7. The method of claim 1 or 2 wherein the at least one gene comprises an open reading frame comprising one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737.
8. The method of claim 1 or 2 wherein the pathogen is a bacterium.
- 5 9. The method of claim 1 or 2 wherein the pathogen is a fungus.
- 10 10. The method of claim 1 or 2 wherein the pathogen is a virus.
11. The method of claim 1 or 2 wherein gene expression is detected or determined using a gene chip, a cDNA array, cDNA-AFLP, or differential display PCR
12. The method of any one of claims 1 to 6 or 8 to 11 wherein the plant is a dicot.
- 15 13. The method of any one of claims 1 to 6 or 8 to 11 wherein the plant is a monocot.
14. The method of any one of claims 1 to 13 further comprising isolating the at least one gene or a portion thereof which includes the open reading frame or promoter for the gene.
- 20 15. A method to identify at least one gene, the expression of which is altered by pathogen infection in a wild-type plant relative to a plant having a mutation that decreases jasmonic acid or ethylene-dependent signaling, comprising:
  - a) contacting a plurality of isolated nucleic acid samples on a solid substrate each comprising isolated nucleic acid with a probe comprising plant nucleic acid  
25 corresponding to RNA from a wild-type plant infected with the pathogen, so as to form a complex, wherein each sample comprises a plurality of oligonucleotides corresponding to at least a portion of one plant gene; and
  - b) comparing complex formation in a) with complex formation between a second plurality of isolated nucleic acid samples on a solid substrate with a second probe comprising  
30 nucleic acid corresponding to RNA from the plant having the mutation and infected with the pathogen, so as to identify a gene, the expression of which is altered by pathogen infection in a wild-type plant relative to the mutant plant.

16. A method to identify at least one gene, the expression of which is altered by pathogen infection in a wild-type plant relative to a plant having a mutation in a gene that interferes with salicylic acid dependent signaling, comprising:
- a) contacting a plurality of isolated nucleic acid samples on a solid substrate each comprising isolated nucleic acid with a probe comprising plant nucleic acid corresponding to RNA from a wild-type plant infected with the pathogen, so as to form a complex, wherein each sample comprises a plurality of oligonucleotides corresponding to at least a portion of one plant gene; and
  - b) comparing complex formation in a) with complex formation between a second plurality of isolated nucleic acid samples on a solid substrate with a second probe comprising nucleic acid corresponding to RNA from the plant having a mutation and infected with the pathogen, so as to identify a gene, the expression of which is altered by pathogen infection in a wild-type plant relative to the mutant plant.
17. A method to identify at least one gene, the expression of which is altered by pathogen infection in a wild-type plant relative to a plant having a mutation that results in enhanced susceptibility to bacterial infection, comprising:
- a) contacting a plurality of isolated nucleic acid samples on a solid substrate each comprising isolated nucleic acid with a probe comprising plant nucleic acid corresponding to RNA from a wild-type plant infected with the pathogen, so as to form a complex, wherein each sample comprises a plurality of oligonucleotides corresponding to at least a portion of one plant gene; and
  - b) comparing complex formation in a) with complex formation between a second plurality of isolated nucleic acid samples on a solid substrate with a second probe comprising nucleic acid corresponding to RNA from the plant having a mutation and infected with the pathogen, so as to identify a gene, the expression of which is altered by pathogen infection in a wild-type plant relative to the mutant plant.
18. A method to identify at least one gene, the expression of which is altered by infection with at least one virus, comprising:
- a) contacting a plurality of isolated nucleic acid samples on a solid substrate each comprising isolated nucleic acid with a probe comprising plant nucleic acid corresponding to RNA from a wild-type plant infected with a virus, so as to form a

- complex, wherein each sample comprises a plurality of oligonucleotides corresponding to at least a portion of one plant gene; and
- 5       b) comparing complex formation in a) with complex formation between a second plurality of isolated nucleic acid samples on a solid substrate with a second probe comprising nucleic acid corresponding to RNA from an uninfected plant, so as to identify a gene, the expression of which is altered by virus infection.
19. A method to identify at least one gene, the expression of which is altered by infection with at least one pathogen comprising:
- 10       a) contacting a plurality of isolated nucleic acid samples on a solid substrate each comprising isolated nucleic acid with a probe comprising plant nucleic acid corresponding to RNA from an incompatible interaction so as to form a complex, wherein each sample comprises a plurality of oligonucleotides corresponding to at least a portion of one plant gene; and
- 15       b) comparing complex formation in a) with complex formation between a second plurality of isolated nucleic acid samples on a solid substrate with a second probe comprising nucleic acid corresponding to RNA from a corresponding compatible interaction so as to identify a gene, the expression of which is altered by the pathogen.
- 20       20. The method of any one of claims 15 to 19 wherein the at least one gene is upregulated.
21. The method of any one of claims 15 to 19 wherein the at least one gene is downregulated.
22. The method of any one of claims 15 to 19 wherein the gene which is identified encodes a polypeptide that is substantially similar to a polypeptide encoded by an open reading
- 25       frame comprising any one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737.
23. The method of any one of claims 15 to 19 wherein the probes comprise nucleic acid from a dicot.
- 30       24. The method of any one of claims 15 to 19 wherein the probes comprise nucleic acid from a cereal plant.

25. The method of any one of claims 15 to 19 wherein the probes comprise nucleic acid from a monocot.
26. The method of any one of claims 15 to 19 further comprising identifying the promoter for the at least one gene.
27. An isolated polynucleotide comprising a plant nucleotide sequence that alters transcription of an operatively linked nucleic acid segment in a plant cell after pathogen infection, which plant nucleotide sequence is from a gene encoding a polypeptide that is substantially similar to a polypeptide encoded by a gene comprising a promoter selected from the group consisting of SEQ ID NOs:2137-2661 or SEQ ID NOs:4738-6813.
28. An isolated polynucleotide comprising a plant nucleotide sequence that alters transcription of an operatively linked nucleic acid segment in a plant cell after pathogen infection, which plant nucleotide sequence hybridizes under high stringency conditions to the complement of any one of SEQ ID NOs:2137-2661 or SEQ ID NOs:4738-6813.
29. An isolated polynucleotide comprising a plant nucleotide sequence that alters transcription of an operatively linked nucleic acid segment in a plant after pathogen infection, which plant nucleotide sequence hybridizes under very high stringency conditions to the complement of any one of SEQ ID NOs:2137-2661 or SEQ ID NOs:4738-6813.
30. An isolated polynucleotide comprising a plant nucleotide sequence that directs transcription of an operatively linked nucleic acid segment in a plant cell, which plant nucleotide sequence is selected from the group consisting of SEQ ID NOs:2137-2661 and or SEQ ID NOs:4738-6813.
31. The polynucleotide of any one of claims 27 to 30 wherein the plant nucleotide sequence is 25 to 2000 nucleotides in length.
32. The polynucleotide of any one of claims 27 to 30 wherein the plant nucleotide sequence has at least 90% nucleotide sequence identity to one of SEQ ID NOs:2137-2661 or SEQ ID NOs:4738-6813.

33. The polynucleotide of any one of claims 27 to 30 wherein the plant nucleotide sequence has at least 98% nucleotide sequence identity to one of SEQ ID NOs:2137-2661 or SEQ ID NOs:4738-6813.
- 5
34. The polynucleotide of any one of claims 27 to 29 and 31 to 33 wherein the plant nucleotide sequence is from a dicot.
35. The polynucleotide of any one of claims 26 to 29 and 31 to 33 wherein the plant nucleotide sequence is from a monocot.
- 10
36. The polynucleotide of any one of claims 26 to 29 and 31 to 33 wherein the plant nucleotide sequence is a maize, soybean, barley, alfalfa, sunflower, canola, soybean, cotton, peanut, sorghum, tobacco, sugarbeet, rice or wheat sequence.
- 15
37. The polynucleotide of any one of claims 27 to 36 which comprises a TATA box, a CAAT box, or both.
38. A composition comprising the polynucleotide of any one of claims 27 to 37.
- 20
39. A recombinant vector comprising the polynucleotide of any one of claims 27 to 38.
40. The vector of claim 39 which is selected from the group consisting of a plasmid, phagemid, cosmid, virus, F-factor and phage.
- 25
41. An expression cassette comprising the polynucleotide of any one of claims 27 to 37 operatively linked to an open reading frame.
42. The expression cassette of claim 41 operably linked to other suitable regulatory sequences.
- 30
43. The expression cassette of claim 41 wherein the open reading frame is in an antisense orientation relative to the nucleotide sequence which alters transcription.

44. The expression cassette of claim 41 wherein the open reading frame is in a sense orientation relative to the nucleotide sequence which alters transcription.
45. A recombinant vector comprising the expression cassette of claim 41.
- 5 46. The vector of claim 45 wherein the vector is selected from the group consisting of a plasmid, phagemid, cosmid, virus, F-factor or phage.
47. A host cell comprising the expression cassette of claim 41.
- 10 48. The host cell of claim 47 wherein the cell is selected from the group consisting of a yeast, a bacterium, a cereal plant cell, and an *Arabidopsis* cell.
49. A plant cell containing the expression cassette of claim 41.
- 15 50. The plant cell of claim 49 which is a monocot cell.
51. The plant cell of claim 49 which is a dicot cell.
- 20 52. A transformed plant, the genome of which is augmented with the expression cassette of claim 41.
53. A transformed plant comprising transformed plant cells, which cells contain the expression cassette of claim 41.
- 25 54. The transformed plant of claim 52 or 53 which is a dicot.
55. The transformed plant of claim 52 or 53 which is a monocot.
- 30 56. The transformed plant of claim 52 or 53 which is selected from the group consisting of maize, soybean, barley, alfalfa, sunflower, canola, soybean, cotton, peanut, sorghum, tobacco, sugarbeet, rice, wheat and *Arabidopsis*.

57. A method for augmenting a plant genome, comprising:
- a) contacting plant cells with the expression cassette of claim 41 so as to yield a transformed plant cell; and
  - b) regenerating the transformed plant cell to provide a differentiated transformed plant,  
5        wherein the differentiated transformed plant expresses the open reading frame in the cells of the plant.
58. A method to alter the phenotype of a plant cell comprising: introducing the expression cassette of claim 41 into a plant cell and expressing that open reading frame in the cell so  
10        as to alter a characteristic of that cell relative to a plant cell that does not comprise the expression cassette.
59. The method of claim 57 or 58 wherein the plant cell is a dicot cell.
- 15    60. The method of claim 57 or 58 wherein the plant is a monocot cell.
61. The method of claim 57 or 58 wherein the plant cell a cereal cell.
62. The method of claim 57 or 58 wherein the plant cell is selected from the group consisting  
20        of a cell of maize, soybean, barley, alfalfa, sunflower, canola, soybean, cotton, peanut, sorghum, tobacco, sugarbeet, rice, wheat and *Arabidopsis*.
63. The method of claim 57 or 58 wherein the open reading frame is in an antisense orientation relative to the nucleotide sequence which alters transcription.  
25
64. The method of claim 57 or 58 wherein the expression inhibits transcription or translation of endogenous plant nucleic acid sequences corresponding to the open reading frame.
65. The method of claim 57 or 58 wherein the open reading frame is in a sense orientation  
30        relative to the nucleotide sequence which alters transcription.
66. The method of claim 57 wherein the open reading frame is expressed in an amount that is greater than the amount in a plant which does not comprise the expression cassette.

67. The method of claim 57 or 58 wherein the open reading frame encodes a protein.
68. The method of claim 67 wherein the protein encodes a regulatory product.
- 5 69. The method of claim 67 wherein the expression of the open reading frame confers insect resistance, bacterial resistance, fungal resistance, viral resistance, or nematode resistance.
70. A transformed plant prepared by the method of claim 57.
- 10 71. A product of the plant of claim 70 which comprises the expression cassette or the gene product encoded by the open reading frame.
72. The product of claim 71 which is selected from the group consisting of a seed, fruit,  
15 vegetable, transgenic plant, and a progeny plant.
73. A computer-readable medium having stored thereon a data structure comprising:  
a) a nucleic acid molecule that has at least 70% nucleic acid sequence identity to a  
nucleotide molecule selected from the group consisting of SEQ ID NOs:1-953, 2137-  
20 2661, 1954-1966, 2000-2129, 2662-4737, 4738-6813 or the complement thereof; and  
b) a module receiving the nucleic acid molecule which compares the nucleic acid  
sequence of the molecule to at least one other nucleic acid sequence.
74. The computer readable medium of claim 73 wherein the medium is selected from the  
25 group consisting of magnetic tape, optical disk, CD-ROM, random access memory,  
volatile memory, non-volatile memory and bubble memory.
75. A computer-readable medium having stored thereon computer executable instructions for  
performing a method comprising:  
30 a) receiving a nucleic acid molecule having at least 70% nucleic acid sequence identity to  
a nucleotide sequence selected from the group consisting of SEQ ID NOs:1-953, 2137-  
2661, 1954-1966, 2000-2129, 2662-4737, 4738-6813 or the complement thereof; and



b) comparing the nucleic acid sequence of the molecule to at least one other nucleic acid sequence.

76. The computer readable medium of claim 75 wherein the medium is selected from the group consisting of magnetic tape, optical disk, CD-ROM, random access memory, volatile memory, non-volatile memory and bubble memory.
77. The computer readable medium of any one of claims 73 to 76 wherein the nucleotide sequence is not SEQ ID NOs. 1-208, 210-215, 217-261, 263-266, 268-316, 318-385, 387-424, 426-439, 441-799, or 801-953.
78. A method to confer resistance or tolerance to a plant to a pathogen, comprising:
- a) contacting plant cells with an expression cassette comprising a polynucleotide encoding a polypeptide that is substantially similar to a polypeptide encoded by an open reading frame comprising any one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737 so as to yield transformed cells; and
  - b) regenerating the transformed plant cells to provide a differentiated transformed plant, wherein the differentiated transformed plant expresses the polynucleotide in the cells of the plant in an amount effective to confer resistance or tolerance to the plant to a pathogen relative to a corresponding plant which does not comprise the expression cassette.
79. The method of claim 78 wherein the polynucleotide hybridizes under moderate stringency conditions to the complement of any one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737.
80. The method of any one of claims 78 to 79 wherein the cells are monocot cells.
81. The method of any one of claims 78 to 79 wherein the cells are dicot cells.
82. The method of claim 78 wherein the open reading frame encodes a DNA binding protein, hormone response protein, membrane protein, metabolic protein, transposon, receptor/kinase, phosphatase, stress protein, cell wall protein, lipid transfer protein, heat

shock protein, protein processing protein, RNA processing protein, non-cell wall structural protein or a non-kinase signaling protein.

83. A transformed plant prepared by the method of any one of claims 78 to 82.

5

84. A seed of the plant of claim 83.

85. A progeny plant of the plant of claim 83.

10 86. A method to identify a plant cell infected with a pathogen, comprising:

a) contacting isolated nucleic acid obtained from a plant cell suspected of being infected with a pathogen with at least one oligonucleotide under conditions effective to specifically amplify a nucleotide sequence corresponding to one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737 or a portion thereof, so as to yield an amplified product; and

15

b) detecting or determining the presence or amount of the amplified product, wherein the presence or amount of the amplified product is indicative of pathogen infection.

87. A method to identify a plant cell infected with a pathogen, comprising:

20 a) contacting a sample comprising polypeptides obtained from a plant cell suspected of being infected with a pathogen with an agent that specifically binds to a polypeptide that is substantially similar to a polypeptide encoded by an open reading frame comprising one of SEQ ID NOs:1-953, 1954-1966, 2000-2129 or 2662-4737 so as to form a complex, wherein the; and

25 b) detecting or determining the presence or amount of the complex, wherein the presence or amount of the complex is indicative of pathogen infection.

88. A method to identify a plant cell infected with a pathogen, comprising:

30 a) contacting nucleic acid obtained from a plant cell suspected of being infected with a pathogen with a probe corresponding to a sequence selected from the group consisting of SEQ ID Nos. 1-953, 1954-1966, 2000-2129 or 2662-4737 or a portion thereof, under stringent hybridization conditions to form a duplex, and

- b) detecting or determining the presence or amount of the duplex, wherein the presence of a duplex is indicative of infection.

89. A method for marker-assisted breeding to select for plants having altered

5 resistance to a pathogen comprising:

- a) contacting plant DNA or cDNA with a probe comprising a sequence selected from the group consisting of SEQ ID Nos. 1-953, 1954-1966, 2000-2129 or 2662-4737 or a portion thereof which hybridizes under moderate stringency conditions to a gene corresponding to one of of SEQ ID Nos. 1-953, 1954-1966, 2000-2129 or 2662-4737

10 so as to form a duplex; and

- b) detecting or determining the presence or amount of the duplex, wherein the amount or presence of the duplex is indicative of the presence of a gene, the expression of which alters the resistance of the plant to a pathogen.

15

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 01/01105

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/29 C12N15/82 C12Q1/68 A01H5/00 G06F17/00  
C07K14/415

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, SEQUENCE SEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages   | Relevant to claim No.  |
|------------|--|------------------------|
| X          | SCHENK P M ET AL: "Coordinated plant defense responses in Arabidopsis revealed by microarray analysis"<br>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US,<br>vol. 97, no. 21,<br>10 October 2000 (2000-10-10), pages<br>11655-11660, XP002153163<br>ISSN: 0027-8424<br>the whole document | 1-5,<br>8-21,<br>23-26 |
| X          | WO 97 49822 A (CIBA GEIGY AG ;ELLIS DANIEL MURRAY (US); FRIEDRICH LESLIE BETHARDS)<br>31 December 1997 (1997-12-31)<br>the whole document<br>---<br>-/--   | 1-5,<br>8-21,<br>23-26 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

9 July 2002

Date of mailing of the international search report

25. 09. 2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 01/01105

| C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT |   |                           |
|--|---|---------------------------|
| Category *   | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No.     |
| X  | WO 01 07603 A (TUZUN SADIK ;UNIV AUBURN<br>(US); ABDULLAH MALIKAH (US); SINGH NAREND)<br>1 February 2001 (2001-02-01)<br>the whole document | 1-5,<br>8-21,<br>23-26    |
| X  | EP 1 033 405 A (CERES INC)<br>6 September 2000 (2000-09-06)   | 73-76,<br>78-86,<br>88,89 |
| A  | page 1 -page 26<br>see SEQ IN NO: 38097<br>page 325 -page 341; claims 1-34<br>-----   | 6,7,22                    |

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/IB 01/01105

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 87  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:  
Claims 1-26, 73-89 partially.
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 87

Present claim 87 relates to a product/compound defined by reference to a desirable characteristic or property, namely an agent that binds to a polypeptide encoded by an open reading frame.

The claims cover all products/compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such products/compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, no search has been carried out.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

**Invention 1: claims 1-26, 73-89 partially**

A method to identify at least one gene involved in plant resistance or response to infection by at least one pathogen wherein the gene encodes a polypeptide encoded by an open reading frame comprising SEQ ID NO: 1. A method to identify at least one gene, the expression of which is altered by pathogen infection, a computer-readable medium, a method to confer resistance or tolerance to a plant, a transformed plant, a seed, a method to identify a plant cell infected with a pathogen, a method for marker-assisted breeding comprising said nucleic acid.

**Invention 2-3173: claims 1-26, 73-89 partially**

same as invention 1 but comprising a gene in the order as given in claim 6 (invention 2 is limited to SEQ ID NO: 2 and invention 3173 is limited to SEQ ID NO: 4737).

**Invention 3174: claims 26-77 partially**

An isolated polynucleotide comprising a plant nucleotide sequence that alters transcription of an operatively linked nucleic acid comprising SEQ ID NO: 2137. A composition, a vector, an expression cassette, a host cell, a plant cell, a transformed plant, a method for augmenting a plant genome, a method to alter the phenotype of a plant cell, a computer-readable medium comprising said polynucleotide.

**Invention 3174-5774: claims 26-77 partially**

same as invention 3173 but comprising a promoter sequence in the order as given in claim 27 (invention 3174 comprises SEQ ID NO: 2138 and invention 5774 comprises SEQ ID NO: 6813).



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 01/01105

| Patent document<br>cited in search report |   | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---|---------------------|----------------------------|---------------------|
| WO 9749822                                | A | 31-12-1997          | AU 719639 B2               | 11-05-2000          |
|   |   |                     | AU 2026197 A               | 14-01-1998          |
|   |   |                     | BR 9709925 A               | 10-08-1999          |
|   |   |                     | CA 2258576 A1              | 31-12-1997          |
|   |   |                     | CN 1228813 A               | 15-09-1999          |
|   |   |                     | WO 9749822 A1              | 31-12-1997          |
|   |   |                     | EP 0923648 A1              | 23-06-1999          |
|   |   |                     | HU 9901749 A2              | 28-09-1999          |
|   |   |                     | JP 2000512502 T            | 26-09-2000          |
|   |   |                     | PL 330599 A1               | 24-05-1999          |
|   |   |                     | TR 9802660 T2              | 21-04-1999          |
|   |   |                     | US 6091004 A               | 18-07-2000          |
|   |   |                     | AU 727179 B2               | 07-12-2000          |
|   |   |                     | AU 5663198 A               | 03-07-1998          |
|   |   |                     | BR 9714398 A               | 02-05-2000          |
|   |   |                     | WO 9826082 A1              | 18-06-1998          |
|   |   |                     | EP 0944728 A1              | 29-09-1999          |
|   |   |                     | FR 2757875 A1              | 03-07-1998          |
|   |   |                     | IT MI972741 A1             | 15-06-1998          |
|   |   |                     | JP 2001505774 T            | 08-05-2001          |
|   |   |                     | NL 1007779 C2              | 22-07-1998          |
|   |   |                     | NL 1007779 A1              | 17-06-1998          |
|   |   |                     | US 5986082 A               | 16-11-1999          |
|   |   |                     | US 5955484 A               | 21-09-1999          |
|   |   |                     | US 5780469 A               | 14-07-1998          |
|   |   |                     | US 6031153 A               | 29-02-2000          |
| -----                                     |   |                     |                            |                     |
| WO 0107603                                | A | 01-02-2001          | AU 6227200 A               | 13-02-2001          |
|   |   |                     | WO 0107603 A1              | 01-02-2001          |
| -----                                     |   |                     |                            |                     |
| EP 1033405                                | A | 06-09-2000          | CA 2300692 A1              | 25-08-2000          |
|   |   |                     | EP 1033405 A2              | 06-09-2000          |
| -----                                     |   |                     |                            |                     |